Dr Anna C. Tottman, MB BS, PhD - Neonatal Fellow

(Corresponding author)

a.tottman@auckland.ac.nz

primary affiliation

University of Auckland Liggins Institute
Park Road Grafton Auckland 1142
New Zealand

secondary affiliation

Royal Women's Hospital -
Cnr Gratton Street and Flemington Road Parkville Victoria 3052
Australia

Dr Brett J. Manley, MB BS, PhD

Brett.Manley@thewomens.org.au

Neonatologist

Royal Women's Hospital -
Cnr Gratton Street and Flemington Road Parkville Victoria 3052
Australia

Dr Louise S. Owen, MB ChB, MD

Louise.Owen@thewomens.org.au

Neonatologist

Royal Women's Hospital -
Cnr Gratton Street and Flemington Road Parkville Victoria 3052
Australia

Professor Mark P. Umstad AM, MB BS, MD, FRCOG, FRANZCOG

Mark.Umstad@thewomens.org.au

Director of Maternity Services

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/AJO.12961](https://doi.org/10.1111/AJO.12961)

This article is protected by copyright. All rights reserved

Royal Women's Hospital -
Cnr Grattan Street and Flemington Road Parkville Victoria 3052
Australia

Professor Peter G. Davis, MB BS, MD

pgd@unimelb.edu.au

Director of Newborn Research

Royal Women's Hospital -
Cnr Grattan Street and Flemington Road Parkville Victoria 3052
Australia

DR. ANNA TOTTMAN (Orcid ID : 0000-0001-9944-5402)

Article type : Letter to the Editor

Deliveries at early term gestation: A view from the NICU

Short Title: Early-term morbidity

Word Count: Main text 500.

Keywords: Neonatal, NICU, Respiratory Distress Syndrome

Wilmink et al¹ report an increased risk of neonatal respiratory morbidity accompanying caesarean birth at early term gestations. Our recent experience of three infants admitted to our tertiary neonatal intensive care unit (NICU) within a two-week period, all born via elective caesarean section under regional anaesthesia without antenatal steroid administration, highlights that this morbidity may be severe:

Case 1

A G3P2 37-year old woman underwent elective repeat lower uterine segment caesarean (LUSCS) at 37+1 weeks' gestation. The pregnancy was complicated by a low-lying placenta on imaging at 31 weeks, but with no antepartum bleeds. A 2.8 kg male infant was delivered in good condition, but became hypoxic and tachypnoeic after one hour. There was radiological evidence of both retained fetal lung fluid and respiratory distress syndrome

(RDS). The infant required endotracheal intubation, mechanical ventilation, two doses of surfactant, umbilical line placement, sedation, and bladder catheterisation. He remained mechanically ventilated for two days, with a nine day admission to the neonatal unit.

Case 2

A G3P1 32-year old woman underwent elective repeat LUSCS in a regional hospital at 38+3 weeks' gestation. A 2.8 kg male infant was delivered in good condition, but developed respiratory distress 20 minutes later. He deteriorated rapidly, requiring endotracheal intubation, surfactant and ongoing mechanical ventilation, and air retrieval to our NICU. He developed a large pneumomediastinum, hypoxia and acidosis, requiring high-frequency oscillatory ventilation, sedation and muscle relaxation. A peripheral arterial line, umbilical venous line, bladder catheterisation, fluid boluses and inotropes were required to manage hypotension. He remained mechanically ventilated for four days, and was transferred back to his local hospital after eight days.

Case 3

A G2P1 41-year old woman underwent elective repeat LUSCS in a private metropolitan hospital at 38+0 weeks' gestation. A 3.4 kg female infant was born crying, but became apnoeic at two minutes of age. She was resuscitated, admitted to her local nursery in ambient oxygen, and subsequently transported to our NICU with RDS. She developed a pneumothorax requiring needle thoracocentesis, endotracheal intubation, mechanical ventilation, intercostal chest drain insertion, two doses of surfactant, umbilical venous catheterisation, and bladder catheterisation. She remained mechanically ventilated for three days and was transferred back to her local hospital after six days.

Although antenatal corticosteroids prior to early-term pre-labour caesarean section are associated with reduced neonatal respiratory morbidity^{1,2}, concerns remain regarding associations between antenatal corticosteroid administration and increased incidence of neonatal hypoglycaemia in the late-preterm population³, and with long-term impacts on educational outcomes in term infants⁴. These concerns have resulted in recommendations that antenatal corticosteroids be used with caution beyond 34 weeks' gestation until safety is proven⁵. Thus, timing of delivery is currently the most important modifiable risk factor for poor neonatal respiratory outcomes at early-term gestations. In light of the increasing evidence that birth at early-term gestation is itself associated with higher risks of poor long-

term neurodevelopmental⁶ and metabolic⁷ outcomes, we would urge practitioners planning caesarean deliveries before 39 weeks' gestation in the **absence** of a medical indication, to consider aligning their practice with current RANZCOG guidance⁸.

References

1. Wilmink F.A., Pham C.T., Edge N., et al. Timing of elective pre-labour caesarean section: A decision analysis. *Aust N Z J Obstet Gynaecol* 2018; 1-7.
DOI:10.1111/ajo.12821 [Epub ahead of print]
2. Srinivasjois R., and Silva D. Antenatal steroid administration on medically uncomplicated pregnancy beyond 37 weeks of gestation for the prevention of neonatal morbidities prior to elective caesarean section: a systematic review and meta-analysis of randomised controlled trials. *J Matern Fetal Neonatal Med* 2017;30(10):1151-7
3. Gyamfi-Bannerman C., Thom E.A, Blackwell S.C., et al. Antenatal betamethasone for women at risk of late preterm delivery. *N Engl J Med* 2016;374:1311-20
4. Behavioural, educational and respiratory outcomes of antenatal betamethasone for term caesarean section (ASTECS trial). Stutchfield P.R., Whitaker R., Gliddone A.E., et al. *Arch Dis Child Fetal Neonatal Ed* 2013;98(3):F195-200
5. Kamatkar S., Jobe A. Antenatal late preterm steroids (ALPS): are we ready to accept it? *J Perinatol* 2017;37:624-5
6. Dong Y., Chen S.J., and Yu J.L. A systematic review and meta-analysis of long-term development of early term infants. *Neonatology* 2012;102(3):212-21
7. Paz Levy D., Sheiner E., Wainstock T., et al. Evidence that children born at early term (37-38 6/7 weeks) are at increased risk for diabetes and obesity related disorders. *AJOG* 2017;217(5):588.e1-588.e11
8. Timing of elective caesarean section at term. C-Obs 23. The Royal Australian and New Zealand College of Obstetricians and Gynaecologists.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Tottman, AC;Manley, BJ;Owen, LS;Umstad, MP;Davis, PG

Title:

Deliveries at early term gestation: A view from the NICU

Date:

2019-04-01

Citation:

Tottman, A. C., Manley, B. J., Owen, L. S., Umstad, M. P. & Davis, P. G. (2019). Deliveries at early term gestation: A view from the NICU. AUSTRALIAN & NEW ZEALAND JOURNAL OF OBSTETRICS & GYNAECOLOGY, 59 (2), pp.E7-E7. <https://doi.org/10.1111/ajo.12961>.

Persistent Link:

<http://hdl.handle.net/11343/285613>