Aleksandr Voskoboinik ORCID iD: 0000-0001-6990-302X

Author reply

Aleksandr Voskoboinik MBBS¹², Peter M Kistler MBBS PhD¹²

¹Heart Centre, Alfred Hospital, Melbourne, Australia

²Baker Heart & Diabetes Research Institute, Melbourne, Australia

Correspondence to:

Professor Peter Kistler, Director of Cardiac Electrophysiology, Alfred Hospital, Melbourne, Australia.

Phone: +61 3 90762000

Email: peter.kistler@baker.edu.au

Disclosures: None

Key Words

Clinical: Electrophysiology – cardioversion

The high cardioversion success rates for atrial fibrillation reported using the Ottawa-AF cardioversion protocol¹ coupled with our study's findings² send an important message to physicians: that manual pressure augmentation (MPA) and the use of 360J defibrillators are alternative techniques rather than accepting 'permanent AF' after failed cardioversion using 200J with adhesive patches (as is standard practice at many centers). Of note, some of the patients in our study were referred for inclusion after failed cardioversion elsewhere, and perhaps this more selected population may

This is the author manuscript accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/jce.13909](https://doi.org/10.1111/jce.13909).

This article is protected by copyright. All rights reserved.

explain the higher failure rates with 200J patches observed in our randomised study compared with the observational results reported by Ramirez et al¹. With respect to the methodology of manual pressure augmentation (MPA), this was performed in either the AA or AP configuration (50% each) and a single operator performed this on 75% of occasions. Hand held paddles are not routinely available at many Australian hospitals, and hence it was important to report the MPA technique (using gloves) as a safe and effective technique. In addition many centres do not have external defibrillators to provide shocks up to 360J. We have observed in some instances that a second operator applying MPA is required in morbidly obese patients (depending on the body size of the first operator) to achieve cardioversion success.

References

1. Ramirez FD, Sadek MM, Boileau I, et al. Evaluation of a novel cardioversion intervention for atrial fibrillation: the Ottawa AF cardioversion protocol. *Europace*. 2018 Dec 7. doi: 10.1093/europace/euy285. [Epub ahead of print]
2. Voskoboinik A, Moskovitch J, Plunkett G, et al. Cardioversion of atrial fibrillation in obese patients: Results from the Cardioversion-BMI randomized controlled trial. *J Cardiovasc Electrophysiol*. 2019 Feb;30(2):155-161.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Voskoboinik, A;Kistler, PM

Title:

Letter to the Editor regarding the paper "Cardioversion of atrial fibrillation in obese patients: Results from the Cardioversion-BMI randomized controlled trial" RESPONSE

Date:

2019-09-01

Citation:

Voskoboinik, A. & Kistler, P. M. (2019). Letter to the Editor regarding the paper "Cardioversion of atrial fibrillation in obese patients: Results from the Cardioversion-BMI randomized controlled trial" RESPONSE. JOURNAL OF CARDIOVASCULAR ELECTROPHYSIOLOGY, 30 (9), pp.1764-1764. <https://doi.org/10.1111/jce.13909>.

Persistent Link:

<http://hdl.handle.net/11343/285650>