Introduction

Hepatitis C virus (HCV) infection is a major public health burden and a leading cause of cirrhosis globally^[1]. The introduction of new direct acting antiviral (DAA) therapies onto PBS in March 2016 offered patients an alternative to interferon-based therapies with clinical trials suggesting efficacy of greater than 95%^{[2][3]}.

The primary aim of this study was to compare the efficacy and safety of DAA therapy in HCV patients of all genotypes with and without cirrhosis. A secondary aim was to ascertain likelihood of sustained viral response at 12 weeks after treatment (SVR12) in those patients without an end-of-treatment response (EoTR).

Materials and Methods

All patients commenced on DAA therapy and seen in hepatology clinics across two metropolitan hospitals in Victoria from October 2016 and April 2017 were included. Ethical and governance approval was obtained from the Eastern Health Human Research Ethics Committee and carried out in accordance with the principles of the Helsinki Declaration. Baseline demographic including age, gender, body mass index (BMI), recreational substance abuse, prior treatment and response were collected from electronic medical records. Recreational substance abuse was defined as active illicit drug use, regular alcohol intake of three or more standard drinks per day or binge drinking five or more standard drinks in a single sitting at least once a month. Pre- and post-treatment (EoT and 12-weeks post-treatment) alanine transaminase (ALT), viral load and transient elastography results were compiled. Adverse outcomes (including serious adverse events) were recorded by treating physicians and nurse consultants.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: [doi:10.1111/liv.14279](#)

of quantification. EoTR was defined as possessing HCV viral load not detected on qualitative titre and below level of quantification at completion of DAA therapy. The diagnosis of cirrhosis was determined based on combination of liver biopsy, transient elastography (FibroScan), serum markers and clinical evaluation. Categorical data was compared using chi-square test for independence and non-parametric data was compared using Wilcoxon signed-rank test with a p value <0.05 considered significant.

Results

There were 376 patients identified, of which 42 did not have clinical follow-up and therefore were not assessed for SVR12. Analysis of 334 patients was conducted with the patient disposition shown in Table 1.

Overall sustained virologic response at 12 weeks post-treatment (SVR12) was 96.7% with minimal differences in SVR12 between the cirrhosis and non-cirrhosis groups (95.7% and 97.3%). SVR12 in genotypes 1, 2 and 3 was 98.2%, 93.8% and 94.9% respectively. There were 20 patients (5.99%) that failed to achieve an EoTR of which 80.0% (n=16) went on to achieve SVR12. Comparison between patients with (n=20) and without a positive EoT viral titre (n=239) is displayed in Table 2. Of the patients with a positive EoT viral titre that achieved SVR12, 93.8% (n=15) had a detected viral load below the level of quantification. In those without EoT viral titre that relapsed after 12 weeks, 50% (n=2) had detected viral load below the level of quantification.

The mean ALT at baseline was 97.5 IU/L, and this reduced significantly to 22.4 IU/L 12 weeks after treatment was completed. Of patients with both baseline and post-treatment liver stiffness results (n=90, 26.9%) overall liver stiffness improved post-treatment from a median baseline of 10.5 kPa (IQR 6.0 – 16.9) to 7.3 kPa (IQR 5.5 – 11.1, $p<0.0001$) with cirrhotic patients experiencing greater change (16.9 kPa to 10.9 kPa, $p<0.0001$).

Mild symptoms (most commonly headache and fatigue) occurred in 51.8% of patients during treatment with 2.7% requiring hospital admission. Post-treatment, 6 patients (1.8%) experienced significant adverse events with 3 deaths (0.9%) and 3 patients with hepatic decompensation (0.9%, variceal haemorrhage).

The patients lost to clinical follow-up (n=42) were compared with patients with 12-weeks post-treatment follow-up (n=334) in Table 3. In patients without follow-up, 26.2% had active substance abuse compared with 12.0% of patients with 12-week post-treatment follow-up (p=0.0112).

Discussion

Our results revealed that DAA therapy in an Australian real world setting had similar efficacy to that of international clinical trials. HCV genotype 1 which typically was treated with a course of sofosbuvir and ledipasvir was found to be more effectively treated than genotype 3 which typically underwent a combination of sofosbuvir and daclatasvir; this is consistent with previous studies that have found genotype 3 more difficult to treat with DAA therapy and an area for further investigation with the advent of second generation DAA regimens^[4].

Patients with positive EoT viral titre post completion of DAA therapy were found to have high likelihood of achieving SVR12 especially when viral titre was below the level of quantification (viral load <12). This presents a deviation from past thinking where a positive EoT viral titre post interferon-based therapy was suggestive of treatment failure^[5]. Historically patients would undergo multiple viral titre measurements during the post treatment period to ascertain response and relapse. Given the efficacy of DAA therapy the indication for regular virological monitoring is significantly reduced. Routine assessment of EoT viral titre is no longer considered necessary post DAA therapy in Australia^[6]; a positive result is not indicative of failure while a negative result does

not offer definitive confirmation of clearance as SVR12 remains the standard for determining virological cure of HCV. When compared with the sub-group of patients with EoT negative viral titre, patients with EoT positive titre were shown to typically have a higher BMI. Although clinically significant with respect to EoTR, BMI was not found to be significant as a negative predictive factor for achieving SVR12.

People who inject drugs (PWID) and have achieved clearance of HCV either spontaneously or through treatment are recommended to undergo HCV RNA testing every 12 months or following an episode of high-risk injecting^[7]. Given the imperative for regular follow-up and the high rate of SVR12 with DAA therapy shown, the necessity for routine HCV RNA testing at 12 weeks post therapy in PWID can be contended as delay in investigation holds little bearing on future management. Despite this, confirming SVR12 can prove opportunistic depending on an individual's level of engagement with healthcare.

A barrier to effective management of HCV identified during the study was the high rates of poor attendance and compliance. Greater than 11% of patients initially included in the study were lost to follow-up despite commencement and at least partial completion of their DAA therapy. This is a confounding factor for this study and can be attributed in part to multiple factors within the patient population contributing to difficulty with treatment adherence including limited financial resources, lack of transport, mental health and low levels of social support. This study reflects the importance of substance abuse as a predictive factor for poor healthcare engagement and represents an area for improvement with early involvement of support services during treatment if the individual is receptive. Despite patients frequently experiencing mild side effects during treatment this did not prove to be a significant limitation of care with only two patients ceasing DAA therapy early due to effects directly related to treatment. All significant adverse events experienced during this study were secondary to pre-existing comorbidities.

Conclusion

This large real-world cohort of HCV patients shows treatment with DAA is highly effective and similar to clinical trials with high rates of SVR12. Minimal difference in efficacy between non-cirrhotic and cirrhotic populations was observed. Patients with active substance abuse are less likely to engage with their healthcare service and attend routine follow-up. Patients with detected EoT viral load have a high chance of achieving SVR12 especially when detected but below level of quantification.

Acknowledgements

We would like to thank the staff at Eastern Health who made this project possible.

References

1. Hajarizadeh B, Grebely J, Dore GJ. Epidemiology and natural history of HCV infection. *Nat Rev Gastroenterol Hepatol*. 2013;10(9):553-62.
2. Lawitz E, Poordad FF, Pang PS, Hyland RH, Ding X, Mo H et al. Sofosbuvir and ledipasvir fixed-dose combination with and without ribavirin in treatment-naïve and previously treated patients with genotype 1 hepatitis C virus infection (LONESTAR): an open-label, randomised, phase 2 trial. *Lancet*. 2014;383(9916):515.
3. Sulkowski MS, Gardiner DF, Rodriguez-Torres M, Reddy KR, Hassanein T, Jacobson I et al. Daclatasvir plus sofosbuvir for previously treated or untreated chronic HCV infection. *N Engl J Med*. 2014;370(3):211-21.
4. Tapper EB, Afdhal NH. Is 3 the new 1: perspectives on virology, natural history and treatment for hepatitis C genotype 3. *J Viral Hepat*. 2013 Oct; 20(10):669-77.
5. Emilio Palumbo. Pegylated Interferon and Ribavirin Treatment for Hepatitis C Virus Infection. *Ther Adv Chronic Dis*. 2011 Jan; 2(1): 39-45.

6. Thompson et al. Australian recommendations for the management of hepatitis C virus infection: a consensus statement 2016. Gastroenterological Society of Australia. 2016 Mar.
7. Grebely et al. Recommendations for the management of hepatitis C virus infection among people who inject drugs. The International Journal of Drug Policy. Volume 26, Issue 10, Pages 1028–1038

Figure Legend

Figure 1 – Outcomes of patients without end-of-treatment response

Table 1 – Demographics of patients who underwent DAA therapy

Demographics	Non-cirrhotic (n=218)	Cirrhotic (n=116)	p value
Male gender, n (%)	129 (59.17%)	83 (71.55%)	0.0315
Mean age (range), years	51.19 (22 – 85)	51.21 (30 – 81)	0.3005
Median BMI (IQR), kg/m ²	25.52 (22.44 – 28.95)	27.30 (23.95 – 29.85)	0.0101
Prior treatment naïve, n (%)	183 (83.94%)	90 (77.58%)	0.1806
HCV genotypes, n (%)			
▪ 1	151 (69.26%)	65 (56.03%)	0.0083
▪ 2	10 (4.59%)	6 (5.17%)	1.0000
▪ 3	55 (25.23%)	43 (37.07%)	0.0137
▪ 4	1 (0.46%)	1 (0.86%)	1.0000
▪ 6	1 (0.46%)	1 (0.86%)	1.0000
Pre-treatment median liver stiffness measurement (IQR), kPa	5.6 (4.7 – 6.8)	16.9 (11.1 – 26.3)	<0.0001
Active substance abuse, n (%)	29 (13.30%)	11 (9.48%)	0.8328

Table 2 – Patients with negative and positive End of Treatment viral titres

Demographics	Negative EoT titre (n=239)	Positive EoT titre (n=20)	p value
Male gender, n (%)	151 (63.18%)	10 (50.00%)	0.2430
Mean age (range), years	53.52 (22 – 85)	50.90 (25 – 69)	0.2778
Non-cirrhotic, n (%)	163 (68.20%)	18 (90.00%)	0.0412
Median BMI (IQR), kg/m ²	25.81 (22.67 – 28.93)	31.31 (26.22 – 34.34)	0.0102
Prior treatment naïve, n (%)	192 (80.33%)	20 (100.00%)	-
HCV genotypes, n (%)			
▪ 1	157 (65.69%)	15 (75.00%)	0.3971
▪ 2	6 (2.51%)	0 (0%)	-
▪ 3	72 (30.01%)	5 (25.00%)	0.6300
Pre-treatment median liver stiffness measurement (IQR), kPa	6.7 (5.0 – 13.6)	5.4 (4.0 – 7.2)	0.1436
Mean baseline ALT, IU/L	105.59	66.25	0.1889

Table 3 – Patients with and without 12-week follow-up

Demographics	Patients with 12-week follow-up (n=334)	Patients lost to follow-up (n=42)	p value
Male gender, n (%)	212 (63.47%)	30 (71.4%)	0.3103
Mean age (range), years	53.27 (22 – 85)	47.45 (25 – 68)	0.0008
Non-cirrhotic (%)	216 (64.67%)	34 (80.95%)	0.0351
Median BMI (IQR), kg/m ²	26.31 (22.63 – 29.33)	26.97 (22.94 – 30.92)	0.7719
Prior treatment naïve, n (%)	273 (81.74%)	37 (88.10%)	0.3072
HCV genotypes, n (%)			-
▪ 1	216 (64.67%)	21 (50.00%)	0.0634
▪ 2	16 (4.79%)	1 (2.38%)	0.4787
▪ 3	98 (29.34%)	20 (47.62%)	0.0161
Pre-treatment median liver stiffness measurement (IQR), kPa	6.6 (4.9 – 12.9)	6.9 (5.5 – 10.8)	0.3231
Mean baseline ALT, IU/L	96.63	85.15	0.5521
Adverse outcomes including mild symptoms, n (%)	173 (51.80%)	16 (38.10%)	0.0941
Active substance abuse, n (%)	40 (11.98%)	11 (26.19%)	0.0112

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Williams, J;Lucarelli, N;Nicoll, A;Lubel, J

Title:

Real-world Australian data reflect very high sustained virologic response at 12 weeks with direct acting antiviral therapy for hepatitis C and suggests highly achievable even in those without an end-of-treatment response.

Date:

2019-05

Citation:

Williams, J., Lucarelli, N., Nicoll, A. & Lubel, J. (2019). Real-world Australian data reflect very high sustained virologic response at 12 weeks with direct acting antiviral therapy for hepatitis C and suggests highly achievable even in those without an end-of-treatment response.. Intern Med J, 49 (5), pp.666-669. <https://doi.org/10.1111/imj.14279>.

Persistent Link:

<http://hdl.handle.net/11343/285823>