TITLE PAGE

Title: Clinical characteristics and outcomes of patient presentations to the emergency department via police: A scoping review

Running title: Patient presentations to ED via police

Authors:

^{1,2} Prof Julia Crilly; RN, MEmergN(Hons), PhD; Associate Professor, Emergency Care, Gold Coast Health and Menzies Health Institute Queensland, Griffith University

^{1, 2, 14} Dr Amy N. B. Johnston; RN, BSc(Hons), PhD; Conjoint Senior lecturer, Emergency Care, now at Metro South Health, Menzies Health Institute Queensland, Griffith University and University of Queensland

^{2, 3} Prof Marianne Wallis; RN, BSc(Hons), PhD; Associate Dean of Health, Faculty of Science, Health, Education and Engineering, University of the Sunshine Coast

¹ Josea Polong-Brown; RN, Research Nurse, Robina Department of Emergency Medicine, Robina Hospital, Gold Coast Health

⁴ A/Prof Ed Heffernan; BSc (Hons), MB BS, MPH, PhD, FRANZCP, Queensland Centre for Mental Health Research, University of Queensland

⁵ Prof Gerard Fitzgerald, MB BS, BHealthAdmin, PhD, Professor of Public Health, School of Public Health and Social Work, Queensland University of Technology

^{6,7,8} Jesse T. Young, MPH; Research Fellow, Melbourne School of Population and Global Health, University of Melbourne

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/1742-6723.13300](https://doi.org/10.1111/1742-6723.13300)

9, 10, 11, 12, 13 Prof Stuart Kinner; PhD, NHMRC Senior Research Fellow, Centre for Adolescent Health, Murdoch Children's Research Institute

1. Department of Emergency Medicine, Gold Coast Health, QLD
2. Menzies Health Institute, Queensland, Griffith University, QLD
3. Sunshine Coast Health Institute and University of the Sunshine Coast, QLD
4. Queensland Forensic Mental Health Services, Brisbane, QLD
5. School of Public Health and Social Work, Queensland University of Technology, Brisbane, QLD
6. Melbourne School of Population and Global Health, University of Melbourne, VIC
7. School of Population and Global Health, The University of Western Australia, WA
8. National Drug Research Institute, Curtin University, WA
9. Centre for Adolescent Health, Murdoch Children's Research Institute, VIC
10. Griffith Criminology Institute, Griffith University, QLD
11. Melbourne School of Population and Global Health, University of Melbourne, VIC
12. Mater Research Institute-UQ, University of Queensland, QLD
13. School of Public Health and Preventive Medicine, Monash University, VIC
14. School of Nursing, Midwifery and Social Work, University of Queensland, QLD

Corresponding author:

Julia Crilly, Department of Emergency Medicine, D Block, LG096 1 Hospital Blvd, Southport, QLD, 4218 Australia.

Email: Julia.Crilly@health.qld.gov.au

Telephone: +61 (0) 7 56875359

Author contribution: JC and AJ contributed to the project concept and design, supported literature search, article limiting, data extraction and manuscript preparation and revision; J P-B and MW contributed to article limiting, data extraction and manuscript development; SK, EH, GF contributed to project concept and critical revision of the manuscript; and JY contributed to critical revision of the manuscript. All authors approved the final version of the manuscript for submission.

This work was carried out at the following institutions:

Department of Emergency Medicine, Gold Coast University Hospital, 1 Hospital Blvd, Southport, QLD, 4215, Australia.

School of Nursing and Midwifery, Menzies Health Institute Queensland, Griffith University, Parklands Drive, Southport, QLD, 4215, Australia.

Title: Clinical characteristics and outcomes of patient presentations to the emergency department via police: A scoping review

ABSTRACT

People brought in by police (BIBP) to the emergency department (ED) are a potentially vulnerable group. This narrative scoping review aimed to identify, evaluate and summarise current literature regarding the frequency of presentation, demographic and clinical profile of patients (including reason for presentation), care delivery, and outcomes for people BIBP to the ED, and identify current gaps in knowledge. The review involved searching EMBASE, CINAHL, and PUBMED using a combination of terms: emergency/emergency department coupled with police custody/watch house or police presentation, for papers published in English language from January 2006 to November 2017. A total of 20 studies met the inclusion criteria. These included 17 observational (non-Randomised Controlled Trials) quantitative studies and 3 descriptive case reports. The proportion of presentations to ED that were BIBP varied depending on the study design and sampling frame. People BIBP often presented with aggressive behaviour, mental health problems, and/or substance use problems, and/or aggressive behaviour and injury caused by self or others. Of studies focused specifically on patients arriving to the ED in mental health crisis (i.e., suicidal ideation or self-harm), approximately 20-30% 18-27% were BIBP. ED presentations BIBP were mostly male and typically younger than people arriving by other means. The nature of care provided in the ED and outcomes of the acute episode of care were typically not well-described. Limited research regarding people BIBP to the ED limits the ability to comprehensively

understand their demographic and clinical profile and outcomes of emergency care. Further research is required to inform if and where in the patient's journey further improvements may be targeted.

Key words: emergency department, police, scoping review

Introduction

Within Australia, the number of patient presentations made to emergency departments (EDs) via police or correctional services increased from 45,452 in 2008-09¹ to 48,307 in 2014-15,² despite remaining a relatively consistent proportion (0.79%¹ and 0.75%²) of all ED presentations. Although this increase (of 6.3%) is below the rate of increase of the Australian population as a whole (8.7% for the same timeframe)^{3,4}, people arriving to the ED via police are a vulnerable group of healthcare users who often have healthcare needs reflective of their reason(s) for police contact including substance use and other mental health problems.

The health profile of people in police or correctional custody reflects a high rate of mental illness and substance use disorders⁵⁻⁷, and it is unclear if this profile is similar for those brought in by police (BIBP) to the ED. In many western jurisdictions the police are becoming the default lead agency in addressing acute social and/or mental health crises, many of which are unlikely to result in criminal prosecution. Reflecting papers included in this review, presentation to ED with police can occur when police suspect someone is in need of mental health assessment and treatment and is at risk of harm to themselves or others or when someone encountered (e.g. family violence victim) or in custody is injured or unwell and requires medical treatment. Police also often transport someone to hospital when the presence of substance intoxication or medical factors such as delirium or disorientation impacts the ability to determine whether acute mental illness is present. ~~is often different to that of their community peers, with high rates of communicable and non-communicable disease, mental health disorders and harmful substance use.~~³⁻⁵ Other chronic and under-treated physical health conditions such as heart disease and diabetes are also common in this

population.⁶ Furthermore, While there is a body of research describing people brought in by police (BIBP) to EDs, this group is typically subsumed within other ED user groups such as mental health consumers^{7,8,9} or those with drug^{9,10} and alcohol-related presentations.^{8,10,11,12}

Currently, little is known about the characteristics of people BIBP to the ED, restricting development of evidence-informed approaches to quality service delivery. In order to optimise service delivery and inform future research directions for people BIBP, this narrative scoping review aimed to provide a comprehensive understanding of the current literature related to the prevalence, demographic and clinical profile (including reason for presentation), care delivery and outcomes for this group.

Methods

~~This narrative scoping review used an integrative multistage approach. Data synthesis was based on an existing framework¹² which informed the mapping of evidence patterns to understand the scope of available evidence.¹³~~

~~The review largely followed the Johanna Briggs Institute scoping review process¹³ in which data synthesis was based on an existing existing framework.^{12,14}~~

Search strategy

~~Our search strategy process followed PRISMA guidelines¹⁵ (see Figure 1). We searched EMBASE, CINAHL and PUBMED for relevant literature. Title/abstract search terms included: ED/EDs, emergency department/s, emergency room/s, accident and emergency,~~

ER/s OR A&E; AND police, custody, watch house, correctional services OR police presentation. Activation of 'smart text' and automatic word variation options (unlimited truncation operators such as emergenc*) during searches ensured that word combination options including US and UK spelling variations and plural terms were detected. Reference chaining was undertaken to supplement the title search strategy in an effort to optimise the number of relevant records.^{14 16}

Inclusion criteria applied to the initial literature library were: full-text, peer-reviewed original research articles (inclusive of experimental and non-experimental studies, quantitative and qualitative studies, and systematic reviews) published in English between 1 January 2006 and 3 November 2017 that used two or more of the terms listed above in combination. This time frame was selected based on major changes to mental health acts and mental healthcare (including drugs to control psychosis) in the last decade. Our review included studies that were not exclusively focussed on people BIBP, provided it was possible to extract data that pertained to people BIBP. Exclusion criteria were: editorials, conference abstracts, anonymous articles, articles where authorship was unclear, and publications that focused solely on updated datasets. **Studies focussed solely on transfer from prison (i.e. escorted by correctional health staff) were also excluded.** Two researchers (VA, AJ) screened titles and abstracts for inclusion based on criteria. A third researcher (JP-B) moderated where agreement was not initially achieved. Full-text articles of studies that appeared to meet inclusion criteria were screened by three researchers (VA, AJ, JP-B). Disagreements on full-text inclusion arising during screening were moderated by two other researchers (JC, MW).¹⁵

Data extraction and synthesis

Data extracted from included studies included: study authors, year of publication, country of study, research design, sample, methods of data collection, and main results pertaining to this review (incidence of presentation, demographic profile, reason for presentation, patient and service characteristics, ED care delivery and outcomes pertaining to people BIBP to the ED) (see Table 1 and Supplementary Table 1). Initial extraction was performed by a research assistant (VA) in collaboration with an author (AJ) and checked by a second author (JP-B). The level of evidence for each included study was independently assessed by two authors (AJ, JP-B) using the National Health and Medical Research Council (NHMRC) Level of Evidence Hierarchy body of evidence matrix.^{46,17} This matrix allocates scores for levels of quantitative evidence on the basis of study design. Assessment differences were resolved by consensus with additional reviewers (JC, MW).

Results

A total of 1,946 titles and abstracts were screened for inclusion; 20 studies met the inclusion criteria (see Figure 1, Table 1 and Supplementary Table 1). Methodologies of included studies were 17 observational (non-RCT) quantitative studies and 3 descriptive case reports. Most of these studies were undertaken in high income countries; ten reports were from Australia, four from the USA, two from Canada, and one each from UK, Portugal, France, and Hong Kong. The studies varied in evidence level from III-3 to IV on the NHMRC level of evidence continuum.^{46,17} Two studies were based on the same dataset but focused on different aspects of the dataset and so are both included in this review.^{9,18}

[Insert Figure 1 about here]

[Insert Table 1 about here]

[Insert Supplementary Table 1 about here]

Prevalence of ED presentations BIBP

The proportion of people BIBP to ED varied widely depending on the nature and focus of the study. In the two studies that considered those BIBP as part of a complete ED cohort, the proportion was low (~0.8%^{17,18} - 1.3%^{11,12}). When specified subgroup populations within the ED were considered, frequent presenters (i.e. >8 presentations/year) were BIBP at three times the rate (1.7%) of infrequent presenters (<5 presentations/year).^{18,19} Almost half of the included studies (n=8) reported on people with mental illness. Within these studies, approximately 18-86% were BIBP.^{8,17,19-23} 9, 18, 20-24 However, the prevalence range of people BIBP with mental illness, when studies that explicitly examined patients transported under legal orders are excluded, was much lower, between 18-27%.^{9,18, 20-22} Other studies explored issues such as people BIBP from alcohol and other drugs and intimate partner violence, ~~and excessive use of force in custody~~. In these narrowed samples, the prevalence rates of people BIBP varied (e.g. amphetamine use: 16-24%, other toxicology: 13%, family violence victims: 7%), and are higher than the general ED police presentation rate (0.8-1.3%).^{10,12,18,26}

The demographic profile, clinical profile (reasons for presentation, urgency of condition, day and time of presentation to ED), and ED care delivery (i.e. procedure/treatment) of people BIBP varied depending on the study focus. The majority of

people BIBP were male, with a median age in the early 30s.^{8, 9, 17, 19, 21, 23-27 9,10,18, 20, 25-29}

Where socio-demographic profile was considered, people BIBP had disproportionate representation of people who identified as an ethnic minority, typically described as ‘first people’ or ‘black’ people.^{20, 23-25 21, 25-27} Studies also reported a higher proportion of people who were unemployed^{8, 17, 28 9,18,30} and lived in unstable housing^{19, 25 20,27} when compared with those in other patient populations.

Regarding the clinical profile of people BIBP, ~~the two most~~ common presenting problems ~~noted in included studies (most of which had narrowed samples) included were:~~ 1) violent or aggressive behaviour^{8,9,17,19,21,29-32 9,10,18,20,22,24,31-33} and 2) mental health problems;^{8, 17, 19, 26, 30 9,18,20,24,28} presentation to the ED with both at the same time ~~was not uncommon~~ was reported in several studies.^{8, 17, 19, 30 9,18,20,24} Substance use (alcohol and/or other drug) was also a common reason for presentation, ~~associated~~ co-reported in some cases with both aggression and/or mental health crisis.^{8, 17, 20, 28 30, 33, 34 9,18,21,24,30,31,34,35} ~~People BIBP with violent or aggressive behaviour were more likely than other patients to have a history of previous police contact, violent ideation and violence towards others.~~^{21, 25, 29, 31, 35 22,27,31,32,36} Patients BIBP who were experiencing a mental health problem were often experiencing a situational crisis (e.g. homelessness or family crisis),^{21,25 22,27} with some presenting with suicidal ideation,^{7, 30 8,24} self-harm,^{21, 26, 30 22,24,28} psychosis,^{17, 20, 23 18,21,25} or acute depression and anxiety.^{8 9} Other clinical profiles reported included physical trauma ~~from intimate partner violence~~^{24, 35 26,36} or from injuries that occurred in custody.^{27, 31, 34, 36 29,32,35,37} ~~and from intimate partner violence~~^{24, 35 26,36} Australasian studies reported on the triage category (i.e. urgency of presentation based on a 1-5 scale where 1 is most urgent and should be seen

immediately, and 5 is less urgent but should be seen within 2 hours).^{37,38} People BIBP were typically allocated a triage category of '3' reflecting the need for assessment and treatment within 30 minutes of arrival.^{8,9,17,30,9,10,18,24} Of the studies that reported time of arrival to the ED, most presentations BIBP occurred outside business hours,^{8,17,26,27,9,18,28,29} the definition of which was not specified or varied from 1700-0830, 1700-0800 and or weekend presentations.

Care delivery for people BIBP to the ED

Emergency care delivered for people BIBP was rarely reported, but where cited, fell into three broad categories: health assessment; investigations; and treatment. Health assessment included initial triage, baseline observations, and medical clearance prior to further specialist assessment (such as mental health assessment).^{31,32} Investigations included: pathology; toxicology; and x-ray; whilst treatment included plaster; sutures; wound dressings, and medication (including chemical restraint)^{17,19,21,26,27,30,31,33,35,38,39,18,22,24,28,29,32,34,36,39,40} although these varied based on the reason for presentation and were often alluded to rather than specifically reported.^{19,20,29,36,20,21,31,37} The use of physical restraint (for patients who were aggressive and posed a risk to others) was also reported in some studies that included patients BIBP.^{19,30,20,24}

Outcomes of people BIBP to the ED

Outcomes for people BIBP were infrequently reported and often subsumed within other population groups (such as mental health presentations). When outcomes were examined they

included: ED length of stay (LOS); admission requirement; and post ED referral to specialist, community and outpatient services. Studies focussed on mental health-related presentations and people BIBP reported that people BIBP tended to have longer hospital LOS^{8,17,30 9,18,24} compared to those presenting by other means. ED LOS for those BIBP was reportedly longer^{30,24} or no different to other ED presenters.^{17 18} Hospital admission rate for those BIBP also varied – lower in some studies^{8,30 9,24} but higher in others.¹⁷⁻¹⁸ Commonly reported referral services for people BIBP included specialist mental health, social work, and alcohol and other drugs services.^{7,19,21 8,20,22} Actual attendance at these referral appointments and/or outcomes of those appointments was not reported, possibly confounded by patient return to police custody^{9,23 10,25} or complications associated with custodial residence,^{29 31} although this cannot be generalised to the whole group. Other outcomes post-discharge were rarely noted.

~~Thus study recommendations relevant to those BIBP, often included the need for greater engagement between services (such as police and health) during and/or subsequent to health care episodes.^{18,22,24} Such needs were particularly evident in studies of patients BIBP to ED with a mental health problem or in crisis.^{8,23}~~

Discussion

The prevalence of ED presentations BIBP overall was small. In a number of sub-populations such as mental health, substance affected, family violence victims, prevalence rates were higher. ED presentations BIBP were mostly male and typically younger than people arriving by other means. Mental health, substance intoxication or medical factors were common reasons for transport by police to ED. Care delivery generally consisted of health assessment,

investigations, and treatment. With regards to outcomes, people BIBP have longer ED LOS than people arriving by other means; admission rates varied based on the sample and referral services reflected reason for presentation. These findings should be taken with caution given the paucity of some data and limitations such as study design and narrow focus. This summary of findings in relation to the posed aims does however provide some context for examining the consistency of these findings and their practice implications.

~~Our findings have identified the proportion of people BIBP to the ED is infrequently reported and the profile of people BIBP tends to reflect a group who present with aggressive behaviour, mental health problems and/or substance use problems. The nature of care provided in the ED and outcomes of the acute episode of care were rarely reported. While we have highlighted some key study results and common foci of these studies, most research we examined had a narrow scope of focus, with few studies providing holistic examination across the presentation, care and outcome spectrum, either cross-sectionally or longitudinally.~~

Meaningful data around frequency of presentations for people BIBP was limited and prevalence was difficult to ascertain because for some studies the sampling frame was people transported by police, for some it was specific diagnostic groupings of patients and for others it was all ED presentations. While the rate of presentations BIBP to ED is generally low (under 2%), for mental health presentations it is reported at 18-86%; lower (18-27%) when studies of patients transported under involuntary orders are excluded. This large range possibly reflects: i) different sampling approaches used in different studies; ii) the mental health definition used (mental health problem / mental health illness) or iii) the high proportion of this population with mental health problems and limited access to or utilisation

of community mental health services. ~~and or ii) varied inter-agency local arrangements in responding to people in crisis.~~ It is also likely that the frequency is highly contextual and influenced by community and police culture, legislative and reporting requirements and health system design considerations. Given that mode of arrival by police is often a characteristic of frequent presenters (defined as those presenting ≥ 8 times/year),^{18 19} and that the costs associated with frequent presentations are high,^{40 41} there appears to be scope for improvement in the medium to long term management of this vulnerable group. Collaborative care and continuity of care for this group is limited between first responders, hospital-based primary care and social support agencies, possibly increasing the likelihood of negative outcomes.^{41, 42-42, 43}

We identified literature spanning countries that included Australia, the UK, USA, Canada, Europe, and Hong Kong, highlighting the perceived importance of this population internationally. No studies were identified arising from low- and middle-income countries in South America, Africa or Asia, possibly reflecting our search strategy which only included studies published in English. The apparent dearth of evidence from such countries, which account for over two-thirds of the world's incarcerated population,⁴³⁻⁴⁴ highlights the need for further research to understand whether reported findings are relevant in other regions (including those that are resource-poor). ~~Study recommendations relevant to those BIBP, often included the need for greater engagement between services (such as police and health) during and/or subsequent to health care episodes.~~^{18, 22, 24 19, 23, 26} Such needs were particularly evident in studies of patients BIBP to ED with a mental health problem or in crisis.^{8, 23 9, 25} Further research is also required to understand if and how changes in police and social and

community healthcare standards have impacted on the incidence of people BIBP to the ED, the nature of care delivery, and outcomes for this vulnerable group.

Our paper focused on people BIBP to ED, a cohort where mental health problems, ~~violent / aggressive behaviour~~ and substance use is often reported. Critically, many aspects of clinical response may be required for patients BIBP, which includes assessment and management of substance misuse, injury, mental health problems and psychosocial stressors in addition to behaviour management. Difficulties for police in managing people with mental health and substance use problems in the community have led some police jurisdictions to adopt a co-response (police and mental health) strategy.^{44, 45 45-47} Whilst such programs have strong linkages with community services and reduce pressure on the justice system, there is limited evidence on other impacts.^{45 46} A recent mixed-methods study undertaken in Canada evaluated a police-mental health co-response team.^{44 45} The co-response team had low rates of injury and arrest and when compared to a police only team response, had higher overall rates of voluntary escorts to hospital and lower rates of involuntary escorts. Police in co-response teams also spent less time on hospital handovers. Interviews with 15 service users indicated an appreciation of specialised mental health response, use of de-escalation skills, compassion, empowerment and non- criminalizing approach.⁴⁴⁻⁴⁵ A similar model, termed Police Ambulance Crisis Emergency Response (PACER), exists in Australia.⁴⁷ With this model, the provision of mental health evaluations ‘in the field’ by mental health personnel and police meant that some consumers could be diverted away from the ED. Similar models warrant rigorous evaluation in other jurisdictions. Although the majority of people BIBP to the ED will not be placed in long-term police custody, there is a need for research targeting

the subset of those who are under current custodial orders and/or progress to incarceration. The WHO^{47,48 48,49} has argued for the importance of understanding prisoners' health needs to inform delivery of evidence-based health services for this vulnerable population. Similarly, our review highlights that substance misuse and mental illness are key areas for care planning for people who arrive to the ED by police.

Supporting guidelines, clinical frameworks, policy and legislative requirements cited in the reviewed studies did not always appear to meet the needs of police and/or consumers⁴⁹⁵⁰ or assist in the provision of quality care in the ED.^{30,39 24,40} Legislatively required documentation associated with presentations to ED that were BIBP^{50 51} was rarely mentioned or reported in studies. Actual or potential medico-legal consequences/outcomes were sometimes, but not always referred to in the reviewed studies even though medico-legal considerations were important factors that guided processes such as detention and restraint applied by police and clinical staff. While a minority of studies cited injury that **apparently** **allegedly** occurred during police pursuit, arrest or whilst in police/correctional officer custody,^{27,31 29,32} the mechanism of injury was mostly not explicitly reported. Although these studies highlighted the need for effective reporting processes, there was limited discussion of the outcomes from the injury.^{27,31,36 29,32,37} This lack of reporting, guidelines and frameworks restricts the ability to empirically evaluate the impact of these processes and related outcomes.

The relative dearth of studies exploring the specific needs of patients arriving to the ED by police limits the capacity to recommend evidence-based interventions to improve their management and care in the ED and beyond. In studies where the patient profile and the

nature of presentations were considered, presentation for mental illness, complicated by intoxication with alcohol or other drugs, was typical. Further research, including longitudinal studies, using multi-sectoral data linkage to articulate care trajectories and identify when and where to intervene to improve outcomes and reduce harm for this vulnerable group of people is warranted. Additionally, increased effort to improve consistency in data collection and reporting by clinicians and health (including ED) services and researchers, is recommended. The impact of this patient group on clinical staff and other patients also warrants further research. Despite the minimal evidence available, there was strong consensus in the literature that further improvement in healthcare delivery for people BIBP to the ED is required. Although people BIBP were more likely to be referred to a mental health service, and other community services or resources, there was a noted lack of literature examining the uptake and outcomes of these referrals. This presents a critical opportunity for further research, especially in the context of national mental health and suicide prevention priority areas that include: coordinated treatment and support for people with severe and complex mental illness; improving physical health and reducing early mortality; and enabling effective system performance and system improvement strategies.⁵⁰⁻⁵¹ By virtue of their presentation, people presenting to the ED by police are a vulnerable population who require co-ordinated multi-agency, multi-sectoral responses. These responses would include, but are not limited to, emergency care, primary care, justice stakeholders, mental health services, and alcohol and other drug services, to ensure continuity of care between services and over time.

Based on this review, key areas for future research include: i) using multi-sectoral data linkage to articulate care trajectories and identify when and where to intervene to

improve outcomes and reduce harm for people BIBP (e.g., frequent attenders or people recently released from prison who may benefit from specific care plans, examining the uptake and outcomes of referrals); ii) providing a consensus on structure, process and outcome indicators for people BIBP to the ED that include post discharge services; iii) describing the impact of people BIBP to ED on clinical staff and other patients; iv) trialling evidenced-based models of care that aim to provide the most appropriate care in the most appropriate place for people in police custody or in a situational crisis in the community; v) evaluating the impacts on the ED of broader changes in police and social and community healthcare standards, practices or policies; and vi) describing if and how ED care delivery may need to vary based on geographical location (remote, regional, metropolitan areas) or low- and middle-income countries.

Limitations

The limitations of this review reflect the paucity of literature about people BIBP to ED. Many of the included studies were limited by sample size, which impacted on their utility to inform clinical practice. Furthermore, ~~with limited information was presented regarding patients BIBP, including their prevalence within the entire ED population. Methodological limitations in the available data preclude robust ascertainment of the actual prevalence such as violent and aggressive behaviour as a cause for police escort to the ED. This restricted for each of the samples presented, there are associated restrictions on~~ Limited information also restricted our ability to discuss important aspects of services provision for this group including: i) safety aspects around people BIBP in the ED (i.e. security of person in custody and safety of others

in the ED), ii) proportion of those BIBP with intellectual disability; iii) acute health and legal implications of those BIBP following suspected body packing; and iv) involuntary detention orders (e.g. for treatment of severe alcohol and other drug dependence) and other similar statutes in other ~~across a number~~ jurisdictions. ~~that impact on the numbers and management of people BIBP.~~ This study was also limited to literature published in English, potentially limiting ascertainment of studies in low-to-middle-income countries and non-English speaking high-income countries. The review was also limited to peer-reviewed publications from the last decade and did not explore grey literature such as governmental reports.

Conclusion

Most people BIBP to the ED were young males. Reasons for presentation ~~included pertained to violent/aggressive behaviour and~~ mental health problems, ~~injury, and violent/aggressive behaviour~~ and these were often underpinned by substance misuse. The specific care delivery provided in the ED was inconsistently reported, and outcomes for this patient group are not well described. A lack of continuity of care from hospitals into the community services or correctional facilities was often noted although few studies examined this empirically. Further research is required to improve our understanding of trajectories for this patient group beyond the ED, and the inter-agency nature of care delivery that they require.

Acknowledgements: We thank Vinah Anderson for her research assistance with this review.

Disclosure of competing interests: There are no known conflicts of interest in the publication of this article. Stuart Kinner receives salary support from the NHMRC (APP1078168). Jesse Young is supported by a Melbourne International Research Scholarship (PhD) from The University of Melbourne.

Funding: This review was supported by a Griffith University Health Practice Innovation seed grant.

References

1. AIHW. Australian hospital statistics 2008-09. Health services series no. 34. Cat. no. HSE 84. Australian Institute of Health and Welfare, Canberra 2010.
2. AIHW. Australian hospital statistics 2014-15: emergency department care. Health services series no. 65. Cat. no. HSE 168. Australian Institute of Health and Welfare, Canberra 2015.
3. ABS. 3201.0 - Population by Age and Sex, Australian States and Territories, Jun 2009 [cited 26 Oct 2018] Available from:
<http://www.abs.gov.au/AUSSTATS/abs@.nsf/Lookup/3201.0Main+Features1Jun%202009?OpenDocument>
4. ABS. 3101.0 - Australian Demographic Statistics, Jun 2015. [Cited 26 Oct 2018] Available from: URL:
<http://www.abs.gov.au/AUSSTATS/abs@.nsf/Previousproducts/3101.0Main%20Features1Jun%202015>
5. ~~3~~-AIHW. The health of Australia's prisoners. Cat No. PHE 170. Australian Institute of Health and Welfare, Canberra 2013.
6. ~~4~~-Fazel S, Bains P, Doll H. Substance abuse and dependence in prisoners: a systematic review. *Addiction*. 2006;101(2):181-191.

7. ~~5.~~ Herbert K, Plugge E, Foster C, Doll H. Prevalence of risk factors for non-communicable diseases in prison populations worldwide: a systematic review. *The Lancet*. 2012;379(9830):1975-1982.

6. ~~Australian Bureau of Statistics. Prisoners in Australia. ABS cat. no. 4517.0. Canberra: Australian Bureau of Statistics 2013.~~

8. ~~7.~~ McKenna B, Furness T, Brown S, Tacey M, Hiam A, Wise M. Police and clinician diversion of people in mental health crisis from the Emergency Department: a trend analysis and cross comparison study. *BMC Emerg Med*. 2015;15(1):1.

9. ~~8.~~ Lee S, Brunero S, Fairbrother G, Cowan D. Profiling police presentations of mental health consumers to an emergency department. *Int J Mental Health Nurs*. 2008;17(5):311-316

10. ~~9.~~ Gray SD, Fatovich DM, McCoubrie DL, Daly FF. Amphetamine-related presentations to an inner-city tertiary emergency department: a prospective evaluation. *Med J Aust*. 2007;186(7):336-339.

11. ~~10.~~ Miller P, Curtis A, Palmer D, et al. Changes in injury -related hospital
department presentations associated with the imposition of regulatory versus voluntary licensing conditions on licensed venues in two cities. *Drug Alcohol Rev*. 2014;33(3):314-322.

12. ~~11.~~ McLay SV, MacDonald E, Fatovich DM, Alcohol Harm in Emergency Departments I. Alcohol-related presentations to the Royal Perth Hospital Emergency Department: A prospective study. *Emerg Med Australas*. 2017;29(5):531-538.

13. Peters MDJ, Godfrey CM, McInerney P, Baldini Soares C, Khalil H, Parker D. Methodology for JBI scoping reviews. *The Joanna Briggs Institute Reviewers' Manual*. 2015:1-24.
14. ~~12.~~ Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Social Res Methodol*. 2005;8(1):19-32.
15. PRISMA. PRISMA: Transparent reporting of systematic reviews and meta-analyses. [Cited 26 Oct 2018] Available from URL: <http://www.prisma-statement.org/PRISMAStatement/>.
16. ~~14.~~ Sampson M, Zhang L, Morrison A, et al. An alternative to the hand searching gold standard: validating methodological search filters using relative recall. *BMC Med Research Methodol*. 2006;6(1):33.
17. ~~16.~~ NHMRC. Additional levels of evidence and grades for recommendations for developers of guidelines. *National Health and Medical Research Council*. [Cited 26 Oct 2018] Available from URL: <https://www.mja.com.au/sites/default/files/NHMRC.levels.of.evidence.2008-09.pdf>
18. ~~17.~~ Lee S. The characteristics of police presentations to an emergency department in a community hospital. *Austr Emerg Nurs J*. 2006;9(2):65-72 68p.
19. ~~18.~~ Markham D, Gaudins A. Characteristics of frequent emergency department presenters to an Australian emergency medicine network. *BMC Emerg Med*. 2011;11:21-21.
20. ~~19.~~ Charette Y, Crocker AG, Billette I. The judicious judicial dispositions juggle: characteristics of police interventions involving people with a mental illness. *Canad J Psychiatry*. 2011;56(11):677-685.

21. ~~20.~~ Downey LVA, Zun LS, Burke TM. Differences in How Psychiatric Patients Come Into the Emergency Department. *Primary Psychiatry*. 2010;17(8):53-61 59p.
22. ~~21.~~ Knott JC, Pleban A, Taylor D, Castle D. Management of mental health patients attending Victorian emergency departments. *Aust NZ J Psych*. 2007;41(9):759-767
23. ~~22.~~ Meehan T, Stedman T. Trends in the use of Emergency Examination Orders in Queensland since the implementation of the Mental Health Intervention Project. *Austral Psychiatry*. 2012;20(4):287-290.
24. ~~30.~~ Llewellyn P, Arendts G, Weeden J, Pethebridge A. Involuntary psychiatric attendances at an Australasian emergency department: A comparison of police and health-care worker initiated presentations. *Emerg Med Aust*. 2011;23(5):593-599.
25. ~~23.~~ Strote J, Walsh M, Auerbach D, Burns T, Maher P. Medical conditions and restraint in patients experiencing excited delirium. *Am J Emer Med*. 2014;32(9):1093-1096.
26. ~~24.~~ Kothari CL, Rhodes KV. Missed opportunities: emergency department visits by police-identified victims of intimate partner violence. *Ann Emer Med*. 2006;47(2):190-199
27. ~~25.~~ Raina P, Arenovich T, Jones J, Lunskey Y. Pathways into the Criminal Justice System for Individuals with Intellectual Disability. *J Applied Res Intellect Disabil*. 2013;26(5):404-409
28. ~~26.~~ Al-Khafaji K, Loy J, Kelly A-M. Characteristics and outcome of patients brought to an emergency department by police under the provisions (Section 10) of the Mental Health Act in Victoria, Australia. *Int J Law Psychiatry*. 2014;37(4):415-419.
29. ~~27.~~ Strote J, Verzemnieks E, Walsh M, Hutson HR. Use of force by law enforcement: an evaluation of safety and injury. *J Trauma*. 2010;69(5):1288-1293.

30. ~~28.~~ Richoux C, Ferrand I, Casalino E, Fleury B, Ginsburg C, Lejoyeux M. Alcohol use disorders in the emergency ward: choice of the best mode of assessment and identification of at-risk situations. *Int J Emer Med*. 2011;4(1):27-27.
31. ~~29.~~ Bunting PJ, Fulde GWO, Forster SL. Comparison of crystalline methamphetamine ("ice") users and other patients with toxicology-related problems presenting to a hospital emergency department. *Med J Aust*. 2007;187(10):564-566.
32. ~~31.~~ Stevenson RJ, Clark K, Kelliher T. Elbow dislocation secondary to fall during police arrest. *J Forensic Legal Med*. 2014;22:125-126.
33. ~~32.~~ Akar FA, Arbel R, Benninga Z, Dia MA, Steiner-Birmanns B. The Istanbul protocol (manual on the effective investigation and documentation of torture and other cruel, inhuman or degrading treatment or punishment): implementation and education in Israel. *Israel Med Ass J*. 2014;16(3):137-141.
34. ~~33.~~ Wang J-P, Chiu C-C, Yang T-H, Liu T-H, Wu C-Y, Chou P. The Low Proportion and Associated Factors of Involuntary Admission in the Psychiatric Emergency Service in Taiwan. *Plos One*. 2015;10(6):e0129204-e0129204.
35. ~~34.~~ McGarvey EL, Leon-Verdin M, Wanchek TN, Bonnie RJ. Decisions to initiate involuntary commitment: the role of intensive community services and other factors. *Psychiatric Services*. 2013;64(2):120-126
36. ~~35.~~ Puentes K, Ribeiro C, Jardim P, Santos A, Magalhães T. Non-fatal gunshot wounds in the context of intimate partner violence. The importance of a multidisciplinary approach: a case report. *J Forensic Legal Med*. 2011;18(5):221-224.

37. ~~36.~~ Strote J, Verzemnieks E, Walsh M. Emergency department documentation of alleged excessive use of force. *Am J Foren Med Pathol.* 2013;34(4):363-365.
38. ~~37.~~ ACEM. *Guidelines on the implemetation of the Australasian triage scale in Emergency Departments.* [Cited: 26 Oct 2018] Available from URL: https://acem.org.au/getmedia/51dc74f7-9ff0-42ce-872a-0437f3db640a/G24_04_Guidelines_on_Implementation_of_ATS_Jul-16.aspx Australasian College of Emergency Medicine; 2016.
39. ~~38.~~ Chung CH, Fung WT. Detection of gastric drug packet by ultrasound scanning. *Eur J Emerg Med.* 2006;13(5):302-303.
40. ~~39.~~ Lunsky Y, Paquette-Smith M, Weiss JA, Lee J. Predictors of emergency service use in adolescents and adults with autism spectrum disorder living with family. *Emerg Med J.* 2015;32(10):787-792.
41. ~~40.~~ Heslin M, Callaghan L, Barrett B, et al. Costs of the police service and mental healthcare pathways experienced by individuals with enduring mental health needs. *Br J Psychiatry.* Feb 2017;210(2):157-164.
42. ~~41.~~ Wood JD, Watson AC, Fulambarker AJ. The "Gray Zone" of Police Work During Mental Health Encounters: Findings from an Observational Study in Chicago. *Police Q.* Mar 2017;20(1):81-105.
43. ~~42.~~ Green S, Foran J, Kouyoumdjian FG. Access to primary care in adults in a provincial correctional facility in Ontario. *BMC Res Notes.* 2016;9(1):1.

- ~~44.~~ ~~43.~~ ICPR. *World Prison Population List*. [Cited 26 Oct 2018] Available from URL: http://www.prisonstudies.org/sites/default/files/resources/downloads/world_prison_population_list_11th_edition_0.pdf Institute for Criminal Policy Research; 2015.
- ~~45.~~ ~~44.~~ Lamanna D, Shapiro GK, Kirst M, Matheson FI, Nakhost A, Stergiopoulos V. Co-responding police-mental health programmes: Service user experiences and outcomes in a large urban centre. *Int J Ment Health Nurs*. 2018; 27(2):891-900.
- ~~46.~~ ~~45.~~ Shapiro G, Cusi A, Kirst M, O'Campo P, Nakhost A, Stergiopoulos V. Co-responding police-mental health programs: a review. *Admin Policy Mental Health*. 2015;42(5):606-620.
- ~~47.~~ ~~46.~~ Huppert D, Griffiths M. Police mental health partnership project: Police Ambulance Crisis Emergency Response (PACER) model development. *Australasian Psychiatry*. 2015;23(5):520-523.
- ~~48.~~ ~~47.~~ WHO. *Good governance for prison health in the 21st century*. World Health Organization Copenhagen: Organization Regional Office for Europe: World Health Organization; 2013.
- ~~49.~~ ~~48.~~ Møller L, Gatherer A, Jürgens R, Stöver H, Nikogosian H. *Health in prisons: a WHO guide to the essentials in prison health*: WHO Regional Office Europe; 2007.
- ~~50.~~ ~~49.~~ Martin T, Thomas S. Police officers' views of their encounters with people with personality disorder. *J Psychiat Ment Health Nurs*. 2015;22(2):125-132.
- ~~51.~~ ~~50.~~ Hoffman R, Hirdes J, Brown GP, Dubin JA, Barbaree H. The use of a brief mental health screener to enhance the ability of police officers to identify persons with serious mental disorders. *Int J Law Psychiatry*. 2016;47:28-35.

EMM_13300_EMA-2018-204.R2-Figure1.JPG

Table 1. Summary of included studies regarding presentation, demographic and clinical profile, care delivery and outcomes for people brought into EDs by police (BIBP)

Author (Year)	Design	Number BIBP	Demographics of those BIBP Sex, age (median/ mean)	Reason(s) for ED presentation	Clinical characteristics reported for those BIBP	ED characteristics reported for those BIBP	ED care delivery described for those BIBP	Outcomes reported for those BIBP
Al-Khafaji et al., 2014 ²⁶⁻²⁸	Observational study	164 people BIBP	58% Male, 35yrs	Mental health problem	Yes	Yes	Yes	Admitted/ discharged ED LOS
Bunting et al., 2007 ²⁹⁻³¹	Observational study	57 people BIBP	NS	Toxicology problem	NS	NS	NS	NS
Charette et al., 2011 ¹⁹⁻²⁰	Observational study	98 people BIBP	NS	Mental health problem	NS	NS	NS	NS
Chung & Fung, 2006 ³⁸⁻³⁹	Case report	1 person BIBP	Male, 53yo	Drug related	Yes	NS	Yes	Admitted Procedure
Downey et al., 2010 ²⁰⁻²¹	Observational study	82 BIBP	61%Male, 27% aged 40-49yrs	Mental health problem	NS	NS	Yes	NS
Gray et al., 2007 ⁹⁻¹⁰	Observational study	25 people BIBP	NS	Drug (amphetamine) related	NS	NS	NS	NS

Knott et al., 2007 ^{24 22}	Observational study	651 people BIBP	NS	Mental health problem	NS	NS	NS	NS
Kothari & Rhodes, 2006 ^{24 26}	Observational study	53 women BIBP	100%Female, Age NS	Victims of intimate partner violence	NS	NS	NS	NS
Lee et al., 2008 ^{8 9}	Observational study	460 people BIBP	66%Male, mean age: 33yrs	Mental health problem	Yes	Yes	Yes	Admitted
Lee, 2006 ^{17 18}	Observational study	452 patient presentati ons BIBP	67%Male, mean age: 33yrs	Mental health problem	Yes	Yes	Yes	ED LOS Hospital LOS Admitted/ discharged Follow up by community mental health
Llewellyn et al., 2011 ^{30 24}	Observational study	205 people BIBP	47%Male, mean age: 35yrs	Mental health problem	Yes	No	Yes	Admitted
Markham & Graudins 2011 ^{18 19}	Case-control study	447 people BIBP	NS	Frequent ED presenters	NS	NS	NS	NS
McLay et al.,	Observational	18 people	NS	Alcohol related	NS	NS	NS	NS

2017- ^{11 12}	study	BIBP		presentation				
Meehan & Stedman 2012 ²²⁻²³	Observational study	Approx: 21,097 people BIBP	NS	Mental health problem	NS	NS	NS	NS
Puentes et al., 2011- ^{35 36}	Case report	1 person BIBP	Female, 37yo	Intimate partner violence	Yes	NS	Yes	Admitted Follow-up
Raina, et al., 2013 ^{25 27}	Observational study	76 people BIBP	58%Male, Mean age: 32yrs	Intellectual disability	Yes	NS	NS	NS
Richoux et al., 2011- ^{28 30}	Cross-sectional study	NS	NS	Alcohol related presentation	NS	NS	NS	NS
Stevenson et al., 2014 ³¹⁻³²	Case report	1 person BIBP	Male, 24yo	Injury	NS	NS	Yes	Admitted
Strote et al., 2014 ²³⁻²⁵	Observational study	43 people BIBP	95%Male, 30yrs	Excited delirium	Yes	No	Yes	Admitted / discharged
Strote et al., 2010 ^{27 29}	Observational study	187 people BIBP	NS	Required police use of force	Yes	No	Yes	Admitted / discharged

NS: not specified; LOS: length of stay; ED: emergency department; yo: years old; yrs: years; BIBP: brought in by police

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Crilly, J;Johnston, ANB;Wallis, M;Polong-Brown, J;Heffernan, E;Fitzgerald, G;Young, JT;Kinner, S

Title:

Review article: Clinical characteristics and outcomes of patient presentations to the emergency department via police: A scoping review

Date:

2019-08-01

Citation:

Crilly, J., Johnston, A. N. B., Wallis, M., Polong-Brown, J., Heffernan, E., Fitzgerald, G., Young, J. T. & Kinner, S. (2019). Review article: Clinical characteristics and outcomes of patient presentations to the emergency department via police: A scoping review. EMERGENCY MEDICINE AUSTRALASIA, 31 (4), pp.506-515. <https://doi.org/10.1111/1742-6723.13300>.

Persistent Link:

<http://hdl.handle.net/11343/285881>