DR. ADAM MILLER (Orcid ID : 0000-0002-1632-7206)
 DR. COLLIN W AHRENS (Orcid ID : 0000-0002-0614-9928)

Article type : Original Article

Local and regional scale habitat heterogeneity contribute to genetic adaptation in a commercially important marine mollusc (*Haliotis rubra*) from southeastern Australia

A. D. Miller^{1,2*}, A. A. Hoffmann³, M. H. Tan^{1,2}, M. Young¹, C. Ahrens⁴, M. Cocomazzo¹, A. Rattray¹, D. A. Ierodiasconou¹, E. Trembl¹, C. D. H Sherman^{1,2}

¹ Deakin University, Geelong, School of Life and Environmental Sciences, Centre for Integrative Ecology, Warrnambool, Victoria, Australia

² Deakin Genomics Centre, Deakin University, Geelong, VIC 3220, Australia

³ School of BioSciences, Bio21 Institute, The University of Melbourne, Parkville, Victoria, Australia

⁴ Hawkesbury Institute for the Environment, Western Sydney University, Hawkesbury, New South Wales, Australia

*Corresponding author: Adam D. Miller (a.miller@deakin.edu.au)

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/MEC.15128](https://doi.org/10.1111/MEC.15128)

This article is protected by copyright. All rights reserved

34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62

Abstract

Characterising adaptive genetic divergence among conspecific populations is often achieved by studying genetic variation across defined environmental gradients. In marine systems this is challenging due to a paucity of information on habitat heterogeneity at local and regional scales and a dependency on sampling regimes that are typically limited to broad longitudinal and latitudinal environmental gradients. As a result, the spatial scales at which selection processes operate and the environmental factors that contribute to genetic adaptation in marine systems are likely to be unclear. In this study we explore patterns of adaptive genetic structuring in a commercially-harvested abalone species (*Haliotis rubra*) from south-eastern Australia, using a panel of genome-wide SNP markers (5,239 SNPs), and a sampling regime informed by marine LiDAR bathymetric imagery and 20-year hindcasted oceanographic models. Despite a lack of overall genetic structure across the sampling distribution, significant genotype associations with heterogeneous habitat features were observed at local and regional spatial scales, including associations with wave energy, ocean current, sea surface temperature, and geology. These findings provide insights into the potential resilience of the species to changing marine climates and the role of migration and selection on recruitment processes, with implications for conservation and fisheries management. This study points to the spatial scales at which selection processes operate in marine systems and highlights the benefits of geospatially-informed sampling regimes for overcoming limitations associated with marine population genomic research.

Keywords: abalone, population genomics, genetic adaptation, environmental heterogeneity, LiDAR, hindcasted oceanographic models, fisheries management, marine conservation

Introduction

Modern genomic approaches have revolutionized the field of population genetics, providing unprecedented sensitivity for resolving patterns of genetic variation between individuals and populations (Davey *et al.* 2011; Eklom & Galindo 2011). Notably, these technologies help characterise patterns of adaptive genetic diversity among conspecific populations (Stapley *et al.* 2010; Savolainen *et al.* 2013). Reduced genome representation, whole genome, and transcriptome sequencing approaches are being used routinely for describing signatures of selection and

identifying putative candidate genes that underlie adaptive differences between species populations (Stapley *et al.* 2010; Ahrens *et al.* 2018). Studies of this nature ideally require sampling regimes that allow genetic variation to be contrasted among conspecific populations distributed across defined environmental gradients (Frichot *et al.* 2013). Most have been carried out in terrestrial systems where habitat heterogeneity is relatively easy to measure and is often well described, with many demonstrating evidence of selection resulting from environmental variation at various spatial scales (Buehler *et al.* 2013; Jordan *et al.* 2017; Termignoni-Garcia *et al.* 2017; Pfeifer *et al.* 2018). In contrast, information on the heterogeneity of marine habitats, particularly benthic habitats, is limited for many coastlines, hindering effective sampling to test for adaptive genetic diversity within marine species. Instead, sampling regimes for marine studies are typically limited to broad-scale environmental gradients (Nielsen *et al.* 2009; Hess *et al.* 2013; Milano *et al.* 2014), and tend to overlook habitat variation at local scales. As a result, the spatial scales at which selection processes operate and the environmental factors that contribute to genetic adaptation in marine systems can be unclear.

Advances in marine geospatial science offer new opportunities for overcoming these limitations (Grummer *et al.* 2019). LiDAR and SONAR technologies are being increasingly used for imaging seabeds around the world and characterising the complexity and connectivity of benthic marine habitats (Brown *et al.* 2011; Ierodiaconou *et al.* 2011; Lecours *et al.* 2015; Young *et al.* 2016) (Figure S3). These types of datasets are important for marine navigation, ecological research, and conservation and fisheries management (Pickrill & Todd 2003; Brock & Purkis 2009; Cogan *et al.* 2009), and as a result, a variety of international and national seafloor mapping projects have been initiated in recent years. These include the GEBCO Seabed 2030 Project which has an ambitious target of mapping the world ocean by 2030 (Mayer *et al.* 2018). In addition to information on the benthic structure of the seafloor, an understanding of the dynamic nature of the physical marine environment is being enhanced by contemporary and hindcasted models of physical oceanographic variables (e.g. hydrodynamics and sea surface temperature) for coastal and offshore marine environments around the world (Burrows *et al.* 2011; Young *et al.* 2011; Hemer *et al.* 2013). The integration of physical oceanographic models with seafloor maps can be used to effectively quantify habitat heterogeneity across marine systems. In turn, there is an opportunity to overcome sampling constraints associated with marine population genomic research, and to develop efficient strategies to quantify genetic diversity across environmental gradients (Grummer *et al.* 2019).

Understanding patterns of contemporary genetic structure and the presence of standing genetic variation for adaptation to changing environmental conditions helps direct effective management approaches for threatened, invasive and commercially important marine species (Hoffmann & Sgro 2011; Weeks *et al.* 2017; Miller *et al.* 2018). In particular, research geared toward assessing the extent of genetic structure across broad geographical gradients, and genetic

variation available for adaptation to more acidic and higher temperature environments, can quantify the resilience of marine species under climate change (Hoegh-Guldberg & Bruno 2010; Torda *et al.* 2017; Miller *et al.* 2018). Additionally, adaptive genetic divergences among populations, particularly on finer spatial scales, can assist in deciphering the influence of natural selection on recruitment processes (Marshall *et al.* 2010). Studies on marine fish and invertebrates have demonstrated that adaptive variation can be maintained despite high levels of gene flow (Galindo *et al.* 2009; Martinez-Fernandez *et al.* 2010; Hess *et al.* 2013; Solas *et al.* 2013; Milano *et al.* 2014; Sandoval-Castillo *et al.* 2018), suggesting that selection can influence recruitment processes as some potential recruits are expected to be maladapted to certain habitat types (Marshall *et al.* 2010). Information on gene flow and population connectivity informs whether populations can replenish naturally following disturbance and depletion, and when management intervention is required (Whiterod *et al.* 2017). To assist intervention activities, such as translocations and reseedling, information on neutral and adaptive genetic differentiation among populations is needed to help identify environmentally compatible genotypes (Ward 2006; Weeks *et al.* 2011). This is pertinent to commercial fisheries (Hauser & Carvalho 2008; Waples *et al.* 2008), where stock augmentation activities are being adopted to assist recovery after depletion (Bell *et al.* 2005; Roodt-Wilding 2007).

Abalone are a group of marine benthic molluscs belonging to the family Haliotidae that are targeted commercially in eleven countries, forming an important global industry worth approximately \$US180 million annually (Gordon & Cook 2013). In recent decades many abalone fisheries have collapsed due to over-exploitation, extreme weather events, environmental change and disease, with a number of target species now listed as endangered or considered ‘species of concern’ (Leiva & Castilla 2001; Gruenthal & Burton 2005; Kashiwada & Taniguchi 2007). Current management of abalone fisheries assumes limited connectivity, with stock viability dictated by recruitment from local sources. This assumption is based on ecological studies suggesting that the planktonic larval dispersal is limited for many abalone species (Hahn 1988; Prince *et al.* 1987; Mcshane *et al.* 1988; Prince *et al.* 1988). However, genetic studies of various Haliotids indicate conflicting patterns of population connectivity and genetic structure (Chambers *et al.* 2006; Gruenthal *et al.* 2007; Gruenthal & Burton 2008; Piggott *et al.* 2008; Miller *et al.* 2014), suggesting that larval dispersal and population connectivity are spatially and temporally variable and influenced by abiotic factors such as ocean currents and wave energy (Miller *et al.* 2016). While these studies have provided novel insights into neutral genetic structuring and population connectivity within abalone species, there is little information on adaptive genetic differentiation, the influence of selection on recruitment processes, and the likely resilience of Haliotids to environmental change.

141 Recent genomic analyses have revealed broad geographic patterns of putatively adaptive
142 genetic structure in connected metapopulations of *Haliotis* species from Australia and North
143 America. Analyses of reduced genome representation data from Western Australian greenlip
144 abalone (*Haliotis laevis*) provided evidence of adaptive genetic variation correlated with sea
145 surface temperature and oxygen concentrations (Sandoval-Castillo *et al.* 2018). Genotype by
146 environment association analyses performed by Sandoval- Castillo *et al.* (2018) revealed genetic
147 variants under selective pressure, including variants in annotated genes with functions related to
148 high temperature and/or low oxygen tolerances. Similarly, analyses of transcriptome-wide
149 polymorphisms in red abalone (*Haliotis rufescens*) from the west coast of the United States
150 indicated signatures of local genetic adaptation correlated with carbon dioxide (CO₂) and oxygen
151 concentrations, and involving genes associated with biomineralization, energy metabolism, heat-,
152 and disease- or hypoxia-tolerance (De Wit & Palumbi 2013). These studies provide insights into
153 standing genetic variation within Haliotid species for adaptation to more acidic and warmer
154 environments emerging under climate change. They also indicate that adaptive genetic differences
155 can persist in Haliotid species despite potential for panmixia, at least along broad geographic
156 gradients. Selection processes may also operate at fine spatial scales in Haliotids given the
157 heterogeneous nature of their benthic habitats, but this remains to be considered.

158 The current study builds on previous population genomic research on blacktip abalone (*H.*
159 *rubra*) which showed a lack of overall genetic structure and evidence of a connected
160 metapopulation spanning ~1400 km of the south-eastern Australian coastline (Miller *et al.* 2016).
161 Here we use a hierarchical sampling regime guided by bathymetric LiDAR imagery and hindcasted
162 oceanographic datasets to explore adaptive genetic differentiation in *H. rubra* across the same study
163 area. Using a panel of genome wide single nucleotide polymorphism (SNP) markers derived from
164 genomic sequencing, we contrast genetic variation among animals distributed across environmental
165 gradients spanning local (10s to 100s of meters) to regional (100s of kilometers) spatial scales,
166 using novel geospatial datasets. We hypothesize that selection processes operate on finer spatial
167 scales than currently assumed in abalone species, considering: 1) the highly heterogeneous nature of
168 abalone habitats (potential drivers of selection; Jalali *et al.* 2015), 2) the high levels of genetic
169 polymorphism present in the species (potential gene variants for selection to act upon; Miller *et al.*
170 2016), and 3) pre-existing evidence for standing genetic variation associated environmental factors
171 that vary on regional scales (evidence of existing adaptive genetic differentiation; De Wit &
172 Palumbi 2013; Sandoval-Castillo *et al.* 2018). This study provides insights into the spatial scales of
173 selection processes in abalone species, and identifies putative environmental factors contributing to
174 genetic adaptation during recruitment, with implications for fisheries management.

175
176

177

178 **Methods**

179 *Experimental design and sample collection*

180 Using a combination of temperature observations from satellite data (sea surface temperature),
181 modelled oceanographic data (wave energy, ocean currents), and LiDAR-derived seafloor structure
182 (substrate type, VRM), we developed a hierarchical and spatially replicated sampling design
183 comprising 29 reef locations representing different habitat types and spanning the entire Victorian
184 coastline, encompassing the Western, Central and Eastern Zone abalone fisheries. The selection of
185 reef locations was optimised to include populations from a range of habitat types varying in
186 seasonal sea surface temperature (SST), local geology (substrate type and vector ruggedness
187 measure), exposure to wave energy and ocean currents, and sea surface temperature (Table 2).
188 Habitat maps and hindcasted oceanographic datasets indicate that each of these variables show
189 significant heterogeneity at both local (10s to 100s of meters) and regional (100s of kilometres)
190 scales (Ierodionou *et al.* 2011; Young *et al.* 2016). Temperature was a key environmental
191 parameter as the Victorian coastline is characterised by a strong longitudinal thermal gradient
192 driven by upwelling systems and converging ocean currents (Sandery & Kaempf 2007; Colton &
193 Swearer 2012). Processed sea surface temperature (SST) datasets were sourced from the Integrated
194 Marine Observing System (IMOS) – IMOS is a national collaborative research infrastructure,
195 supported by the Australian Government. Mean winter and summer SST, and annual temperature
196 range for the years 1995-2015 were evaluated, after which mean winter SST was omitted due to a
197 high correlation with summer estimates. Wave energy and ocean current information were derived
198 from a coupled hydrodynamic and wave model developed by Water Technology (more detailed
199 information on the development of these hydrodynamic models can be found in the supplementary
200 material). As the 500 m resolution of the hydrodynamic model did not resolve areas that were
201 sheltered due to fine-scale coastal geometry (such as headlands and exposed offshore reefs), we
202 calculated fetch distances in 1 degree increments from each site in a 30 degree sector, 15 degrees
203 either side of annual mean swell direction using a custom 1:10000 scale digital coastline. Mean
204 fetch values for each site were used to weight local estimates of wave power and orbital velocity to
205 account for local variability in swell protection. The geology of benthic reef habitats is variable
206 within and across regions of the Victorian coastline, and consists of basalt, calcareous and granite
207 reef complexes. Information on substrate type for each location was inferred from existing
208 terrestrial geology maps and inferences from bathymetric LiDAR intensity data and expert
209 knowledge. Structure of these geologic classes was derived from the 5 m resolution LiDAR
210 bathymetry dataset using the vector ruggedness (VRM) tool (Walbridge *et al.* 2018) within ArcGIS
211 10.5. VRM is a measure of surface roughness that is calculated using a 3 x 3 moving window where
212 each cell in the window is assigned a VRM value based on three-dimensional location of the cell

213 centre, slope, and aspect (Wallbridge et al. 2018). Correlation analyses implemented in the ggpubr
214 R package (<https://cran.r-project.org/web/packages/ggpubr/index.html>) confirmed the independence
215 of all environmental variables except summer sea surface temperature and seasonal temperature
216 range which correlated moderately (Pearson's $r = 0.6$, $P < 0.01$).

217 Field sampling was conducted in 2016 with the assistance of commercial survey divers
218 contracted by the Victorian Fisheries Authority. A total of 870 abalone tissue biopsies were
219 collected from 29 localities (30 samples per location) spanning the Victorian Western, Central and
220 Eastern Zone fisheries and provided for genomic analysis (Table 1 and Figure 2). At each location,
221 individual abalone were collected within a 25 m radius of an anchored vessel and were assumed a
222 random sample of the site. Multiple size classes were intentionally sampled to avoid sampling
223 single cohorts (potentially related individuals) that might drive ambiguous estimates of genetic
224 structure. From these samples, we collected 50 mg tissue biopsies from each individual using sterile
225 dissection tools to avoid cross-contamination. The biopsied material was transferred to 2-mL
226 microcentrifuge tubes containing either 100% ethanol or RNAlater (QIAGEN) and transported to
227 the laboratory. A total of 25 samples per location were used for genomic analysis.

228

229 *DNA extraction and Genotyping by Sequencing library construction*

230 Total genomic DNA was extracted from 10 mg of muscle tissue using the Qiagen DNeasy 96 Blood
231 and Tissue Kit (Venlo, Limburg, NL), and reduced representation genome libraries were prepared
232 with a modified genotyping by sequencing (GBS) protocol of Elshire *et al.* (2011). Three hundred
233 nanograms of genomic DNA from each individual were digested in 20 μ L reaction containing four
234 units of restriction enzymes MseI and SbfI for 2 h at 37 °C. Digestion products were then ligated to
235 modified P1 and P2 adapters with unique barcode combinations to allow for subsequent
236 multiplexing of all individuals. Fifty μ L ligations were performed containing the enzyme digested
237 DNA, 1.125 ng of P1 and P2 adapters, 400 units of T4 ligase and T4 buffer (New England Biolabs,
238 Beverly MA, USA). Ligations were incubated at 16 °C for 90 min followed by a 30 min of
239 denaturation at 80 °C. Adapter ligated DNA fragments were purified using a Qiagen MinElute PCR
240 purification kit (Redwood City, CA, USA), eluted in 20 μ L of ddH₂O and subsequently used for
241 PCR amplification. Fifty μ L PCRs were performed using 29 MyTaq™ HS Mix (Bioline, Taunton,
242 MA, USA), and containing 0.2 μ L each of Illumina Dual Index Sequencing Primers 1 & 2 (Illumina
243 Inc., San Diego, CA, USA) and 10 μ L of above purified DNA. PCR conditions were as follows: 95
244 °C for 1 min, 24 cycles of 95 °C for 30 s, 65 °C for 30 s, 72 °C for 30 s and a final extension step of
245 72 °C for 5 min. DNA quantitation and qualitative analysis of individual PCR products were
246 performed on a MCE_-202 MultiNA with a DNA-1000 kit (Shimadzu, Kyoto). Samples were then
247 pooled equimolar into groups of 71 samples (11 pooled libraries in total), with each pooled library

being sequenced on a single Illumina HiSeq 4000 (Illumina, San Diego) lane by MacroGen (Seoul, Korea).

Bioinformatics processing and genotyping

Illumina HiSeq 4000 runs yielded an average of 380 million assigned 100 base pair (bp) paired-end reads per lane, providing an average of 5.2 million paired-end reads per sample for genomic analysis. Raw sequences were first processed using the Trimmomatic V0.36 program (Bolger *et al.* 2014) by trimming the raw reads to 80 bp length and discarding all reads that had a Phred score below 20. We used the *de novo* program from *Stacks* 1.44 (Catchen *et al.* 2013) to create a catalogue of SNPs and genotypes for all individuals. Because we expected a high level of intra-population genetic diversity based on microsatellite and SNP data from a previous study (Miller *et al.* 2016), we tested several combinations of parameters that allowed *de novo* assembly of GBS loci from orthologous sequences while rejecting paralogous sequences. The final parameter settings included the maximum distance allowed between stacks ($M = 3$), and the distance between loci in the catalogue ($n = 3$), while the minimum depth of coverage to form a stack was kept constant ($m = 3$). SNPs and genotypes at each locus were called using a maximum likelihood framework with default *Stacks* 1.44 parameters at the significance level of 5%. A subset of polymorphic GBS loci present in at least 70% of individuals at each sampling location (Boehm *et al.* 2015; Dierickx *et al.* 2015) and having SNPs with a global minor allele frequency of ≥ 0.05 were used for downstream genomic analyses.

Estimating overall genetic structure

SNP frequencies over all loci were initially contrasted between all 29 sampled locations to determine patterns of overall genetic structure and population connectivity. Analyses were repeated following the removal of candidate loci potentially under directional selection (refer to section immediately below), but had little effect. The software GenoDive (Meirmans & Van Tienderen 2004) was used to calculate the average number of allele per locus (N_a), observed (H_o) and expected (H_e) heterozygosity across loci, for each sample location, as well as global and pairwise measures of population differentiation including both F_{ST} and D_{EST} (Weir & Cockerham 1984; Jost 2008). Correlations between genetic and geographic distances were assessed using Mantel tests implemented in the adegenet package for R (Jombart 2008; Jombart & Ahmed 2011), with significance determined by permutation (10,000 randomizations). A discriminant analysis of principal components (DAPC) was implemented in the adegenet package for R (Jombart 2008; Jombart & Ahmed 2011), to obtain a graphical depiction of patterns of genetic structure. The number of genetic clusters was then defined using k-means, a clustering algorithm that looks for the value of K that maximizes the variation between groups. The Bayesian Information Criterion (BIC)

was calculated for $K = 1-29$ and the K value with the lowest BIC was selected as the optimal number of clusters. A discriminant analysis was then performed using the function DAPC, implemented in R, to describe the genetic clusters.

Detection of candidate SNPs

To detect putative genomic signatures of selection, we conducted genotype by environment association analyses (GEAs) using three complementary models; Latent Factor Mixed Models (LFMM; (Frichot *et al.* 2013)), the Bayesian method available in BayPass V1.01 (Gautier 2015) and redundancy analysis (RDA) (Forester *et al.* 2018). LFMM and BayPass accounted for potentially confounding allele frequency differences due to population structure in a mixed linear model framework, but in different ways: LFMM estimates GEAs when simultaneously correcting for population structure with latent factors, while BayPass uses a neutral covariance matrix constructed from population allele frequencies.

We first used LFMMs (LEA 1.3; (Frichot *et al.* 2013)) to detect genotype–environment correlations with the continuous variables described above. The number of latent factors was set to two ancestral clusters based on the cross-entropy score results using the *snmf* function in the LEA package. Cross-entropy scores evaluate the fit of the statistical model to the data using a cross-validation technique (Frichot & Francois 2015). The lowest value returned from the cross-entropy scores represents the optimal k -value and was used to correct for confounding effects from population structure in LFMM. LFMM naively imputes missing data, therefore we imputed missing data from the *snmf* run with the lowest cross-entropy score with two ancestral clusters, as described in the LFMM manual. The *lfmm* function was used in the LEA package to identify SNP–environment associations with the settings set to 10,000 iterations of the Gibbs Sampling algorithm with 5,000 burn-in iterations, outputs were averaged over 10 independent runs. The p -values were re-adjusted in R using the Benjamini-Hochberg method for multiple tests on the combined z -scores from all runs, as recommended in the LFMM manual (Frichot *et al.* 2013). Significance was assessed with a false discovery rate (FDR) threshold of 0.001 (i.e. q -value in LFMM terminology), as advised for these kinds of analyses (De Villemereuil *et al.* 2014).

Second, we explored GEAs with BayPass V1.01 (Gautier 2015) under the auxiliary (AUX) covariate mode (-covmcmc and -auxmode flags), after scaling the variables with the -scalecov flag. The underlying models explicitly account for the covariance structure among the population allele frequencies that originates from the shared history of the populations under study, through the estimation of the population covariance matrix Ω , which renders the identification of SNPs subjected to selection less sensitive to the confounding effect of demography (Bonhomme *et al.* 2010; Gunther & Coop 2013). The auxiliary covariate model specifically involves the introduction of a binary auxiliary variable to classify each locus as associated or not. This allows the

320 computation of posterior inclusion probabilities (and Bayes Factors) for each locus while explicitly
321 accounting for multiple testing issues. For each SNP, the Bayes factor was expressed in deciban
322 units (dB) via the transformation $10 \log_{10}(\text{BF})$. Significance was assessed based on the Bayes
323 Factor (BF) between models and according to Jeffrey's rule (Jeffrey 1961); that is, markers with
324 moderately ($3 < \text{BF} < 10$), strong ($10 < \text{BF} < 20$) to decisive ($\text{BF} > 20$) evidence were retained as
325 potential candidates under positive selection.

326 Finally, we used redundancy analysis (RDA) to identify candidate SNPs under selection, a
327 multivariate approach known to reduce false positive and identify unique sets of adaptive loci under
328 weak selection (Forester *et al.* 2018). We modified R script from Forester *et al.* (2018) to test for
329 correlations between allele frequencies and the environment by using the function *rda* implemented
330 in the package *vegan*, version 2.3-5 package (Oksanen *et al.* 2013). Components of the RDA
331 significantly correlated to the constrained axis ($\alpha = 0.05$) were retained and a Fisher transformation
332 was applied to correct for sample size (Bourret *et al.* 2014). SNPs were retained and considered
333 candidates only if correlated with the retained components. In order to compare the outputs from the
334 three methods, Venn diagrams were generated using the *VennDiagram* package for R (Chen &
335 Boutros 2011).

336 Discriminant analysis of principal components (DAPC) were implemented in the *adegenet*
337 package for R (Jombart 2008; Jombart & Ahmed 2011) to obtain a graphical depiction of patterns
338 of genetic structure based on datasets consisting of candidate SNPs for each environmental variable
339 identified by BayPass (BFs > 3), LFMM (q -values < 0.001), and RDA. As outlined previously, the
340 number of genetic clusters was defined using k-means, with Bayesian Information Criterion (BIC)
341 calculated for $K = 1-29$ and the K value with the lowest BIC selected as the optimal number of
342 clusters. A discriminant analysis was then performed using the function *DAPC* to efficiently
343 describe the genetic clusters.

344 Generalized Dissimilarity Modeling (GDM; (Ferrier *et al.* 2007)) was subsequently
345 performed on all SNPs found to be significantly correlated with the environmental variables
346 (derived from LFMM, BayPass and RDA analyses) following the methods described in Fitzpatrick
347 & Keller (2015). First, F_{ST} matrices were created separately for each of the putatively adaptive
348 SNPs using the web version of GenePop (Raymond & Rousset 1995). A GDM analysis, using the
349 *gdm* package v 1.3.7 in R (Manion *et al.* 2018) was independently performed on the F_{ST} matrix for
350 each candidate SNP to estimate allelic turnover through climatic space (deviations in allele
351 frequency associated with environment type). GDM analysis removes variation that is associated
352 with geographic distance and the remaining genetic variation is compared to climate (the y-axis on
353 the spline plots generated by GDM are therefore labeled partial genetic distance). SNPs explaining
354 zero deviance or with missing data from at least five populations were not plotted. Spline plots
355 showed the relative importance of climate on allelic turnover, and assist in identifying

environmental thresholds where selection pressure is expected to be greatest. The significance of climate variables on the dependent variable (in this case a SNP) is calculated by the difference between two models, with and without the climate variable, and is termed deviance (Ferrier et al. 2007), which is analogous to percentage of variation explained. Note that GDM analyses could only be performed on SNPs associated with continuous environmental variables. Allelic turnover was visualised in geographic space for environmental variables SST and VRM by using pairwise F_{ST} matrices generated from the combination of SNPs deemed to be significant with its respective environmental factor based on initial GDM analyses. GDM analyses were performed on each matrix with outputs being transformed with the respective environmental raster layer using the function *gdm.transform*. Analysis of principal components was then performed on the transformed data using *prcomp*, and subsequently predicted across geographic space using the predict function (raster package) to create a raster layer comprising of allelic turnover. The results were saved as a geotiff file and visualised in ArcGIS (ESRI 2018). Analyses could not be performed for waver power or orbital velocity environmental variables as fetch corrected exposure estimates are not available for near-shore habitats at a statewide scale.

Functional annotation

Functional annotations were performed for SNP loci showing strong and significant correlations with each environmental covariable based on LFMM and BayPass (BFs > 10 and q-values < 0.00001). Tag sequences containing candidate SNP loci of interest were searched against five *Haliotis* transcriptomes: *H. laevigata* (Shiel et al. 2015), *H. midae* (Van Der Merwe et al. 2011), *H. rubra* (Deakin University, unpublished), *H. rufescens* (De Wit & Palumbi 2013), and *H. tuberculata* (Harney et al. 2016) using BLAST+ v2.6.0 (*blastn*, e-value 1e-5) (Camacho et al. 2009). Sequences with hits to any transcriptome were further annotated against the UniProtKB/Swiss-Prot protein database (Bateman et al. 2017) based on sequence homology (*blastx*, e-value 1e-5). Additional gene ontology term analysis was performed through InterProScan v5.28.67 (Jones et al. 2014) and visualized with WEGO (Ye et al. 2006).

Results

Genotyping and overall population structure

Genome scans of 725 *Haliotis rubra* specimens from 29 locations distributed across the Victorian western, central and eastern zone abalone fisheries were performed by reduced genome representation sequencing. Illumina sequencing yielded a total of 3,794,534,348 base paired reads, providing an average of 5.2 million base-paired reads per sample (range = 0.5 to 20.2 million). *De*

392 *novovo* assembly using the STACKS bioinformatics pipeline yielded a total of 4,665,377 polymorphic
393 GBS tags, and 1,108,067 tags containing 5,239 SNPs. Levels of diversity were largely consistent
394 across sites with the percentage of polymorphic loci ranging from 72.7% to 96.8% (mean %P =
395 86.0%) (Table S1) and expected heterozygosities ranging from 0.24 to 0.28 (mean H_E = 0.27).

396 Global F_{ST} was weak but significantly different from zero (F_{ST} = 0.009, 95% CIs = 0.008 –
397 0.009; P < 0.001; D_{EST} = 0.003, 95% CIs = 0.003 – 0.004, P < 0.001) indicating some genetic
398 structuring among sample locations. Significant genetic structuring appears to be associated with
399 the Victorian eastern zone and Port Phillip Bay fishing stocks, as all pairwise F_{ST} comparisons with
400 sites from these regions were significantly different from zero (Table S2). In contrast all pairwise
401 estimates of F_{ST} between western zone and central zone fishing stocks (except for Port Phillip Bay)
402 did not differ significantly from zero. Excluding the eastern zone and Port Phillip Bay fishing
403 stocks, global F_{ST} did not differ significantly from zero (F_{ST} = 0.002, 95% CIs = 0 – 0.002; P >
404 0.050; D_{EST} = 0.001, 95% CIs = 0 – 0.001, P > 0.050) indicating a lack of genetic structuring
405 between western and central zone fishing stocks. These patterns of population genetic
406 differentiation are graphically depicted by DAPC analyses which retained 300 principal
407 components, and the first two discriminant functions, capturing 75% of the total variance within the
408 SNP dataset (Figure S4A), and with k-means identifying three population clusters. When plotted
409 across the x- and y-axes, individuals from the eastern zone and Port Phillip Bay sample locations
410 cluster separately from each other and from the main population cluster consisting of admixed
411 individuals from the western and central zone fisheries. After removing the eastern zone and Port
412 Phillip Bay sample locations (retaining 250 principal components, and the first two discriminant
413 functions capturing 65% of the total variance), no genetic structuring between the western and
414 central zone fisheries is apparent. (Figure S4B). Mantel tests showed a moderately strong
415 relationship between genetic and geographic distance (Mantel r = 0.43, P < 0.01). This relationship
416 is no longer significant once the sites from Port Phillip Bay and the Eastern Zone are removed from
417 the analysis (Mantel r = 0.01, P > 0.05) consistent with high levels of gene flow across the Western
418 and Central Zone fisheries (Figures S5A and B).

419

420 *Detection of candidate SNPs*

421 BayPass, LFMM and RDA each identified a number of candidate SNP loci exhibiting significant
422 genotype-by-environment associations for each of the environmental covariables tested (Table 3).
423 BayPass identified between 14 (0.26%) and 76 (1.45%) SNPs with significant correlations
424 ($\log_{10}(BF) > 3$) for each of the environmental covariables (average = 49.28 SNPs; 0.94%), while a
425 larger number of SNP loci showed significant correlations with summer SST (658 SNPs; 12.60%).
426 LFMM identified between 12 (0.2%) and 60 SNPs (2.8%) showing significant correlations ($q <$
427 0.001) with each of the environmental covariables (average = 28.42 SNPs; 0.65%). Note that

428 because LFMM analyses could not be performed on categorical data, estimates for reef geology are
429 not provided. RDA identified between 0 (0%) and 64 SNPs (1.2%) showing significant correlations
430 with each of the environmental covariables, except for seasonal temperature range (average = 28.43
431 SNPs; 0.54%; Figure 2). For each of the environmental covariables, between 0 and 22 SNPs were
432 identified as exhibiting significant correlations by both BayPass, LFMM and /or redundancy
433 analyses. Venn diagrams displaying the number of candidates identified by each GEA method, and
434 overlaps between methods, are provided in Figures S6 – 9. Overlap of candidate loci between
435 methods was low which was expected for several reasons. Specifically, we chose GEA methods that
436 identify candidate loci with varying sensitivities to detecting loci under weak or strong selection,
437 use different methods for controlling for overall population structure, and adopt different yet
438 complementary mathematical algorithms for identifying genotype-environment associations.
439 Candidates from each GEA method were used in post-hoc analyses to better identify true positives,
440 particularly ones that occur in genic regions (annotation) and have strong relationships with
441 environments that cannot be explained by geography (GDM).

442 Discriminant Analysis of Principle Components (DAPC) was performed on SNP loci
443 showing significant correlations with each environmental covariable based on LFMM, BayPass and
444 RDA analyses to provide a graphical depiction of genetic structuring among environmental
445 categories. On each occasion genetic structure was captured by the first two principal components
446 indicated by eigenvalues in Figures 2-5, with each analysis identifying a single genetic cluster using
447 k-means. Patterns of genetic differentiation for these loci are graphically depicted following the
448 retention of principal components and discriminant functions that captured between 60 and 85% of
449 the total variance. To visualise patterns of genetic structuring, populations were colour coded based
450 on their respective habitat classifications (Figures 3 – 6). For geology, which is a categorical
451 variable, this involved colour coding populations sampled from the various substrate types (blue =
452 basalt, black = calcareous, red = granite; Figure 3A). For the continuous environmental variables,
453 populations were colour coded based on relative environmental scores (low, intermediate, high;
454 Table 2). When plotted across the x- and y-axes, patterns of genetic structuring can be seen
455 differentiating populations from basalt, calcareous and granite reef systems. Likewise, there is
456 genetic structuring between populations from reef complexes of low, intermediate and high VRM
457 values (Figure 3B). For summer wave power and orbital velocity, populations with low,
458 intermediate and high environmental values separate across the axes, but patterns associated with
459 winter are less clear (Figures 4 and 5). DAPC plots indicate genetic structuring between
460 populations of varying seasonal temperature ranges, while patterns associated with summer SST are
461 not clear despite the large number of SNP loci having significant correlations with this covariate
462 (Figure 6).

All SNPs identified as having significant associations with environmental variables were used for GDM analysis to quantify allelic turnover through climatic space (Figure 7; Table 4). The three environmental variables that explained the greatest deviance on average were winter wave power (5.31%), VRM (6.42%), and summer wave power (9.55%). In contrast, seasonal temperature range explained the least deviance, with only one SNP explaining a very low (0.28%) level of variation, as presented in the spline plot in Table 4. The majority of SNPs plotted against each environmental variable showed small to moderate responses to climate (i.e. deviance explained; Table 4). However, several SNPs explained greater than 10% of the variation, suggesting directional selection. For example, winter wave power associated with SNPs 94891_79 and 29000_33 and showed strong clines with deviances of 13.47% and 13.2%, respectively; summer wave power associated with SNPs 12085_77, 14150_22, and 29000_33 and had deviances of 17.68%, 24.21%, and 14.88%, respectively; summer sea surface temperature associated with SNP 15442_53 had a deviance of 22.03%; and VRM associated SNPs 15487_52 and 35093_14 and showed strong clines with deviances of 18.96% and 18.12% (Table 4). In contrast, SNPs associated with summer and winter orbital velocity explained little variation (< 5% deviance). The GDM spline plots also demonstrate defined points of allelic turn-over for many SNPs in climate space (Figure 7). Specifically, allelic turn-over occurs at five and three kW/m for winter and summer wave power (Figure 7A and B), approximately 18 °C for summer sea surface temperature (Figure 7C), and 0.0009 for VRM (Figure 7G). These values may be interpreted as environmental thresholds influencing selection and separating adaptive genotypes but should be treated with caution due to the shallow genome coverage dataset used in this study. These thresholds are further depicted by allelic turnover maps which show two very distinct patterns of selection: broad-scale turnover in SST where allelic changes occur at regional scales (Figure 8A) and fine-scale turnover in VRM where allelic changes occur at local spatial scales (Figure 8B).

Functional Annotation

Tag sequences containing candidate SNP loci which showed strong and significant correlations with each environmental covariable were searched against five *Haliotis* transcriptomes. An average of 36% of tags (range 25 – 57%) were confidently aligned to *Haliotis* transcripts, with an average of 16% (range 5 – 27%) having annotations for genes associated with molecular function, biological processes and cellular components. All functional annotations are provided in Table S3.

Discussion

Evidence of adaptive genetic structuring despite gene flow

Using a sampling regime informed by high-resolution bathymetric and hindcasted oceanographic datasets, we contrasted patterns of genetic variation in *H. rubra* across heterogeneous habitats and

499 ~1,400 km of Australia's south eastern coastline. Estimates of overall genetic structure are
500 consistent with previous studies (Brown 1991; Conod *et al.* 2002; Li *et al.* 2006; Miller *et al.* 2016),
501 indicating a lack of genetic differentiation and potential panmixia across almost 1,000 km's of the
502 sampling range, extending from the South Australian border to Wilsons Promontory located on the
503 east-central coast of Victoria. Populations from the eastern extreme of the sampling range (east of
504 Wilson Promontory) and Port Phillip Bay appear weakly differentiated, as demonstrated by
505 previous studies (Brown 1991; Conod *et al.* 2002; Miller *et al.* 2016). Biogeographic structuring of
506 marine communities and limitations to gene flow among populations separated by Wilson
507 Promontory is well-recognised (York *et al.* 2008; Ayre *et al.* 2009; Colton & Swearer 2012; Miller
508 *et al.* 2013), and has been attributed to historical (Lambeck & Chappell 2001) and contemporary
509 physical oceanographic factors, such as converging ocean currents, environmental gradients
510 (temperature and salinity) and habitat discontinuities that persist in the region (Baines *et al.* 1983;
511 Ridgway & Godfrey 1997; Ridgway & Condie 2004; Sandery & Kaempf 2007; Colton & Swearer
512 2012). Weak differentiation of Port Phillip Bay is also consistent with the oceanography of the
513 region, with the low flushing rates of Port Phillip Bay (~270 days; (Walker 1999) expected to limit
514 gene flow between local populations and those from outside coastal waters. Despite a lack of
515 genetic structure across much of the Victorian coastline, we provide evidence for genotype
516 associations with heterogeneous habitat features occurring at local and regional spatial scales which
517 are likely driving patterns of adaptive genetic structuring within *H. rubra*.

518 Drift processes leading to neutral genetic structure are often suppressed in broadcast
519 spawning marine organisms with large populations sizes (Waples 1998; Whitlock & McCauley
520 1999; Hauser & Carvalho 2008), but numerous studies have shown that adaptive genetic
521 divergences can still be established and maintained under strong selection pressure (Marshall *et al.*
522 2010; Milano *et al.* 2014; Torda *et al.* 2017). This appears applicable to *Haliotis* species where
523 studies on Western Australian greenlip and Californian red abalone indicate selection contributing
524 to genetic adaptations associated with broad regional temperature, oxygen, and acidity gradients,
525 despite a lack of overall genetic structure (De Wit & Palumbi 2013; Sandoval-Castillo *et al.* 2015;
526 Sandoval-Castillo *et al.* 2018). We provide additional support, with evidence of adaptive divergence
527 associated with a temperature gradient in *H. rubra*. We are cautious with the interpretation of these
528 findings due to the high number of SNP loci found to be associated with sea surface temperature
529 (~10% of total SNPs). The possible influence of artefactual associations (i.e. false positives) and
530 SNP associations with other environmental factors varying with longitude cannot be ruled out. It is
531 also possible that adaptive loci are contained within inversions, which create genomic areas of
532 linkage disequilibrium (Hoffmann & Rieseberg, 2008) and have been implicated in many recent
533 examples of adaptive divergence (Twyford & Friedman 2015; Kirubakaran *et al.* 2016). Yet, the
534 previous reports of temperature as a driver of local adaptation in other Haliotid species suggests our

results may at least partly reflect temperature-related genetic structuring which warrants further investigation.

Our data also point to the possibility of selection associated with heterogeneous habitat features occurring on more local spatial scales. Specifically, our genotype-environmental association analyses suggest genotype associations with habitat features such as local reef geology, reef complexity, and exposure to wave power and ocean currents that vary over scales of 10s to 100s of meters (Ierodiaconou *et al.* 2011; Young *et al.* 2016). Evidence of fine scale adaptive genetic structuring is unusual in dispersive marine species, with a few exceptions (Sherman *et al.* 2008; Babin *et al.* 2017). While we cannot completely rule out the influence of false positives, each genotype–environment association (GEA) detected was corrected for spatial structure and distance, and our sampling regime contrasted genetic variation across distinct habitat types present both within and between western, central, and eastern regions of the state. Also, the largely panmictic nature of the *H. rubra* population spanning our sampling area suggests that artefactual demographic factors capable of producing ‘selection-like’ signatures across the genome (i.e. such as allelic surfing (Excoffier & Ray 2008)), are unlikely to influence our results.

At this stage it is uncertain if habitat features are directly or indirectly influencing genotype distributions. For example, substrate type (i.e. basalt, calcareous, granite) can influence benthic community composition, biological interactions and resource availability (Edmunds *et al.* 2000). Likewise, VRM is a physical measure of seabed complexity which can also influence benthic community compositions and biological interactions, as well as wave energy exposure, local oxygen saturation levels, and nutrient and resource availability (Sappington *et al.* 2007; Young *et al.* 2010; Martin-Garcia *et al.* 2013). This is also true for subtidal reefs that vary with respect to exposure to wave power and ocean currents (Young *et al.* 2015). As a result, the exact drivers of selection and potential adaptive divergence in *H. rubra* are unclear and warrant further investigation. Interestingly, our results suggest that variation in summer wave power and orbital velocity may lead to stronger patterns of local adaptation compared with winter months. This may reflect the *H. rubra* life cycle, where spawning typically occurs during the warmer months of the year (between February and April and again between October and December (Mcshane & Smith 1991)) when selection is likely to be strongest, acting on early life stages (Fraser *et al.* 2008; Postma & Ågren 2016). Experimental manipulations (i.e. via thermal stress, transplant, and common garden experiments) coupled with genome resequencing will assist in identifying environmental factors directly influencing selection and the genes and genomic features underpinning local adaptations (Jones *et al.* 2012; Berg *et al.* 2017; Price *et al.* 2018).

Insights into environmental resilience and recruitment processes

570 Results from this study are relevant to the potential resilience of *H. rubra* populations to
571 continued exploitation and changing environmental conditions. Marine ecosystems around the
572 world are experiencing rapid environmental transformations with climate change contributing to
573 rising ocean temperatures, the build-up of anthropogenic CO₂, changing ocean currents, and major
574 shifts in seasonal weather patterns (Hoegh-Guldberg & Bruno 2010; Burrows *et al.* 2011; Wernberg
575 *et al.* 2011). This is particularly pertinent in south-eastern Australia where ocean temperatures are
576 warming at four times the global average and ocean currents are strengthening and extending
577 southward (Ridgway 2007). Our results suggest potential genetic adaptations to temperature, wave
578 energy and current intensity, as well as connectivity between locations that show geographical
579 variation in them. This suggests standing genetic variation may be available for selection to act on
580 to counter future environmental change, assisted by wide spread gene flow and a short generation
581 time in this species (~4 years; (Andrews 1999)). This is a timely finding given abalone populations
582 around the world are showing signs of climate change stress, with depletion events in recent years
583 associated with exposure to extreme temperatures (Pearce *et al.* 2011; Micheli *et al.* 2012; Boch *et*
584 *al.* 2018). Quantitative tests via thermal stress, transplant, and common garden experiments, will
585 help to validate our findings, and provide information on the potential role of plasticity and genetic
586 adaptation in assisting *Haliotis* species counter climate change.

587 Abiotic factors such as local current intensities and wave energy are expected to determine
588 distances and directionalities of larvae movement (Trembl *et al.* 2015). However, selection is
589 potentially playing a role in determining habitats to which larvae can recruit depending on their
590 adaptive or maladaptive genetic background. For example, larvae released into the water column
591 from low energy environments are expected to be maladapted to high energy environments, and
592 may not successfully recruit to high energy habitat patches despite dispersal potential. This is not a
593 novel finding; evidence of phenotype-environment mismatches impacting recruitment dynamics
594 and reducing population connectivity in dispersive marine species is well documented and
595 summarised in Marshall *et al.* (2010). Natural selection could potentially occur at pre- or post-
596 settlement stages in *H. rubra*, as seen in other marine and diadromous taxa (Sherman *et al.* 2008;
597 Babin *et al.* 2017). In some marine species, environmental cues influence the settlement of larvae
598 from the water column (Rodriguez *et al.* 1993; Harrington *et al.* 2004; Sneed *et al.* 2014), with
599 encrusting coralline algae, macroalgae, and mucus trails reported to be potential settlement cues for
600 Haliotids (Morse *et al.* 1984; Takami *et al.* 1997; Huggett *et al.* 2005). Consequently, it is possible
601 that abalone larvae with different genotypes respond to specific cues, maximising the chance of
602 successful recruitment by settling upon suitable habitats, as shown in some corals (Meyer *et al.*
603 2009). Conversely, selection may act post-settlement, where maladapted genotypes are selected
604 against. For example, genetic inviability driven by genotype-environment interactions may be
605 responsible for high juvenile mortality in oysters (Plough 2012; Plough *et al.* 2016). As post-

606 settlement juvenile mortality is extremely high in *Haliotis* species (Mcshane & Smith 1991;
607 Kiyomoto 2007), post-settlement selection warrants further investigation. Overall, these findings
608 suggest that recruitment processes in Haliotids are likely to be complicated, with a combination of
609 exogenous and endogenous factors, including selection processes, having an influence.

610

611 *Further implications for fisheries management*

612 Translocation and reseedling activities are common management tools used for establishing new
613 fishing stocks and to catalyse the recovery of depleted stocks, including abalone fisheries (Bell *et al.*
614 *et al.* 2005; Roberts *et al.* 2007; Taniguchi *et al.* 2013). In recent years translocations have been
615 undertaken across *H. rubra* fisheries to assist the recovery of virus and urchin affected stocks in
616 south-eastern Australia (Department of Primary Industries 2012). Our results suggest that managers
617 should pay careful attention to the selection of genetically suitable animals for translocation
618 purposes, either through prior genotyping or habitat matching, to maximise return on investment.
619 The environmental thresholds driving allelic turn-over in our GDM spline plots could be used for
620 such purposes. Failure to consider the genetic profiles of fishing stocks could lead to the movement
621 of animals that are maladapted to their new habitat and lead to stocking failure. Indeed, the failure
622 of many restoration programs within the native ranges of Pacific salmon has been suggested to
623 partly result from inadequate adaptive ‘matching’ of translocated populations to their new
624 environments (Allendorf & Waples 1996).

625

626

627

628

629 **Conclusion**

630 Population genomic studies offer the opportunity to characterise selection processes and
631 environmental factors that contribute to genetic adaptation at species and population levels. Such
632 studies can assist management by providing insights into the likely resilience of species to
633 environmental change, and the influences of migration and selection on recruitment processes. The
634 current study demonstrates how emerging geospatial datasets offer new opportunities for
635 quantifying patterns of habitat heterogeneity in marine systems, and contrasting genetic variation
636 across environmental gradients to identify signatures and environmental drivers of selection with
637 unprecedented sensitivity. This research is becoming increasingly important as marine ecosystems
638 around the world are facing increasing threats from climate change, invasive pests, disease and
639 resource exploitation. Effective marine conservation planning and fisheries management, based on
640 population genomic studies with geospatially informed sampling regimes, will assist in minimizing
641 biodiversity loss and sustainable exploitation of commercially targeted species into the future.

642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676

Acknowledgements

This project was funded by the Fisheries Research and Development Corporation (FRDC) project ‘2015-025: Patterns of interaction between habitat and oceanographic variables affecting the connectivity and productivity of invertebrate fisheries’ on behalf of the Australian Government, plus contributions from the Western Abalone Divers Association. We thank Nick Murphy (La Trobe University), Steve Swearer (The University of Melbourne), and Jan Strugnell (James Cook University) for their contributions to the original project proposal. We would also like to thank Harry Gorfine from the Victorian Fisheries Authority, and government contract divers David Forbes and Johnno Rudge and their respective crews, for their assistance with the genetic sampling. Additional thanks to Chris Austin, Han Ming Gan and Yin Peng Lee (Deakin University), for assistance with generation of genomics libraries and sequencing.

This project was conceived by A.D.M and C.D.H, and experimental design was assisted by AAH. A.D.M, C.D.H., M.H.T., and M.C., were responsible for generating the data, and the analysis was led by A.D.M with assistance from M.H.T., C.A., C.D.H., and A.A.H. Geospatial datasets were provided by M.Y., A.R., D.A.I., and E.T. Writing of the manuscript was led by A.D.M. with assistance from all authors.

Data accessibility

Genomic and environmental data sets, and scripts for GEA analyses are available in the DRYAD archives under accession doi:10.5061/dryad.kb401ch

Table 1 Sites included in genomic analysis

Table 2 Summary of environmental statistics for all 29 collection sites: geology, Vector Ruggedness Measure (VRM), average winter and summer wave power (kW/m); average winter and summer orbital velocity (m/s); average summer and seasonal range in sea surface temperature (°C)

Table 3 Numbers of candidate loci identified as having significant correlations with environmental covariables by BayPass and Latent Factor Mixed Models (LFMM) from a total dataset of 5,239 SNPs.

677 **Figure 1** Sampling locations for population genomic analysis from south-eastern Australia,
678 including the western (inset figure ‘a’), central (inset figures ‘b, c, d and e’), and eastern (inset
679 figure ‘f’) zone fisheries. Refer to Table 1 for sample codes. Wilsons Promontory visible within
680 inset ‘e’.

681
682 **Figure 2** Identification of SNPs under putative selection using redundancy analysis. All points are
683 SNPs, off-white points are SNPs not associated with the 8 environmental variables. Colours signify
684 significant correlation with environmental variables. The “GRA” inertia is not shown in the RDA
685 plot because it was used as the ‘dummy’ variable for the remaining two geological states (“BAS”
686 and “CAL”). SWP = summer wave power; VRM = vector ruggedness measure; WWP = winter
687 wave power; WOV = winter orbital velocity; SOV = summer orbital velocity; SST = summer
688 seasonal temperature; STR = seasonal range sea surface temperature; BAS = geology (basalt); CAL
689 = geology (calcareous); GRA = geology (granite).

690
691 **Figure 3** Discriminant analysis of principal components plots based on candidate SNPs putatively
692 under selection and associated with habitat geology (substrate and VRM). Inset eigenvalues indicate
693 the number of two principal components that captured the observed genetic structure.

694
695 **Figure 4** Discriminant analysis of principal components plots based on candidate SNPs associated
696 with winter and summer wave power. Inset eigenvalues indicate the number of two principal
697 components that captured the observed genetic structure.

698
699 **Figure 5** Discriminant analysis of principal components plots based on candidate SNPs associated
700 with winter and summer orbital velocity. Inset eigenvalues indicate the number of two principal
701 components that captured the observed genetic structure.

702
703 **Figure 6** Discriminant analysis of principal components plots based on candidate SNPs associated
704 with summer sea surface temperature and seasonal temperature range. Inset eigenvalues indicate the
705 number of two principal components that captured the observed genetic structure.

706
707 **Figure 7** Generalized Dissimilarity Modeling spline plots quantifying allelic turnover through
708 climatic space (deviations in allele frequency associated with environment type) for candidate SNPs

709 putatively associated with environmental variables a) winter wave power, b) summer wave power,
710 c) summer seasonal temperature, d) seasonal range sea surface temperature, e) winter orbital
711 velocity, f) summer orbital velocity, g) vector ruggedness measure. Individual lines represent allelic
712 turn over for single loci.

713

714 **Figure 8** Predicted spatial turnover in allele frequencies of *H. rubra* based on generalised
715 dissimilarity modeling analyses for all SNPs associated with a) summer sea surface temperature,
716 and b) vector ruggedness measure. Regions with similar colours are expected to harbor populations
717 with similar genomic compositions. Sediment habitats which are uninhabited by abalone and shown
718 in grey.

719

720

721 **References**

- 722 Ahrens CW, Rymer PD, Stow A, Bragg J, Dillon S *et al.* (2018) The search for loci under selection:
723 trends, biases and progress. *Molecular Ecology* **27**: 1342-1356.
- 724 Allendorf FW, and Waples RS (1996) Conservation and genetics of salmonid fishes, pp. 238-280 in
725 *Conservation Genetics: Case Histories from Nature*, edited by Avise JC and Hamrick JL.
726 Chapman and Hall, New York.
- 727 Andrews NL (1999) *Under Southern Seas: The Ecology of Australia's Rocky Reefs*. University of
728 New South Wales Press, Sydney.
- 729 Ayre DJ, Minchinton TE and Perrin C (2009) Does life history predict past and current connectivity
730 for rocky intertidal invertebrates across a marine biogeographic barrier? *Molecular Ecology*
731 **18**: 1887-1903.
- 732 Babin C, Gagnaire PA, Pavey SA, Bernatchez L (2017) RAD-Seq reveals patterns of additive
733 polygenic variation caused by spatially-varying selection in the American Eel (*Anguilla*
734 *rostrata*). *Genome Biology and Evolution* **9**: 2974–2986.
- 735 Baines PG, Edwards RJ and Fandry CB (1983) Observations of a new baroclinic current along the
736 western continental slope of Bass Strait. *Australian Journal of Marine and Freshwater*
737 *Research* **34**: 155-157.
- 738 Bateman A, Martin MJ, O'Donovan C, Magrane M, Alpi E *et al.* (2017) UniProt: the universal
739 protein knowledgebase. *Nucleic Acids Research* **45**: D158-D169.
- 740 Bell JD, Rothlisberg PC, Munro JL, Loneragan NR, Nash WJ *et al.* (2005) Restocking and Stock
741 Enhancement of Marine Invertebrate Fisheries. *Restocking and Stock Enhancement of*
742 *Marine Invertebrate Fisheries* **49**: 1-374.

743 Berg PR, Star B, Pampoulie C, Bradbury IR, Bentzen P, Hutchings JA, Trans-oceanic genomic
744 divergence of Atlantic cod ecotypes is associated with large inversions

745 Boch CA, Micheli F, AlNajjar M, Monismith SG, Beers JM *et al.* (2018) Local oceanographic
746 variability influences the performance of juvenile abalone under climate change. *Scientific*
747 *Reports* **8**: 5501.

748 Boehm JT, Waldman J, Robinson JD and Hickerson MJ (2015) Population genomics reveals
749 seahorses (*Hippocampus erectus*) of the western mid-Atlantic coast to be residents rather
750 than vagrants. *Plos One* **10**: e0116219.

751 Bolger AM, Lohse M and Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence
752 data. *Bioinformatics* **30**: 2114-2120.

753 Bonhomme M, Chevalet C, Servin B, Boitard S, Abdallah J *et al.* (2010) Detecting Selection in
754 Population Trees: The Lewontin and Krakauer Test Extended. *Genetics* **186**: 241-U406.

755 Bourret V, Dionne M, & Bernatchez L (2014). Detecting genotypic changes associated with
756 selective mortality at sea in Atlantic salmon: polygenic multilocus analysis surpasses
757 genome scan. *Molecular Ecology* **23**: 4444-4457.

758 Brock JC, and Purkis SJ (2009) The Emerging Role of Lidar Remote Sensing in Coastal Research
759 and Resource Management. *Journal of Coastal Research* **25**: 1-5.

760 Brown CJ, Smith SJ, Lawton P and Anderson JT (2011) Benthic habitat mapping: A review of
761 progress towards improved understanding of the spatial ecology of the seafloor using
762 acoustic techniques. *Estuarine Coastal and Shelf Science* **92**: 502-520.

763 Brown LD (1991) Genetic variation and population structure in the blacklip abalone, *Haliotis*
764 *rubra*. *Australian Journal of Marine and Freshwater Research* **42**: 77-90.

765 Buehler D, Poncet BN, Holderegger R, Manel S, Taberlet P *et al.* (2013) An outlier locus relevant
766 in habitat-mediated selection in an alpine plant across independent regional replicates.
767 *Evolutionary Ecology* **27**: 285-300.

768 Burrows MT, Schoeman DS, Buckley LB, Moore P, Poloczanska ES *et al.* (2011) The Pace of
769 Shifting Climate in Marine and Terrestrial Ecosystems. *Science* **334**: 652-655.

770 Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J *et al.* (2009) BLAST plus :
771 architecture and applications. *Bmc Bioinformatics* **10**: 421.

772 Catchen J, Hohenlohe PA, Bassham S, Amores A and Cresko WA (2013) Stacks: an analysis tool
773 set for population genomics. *Molecular Ecology* **22**: 3124-3140.

774 Chambers MD, VanBlaricom GR, Hauser L, Utter F and Friedman CS (2006) Genetic structure of
775 black abalone (*Haliotis cracherodii*) populations in the California islands and central
776 California coast: Impacts of larval dispersal and decimation from withering syndrome.
777 *Journal of Experimental Marine Biology and Ecology* **331**: 173-185.

778 Chen H, Boutros PC (2011) VennDiagram: a package for the generation of highly-customizable
779 Venn and Euler diagrams in R. *BMC Bioinformatics* **12**:35

780 Cogan CB, Todd BJ, Lawton P and Noji TT (2009) The role of marine habitat mapping in
781 ecosystem-based management. *Ices Journal of Marine Science* **66**: 2033-2042.

782 Colton MA, and Swearer SE (2012) Locating faunal breaks in the nearshore fish assemblage of
783 Victoria, Australia. *Marine and Freshwater Research* **63**: 218-231.

784 Conod N, Bartlett JP, Evans BS and Elliott NG (2002) Comparison of mitochondrial and nuclear
785 DNA analyses of population structure in the blacklip abalone *Haliotis rubra* Leach. *Marine
786 and Freshwater Research* **53**: 711-718.

787 Davey JW, Hohenlohe PA, Etter PD, Boone JQ, Catchen JM *et al.* (2011) Genome-wide genetic
788 marker discovery and genotyping using next-generation sequencing. *Nature Reviews
789 Genetics* **12**: 499-510.

790 Department of Primary Industries (2012) A review of rebuilding options for the Victorian abalone
791 fishery. Fisheries Victoria Management Report, State Government of Victoria.

792 de Villemereuil P, Frichot E, Bazin E, Francois O and Gaggiotti OE (2014) Genome scan methods
793 against more complex models: when and how much should we trust them? *Molecular
794 Ecology* **23**: 2006-2019.

795 De Wit P, and Palumbi SR (2013) Transcriptome-wide polymorphisms of red abalone (*Haliotis
796 rufescens*) reveal patterns of gene flow and local adaptation. *Molecular Ecology* **22**: 2884-
797 2897.

798 Dierickx EG, Shultz AJ, Sato F, Hiraoka T and Edwards SV (2015) Morphological and genomic
799 comparisons of Hawaiian and Japanese Black-footed Albatrosses (*Phoebastria nigripes*)
800 using double digest RADseq: implications for conservation. *Evolutionary Applications* **8**:
801 662-678.

802 Edmunds M, Chidgey S and Willcox S (2000) Environmental Inventory of Victoria's Marine
803 Ecosystems Stage 3, edited by Ferns LW and Hough D. Parks, Flora and Fauna Division,
804 Department of Natural Resources and Environment, East Melbourne, Australia.

805 Ekblom R, and Galindo J (2011) Applications of next generation sequencing in molecular ecology
806 of non-model organisms. *Heredity* **107**: 1-15.

807 Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K *et al.* (2011) A Robust, Simple
808 Genotyping-by-Sequencing (GBS) Approach for High Diversity Species. *Plos One* **6**.

809 Environmental Systems Research Institute (ESRI) 2018. ArcGIS Release 10.5. Redlands, CA.

810 Excoffier L, and Ray N (2008) Surfing during population expansions promotes genetic revolutions
811 and structuration. *Trends in Ecology & Evolution* **23**: 347-351.

- 812 Ferrier S, Manion G, Elith J and Richardson K (2007) Using generalized dissimilarity modelling to
813 analyse and predict patterns of beta diversity in regional biodiversity assessment. *Diversity*
814 *and Distributions* **13**: 252-264.
- 815 Fitzpatrick MC, and Keller SR (2015) Ecological genomics meets community-level modelling of
816 biodiversity: mapping the genomic landscape of current and future environmental
817 adaptation. *Ecology Letters* **18**: 1-16.
- 818 Forester, BR, Lasky JR, Wagner HH, and Urban DL (2018). Comparing methods for detecting
819 multilocus adaptation with multivariate genotype–environment associations. *Molecular*
820 *Ecology* **27**: 2215-2233.
- 821 Fraser DJ, Cook AM, Eddington JD, Bentzen P and Hutchings JA (2008) Mixed evidence for
822 reduced local adaptation in wild salmon resulting from interbreeding with escaped farmed
823 salmon: complexities in hybrid fitness. *Evolutionary Applications* **1**: 501-512.
- 824 Frichot E, and Francois O (2015) LEA: An R package for landscape and ecological association
825 studies. *Methods in Ecology and Evolution* **6**: 925-929.
- 826 Frichot E, Schoville SD, Bouchard G and Francois O (2013) Testing for Associations between Loci
827 and Environmental Gradients Using Latent Factor Mixed Models. *Molecular Biology and*
828 *Evolution* **30**: 1687-1699.
- 829 Galindo J, Martinez-Fernandez M, Suarez P, Moran P, Juan FS *et al.* (2009) The adaptive role of
830 Phosphoglucosyltransferase and other allozymes in a marine snail across the vertical rocky-shore
831 gradient. *Biological Journal of the Linnean Society* **98**: 225-233.
- 832 Gautier M (2015) Genome-wide scan for adaptive divergence and association with population-
833 specific covariates. *Genetics* **201**: 1555-1579.
- 834 Gordon HR, and Cook PA (2013) World abalone supply, markets, and pricing: 2011 update.
835 *Journal of Shellfish Research* **32**: 5-7.
- 836 Gruenthal KM, Acheson LK and Burton RS (2007) Genetic structure of natural populations of
837 California red abalone (*Haliotis rufescens*) using multiple genetic markers. *Marine Biology*
838 **152**: 1237-1248.
- 839 Gruenthal KM, and Burton RS (2005) Genetic diversity and species identification in the endangered
840 white abalone (*Haliotis sorenseni*). *Conservation Genetics* **6**: 929-939.
- 841 Gruenthal KM, and Burton RS (2008) Genetic structure of natural populations of the California
842 black abalone (*Haliotis cracherodii* Leach, 1814), a candidate for endangered species status.
843 *Journal of Experimental Marine Biology and Ecology* **355**: 47-58.
- 844 Grummer JA, Beheregary LB, Bernatchez L, Hand BK, Luikart G, Narum S, Taylor EB (2019)
845 Aquatic landscape genomics and environmental effects on genetic variation. *Trends in*
846 *Ecology and Evolution*. DOI.org/10.1016/j.tree.2019.02.013.

847 Gunther T, and Coop G (2013) Robust identification of local adaptation from allele frequencies.
848 *Genetics* **195**: 205-220.

849 Hahn KO (1988) Larval development of abalone, pp. 71-99 in *The Culture of Abalone and Other*
850 *Marine Gastropods*, edited by Hahn KO. CRC Press, Boca Raton.

851 Harney E, Dubief B, Boudry P, Basuyaux O, Schilhabel MB *et al.* (2016) De novo assembly
852 and annotation of the European abalone *Haliotis tuberculata* transcriptome. *Marine*
853 *Genomics* **28**: 11-16.

854 Harrington L, Fabricius K, De'ath G and Negri A (2004) Recognition and selection of settlement
855 substrata determine post-settlement survival in corals. *Ecology* **85**: 3428-3437.

856 Hauser L, and Carvalho GR (2008) Paradigm shifts in marine fisheries genetics: ugly hypotheses
857 slain by beautiful facts. *Fish and Fisheries* **9**: 333-362.

858 Hemer MA, Fan YL, Mori N, Semedo A and Wang XLL (2013) Projected changes in wave climate
859 from a multi-model ensemble. *Nature Climate Change* **3**: 471-476.

860 Hess JE, Campbell NR, Close DA, Docker MF and Narum SR (2013) Population genomics of
861 Pacific lamprey: adaptive variation in a highly dispersive species. *Molecular Ecology* **22**:
862 2898-2916.

863 Hoegh-Guldberg O, and Bruno JF (2010) The Impact of Climate Change on the World's Marine
864 Ecosystems. *Science* **328**: 1523-1528.

865 Hoffmann AA, and Sgro CM (2011) Climate change and evolutionary adaptation. *Nature* **470**: 479-
866 485.

867 Hoffmann AA, and Rieseberg LH (2008) Revisiting the impact of inversions in evolution: from
868 population genetic markers to drivers of adaptive shifts and speciation? *Annual Review of*
869 *Ecology, Evolution, and Systematics* **39**: 21-42.

870 Huggett MJ, de Nys R, Williamson JE, Heasman M and Steinberg PD (2005) Settlement of larval
871 blacklip abalone, *Haliotis rubra*, in response to green and red macroalgae. *Marine Biology*
872 **147**: 1155-1163.

873 Ierodiconou D, Monk J, Rattray A, Laurenson L and Versace VL (2011) Comparison of automated
874 classification techniques for predicting benthic biological communities using hydroacoustics
875 and video observations. *Continental Shelf Research* **31**: S28-S38.

876 Ierodiconou DA, Young M, Miller AD, Treml E, Swearer S *et al.* (2018) Patterns of interaction
877 between habitat and oceanographic variables affecting the connectivity and productivity of
878 invertebrate fisheries. Fisheries Research and Development Corporation (Final report, 2015-
879 025).

880 Jalali MA, Ierodiconou D, Gorfine H, Monk J and Rattray A (2015) Exploring spatiotemporal
881 trends in commercial fishing effort of an abalone fishing zone: A GIS-based hotspot model.
882 *Plos One* **10**: e0122995.

883   Jeffreys H (1961) Theory of Probability (3rd edition). Oxford: Oxford University Press.

884   Jombart T (2008) adegenet: a R package for the multivariate analysis of genetic markers.

885   *Bioinformatics* **24**: 1403-1405.

886   Jombart T, and Ahmed I (2011) adegenet 1.3-1: new tools for the analysis of genome-wide SNP

887   data. *Bioinformatics* **27**: 3070-3071.

888   Jones FC, Grabherr MG, Chan YF, Russell P, Mauceli E *et al.* (2012) The genomic basis of

889   adaptive evolution in threespine sticklebacks. *Nature* **484**: 55-61.

890   Jones P, Binns D, Chang HY, Fraser M, Li WZ *et al.* (2014) InterProScan 5: genome-scale

891   protein function classification. *Bioinformatics* **30**: 1236-1240.

892   Jordan R, Hoffmann AA, Dillon SK and Prober SM (2017) Evidence of genomic adaptation to

893   climate in *Eucalyptus microcarpa*: Implications for adaptive potential to projected climate

894   change. *Molecular Ecology* **26**: 6002-6020.

895   Jost I (2008) Gst and its relatives do not measure differentiation. *Molecular Ecology* **17**: 4015-4026.

896   Kashiwada JV, and Taniguchi IK (2007) Application of recent red abalone *Haliotis rufescens*

897   surveys to management decisions outlined in the California Abalone Recovery and

898   Management Plan. *Journal of Shellfish Research* **26**: 713-717.

899   Kirubakaran TG, Grove H, Kent MP, *et al.* (2016) Two adjacent inversions maintain genomic

900   differentiation between migratory and stationary ecotypes of Atlantic cod. *Molecular*

901   *Ecology* **25**: 2130-2143.

902   Kiyomoto S (2007) The presence of scavengers increases the vulnerability of juvenile abalone to

903   predatory fish. *Fisheries Science* **73**: 732-734.

904   Kough AS, Claro R, Lindeman KC and Paris CB (2016) Decadal analysis of larval connectivity

905   from Cuban snapper (Lutjanidae) spawning aggregations based on biophysical modeling.

906   *Marine Ecology Progress Series* **550**: 175-190.

907   Lambeck K, and Chappell J (2001) Sea level change through the last glacial cycle. *Science* **292**:

908   679-686.

909   Lecours V, Lucieer VL, Donlan MFJ and Micallef A (2015) An ocean of possibilities: applications

910   and challenges of marine geomorphometry in *Geomorphometry for geosciences*, edited by

911   Zwolinski Z, Mitasova H and Hengl T. International Society for Geomorphometry, Poznan,

912   Poland.

913   Leiva GE, and Castilla JC (2001) A review of the world marine gastropod fishery: evolution of

914   catches, management and the Chilean experience. *Reviews in Fish Biology and Fisheries* **11**:

915   283-300.

916   Li ZB, Appleyard SA and Elliott NG (2006) Population structure of *Haliotis rubra* from South

917   Australia inferred from nuclear and mtDNA analyses. *Acta Oceanologica Sinica* **25**: 99-112.

- 918 Manion G, Lisk M, Ferrier S, Nieto-Lugilde D and Fitzpatrick MC (2018) GDM: Functions for
919 Generalized Dissimilarity Modeling; R Package Version 1.2.3.
- 920 Marshall DJ, Monro K, Bode M, Keough MJ and Swearer S (2010) Phenotype-environment
921 mismatches reduce connectivity in the sea. *Ecology Letters* **13**: 128-140.
- 922 Martin-Garcia L, Gonzalez-Lorenzo G, Brito-Izquierdo IT and Barquin-Diez J (2013) Use of
923 topographic predictors for macrobenthic community mapping in the Marine Reserve of La
924 Palma (Canary Islands, Spain). *Ecological Modelling* **263**: 19-31.
- 925 Martinez-Fernandez M, Bernatchez L, Rolan-Alvarez E and Quesada H (2010) Insights into the role
926 of differential gene expression on the ecological adaptation of the snail *Littorina saxatilis*.
927 *BMC Evolutionary Biology* **10**: 356.
- 928 Mayer L, Jakobsson M, Allen G, Dorschel B, Falconer R *et al.* (2018) The Nippon Foundation-
929 GEBCO Seabed 2030 Project: The Quest to See the World's Oceans Completely Mapped by
930 2030. *Geosciences* **8**: 63.
- 931 McShane PE, Black KP and Smith MG (1988) Recruitment processes in *Haliotis rubra* Leach
932 (Mollusca: Gastropoda) and regional hydrodynamics in southeastern Australia imply
933 localized dispersal of larvae. *Journal of Experimental Marine Biology and Ecology* **124**:
934 175-203.
- 935 McShane PE, and Smith MG (1991) Recruitment variation in sympatric populations of *Haliotis*
936 *rubra* (Mollusca, Gastropoda) in southeast Australian waters. *Marine Ecology Progress*
937 *Series* **73**: 203-210.
- 938 Meirmans PG, and Van Tienderen PH (2004) GENOTYPE and GENODIVE: two programs for the
939 analysis of genetic diversity of asexual organisms. *Molecular Ecology Notes* **4**: 792-794.
- 940 Meyer E, Davies S, Wang S, Willis BL, Abrego D *et al.* (2009) Genetic variation in responses to a
941 settlement cue and elevated temperature in the reef-building coral *Acropora millepora*.
942 *Marine Ecology Progress Series* **392**: 81-92.
- 943 Micheli F, Saenz-Arroyo A, Greenley A, Vazquez L, Montes JAE *et al.* (2012) Evidence That
944 Marine Reserves Enhance Resilience to Climatic Impacts. *Plos One* **7**: e40832.
- 945 Milano I, Babbucci M, Cariani A, Atanassova M, Bekkevold D *et al.* (2014) Outlier SNP markers
946 reveal fine-scale genetic structuring across European hake populations (*Merluccius*
947 *merluccius*). *Molecular Ecology* **23**: 118-135.
- 948 Miller AD, van Rooyen A, Rasic G, Ierodiaconou DA, Gorfine HK *et al.* (2016) Contrasting
949 patterns of population connectivity between regions in a commercially important mollusc
950 *Haliotis rubra*: integrating population genetics, genomics and marine LiDAR data.
951 *Molecular Ecology* **25**: 3845-3864.
- 952 Miller AD, Versace VL, Matthews TG, Montgomery S and Bowie KC (2013) Ocean currents
953 influence the genetic structure of an intertidal mollusc in south-eastern Australia -

954 implications for predicting the movement of passive dispersers across a marine
 955 biogeographic barrier. *Ecology and Evolution* **3**: 1248-1261.

956 Miller DD, Ota Y, Sumaila UR, Cisneros-Montemayor AM and Cheung WWL (2018) Adaptation
 957 strategies to climate change in marine systems. *Global Change Biology* **24**: E1-E14.

958 Miller KJ, Maynard BT and Mundy CN (2009) Genetic diversity and gene flow in collapsed and
 959 healthy abalone fisheries. *Molecular Ecology* **18**: 200-211.

960 Miller KJ, Mundy CN and Mayfield S (2014) Molecular genetics to inform spatial management in
 961 benthic invertebrate fisheries: a case study using the Australian Greenlip Abalone.
 962 *Molecular Ecology* **23**: 4958-4975.

963 Morse ANC, Froyd CA and Morse DE (1984) Molecules from cyanobacteria and red algae that
 964 induce larval settlement and metamorphosis in the mollusk *Haliotis rufescens*. *Marine*
 965 *Biology* **81**: 293-298.

966 Nielsen EE, Hemmer-Hansen J, Poulsen NA, Loeschcke V, Moen T *et al.* (2009) Genomic
 967 signatures of local directional selection in a high gene flow marine organism; the Atlantic
 968 cod (*Gadus morhua*). *BMC Evolutionary Biology* **9**: 276.

969 Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, *et al.* (2013) Package ‘vegan’.
 970 Community ecology package, version, 2(9).

971 Peakall R, Smouse PE (2006) GENALEX 6: genetic analysis in Excel. Population genetics software
 972 for teaching and research. *Molecular Ecology Notes* **6**: 288–295.

973 Pearce A, Lenanton R, Jackson G, Moore J, Feng M *et al.* (2011) The “marine heat wave” off
 974 Western Australia during the summer of 2010/11, in *Fisheries Research Report No. 222*.
 975 Government of Western Australia, Department of Fisheries, Western Australia.

976 Petrik CM, Duffy-Anderson JT, Mueter F, Hedstrom K and Curchitser EN (2015) Biophysical
 977 transport model suggests climate variability determines distribution of Walleye Pollock
 978 early life stages in the eastern Bering Sea through effects on spawning. *Progress in*
 979 *Oceanography* **138**: 459-474.

980 Pfeifer SP, Laurent S, Sousa VC, Linnen CR, Foll M *et al.* (2018) The eolutionary history of
 981 Nebraska deer mice: local adaptation in the face of strong gene flow. *Molecular Biology and*
 982 *Evolution* **35**: 792-806.

983 Pickrill RA, and Todd BJ (2003) The multiple roles of acoustic mapping in integrated ocean
 984 management, Canadian Atlantic continental margin. *Ocean & Coastal Management* **46**:
 985 601-614.

986 Piggott MP, Banks SC, Tung P and Beheregaray LB (2008) Genetic evidence for different scales of
 987 connectivity in a marine mollusc. *Marine Ecology Progress Series* **365**: 127-136.

988 Plough LV (2012) Environmental stress increases selection against and dominance of deleterious
989 mutations in inbred families of the Pacific oyster *Crassostrea gigas*. *Molecular Ecology* **21**:
990 3974-3987.

991 Plough LV, Shin G and Hedgecock D (2016) Genetic inviability is a major driver of type III
992 survivorship in experimental families of a highly fecund marine bivalve. *Molecular Ecology*
993 **25**: 895-910.

994 Postma FM, and Ågren J (2016) Early life stages contribute strongly to local adaptation in
995 *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences* **113**: 7590-7595.

996 Price N, Moyers BT, Lopez L, Lasky JR, Monroe JG *et al.* (2018) Combining population
997 genomics and fitness QTLs to identify the genetics of local adaptation in *Arabidopsis*
998 *thaliana*. *Proceedings of the National Academy of Sciences of the United States of America*
999 **115**: 5028-5033.

1000 Prince JD, Sellers TL, Ford WB and Talbot SR (1987) Experimental evidence for limited
1001 dispersal of haliotid larvae (genus *Haliotis*; Mollusca: Gastropoda). *Journal of Experimental*
1002 *Marine Biology and Ecology* **106**: 243-263.

1003 Prince JD, Sellers TL, Ford WB and Talbot SR (1988) Confirmation of a relationship between the
1004 localized abundance of breeding stock and recruitment for *Haliotis rubra* Leach (Mollusca:
1005 Gastropoda). *Journal of Experimental Marine Biology and Ecology* **122**: 91-104.

1006 Raymond M, and Rousset F (1995) An exact test for population differentiation. *Evolution* **49**: 1280-
1007 1283.

1008 Ridgway KR (2007) Long-term trend and decadal variability of the southward penetration of the
1009 East Australian Current. *Geophysical Research Letters* **34**: L13613.

1010 Ridgway KR, and Condie SA (2004) The 5500-km-long boundary flow off western and southern
1011 Australia. *Journal of Geophysical Research-Oceans* **109**: 1-18.

1012 Ridgway KR, and Godfrey JS (1997) Seasonal cycle of the East Australian Current. *Journal of*
1013 *Geophysical Research-Oceans* **102**: 22921-22936.

1014 Roberts RD, Keys EF, Prendeville G and Pilditch CA (2007) Viability of abalone (*Haliotis iris*)
1015 stock enhancement by release of hatchery-reared seed in Marlborough, New Zealand.
1016 *Journal of Shellfish Research* **26**: 697-703.

1017 Rodriguez SR, Ojeda FP and Inestrosa NC (1993) Settlement of benthic marine-invertebrates.
1018 *Marine Ecology Progress Series* **97**: 193-207.

1019 Roodt-Wilding R (2007) Abalone ranching: a review on genetic considerations. *Aquaculture*
1020 *Research* **38**: 1229-1241.

1021 Rousset F (1997) Genetic differentiation and estimation of gene flow from F-statistics under
1022 isolation by distance. *Genetics* **145**: 1219–1228.

- 1023 Sandery PA, and Kaempf J (2007) Transport timescales for identifying seasonal variation in Bass
1024 Strait, south-eastern Australia. *Estuarine Coastal and Shelf Science* **74**: 684-696.
- 1025 Sandoval-Castillo J, Robinson NA, Hart AM, Strain WS and Beheregaray LB (2018) Seascape
1026 genomics reveals adaptive divergence in a connected and commercially important mollusc,
1027 the greenlip abalone (*Haliotis laevis*), along a longitudinal environmental gradient.
1028 *Molecular Ecology* **27**: 1603-1620.
- 1029 Sandoval-Castillo J, Strain WS, Hart AM, Robinson NA and Beheregaray LB (2015) Use of next
1030 generation DNA technologies for revealing the genetic impact of fisheries restocking and
1031 ranching, Adelaide: The Seafood CRC Company Ltd, the Fisheries Research and
1032 Development Corporation, Flinders University.
- 1033 Sappington JM, Longshore KM and Thompson DB (2007) Quantifying landscape ruggedness for
1034 animal habitat analysis: A case study using bighorn sheep in the Mojave Desert. *Journal of*
1035 *Wildlife Management* **71**: 1419-1426.
- 1036 Savolainen O, Lascoux M and Merila J (2013) Ecological genomics of local adaptation. *Nature*
1037 *Reviews Genetics* **14**: 807-820.
- 1038 Sherman, CDH, Ayre, DJ (2008) Fine-scale adaptation in a clonal sea anemone. *Evolution* **62**:
1039 1373-1380.
- 1040 Shiel BP, Hall NE, Cooke IR, Robinson NA and Strugnell JM (2015) De Novo
1041 Characterisation of the Greenlip Abalone Transcriptome (*Haliotis laevis*) with a Focus
1042 on the Heat Shock Protein 70 (HSP70) Family. *Marine Biotechnology* **17**: 23-32.
- 1043 Sneed JM, Sharp KH, Ritchie KB and Paul VJ (2014) The chemical cue tetrabromopyrrole from a
1044 biofilm bacterium induces settlement of multiple Caribbean corals. *Proceedings of the Royal*
1045 *Society B-Biological Sciences* **281**: 1786.
- 1046 Solas MR, Sepulveda RD and Brante A (2013) Genetic variation of the shell morphology in
1047 *Acanthina monodon* (Gastropoda) in habitats with different wave exposure conditions.
1048 *Aquatic Biology* **18**: 253-260.
- 1049 Stapley J, Reger J, Feulner PGD, Smadja C, Galindo J *et al.* (2010) Adaptation genomics: the next
1050 generation. *Trends in Ecology & Evolution* **25**: 705-712.
- 1051 Takami H, Kawamura T and Yamashita Y (1997) Survival and growth rates of post-larval abalone
1052 *Haliotis discus hannai* fed conspecific trail mucus and/or benthic diatom *Cocconeis*
1053 *scutellum* var. *parva*. *Aquaculture* **152**: 129-138.
- 1054 Taniguchi IK, Stein D, Lampson K and Rogers-Bennett L (2013) Testing translocation as a
1055 recovery tool for pink (*Haliotis corrugata*) and green (*Haliotis fulgens*) abalone in southern
1056 California. *Journal of Shellfish Research* **32**: 209-216.

1057 Temby N, Miller K and Mundy C (2007) Evidence of genetic subdivision among populations of
1058 blacklip abalone (*Haliotis rubra* Leach) in Tasmania. *Marine and Freshwater Research* **58**:
1059 733-742.

1060 Termignoni-Garcia F, Jaramillo-Correa JP, Chable-Santos J, Liu M, Shultz AJ *et al.* (2017)
1061 Genomic footprints of adaptation in a cooperatively breeding tropical bird across a
1062 vegetation gradient. *Molecular Ecology* **26**: 4483-4496.

1063 Trembl EA, Ford JR, Black KP and Swearer SE (2015) Identifying the key biophysical
1064 drivers, connectivity outcomes, and metapopulation consequences of larval dispersal in the
1065 sea. *Movement Ecology* **3**: 17.

1066 Torda G, Donelson JM, Aranda M, Barshis DJ, Bay L *et al.* (2017) Rapid adaptive responses to
1067 climate change in corals. *Nature Climate Change* **7**: 627-636.

1068 Twyford AD, and Friedman J (2015) Adaptive divergence in the monkey flower *Mimulus guttatus*
1069 is maintained by a chromosomal inversion. *Evolution* **69**: 1476-1486.

1070 van der Merwe M, Franchini P and Roodt-Wilding R (2011) Differential growth-related
1071 gene expression in abalone (*Haliotis midae*). *Marine Biotechnology* **13**: 1125-1139.

1072 Walbridge S, Slocum N, Pobuda M, Wright DJ (2008) Unified geomorphological analysis workflows
1073 with benthic terrain modeler. *Geosciences* **8**: 94.

1074 Walker SJ (1999) Coupled hydrodynamic and transport models of Port Phillip Bay, a semi-enclosed
1075 bay in south-eastern Australia. *Marine and Freshwater Research* **50**: 469-481.

1076 Waples RS (1998) Separating the wheat from the chaff: Patterns of genetic differentiation in high
1077 gene flow species. *Journal of Heredity* **89**: 438-450.

1078 Waples RS, Punt AE and Cope JM (2008) Integrating genetic data into management of marine
1079 resources: how can we do it better? *Fish and Fisheries* **9**: 423-449.

1080 Ward RD (2006) The importance of identifying spatial population structure in restocking and stock
1081 enhancement programmes. *Fisheries Research* **80**: 9-18.

1082 Weeks AR, Heinze D, Perrin L, Stoklosa J, Hoffmann AA *et al.* (2017) Genetic rescue increases
1083 fitness and aids rapid recovery of an endangered marsupial population. *Nature*
1084 *Communications* **8**: 1071.

1085 Weeks AR, Sgro CM, Young AG, Frankham R, Mitchell NJ *et al.* (2011) Assessing the benefits
1086 and risks of translocations in changing environments: a genetic perspective. *Evolutionary*
1087 *Applications* **4**: 709-725.

1088 Weir B, and Cockerham C (1984) Estimating F-statistics for the analysis of population structure.
1089 *Evolution* **38**: 1358-1370.

1090 Wernberg T, Russell BD, Moore PJ, Ling SD, Smale DA *et al.* (2011) Impacts of climate change in
1091 a global hotspot for temperate marine biodiversity and ocean warming. *Journal of*
1092 *Experimental Marine Biology and Ecology* **400**: 7-16.

Whitlerod NS, Zukowski S, Asmus M, Gilligan D and Miller AD (2017) Genetic analyses reveal limited dispersal and recovery potential in the large freshwater crayfish *Euastacus armatus* from the southern Murray-Darling Basin. *Marine and Freshwater Research* **68**: 213-225.

Whitlock MC, and McCauley DE (1999) Indirect measures of gene flow and migration: F_{ST} not equal $1/(4Nm+1)$. *Heredity* **82**: 117-125.

Ye J, Fang L, Zheng HK, Zhang Y, Chen J *et al.* (2006) WEGO: a web tool for plotting GO annotations. *Nucleic Acids Research* **34**: W293-W297.

York KL, Blacket MJ and Appleton BR (2008) The Bassian Isthmus and the major ocean currents of southeast Australia influence the phylogeography and population structure of a southern Australian intertidal barnacle *Catomerus polymerus* (Darwin). *Molecular Ecology* **17**: 1948-1961.

Young IR, Zieger S and Babanin AV (2011) Global Trends in Wind Speed and Wave Height. *Science* **332**: 451-455.

Young M, Ierodiaconou D and Womersley T (2015) Forests of the sea: Predictive habitat modelling to assess the abundance of canopy forming kelp forests on temperate reefs. *Remote Sensing of Environment* **170**: 178-187.

Young MA, Iampietro PJ, Kvitek RG and Garza CD (2010) Multivariate bathymetry-derived generalized linear model accurately predicts rockfish distribution on Cordell Bank, California, USA. *Marine Ecology Progress Series* **415**: 247-261.

Young MA, Ierodiaconou D, Edmunds M, Hulands L and Schimel ACG (2016) Accounting for habitat and seafloor structure characteristics on southern rock lobster (*Jasus edwardsii*) assessment in a small marine reserve. *Marine Biology* **163**: 1-13.

1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143

Table 1 Site location details and corresponding codes for 29 collection sites of *Haliotis rubra* from south eastern Australia.

Zone and Location	Code	GPS Location	
		Latitude	Longitude
<i>Western Zone</i>			
Whites Beach	WHI	-38.349	141.384
Bully Cove	BUL	-38.397	141.411
Inside Murrels	INM	-38.404	141.526
Inside Nelson	INE	-38.404	141.560
Lawrence Rocks	LRO	-38.404	141.667
Lady Julia Percy Island	LJP	-38.421	141.995
The Craggs	OSC	-38.390	142.135
Warrnambool	WAR	-38.397	142.459
<i>Central Zone</i>			
Cat Reef	CAT	-38.741	143.188
Moonlight Head	MLH	-38.767	143.266
White Cliffs	WCF	-38.758	143.330
Blanket Bay	BLK	-38.827	143.586
Point Cook, Port Phillip Bay	PCO	-37.946	144.756
Freds Farm, Port Phillip Bay	FFM	-37.870	144.888
Seal Rocks	SRO	-38.528	145.100
The Nobbies	NOB	-38.518	145.110

Cape Woolami	WOO	-38.564	145.346
Arch Rock	ARK	-38.847	145.894
Cape Liptrap	LIP	-38.889	145.906
Cape Liptrap site 2	CCC	-38.901	145.913
Grinder Bay	GRB	-38.902	145.956
Tongue Point	TON	-38.994	146.258
Norman Point	NRM	-39.025	146.242
Great Glennie Island	GGI	-39.087	146.231
Norman Island	NMP	-39.053	146.320

Eastern Zone

Marlo Frenches Narrows	MFN	-37.809	148.596
Point Ricardo	PRC	-37.813	148.619
Cape Conron	CCO	-37.814	148.731
East Cape Deep	ECD	-37.809	148.749

1144

1145

1146

1147

1148

1149

1150 **Table 2** Summary of environmental statistics for all 29 collection sites: geology, Vector

1151 Ruggedness Measure (VRM), average winter and summer wave power (kW/m); average winter and

1152 summer orbital velocity (m/s); average summer and seasonal range in sea surface temperature (°C)

1153

Geology			Wave Power (kW/m)		Orbital Velocity (m/s)		Sea Surface Temperature (°C)	
Code	Substrate	VRM	Winter	Summer	Winter	Summer	Summer	Seasonal Range
<i>Western Zone</i>								
WHI	Basalt	0.0017	8.20	4.42	173.13	126.85	17.14	3.08
BUL	Basalt	0.0028	17.64	10.54	177.84	135.03	17.93	3.07
INM	Calcareous	0.0007	6.51	4.96	189.12	160.40	17.96	3.55
INE	Calcareous	0.0007	5.87	3.89	84.83	63.09	17.96	3.55
LRO	Calcareous	0.0028	4.28	1.81	36.42	17.77	18.30	3.74
LJP	Basalt	0.0000	28.76	15.94	97.71	60.13	18.14	3.67
OSC	Basalt	0.0014	20.96	12.04	171.03	121.20	18.21	3.83

WAR	Calcareous	0.0037	3.31	2.24	56.00	44.62	18.07	3.97

Central Zone

CAT	Calcareous	0.0049	20.66	12.86	206.55	159.15	17.69	3.41
MLH	Calcareous	0.0022	0.00	0.00	0.03	0.02	17.69	3.45
WCF	Calcareous	0.0015	14.52	8.96	168.51	123.96	17.65	3.47
BLK	Calcareous	0.0011	2.36	1.81	39.58	32.55	17.88	3.96
PCO	Basalt	0.0000	0.00	0.01	0.78	0.72	21.12	9.39
FFM	Basalt	0.0001	0.01	0.01	0.66	0.70	21.34	9.61
SRO	Basalt	0.0010	0.00	0.00	0.01	0.01	19.04	6.11
NOB	Basalt	0.0012	7.07	4.77	79.35	62.28	19.04	6.11
WOO	Granite	0.0034	8.82	5.62	22.65	13.86	19.05	5.52
ARK	Calcareous	0.0009	9.17	5.42	110.09	85.43	18.78	5.13
LIP	Calcareous	0.0018	9.58	5.29	107.92	77.24	18.78	5.13
CCC	Calcareous	0.0029	11.34	6.27	89.30	61.95	18.68	4.92
GRB	Granite	0.0019	4.46	3.12	107.46	87.20	18.68	4.92
TON	Granite	0.0008	7.01	3.77	42.15	29.24	18.58	4.99
NRM	Granite	0.0002	8.86	3.54	5.76	2.48	18.43	4.76
GGI	Granite	0.0005	9.85	4.93	4.46	2.08	18.15	4.49
NMP	Granite	0.0035	3.90	2.47	41.60	35.35	18.58	4.99

Eastern Zone

MFN	Calcareous	0.0003	3.27	1.77	26.99	14.92	18.62	5.20
PRC	Calcareous	0.0003	3.15	1.75	25.07	14.19	18.62	5.20
CCO	Calcareous	0.0005	3.63	2.00	21.93	12.16	18.59	4.87
ECD	Calcareous	0.0005	3.65	2.00	33.98	20.03	18.59	4.87

1154

1155

1156

1157

1158

1159

1160 **Table 3** Numbers of candidate loci identified as having significant correlations with environmental covariables by BayPass, Latent Factor Mixed
1161 Modelling (LFMM) and Redundancy Analysis (RDA) from a total dataset of 5,239 SNPs for the blacklip abalone *Haliotis rubra*.

1162

	BayPass			LFMM	RDA	
	3 < BF < 10	10 < BF < 20	BF > 20	q < 0.001	a = 0.05	Total candidates
Geology	60	15	4	N/A	29	106
VRM	48	23	3	12	16	95
Winter wave power	46	12	6	25	11	97
Summer wave power	39	11	5	23	12	88
Winter orbital velocity	13	0	1	22	12	43
Summer orbital velocity	13	7	0	24	8	48
Summer average SST	268	270	120	60	47	718
Seasonal temp range	24	18	0	40	0	76

1163

1164

1165

1166

1167

1168

1169

1170

1171

1173 **Table 4** SNP loci identified by GDM analyses as explaining greater than 0% deviance for each
 1174 environmental variable. Numbers of non-significant candidates for each data set were as follows:
 1175 winter wave power = 88; summer wave power = 80; seasonal temperature range = 74; summer sea
 1176 surface temperature = 705; winter orbital velocity = 39; summer orbital velocity = 43; VRM = 75.

Environmental			Deviance			Environmental			Deviance		
variable		SNP locus	Deviance		(%)	variable		SNP locus	Deviance		(%)
Winter wave Power		29000_33			13.2	Summer wave power		10080_72			0.34
		93527_52			0.07			12085_77			17.68
		94891_79			13.47			12099_54			0.07
		31220_73			7.62			14150_22			24.21
		81807_43			0.77			29000_33			14.88
		11925_63			1.03			93527_52			0.1
		10080_72			0.32			22064_40			10.0
		5351_44			8.8			34419_52			8.4
		8277_7			5.96						
Seasonal temperature range		12934_28			0.28	Summer sea surface temperature		9325_22			2.48
		32210_29			0.19			9379_73			0.6
								10080_72			0.55
Winter orbital velocity		8277_7			1.59			12934_28			0.28
		6784_54			3.15			14745_47			1.23
		25404_6			16.9			15884_37			2.03
		30737_35			27.4			17019_41			1.01
								24395_36			0.01
Summer orbital velocity		5038_65			4.92			30165_46			3.14
		6784_54			3.43			34419_52			1.31
		30165_46			4.4			50488_50			0.31
		30737_35			0.19			2755_22			0.34
		25404_6			0.001			32754_60			0.19
VRM											
		2755_22			5.3						
		8277_7			6.71						

	9824_42	0.03				
	11778_32	7				
	12077_74	2.54				
	13056_29	3.07				
	15442_53	9.75				
	15487_52	18.96				
	22064_40	0.34				
	34419_52	3.43				
	24136_79	6.65				
	24692_77	1.52				
	35093_14	18.12				
	2834_53	15.07				
	4401_40	11.96				
	5356_71	5.04				
	13077_33	17.19				
	13824_41	5.70				
	16880_71	8.53				

1177

Table 1 Site location details and corresponding codes for 29 collection sites of *Haliotis rubra* from south eastern Australia.

Zone and Location	Code	GPS Location	
		Latitude	Longitude
Western Zone			
Whites Beach	WHI	-38.349	141.384
Bully Cove	BUL	-38.397	141.411
Inside Murrels	INM	-38.404	141.526
Inside Nelson	INE	-38.404	141.560
Lawrence Rocks	LRO	-38.404	141.667
Lady Julia Percy Island	LJP	-38.421	141.995
The Craggs	OSC	-38.390	142.135
Warrnambool	WAR	-38.397	142.459
Central Zone			
Cat Reef	CAT	-38.741	143.188
Moonlight Head	MLH	-38.767	143.266
White Cliffs	WCF	-38.758	143.330
Blanket Bay	BLK	-38.827	143.586
Point Cook, Port Phillip Bay	PCO	-37.946	144.756
Freds Farm, Port Phillip Bay	FFM	-37.870	144.888
Seal Rocks	SRO	-38.528	145.100
The Nobbies	NOB	-38.518	145.110
Cape Woolami	WOO	-38.564	145.346
Arch Rock	ARK	-38.847	145.894
Cape Liptrap	LIP	-38.889	145.906
Cape Liptrap site 2	CCC	-38.901	145.913
Grinder Bay	GRB	-38.902	145.956
Tongue Point	TON	-38.994	146.258
Norman Point	NRM	-39.025	146.242
Great Glennie Island	GGI	-39.087	146.231
Norman Island	NMP	-39.053	146.320

Eastern Zone

Marlo Frenches Narrows	MFN	-37.809	148.596
Point Ricardo	PRC	-37.813	148.619
Cape Conron	CCO	-37.814	148.731
East Cape Deep	ECD	-37.809	148.749

Table 2 Summary of environmental statistics for all 29 collection sites: geology, Vector Ruggedness Measure (VRM), average winter and summer wave power (kW/m); average winter and summer orbital velocity (m/s); average summer and seasonal range in sea surface temperature (°C)

Code	Geology		Wave Power (kW/m)		Orbital Velocity (m/s)		Sea Surface Temperature (°C)	
	Substrate	VRM	Winter	Summer	Winter	Summer	Summer	Seasonal Range
Western Zone								
WHI	Basalt	0.0017	8.20	4.42	173.13	126.85	17.14	3.08
BUL	Basalt	0.0028	17.64	10.54	177.84	135.03	17.93	3.07
INM	Calcareous	0.0007	6.51	4.96	189.12	160.40	17.96	3.55
INE	Calcareous	0.0007	5.87	3.89	84.83	63.09	17.96	3.55
LRO	Calcareous	0.0028	4.28	1.81	36.42	17.77	18.30	3.74
LJP	Basalt	0.0000	28.76	15.94	97.71	60.13	18.14	3.67
OSC	Basalt	0.0014	20.96	12.04	171.03	121.20	18.21	3.83
WAR	Calcareous	0.0037	3.31	2.24	56.00	44.62	18.07	3.97
Central Zone								
CAT	Calcareous	0.0049	20.66	12.86	206.55	159.15	17.69	3.41
MLH	Calcareous	0.0022	0.00	0.00	0.03	0.02	17.69	3.45
WCF	Calcareous	0.0015	14.52	8.96	168.51	123.96	17.65	3.47
BLK	Calcareous	0.0011	2.36	1.81	39.58	32.55	17.88	3.96
PCO	Basalt	0.0000	0.00	0.01	0.78	0.72	21.12	9.39
FFM	Basalt	0.0001	0.01	0.01	0.66	0.70	21.34	9.61
SRO	Basalt	0.0010	0.00	0.00	0.01	0.01	19.04	6.11
NOB	Basalt	0.0012	7.07	4.77	79.35	62.28	19.04	6.11
WOO	Granite	0.0034	8.82	5.62	22.65	13.86	19.05	5.52
ARK	Calcareous	0.0009	9.17	5.42	110.09	85.43	18.78	5.13
LIP	Calcareous	0.0018	9.58	5.29	107.92	77.24	18.78	5.13
CCC	Calcareous	0.0029	11.34	6.27	89.30	61.95	18.68	4.92
GRB	Granite	0.0019	4.46	3.12	107.46	87.20	18.68	4.92
TON	Granite	0.0008	7.01	3.77	42.15	29.24	18.58	4.99
NRM	Granite	0.0002	8.86	3.54	5.76	2.48	18.43	4.76
GGI	Granite	0.0005	9.85	4.93	4.46	2.08	18.15	4.49
NMP	Granite	0.0035	3.90	2.47	41.60	35.35	18.58	4.99

Eastern Zone

MFN	Calcareous	0.0003	3.27	1.77	26.99	14.92	18.62	5.20
PRC	Calcareous	0.0003	3.15	1.75	25.07	14.19	18.62	5.20
CCO	Calcareous	0.0005	3.63	2.00	21.93	12.16	18.59	4.87
ECD	Calcareous	0.0005	3.65	2.00	33.98	20.03	18.59	4.87

Table 3 Numbers of candidate loci identified as having significant correlations with environmental covariables by BayPass, Latent Factor Mixed Modelling (LFMM) and Redundancy Analysis (RDA) from a total dataset of 5,239 SNPs for the blacklip abalone *Haliotis rubra*.

	BayPass			LFMM	RDA	Total candidates
	3 < BF < 10	10 < BF < 20	BF > 20	q < 0.001	a = 0.05	
Geology	60	15	4	N/A	29	106
VRM	48	23	3	12	16	95
Winter wave power	46	12	6	25	11	97
Summer wave power	39	11	5	23	12	88
Winter orbital velocity	13	0	1	22	12	43
Summer orbital velocity	13	7	0	24	8	48
Summer average SST	268	270	120	60	47	718
Seasonal temp range	24	18	0	40	0	76

Table 4 SNP loci identified by GDM analyses as explaining greater than 0% deviance for each environmental variable. Numbers of non-significant candidates for each data set were as follows: winter wave power = 88; summer wave power = 80; seasonal temperature range = 74; summer sea surface temperature = 705; winter orbital velocity = 39; summer orbital velocity = 43; VRM = 75.

Environmental			Deviance			Environmental			Deviance		
variable		SNP locus	SNP locus		(%)	variable		SNP locus	SNP locus		(%)
Winter wave Power		29000_33			13.2	Summer wave power		10080_72			0.34
		93527_52			0.07			12085_77			17.68
		94891_79			13.47			12099_54			0.07
		31220_73			7.62			14150_22			24.21
		81807_43			0.77			29000_33			14.88
		11925_63			1.03			93527_52			0.1
		10080_72			0.32			22064_40			10.0
		5351_44			8.8			34419_52			8.4
		8277_7			5.96						
Seasonal temperature range						Summer sea surface temperature					
		12934_28			0.28			9325_22			2.48
		32210_29			0.19			9379_73			0.6
								10080_72			0.55
Winter orbital velocity		8277_7			1.59			12934_28			0.28
		6784_54			3.15			14745_47			1.23
		25404_6			16.9			15884_37			2.03
		30737_35			27.4			17019_41			1.01
								24395_36			0.01
Summer orbital velocity		5038_65			4.92			30165_46			3.14
		6784_54			3.43			34419_52			1.31
		30165_46			4.4			50488_50			0.31

	30737_35	0.19			2755_22	0.34
	25404_6	0.001			32754_60	0.19
VRM	2755_22	5.3				
	8277_7	6.71				
	9824_42	0.03				
	11778_32	7				
	12077_74	2.54				
	13056_29	3.07				
	15442_53	9.75				
	15487_52	18.96				
	22064_40	0.34				
	34419_52	3.43				
	24136_79	6.65				
	24692_77	1.52				
	35093_14	18.12				
	2834_53	15.07				
	4401_40	11.96				
	5356_71	5.04				
	13077_33	17.19				
	13824_41	5.70				
	16880_71	8.53				

mec_15128_f3.png

mec_15128_f4.png

mec_15128_f5.png

mec_15128_f6.png

mec_15128_f8.tif

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Miller, AD;Hoffmann, AA;Tan, MH;Young, M;Ahrens, C;Cocomazzo, M;Ratray, A;Ierodiasconou, DA;Treml, E;Sherman, CDH

Title:

Local and regional scale habitat heterogeneity contribute to genetic adaptation in a commercially important marine mollusc (*Haliotis rubra*) from southeastern Australia

Date:

2019-06-01

Citation:

Miller, A. D., Hoffmann, A. A., Tan, M. H., Young, M., Ahrens, C., Cocomazzo, M., Ratray, A., Ierodiasconou, D. A., Treml, E. & Sherman, C. D. H. (2019). Local and regional scale habitat heterogeneity contribute to genetic adaptation in a commercially important marine mollusc (*Haliotis rubra*) from southeastern Australia. *MOLECULAR ECOLOGY*, 28 (12), pp.3053-3072. <https://doi.org/10.1111/mec.15128>.

Persistent Link:

<http://hdl.handle.net/11343/285938>