Multivisceral, vascular and nodal resection for recurrent rectal cancer involving the left renal tract, left pelvic side wall and abdominal aorta.

Kasmira Wilson MBBS FRACS¹,
Peadar S Waters MD FRCS¹
Oliver Peacock PhD FRCS¹,
Alexander G Heriot MD FRACS¹
Timothy Wagner MBBS FRACS²
Satish K Warriar MS FRACS¹

1. Colorectal Surgery Unit, Peter MacCallum Cancer Centre, Melbourne, Australia.
2. Department of Vascular Surgery, Royal Melbourne Hospital, Melbourne, Australia.

Corresponding Author: Mr Peadar Waters, Colorectal Surgery Unit, Peter MacCallum Cancer Centre, 305 Grattan Street, Melbourne, Victoria 3000, Australia

Telephone Number: 0385595000, Fax Number: 0385597379

Email: peadarwaters@hotmail.com

Word Count: 732

References: 8

No sources of support

Key Words: Pelvic exenteration, rectal cancer, pelvic malignancy, recurrent rectal cancer, vascular reconstruction

Category of Manuscript: Images for surgeons

CASE REPORT

A 47-year old male initially presented with rectal bleeding and was subsequently diagnosed with a poorly differentiated rectal adenocarcinoma at 5cm from the anal verge in July 2016. The patient completed neo-adjuvant long-course chemoradiotherapy (50.4Gy/28f, 5FU), followed by a laparoscopic ultralow anterior resection and defunctioning loop ileostomy in January 2017. Final

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/ans.15281](https://doi.org/10.1111/ans.15281)

histopathological staging was pT3N2 (9/18 lymph nodes positive) mismatch-repair proficient mucinous signet ring adenocarcinoma, with clear resection margins and he completed adjuvant chemotherapy (FOLFOX) in June 2017.

Routine surveillance computed tomography (CT) scan in February 2018, revealed left sided hydronephrosis and an atrophic left kidney, with subsequent magnetic resonance imaging (MRI) demonstrating a 33x23x17 mm infiltrative mass at the bifurcation of the common iliac vessels. A F-18 fluorodeoxyglucose (FDG) positron emission tomography (PET) scan confirmed an FDG avid left pelvic recurrence with an additional FDG avid para-aortic lymph node at the 3rd lumbar vertebral level. Following discussion at the colorectal multidisciplinary team (MDT) meeting, he received four cycles of systemic therapy (FOLFIRI & Bevacizumab) and a repeat PET scan revealed a response to therapy at both sites with no further progressive disease. Additional chemoradiotherapy (39.6Gy/22f, 5FU) was administered to the left pelvic sidewall recurrence. Restaging imaging did not reveal progression of disease (Figure 1), therefore a curative resection with extensive vascular resection was planned. Four weeks prior to definitive surgical intervention, the patient underwent a right to left femoral-femoral artery crossover graft and a left to right femoral-femoral vein crossover graft, without complication in November 2018.

Subsequently the patient underwent an extensive field lymphadenectomy, pelvic exenteration with en-bloc left iliac vessels, abdominoperineal and pre-sacral fascia resection, left nephroureterectomy and excision of the infrarenal aorta with reconstruction with a Dacron tube graft in December 2018 (Figure 2 and Figure 3). Histopathology demonstrated clear resection margins, but recurrent poorly differentiated signet ring adenocarcinoma with direct extension into the left ureter and left iliac vessels, with 1 out of 36 lymph nodes involved (pT4N1). The resected portion of the aorta also had

signet ring adenocarcinoma present, but clear superior and inferior resection margins were achieved, with no luminal surface involvement either. The patient received extended antibiotic prophylaxis due to a graft reconstruction with concomitant bowel resection. The patient was discharged home 9 days after surgery, with no immediate complications.

DISCUSSION

Local recurrence rates after curative surgical resection for colorectal cancer vary between 3 and 38% of patients(1-3). Recurrent disease involving vascular structures, such as the aorta, have previously been deemed as contraindications for resection. Without treatment, these patients have a median survival of 8 months and may experience a poor quality of life, whilst systemic therapy achieves only temporary symptomatic palliation in the majority of patients(4). However, complete surgical resection offers the potential for cure, providing a negative resection margin is achieved. Reluctance to pursue surgical resection for these cases has been based on the assumed significant morbidity, limitations in achieving clear margins and the impact on prognosis(5, 6). Nevertheless, recent small series for vascular resections have presented good outcomes(4, 7, 8).

Pelvic exenteration is an established procedure for recurrent pelvic cancer, when clear resection margins are deemed achievable. Traditionally, pelvic sidewall involvement has been considered a contraindication for surgery, with the lateral compartment typically posing the most challenge(7). However, with improved surgical technique, surgical units with exenterative experience have expanded the boundaries of resectability and redefined what constitutes resectable disease(7). This case demonstrates that clear resection margins can be achieved in selected patients with en-bloc major vascular resections within a specialised centre for exenterative surgery and vascular reconstruction can be performed safely.

Furthermore, given the patient's increasing size of the paraortic node on PET scan, and following a robust discussion in our specialist MDT setting, a decision was made to prioritise systemic relapse. The rationale behind this approach was to assess the disease biology and response and to ensure there was no further progression prior to embarking upon large scale surgery. Systemic control was achieved as shown by lack of progression on subsequent imaging.

Following further discussion, this case was deemed surgically resectable with curative intent, allowing local therapy to be performed and meticulous planning of the surgical intervention with a specialist vascular surgeon. The authors believe this case with aortoiliac and pelvic sidewall nodal recurrence again highlights the importance of multidisciplinary management in complex recurrent rectal cancer patients

References:

1. Pilipshen SJ, Heilweil M, Quan SH, Sternberg SS, Enker WE. Patterns of pelvic recurrence following definitive resections of rectal cancer. *Cancer*. 1984;53(6):1354-62.
2. McDermott FT, Hughes ES, Pihl E, Johnson WR, Price AB. Local recurrence after potentially curative resection for rectal cancer in a series of 1008 patients. *Br J Surg*. 1985;72(1):34-7.
3. Haddock MG, Miller RC, Nelson H, Pemberton JH, Dozois EJ, Alberts SR, et al. Combined modality therapy including intraoperative electron irradiation for locally recurrent colorectal cancer. *Int J Radiat Oncol Biol Phys*. 2011;79(1):143-50.
4. Abdelsattar ZM, Mathis KL, Colibaseanu DT, Merchea A, Bower TC, Larson DW, et al. Surgery for locally advanced recurrent colorectal cancer involving the aortoiliac axis: can we achieve R0 resection and long-term survival? *Dis Colon Rectum*. 2013;56(6):711-6.

5. Boyle KM, Sagar PM, Chalmers AG, Sebag-Montefiore D, Cairns A, Eardley I. Surgery for locally recurrent rectal cancer. *Dis Colon Rectum*. 2005;48(5):929-37.
6. Bouchard P, Efron J. Management of recurrent rectal cancer. *Ann Surg Oncol*. 2010;17(5):1343-56.
7. Brown KG, Koh CE, Solomon MJ, Qasabian R, Robinson D, Dubenec S. Outcomes After En Bloc Iliac Vessel Excision and Reconstruction During Pelvic Exenteration. *Dis Colon Rectum*. 2015;58(9):850-6.
8. Austin KK, Solomon MJ. Pelvic exenteration with en bloc iliac vessel resection for lateral pelvic wall involvement. *Dis Colon Rectum*. 2009;52(7):1223-33.

ANS_15281_ANZ Figure 1.tiff

ANS_15281_ANZ Figure 2 .tiff

ANS_15281_ANZ Figure 3 .tiff

Figure legend

Figure 1A: Sagittal Magnetic resonance imaging demonstrating recurrent rectal tumour invading into left pelvic sidewall (red arrow) with nodal mass left external iliac artery 29mm x 29mm. **Figure**

Figure 1B: FDG avid paraaortic node (red arrow) measuring 31 mm in size proximal to the level of the left common iliac artery to level of left renal hilum.

Figure 2A: Intraoperative imaging demonstrating tumour infiltrating plaque (blue arrow) on left aortic side wall.

Figure 2B: Enbloc resection of paraaortic lymph node invading into aorta with resection of 3cm aorta and Dacron tube graft reconstruction.

Figure 3a: Intraoperative imaging demonstrating resection of presacral fascia from sacral promontory with resection of nodal mass recurrence enbloc with left common iliac artery, left ureter, and gonadals.

Figure 3B: left nephroureterectomy with enbloc retroperitoneal lymphadenectomy

Multivisceral, vascular and nodal resection for recurrent rectal cancer involving the left renal tract, left pelvic side wall and abdominal aorta.

Kasmira Wilson MBBS FRACS¹,
Peadar S Waters MD FRCS¹
Oliver Peacock PhD FRCS¹,
Alexander G Heriot MD FRACS¹
Timothy Wagner MBBS FRACS²
Satish K Warriar MS FRACS¹

1. Colorectal Surgery Unit, Peter MacCallum Cancer Centre, Melbourne, Australia.
2. Department of Vascular Surgery, Royal Melbourne Hospital, Melbourne, Australia.

Corresponding Author: Mr Peadar Waters, Colorectal Surgery Unit, Peter MacCallum Cancer Centre, 305 Grattan Street, Melbourne, Victoria 3000, Australia

Telephone Number: 0385595000, Fax Number: 0385597379

Email: peadarwaters@hotmail.com

Word Count: 732

References: 8

No sources of support

Key Words: Pelvic exenteration, rectal cancer, pelvic malignancy, recurrent rectal cancer, vascular reconstruction

Category of Manuscript: Images for surgeons

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Wilson, K;Waters, PS;Peacock, O;Heriot, AG;Wagner, T;Warrier, SK

Title:

Multivisceral, vascular and nodal resection for recurrent rectal cancer involving the left renal tract, left pelvic side wall and abdominal aorta

Date:

2020-04-01

Citation:

Wilson, K., Waters, P. S., Peacock, O., Heriot, A. G., Wagner, T. & Warrier, S. K. (2020). Multivisceral, vascular and nodal resection for recurrent rectal cancer involving the left renal tract, left pelvic side wall and abdominal aorta. ANZ JOURNAL OF SURGERY, 90 (4), pp.632-634. <https://doi.org/10.1111/ans.15281>.

Persistent Link:

<http://hdl.handle.net/11343/285946>