DR PENELOPE ANNE HANCOCK (Orcid ID : 0000-0001-5531-1082)

Article type : Research Article

Handling Editor: Bret Elderd

PREDICTING THE SPATIAL DYNAMICS OF *WOLBACHIA* INFECTIONS IN *AEDES AEGYPTI* ARBOVIRUS VECTOR POPULATIONS IN HETEROGENEOUS LANDSCAPES

Penelope A. Hancock^{a*}, Scott A. Ritchie^b, Constantianus J. M. Koenraadt^c, Thomas W. Scott^d, Ary A. Hoffmann^e, H. Charles J. Godfray^a

a. *Corresponding author. Department of Zoology, University of Oxford, Oxford, OX1 3PS, UK. Email: hancock.penelope@gmail.com

b. Australian Institute of Tropical Health and Medicine, James Cook University, 4870, Australia.

c. Laboratory of Entomology, Wageningen University, 6700AA, The Netherlands.

d. Department of Entomology and Nematology, University of California, Davis, CA 95616-5270, USA.

e. Bio 21 Molecular Science and Biotechnology Institute, University of Melbourne, 3010, Australia.

ABSTRACT

1. A promising strategy for reducing the transmission of dengue and other arboviral human diseases by *Aedes aegypti* mosquito vector populations involves field introductions of the endosymbiotic bacteria *Wolbachia*.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/1365-2664.13423](https://doi.org/10.1111/1365-2664.13423)

This article is protected by copyright. All rights reserved

Wolbachia infections inhibit viral transmission by the mosquito, and can spread between mosquito hosts to reach high frequencies in the vector population. *Wolbachia* spreads by maternal transmission, and spread dynamics can be variable and highly dependent on natural mosquito population dynamics, population structure and fitness components.

2. We develop a mathematical model of an *Ae. aegypti* metapopulation that incorporates empirically validated relationships describing density-dependent mosquito fitness components. We assume that density dependent relationships differ across subpopulations, and construct heterogeneous landscapes for which model-predicted patterns of variation in mosquito abundance and demography approximate those observed in field populations. We then simulate *Wolbachia* release strategies similar to that used in field trials.
3. We show that our model can produce rates of spatial spread of *Wolbachia* similar to those observed following field releases.
4. We then investigate how different types of spatio-temporal variation in mosquito habitat, as well as different fitness costs incurred by *Wolbachia* on the mosquito host, influence predicted spread rates. We find that fitness costs reduce spread rates more strongly when the habitat landscape varies temporally due to stochastic and seasonal processes.
5. *Synthesis and applications.* Our empirically based modelling approach represents effects of environmental heterogeneity on the spatial spread of *Wolbachia*. The models can assist in interpreting observed spread patterns following field releases and in designing suitable release strategies for targeting spatially heterogeneous vector populations.

KEYWORDS

arbovirus, dengue, gene drive, spatial spread, *wAlbB*, *wMel*, *wMelPop*, Zika

INTRODUCTION

Aedes aegypti mosquitoes transmit several arboviruses, including dengue which infects an estimated 390 million people per year (Bhatt *et al.* 2013), as well as

Zika and chikungunya which have expanded geographically on a global scale in recent years (Nunes *et al.* 2015; Faria *et al.* 2017). Vaccines against these viruses are not yet widely available (Saez-Llorens *et al.* 2017) and disease control strategies rely on interventions that target the mosquito vector. A promising new control strategy involves the field releases of the endosymbiotic bacteria *Wolbachia* into *Ae. aegypti* populations. *Wolbachia* infections in *Ae. aegypti* inhibit virus development and their transmission by the vector (Moreira *et al.* 2009; Walker *et al.* 2011). *Wolbachia* is maternally transmitted in the mosquito host, and utilizes a mechanism known as cytoplasmic incompatibility (CI), which drives the invasion of the bacteria throughout the population. CI works by preventing the development of offspring from matings between *Wolbachia*-infected males and uninfected females (Caspari & Watson 1959), conferring a strong relative fitness advantage to infected females that are able to mate with both infected and uninfected males. Field releases conducted in northeast Australia in 2011 achieved the first successful invasion and stable establishment of a *Wolbachia* strain (*w*Mel) in wild *Ae. aegypti* populations (Hoffmann *et al.* 2011). Current field trials are evaluating the public health impact, as measured by reductions in human disease incidence, of *Wolbachia* infected *Ae. aegypti* (O'Neill *et al.* 2018).

Given the growing need and potential for applying *Wolbachia* to mosquito vector control and disease prevention, it is necessary to understand the dynamics of *Wolbachia* spread in field *Ae. aegypti* populations. The spatial spread of *Wolbachia* is expected to depend on the natural heterogeneity in mosquito abundance and demography that arises from changing ecological and environmental conditions (Hancock *et al.* 2016a; Hancock *et al.* 2016b). Theoretical studies demonstrate that spatial heterogeneity in host abundance is critical to *Wolbachia* spread, slowing and sometimes stopping the invasion (Barton 1979; Hancock & Godfray 2012). Demographic variation amongst individual hosts, and how this is spatially structured, also has important consequences for spread because the rate of maternal transmission of *Wolbachia* depends directly on its hosts' fitness (Hancock *et al.* 2016b).

In mosquito populations, important demographic traits such as female fecundity and larval development rates can vary dramatically depending on the levels of density-dependent competition for limited food resources that are experienced during larval development (Barrera *et al.* 2006; Muriu *et al.* 2013; Hancock *et al.* 2016a). Parameters describing female fecundity and larval development rates are fundamental to models of *Wolbachia* dynamics in mosquito populations, because the bacteria is only transmitted maternally (Turelli 1994). Incorporation of density-dependent demographic variation in these parameters into models of the spatial dynamics of *Wolbachia* has, however, been hindered by the lack of an empirical understanding of the form of density dependence (Legros *et al.* 2009). Models of the spatial spread of *Wolbachia* in mosquito populations developed thus far have typically either assumed constant demographic rates across space and time (Turelli & Hoffmann 1991; Schmidt *et al.* 2017; Turelli & Barton 2017), or restricted the effects of density-dependence to impacts on larval survival (Hancock & Godfray 2012).

Here we develop a spatially explicit model of *Wolbachia*-*Ae. aegypti* dynamics that aims to incorporate realistic patterns of demographic variation in the mosquito population. *Ae. aegypti* is a highly domestic species that is found in or near areas with human habitation and has low rates of dispersal (Harrington *et al.* 2005), which makes population size estimation easier than for other mosquito species (Ritchie *et al.* 2013a). Studies from multiple geographic regions show characteristic strong spatial overdispersion in population size (Getis *et al.* 2003; Focks & Alexander 2006; Koenraadt *et al.* 2008; Jeffery *et al.* 2009; Aldstadt *et al.* 2011; LaCon *et al.* 2014), which may arise from the reliance of *Ae. aegypti* larvae on developing in water filled artificial containers close to human dwellings (Southwood *et al.* 1972; Aldstadt *et al.* 2011; Williams *et al.* 2013). Many of these containers produce low numbers of pupae (and adults), which is thought to be due limited larval food resources (Arrivillaga & Barrera 2004; Focks & Alexander 2006). Adult body size in field populations is typically smaller than those produced from larvae that are fed *ad libitum* in the laboratory (Nasci 1986; Barrera *et al.* 2006), with more crowded field containers typically producing smaller adults (Schneider *et al.* 2004). This indicates that larval development in

field *Ae. aegypti* populations is affected by food limitation arising from density-dependent larval competition, which adversely affects juvenile and adult fitness components (Hancock *et al.* 2016a). Thus, density dependent effects on mosquito production and fitness could be an impediment for *Wolbachia* to establish and spread, especially as releases of additional mosquitoes will lead to increased crowding in containers.

Wolbachia spread may also be impacted by spatio-temporal variation in mosquito habitat (Schmidt *et al.* 2017). Field studies have shown that over time it is not always the same houses that are responsible for the production of high numbers of *Ae. aegypti* (Jeffery *et al.* 2009; LaCon *et al.* 2014), suggesting that the quality of mosquito habitat within a given house may be temporally variable. In some environments, including northeast Australia, *Ae. aegypti* abundance is seasonal and higher in the wet season (November-April) than in the dry season (May-October) (Ritchie *et al.* 2013b). In some other environments, however, *Ae. aegypti* abundance does not show seasonal variation (Scott *et al.* 2000b; Koenraadt *et al.* 2008).

We develop a metapopulation model that describes the spatially heterogeneous demography of *Ae. aegypti* using empirically validated relationships that link density-dependent demographic traits to mosquito abundance. Incorporating an empirically realistic form of density-dependence significantly reduces parameter uncertainty compared to other models of *Ae. aegypti* population dynamics. We ask whether our model can produce patterns of demographic variation approximating those observed in field populations, in terms of variation in both mosquito population size and density-dependent demographic traits. We then ask how rates of spatial spread of *Wolbachia* predicted by our model compare to those observed following field releases. We explore the effects on the rate of spread of different types of spatio-temporal variation in the mosquito habitat landscape, including stochastic and seasonal variation, and of different fitness costs incurred by *Wolbachia* on the mosquito host.

MATERIALS AND METHODS

We develop an age-structured model of an *Ae. aegypti* metapopulation that assumes that variation amongst individuals in two demographic traits, larval development times and per-capita female fecundity, is influenced by varying levels of larval density-dependent competition for limited food resources (Hancock *et al.* 2016b) (Fig. 1). The level of density-dependent competition within each of the subpopulations that make up the metapopulation is assumed to depend on the local larval density as well as the quantity of larval food resources available to the subpopulation (Fig 1A). We conceptualize each subpopulation of the metapopulation as a “house” and define the larval habitat quality in each house as high quality, low quality or empty (no larval habitat present). High quality habitats contain a greater larval food resource quantity than low quality habitats.

In order to estimate the functional forms of density-dependent demographic variation occurring in high and low quality habitats, we utilize a recently developed methodology that is based on frequent observations of the temporal variation in the abundance of the juvenile life stages of *Ae. aegypti* in experimental field-caged populations (see below and Hancock *et al.* (2016b)). Mathematical functions describing the density-dependent co-variation in per-capita female fecundity and larval development times occurring in each population are then parameterized from the observed variation in juvenile abundance (Hancock *et al.* 2016b). Details of how these different forms of density-dependence are quantified are given below.

Our model also includes parameters describing density-independent mosquito demographic traits and dispersal. These parameters are assumed to be constant and are estimated using published data. Density-independent demographic traits also affect the level of density-dependent competition occurring in the metapopulation through their effects on population size (Hancock *et al.* 2016b). We describe below the structure of our model, and detail our methodology for estimating density-dependent and density-independent model parameters. Model parameters are provided in Tables 1 & 2.

Model Structure

Our model of mosquito dynamics within each subpopulation follows that described in Hancock et al. (2016b). In summary, the model subdivides the mosquito subpopulation into a series of juvenile stages (including eggs, larvae and pupae) and male and female adults, where adult females are classified according to whether they have reached the minimum age required for oviposition (T_G ; Fig. 1B and Table 1). For each larval cohort, both the mean larval development time, $\mu_{c,q}$, and the standard deviation of the larval development times, $\sigma_{c,q}$, are assumed to depend on $\bar{L}_{c,q}^P$, the average larval density that the larvae in cohort c in a habitat with quality q experience during the time period from hatching to eclosion of the first pupa from the cohort. The per-capita fecundity of adult females aged T_G days or older on day i , $\lambda_{i,q}$, defined as the number of first instar larvae that hatch per day per adult female, is assumed to depend on $\bar{L}_{n,q}^A$, the average larval density in the subpopulation over a fixed time lag of three weeks ending on day $n=i-T_G-T_P$ (where T_P is the duration of the pupal development stage; see Fig. 1B and Table 1). Mosquito demographic traits that are assumed to be density-independent include the daily larval mortality, μ_L , the daily pupal and adult mortality, μ_A , and the developmental time lags T_P , T_G and the time lag between oviposition and hatching of eggs, T_H (Fig. 1B, Table 1). Adult mosquitoes migrate between subpopulations according to a daily probability, p_D , of moving to an adjacent house within the neighbourhood of the eight adjacent houses (Fig. 1A). Details of the mathematical formulation of the model are provided in Section S4.

Estimating the form of density-dependent demographic variation

We estimated the forms of density-dependent variation in mosquito demographic traits occurring in two independent experimental populations of *Ae. aegypti*. The populations were housed in field-cages designed to simulate the natural habitat of *Ae. aegypti* and the climatic conditions of northeast Australia (Darbro et al. 2012) and maintained as described in Hancock et al. (2016a; 2016b) (see also Section S1). Each population received a different larval food

regime. One population, which we refer to as the HF (high food) Population, received a fixed quantity of food (0.32g ground lucerne) three times per week, and the other, referred to as the LF (low food) Population, received this same food quantity only once per week.

We performed counts of all larvae (3 times per week) and pupae (daily) over approximately six months. We obtained an expression for the likelihood of our observed abundances of the juvenile mosquito lifestages conditional on mathematical functions describing the form of larval density-dependent variation in larval development times and per-capita female fecundity. We used Bayesian Markov Chain Monte Carlo (MCMC) methods, described in Hancock *et al.* (2016b), to estimate the parameters of these mathematical functions for each field-caged population (see Section S2). For both populations, the fitted models accurately describe the major features of the temporal variation in juvenile mosquito abundances in the field-caged populations (Fig S1). The form of density-dependent demographic variation occurring in the high and low quality habitats in our modelled metapopulation are given by the fitted relationships for the HF and LF Populations, respectively. For the HF Population, these model fits have been reported previously (Hancock *et al.* 2016b), while for the LF Population our results are reported here for the first time. We note that incorporating density-dependent variation in additional demographic traits, such as the daily larval mortality rate, may have produced a closer fit to the field cage observations.

Estimating density-independent demographic traits

Values of parameters describing density-independent demographic rates (assumed constant) were estimated from published studies (Table 1). We estimated the daily probability, p_a that adults move to an adjacent house using published data from mark-release-recapture (MRR) experiments performed in Thailand (lines 1 and 2 of Table 2 of Harrington *et al.* (2005)) that record numbers of recaptured *Ae. aegypti* mosquitoes at different distances from the release sites. We ran our metapopulation model to simulate the MRR protocol described in Harrington *et al.* assuming a constant daily rate of adult mosquito

mortality μ_A (Table 1). We calculated the value of p_d that provided the minimum least squares fit to the observed recapture rates. We obtained $p_d = 0.56$ (Table 1), but note that the data describing both mosquito mortality and the recapture rates is insufficient to quantify the uncertainty in this estimate. We assumed that larvae experience a constant daily rate of mortality μ_L similar to the value estimated for our field-cage experiments (Hancock *et al.* 2016b), and that pupae experience the same daily mortality rate as adults. Values of the development time lags T_G , T_H and T_P were estimated from our field-cage experiments (Hancock *et al.* 2016b).

Estimating spatial variation in habitat quality

We denote the proportion of houses in the metapopulation with high habitat quality, low habitat quality and empty as h_H , h_L , and h_E . We assume that the habitat quality type across houses is spatially uncorrelated (Getis *et al.* 2003) and is described by a multinomial distribution $\text{Mult}(h_H, h_L, h_E)$. We then compare the abundance of pupae across subpopulations simulated by our model with observations from field surveys conducted by Koenraadt *et al.* (2008). They recorded the number of pupae found across a set of residential properties distributed over two villages in Kamphaeng Phet province, Thailand, in four surveys conducted over 2 month periods at the end of the dry season (March and April, which is the beginning of the dengue transmission season) and at the end of the wet season (September and October, which is the end of the dengue transmission season) in 2004 and 2005. They made a total of 2123 observations across 604 houses. For each house, pupae in all water-holding containers were counted. Pupal abundances were determined for all potential larval development sites. The observed distribution of pupal numbers across houses across both surveys shows a spatially aggregated pattern that is typical for *Ae. aegypti* field populations (Getis *et al.* 2003; Focks & Alexander 2006; Jeffery *et al.* 2009; LaCon *et al.* 2014).

We ran one-year simulations of our metapopulation model across which the only parameters that varied were the proportions of houses with high quality, low

quality and empty habitats (h_H , h_L and h_E). We calculated the parameters of the multinomial habitat quality distribution for which the simulated distribution of pupal numbers per house best matched that observed (Table 2). This procedure assumed that the habitat quality of each house does not vary over time because the pupal count data cover short time periods but are not temporally explicit within this time frame (see the Discussion). Further details of this methodology are provided in Section S3.

Comparing the predicted intensity of density dependence with observations from field populations

We assessed how the intensity of larval density-dependent competition occurring in our modelled metapopulation compared to observations from field populations using observed wing lengths of adult females produced from field-collected *Ae. aegypti* pupae. Food limitation during larval development greatly affects overall adult body size, which can be estimated using wing length (Briegel 1990; Barrera *et al.* 2006), so field populations that show smaller adult wing lengths are likely to have experienced more intense larval resource competition (Barrera *et al.* 2006). Temperature conditions can also affect *Ae. aegypti* body size (Scott *et al.* 2000a; Mohammed & Chadee 2011), although in our field-cage populations changes in larval water temperature had less influence on demographic traits than variation in larval density (Hancock *et al.* 2016a) (see the Discussion). We used published observations of the average wing length per household of 2316 female *Ae. aegypti* collected as pupae from 2931 houses in Iquitos, Peru (Schneider *et al.* 2004). The wing lengths of the sampled mosquitoes are highly variable, and median wing length is low relative to that occurring when adult females experience plentiful food conditions during larval development (Fig. 1 of (Schneider *et al.* 2004)). Other observed wing lengths distributions obtained from field-sampled *Ae. aegypti* also show similar characteristics (Nasci 1986; Barrera *et al.* 2006).

In order to compare these wing length observations to the output of our model, we used the relationship between wing length and female fecundity obtained by Briegel (1990). Briegel (1990) recorded adult female wing length, w , and the number of eggs oviposited per gonotrophic cycle, $\tilde{\lambda}_g$, for 206 *Ae. aegypti* females

reared in a laboratory under varying larval density conditions and blood-fed on human hosts (Table 3 of Briegel 1990)). He then fitted a linear relationship to these observations:

$$\tilde{\lambda}_G = 2.505w^3 - 8.616 \quad (1)$$

We applied eqn (1) to estimate the distribution of the average $\tilde{\lambda}_G$ per house, $\lambda_{G,h}$, from the observed distribution of average female wing lengths per house. This procedure assumes that individual wing lengths can be approximated by the house average. We then calculated the distribution of the weekly average per-capita female fecundity for each house, λ_h , using observations from our field-cage experiments (see Methods). This procedure makes two further assumptions: (i) $\lambda_{G,h}$ and λ_h are assumed to be directly proportional; and (ii) the adult female wing lengths in our field-caged populations under conditions of plentiful larval food are equal to the highest value observed by Briegel (1990). Having obtained a distribution of per-capita female fecundity across individuals derived from the field wing length data, we compared it with the distribution of per-capita female fecundity values across all subpopulations simulated by our metapopulation model.

Comparing the predicted rate of spatial spread of *w*Mel *Wolbachia* with observations from field populations

We compared the rates of spatial spread of *Wolbachia* predicted by our model to those observed following the *w*Mel field release trials conducted at two sites in the city of Cairns, northeast Australia, in 2013 (Schmidt *et al.* 2017). The two sites, Edge Hill and Parramatta Park, are residential areas close to the center of Cairns. The field trials involved releasing mosquitoes infected with *w*Mel into a restricted area, or “release zone”, with the aim of allowing the *Wolbachia* to spread into surrounding areas. Observed rates of spread of *w*Mel beyond the release zone for both sites over a time period of 20 months are reported in Schmidt *et al.* (2017) (see Methods). We used their Gaussian/logistic model fits to observations of *w*Mel frequencies in pooled samples (Fig. S3 of Schmidt *et al.*

(2017)) to obtain the estimated distance from the edge of the release zone at which the *w*Mel frequency was 0.5 at multiple time points following the releases.

To represent their spatial release strategy, we assume in our model that mosquitoes infected with *w*Mel are liberated within a square-shaped release zone located at the center of a larger area of potential habitat containing 10,000 houses (also square-shaped; 2.5 km²). We set parameters specifying the release protocol to values matching those used in the field trials (these included the area of the release zone, the duration and frequency of releases, and the frequency of *w*Mel inside the release zone at the time of the final release; Table 2). We assumed that a fixed number of infected mosquitoes are released within each house inside the release zone once per week. Because released mosquitoes were reared in the laboratory, we also assumed that the per-capita fecundity of the released females is high and equal to that observed when larval food resources are plentiful (Hancock *et al.* 2016b). Our results assume that houses have dimensions of 25m² (Table 2), which is approximately equal to average area of the housing blocks within the two release zones in Cairns, calculated by randomly sampling 100 houses within each release zone and obtaining the dimensions of each housing block using Google Maps. The available data do not allow estimation of the mosquito habitat quality in the houses within the study site, so we simulated *w*Mel releases for 50 landscapes where the habitat quality in each house (low, high, or empty) was randomly drawn from a multinomial distribution with parameters h_H , h_L , and h_E (Table 1).

Our model of the *Wolbachia* dynamics in each subpopulation of the metapopulation is based on that described in Hancock *et al.* (2016b). The approach subdivides each mosquito life stage into classes infected and uninfected with *Wolbachia*, and describes the effects of cytoplasmic incompatibility on mosquito demography (Fig. 1B); mathematical details are provided in Section S5. We assume high values of the strength of cytoplasmic incompatibility, s_b , and the rate of maternal transmission of *Wolbachia*, ω (Table 1), consistent with observations of the *Wolbachia* strain *w*Mel that has been used in recent field release trials (Hoffmann *et al.* 2011). We begin by

assuming that *Wolbachia* infection does not affect the fitness of its mosquito host (Hancock *et al.* 2016b), and then explore the effects of varying fitness costs on spread dynamics.

Sensitivity analysis

We performed local sensitivity analysis to investigate how changes in mosquito demographic parameters affect three aspects of the dynamics of our metapopulation model: (i) the distance that *Wolbachia* spreads beyond the release zone in 20 months following the final release (using 0.5 *Wolbachia* frequency contour as described above); (ii) the total mosquito population size and (iii) the average per-capita female fecundity across the entire spatial region ((ii) and (iii) are evaluated the equilibrium in the absence of *Wolbachia*). We chose to vary three demographic parameters that are known to be important to the predictions of mosquito population dynamic models, but are poorly quantified for mosquito populations (Schofield 2002; Hancock *et al.* 2016b; Legros *et al.* 2016): the daily larval mortality, μ_L , the daily pupal and adult mortality, μ_A , and the daily probability of adult dispersal, p_D (Table 1). We assume in this analysis that *Wolbachia* infection does not affect mosquito fitness and that the mosquito habitat quality landscape is temporally invariant.

RESULTS

Modelling spatially varying demography in the mosquito population

Modelling spatial variation in mosquito abundance

The fitted distribution of the numbers of pupae across houses is similar in both shape and range to that obtained from the field surveys (Fig. 2A). The predicted pupal numbers are highly aggregated across space due to the small proportion of houses with high quality habitat ($h_H = 0.1$). The fitted proportions of houses with low and empty habitat quality (h_L and h_E) were equal (Fig. 2A). These results demonstrate that the relationships describing density-dependent demography derived from our field-caged population experiments can be applied directly,

without rescaling, to approximate patterns of variation in abundance that occur in field *Ae. aegypti* populations.

Modelling the intensity of density-dependent competition

The median of the field-derived per-capita female fecundity distribution is in the lower part of the per-capita fecundity range and is close to that predicted by our model (Fig. 2B). Thus both field observations and model predictions indicate that density-dependent competition is relatively intense. While both the distributions from the field and model show wide variation in per-capita female fecundity across subpopulations, the variance predicted by our model is larger. The larval development times predicted by our model also vary widely across the metapopulation, with the majority of subpopulations having protracted larval development (Fig. 2B).

Predicting the speed of spatial spread of *Wolbachia*

We simulated *Wolbachia* field releases for 50 landscapes where habitat quality in each house is randomly drawn from a multinomial distribution with parameters h_H , h_L , and h_E (Table 1). We begin by assuming that *Wolbachia* infection has no effect on mosquito fitness. The mean of the simulated rates of spread across all landscapes agrees well with the rates observed at Edge Hill and Parramatta Park (Fig. 3A). This demonstrates that under our default parameter assumptions (Table 1) our model is able to predict *Wolbachia* spread rates similar to those observed in field populations. Spread rates were spatially heterogeneous, however, and spread dynamics varied considerably depending on random differences between the realized habitat quality landscapes (see the credible intervals in Fig. 3A and see also Animation S1).

Effects of stochastic habitat variation

We first extend our analysis to consider landscapes where habitat quality shows stochastic temporal variation. We defined a fixed daily probability t_q that the habitat quality in each house is reassigned to a value chosen randomly from $\text{Mult}(h_H, h_L, h_E)$. We then defined the expected habitat quality turnover period, $\bar{t}_q = 1/t_q$, as the expected time interval between a reassignment of habitat quality

for each house, and considered cases where turnover is slow ($\bar{t}_q = 3$ months) and relatively fast ($\bar{t}_q = 1$ month). For each value of $\bar{t}_q$ we repeated the analysis described in the previous section to obtain predictions of the speed of spatial spread of *Wolbachia* by simulating the release protocol used in the field releases conducted in Edge Hill and Parramatta Park, Cairns (Table 2). We found that the predicted rates of spatial spread of *Wolbachia* for both values of t_q was similar to that obtained when habitat quality was assumed not to vary temporally (Fig. 3B-C).

Effects of seasonal habitat variation

Next we extended our analysis to explore effects of seasonal variation in *Ae. aegypti* abundance on *Wolbachia* spread. We simulated seasonal variation in *Ae. aegypti* abundance by periodically increasing the proportion of high quality habitat, h_H . This involved randomly selecting a set of houses (of size n_s) with either low or empty habitat quality, and reassigning the habitat quality in these houses to high, once every 365 days. This “wet season” habitat quality landscape remained fixed for 182 days, and then a set of high quality habitat houses, also of size n_s , was selected at random and the habitat quality in these houses was reassigned to either low, or empty, with a probability of 0.5. We set the value of n_s such that the proportion of houses with high quality habitat, h_H , increased to 0.175 at the start of the wet season (and returned to 0.1 at the start of the dry season). This produced a simulated pattern of seasonal variation in adult abundance with wet season peak about 4 times higher than the dry season trough (Fig. S3). Published observations of the seasonal variation in the numbers of *Ae. aegypti* caught in Biogents Sentinel traps (BGS) throughout Cairns show a similar relative amplitude across seasons (Fig. 1 of Ritchie et al. (2013b)). We again simulated the above *Wolbachia* field release protocol, starting the releases 40 days into the wet season, and again found that the predicted rate of spatial spread of *Wolbachia* was similar to that obtained when habitat quality was assumed not to vary temporally (Fig. 3D).

Effects of fitness costs

We then investigated how these spread patterns are affected by fitness costs imposed by *Wolbachia* on its mosquito host. Specifically, we assumed that *Wolbachia*-infected individuals experience a reduction in average adult lifespan relative to uninfected individuals by a proportion s . Estimates of fitness costs incurred by *wMel* on *Ae. aegypti* are highly uncertain (Hoffmann *et al.* 2011; Hancock *et al.* 2016b; Schmidt *et al.* 2017) (and see the Discussion) but high costs are unlikely; here we consider values of $s=0.15$ and $s=0.3$. For all four types of temporal variation in mosquito habitat quality considered above (including static landscapes with no temporal variation), we simulated the *Wolbachia* release protocol previously described, but augmented the total release size so that the *Wolbachia* frequency reached the required value at the time of the final release (Table 2).

For environments where the habitat quality landscape is static, either fitness cost resulted in slower rates of *Wolbachia* spread that show less agreement with the rates observed in the field populations (Figs. 3E & 3I). When $s=0.15$ the observed spread rates are above the mean of the simulated rates, but within the 95% quantiles (Fig. 3E), and when $s=0.3$ the observed rates are higher than the quantiles of the simulated rates for times later than six months following the final release (Fig. 3I).

In the presence of fitness costs, predicted *Wolbachia* spread rates also differ across the different types of spatio-temporal variation in mosquito habitat quality. They are slower in landscapes where house habitat quality undergoes stochastic changes over time compared to landscapes where house habitat quality is static, with differences being larger when the fitness cost is higher (Figs. 3F-G & 3J-K). Seasonal variation in mosquito habitat quality also results in slower predicted spread rates, with the difference again being strongest for higher fitness costs (Figs. 3H & 3L). When the *Wolbachia* imposes fitness costs, these types of temporal variation in habitat quality result in slower rates of spatial spread because a reduction in a house's habitat quality causes a localized increase in mosquito mortality. If this occurs before the *Wolbachia* has spread to fixation, the frequency dependent advantage that the *Wolbachia* has gained by

invading and spreading in areas with good habitat quality is lost; thus spread slows down, and may fail if frequencies fall too low. Interestingly, the higher rates of mosquito mortality in temporally variable habitat quality landscapes lead to reductions in the intensity of density-dependent larval competition, which can lead to faster *Wolbachia* spread rates (Hancock *et al.* 2016b). However in the presence of fitness costs this effect did not compensate for the inhibitory effects on rates of spread of temporal changes in house habitat quality.

Sensitivity analysis

The effects of variation in daily larval mortality are relatively straightforward; when larval mortality is lower, the total mosquito population size is higher and average per-capita female fecundity is lower because larval density-dependent competition is more intense (Fig. 4A). Lower values of larval mortality result in slower *Wolbachia* spread, firstly because mosquito generation times are longer when density-dependence is more intense, and secondly because the release of a fixed number of *Wolbachia* infected mosquitoes is less effective at elevating the *Wolbachia* frequency when the abundance of the wild mosquito population is greater. The effects of variation in pupal and adult mortality on mosquito population size and density-dependent demographic rates are the same in direction but stronger in magnitude. Due to these effects, *Wolbachia* spreads more quickly when pupal and adult mortality is higher, even though shorter-lived adults have a reduced lifetime dispersal distance, which inhibits the spatial spread of *Wolbachia* (Fig. 4B,C). Variation in the daily dispersal probability has less effect on mosquito demography, in terms of the total population size and the intensity of density dependence, but effects on the spatial spread of *Wolbachia* across the full range of dispersal probabilities are substantial (Fig. 4C). *Wolbachia* spreads more quickly when the dispersal probability is higher because the bacterium is propagated more quickly through the landscape.

DISCUSSION

Our metapopulation model describes the dynamics of *Ae. aegypti* populations with spatially heterogeneous abundance and demography, incorporating empirically derived relationships that model density-dependent demographic

traits. We demonstrated that the model could approximate the patterns of demographic variation that occur in field populations and predict rates of spatial spread of *Wolbachia* close to those observed following field releases. We discuss below how our representation of spatially structured population dynamics is important to predicting *Wolbachia* spread, and also identify limitations of our model in describing the dynamics of natural populations.

Previous studies have demonstrated several mechanisms through which the dynamics of *Wolbachia* spread are affected by spatial heterogeneity in host abundance (Barton 1979; Hancock & Godfray 2012), and demographic heterogeneity across individuals (Hancock *et al.* 2016a; Hancock *et al.* 2016b). Our modelling approach empirically relates variation in mosquito abundance to density-dependent variation in two demographic traits, namely larval development times and per-capita female fecundity, which are major determinants of the rate of *Wolbachia* transmission through successive (overlapping) mosquito generations. By allowing density dependence relationships to vary across subpopulations, our model is able to represent realistically two important demographic features of *Ae. aegypti* field populations: (i) highly variable, spatially aggregated patterns of abundance; and (ii) a relatively intense level of density-dependent larval resource competition on average across the metapopulation.

However, other aspects of demographic and environmental variation that were not included in our model may also cause variable rates of *Wolbachia* spread. Mosquito mortality and dispersal rates also affect *Wolbachia* spread dynamics (Schofield 2002; Hancock *et al.* 2016b), but these traits are poorly quantified and likely to vary with changing environmental conditions (Harrington *et al.* 2005; Maciel-De-Freitas *et al.* 2007; Hoffmann 2014). Several demographic traits vary due to changes in weather and climate (Scott *et al.* 2000a; Mohammed & Chadee 2011), and predation (Bowatte *et al.* 2013). Moreover, density dependence may impact other demographic traits such as stage-specific daily survival rates (Hancock *et al.* 2016a), and mating success (Segoli *et al.* 2014). Density-dependent dynamics may be further complicated by preferential oviposition

behaviour by adult females, whereby females are more likely to oviposit in containers that are already inhabited by conspecific larvae (Wong *et al.* 2011). Thus our model cannot fully describe the demographic variability in a particular metapopulation. By specifying realistic forms of density dependence, however, our model offers a useful tool for exploring dynamic behaviour across different assumptions about uncertain aspects of mosquito-*Wolbachia* demography and different types of environmental variation, as our results demonstrate. In order to understand the effects of variation in demographic parameters, it is necessary to consider how these demographic rates impact the dynamics of the full system, as demonstrated by our sensitivity analysis (Fig. 4).

A current challenge in forecasting *Wolbachia* dynamics is the difficulty of estimating fitness costs in the field (Hancock *et al.* 2016b), which is exacerbated by the very high levels of demographic variation caused by environmental and genetic effects (Hoffmann *et al.* 2011; Walker *et al.* 2011; Hoffmann 2014; Hancock *et al.* 2016b; Ross *et al.* 2016; Schmidt *et al.* 2017). This is problematic because theory suggests that even small fitness costs strongly affect the spatial spread of *Wolbachia* (Turelli & Hoffmann 1991; Hancock & Godfray 2012). For example, the classical reaction-diffusion models of Barton and Turelli (Barton 1979; Turelli & Hoffmann 1991) predict that a fitness cost of 0.25 halves the equilibrium wave speed of *Wolbachia*. These reaction-diffusion approaches assume spatial and temporal constancy of mosquito abundance and demography (Turelli & Barton 2017). Under plausible sets of demographic and environmental parameters (Tables 1 & 2), and across a range of models of spatio-temporal habitat variation, our model accurately described observed spread rates without assuming that *Wolbachia* incurs fitness costs. It is important to note that, while *Ae. aegypti* abundance varies seasonally in northeast Australia, in many dengue-endemic regions (such as parts of Thailand and Peru) abundance does not show a strong seasonal signal (Morrison *et al.* 2004; Koenraadt *et al.* 2008). Interestingly, however, for cases where the fitness cost incurred by *Wolbachia* was low, our predicted spread rates were relatively robust to the different forms of seasonal and stochastic habitat variation that we considered. While our results predict that fitness costs incurred by *wMel* are usually small, the uncertainty

surrounding the demographic and environmental parameters and processes that we aim to model prevents us from making accurate inferences about the effects of *Wolbachia* on fitness experienced in field *Ae. aegypti* populations.

In summary, we present a mathematical model of the spatial dynamics of *Wolbachia* in *Ae. aegypti* populations that incorporates realistic features of mosquito demographic variation that impact *Wolbachia* spread. Our approach facilitates exploration of the effects of demographic and environmental heterogeneity on the spread of *Wolbachia* following field releases. Our models can be applied to compare the performance of different field release strategies, for example by varying the release size and timing, and the spatial distribution of the release locations. Further, our approach can be extended to explore the dynamics of other mosquito control interventions, such as the release of *Wolbachia*-infected male mosquitoes in order to suppress vector abundance, or the release of mosquitoes carrying gene drive constructs that aim to suppress abundance or lower vectorial capacity (Flores & O'Neill 2018). By developing our understanding of the ecological dynamics of *Ae. aegypti* arbovirus vector populations, and their interactions with vector control interventions, our models can contribute to interpreting observed impacts in field populations and allowing effects of environmental heterogeneity to be considered in designing intervention strategies.

AUTHOR CONTRIBUTIONS

PAH, HCG and SAR conceived the ideas; PAH designed the methodology; PAH collected the data; PAH analysed the data; PAH led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

ACKNOWLEDGEMENTS

This research was supported by a Marie Curie International Outgoing Fellowship within the 7th European Community Framework Programme (Grant no. 326551-WOLBACHIA-MOD).

DATA ACCESSIBILITY

Mosquito age-class abundances for LF Population and observed pupal abundances per house from the pupal suveys conducted by Koenraadt *et al.* (2008) are available at <https://doi.org/10.6084/m9.figshare.8117939.v1> (Hancock et al. 2019)

C++ Computer Code for implementing the metapopulation model is available via Zenodo <http://doi.org/10.5281/zenodo.2784071> (Hancock 2019)

REFERENCES

- Aldstadt, J., Koenraadt, C.J.M., Fansiri, T., Kijchalao, U., Richardson, J., Jones, J.W. & Scott, T.W. (2011) Ecological Modeling of *Aedes aegypti* (L.) Pupal Production in Rural Kamphaeng Phet, Thailand. *PLoS Negl Trop Dis*, **5**.
- Arrivillaga, J. & Barrera, R. (2004) Food as a limiting factor for *Aedes aegypti* in water-storage containers. *J Vector Ecol*, **29**, 11-20.
- Barrera, R., Amador, M. & Clark, G.G. (2006) Ecological factors influencing *Aedes aegypti* (Diptera : Culicidae) productivity in artificial containers in Salinas, Puerto Rico. *J Med Entomol*, **43**, 484-492.
- Barton, N.H. (1979) The dynamics of hybrid zones. *Heredity*, **43**, 341-359.
- Bhatt, S., Gething, P.W., Brady, O.J., Messina, J.P., Farlow, A.W., Moyes, C.L., . . . Hay, S.I. (2013) The global distribution and burden of dengue. *Nature*, **496**, 504-507.
- Bowatte, G., Perera, P., Senevirathne, G., Meegaskumbura, S. & Meegaskumbura, M. (2013) Tadpoles as dengue mosquito (*Aedes aegypti*) egg predators. *Biological Control*, **67**, 469-474.
- Briegel, H. (1990) Metabolic relationship between female body size, reserves, and fecundity of *Aedes aegypti*. *J Insect Physiol*, **36**, 165-172.
- Caspari, E. & Watson, G.S. (1959) On the evolutionary importance of cytoplasmic sterility in mosquitoes. *Evolution*, **13**, 568-570.
- Darbro, J.M., Johnson, P.H., Thomas, M.B., Ritchie, S.A., Kay, B.H. & Ryan, P.A. (2012) Effects of *Beauveria bassiana* on Survival, Blood-Feeding Success,

- 685 and Fecundity of *Aedes aegypti* in Laboratory and Semi-Field Conditions.
 686 *Am J Trop Med Hyg*, **86**, 656-664.
- 687 Faria, N.R., Quick, J., Claro, I.M., Theze, J., de Jesus, J.G., Giovanetti, M., . . . Pybus,
 688 O.G. (2017) Establishment and cryptic transmission of Zika virus in Brazil
 689 and the Americas. *Nature*, **546**, 406-+.
- 690 Flores, H.A. & O'Neill, S.L. (2018) Controlling vector-borne diseases by releasing
 691 modified mosquitoes. *Nature Reviews Microbiology*, **16**, 508-518.
- 692 Focks, D.A. & Alexander, N. (2006) Multicountry study of *Aedes aegypti* pupal
 693 productivity: findings and recommendations. World Health Organization.
- 694 Getis, A., Morrison, A.C., Gray, K. & Scott, T.W. (2003) Characteristics of the
 695 spatial pattern of the dengue vector, *Aedes aegypti*, in Iquitos, Peru. *Am J*
 696 *Trop Med Hyg*, **69**, 494-505.
- 697 Hancock, P.A. & Godfray, H.C.J. (2012) Modelling the spread of *Wolbachia* in
 698 spatially heterogeneous environments. *J Royal Soc Interface*, **9**, 3045-
 699 3054.
- 700 Hancock, P.A., Linley-White, V., Callahan, A.G., Godfray, H.C.J., Hoffmann, A.A. &
 701 Ritchie, S.A. (2016a) Density-dependent population dynamics in *Aedes*
 702 *aegypti* slow the spread of *wMel Wolbachia*. *J Appl Ecol*, **53**, 785-793.
- 703 Hancock, P.A., White, V.L., Ritchie, S.A., Hoffmann, A.A. & Godfray, H.C.J. (2016b)
 704 Predicting *Wolbachia* invasion dynamics in *Aedes aegypti* populations
 705 using models of density-dependent demographic traits. *BMC Biology*, **14**.
- 706 Hancock, Penelope; Ritchie, Scott A.; Koenraadt, Constantianus J. M.; Scott,
 707 Thomas; HOFFMANN, ARY; Godfray, H. Charles J. (2019): Predicting the
 708 spatial dynamics of *Wolbachia* infections in *Aedes aegypti* arbovirus
 709 vector populations in heterogeneous landscapes, 2019, Journal of Applied
 710 Ecology. figshare. Dataset. <https://doi.org/10.6084/m9.figshare.8117939.v1>
- 711 Hancock, P.A. (2019). pahanc/*Wolbachia*-invasion-dynamics-spatial: *Wolbachia*
 712 invasion dynamics spatial (Version v1.0.0). Zenodo.
 713 <http://doi.org/10.5281/zenodo.2784071>
- 714 Harrington, L.C., Scott, T.W., Lerdthusnee, K., Coleman, R.C., Costero, A., Clark,
 715 G.G., . . . Edman, J.D. (2005) Dispersal of the dengue vector *Aedes aegypti*
 716 within and between rural communities. *Am J Trop Med Hyg*, **72**, 209-220.

- Hoffmann, A.A. (2014) Facilitating *Wolbachia* invasions. *Austral Entomology*, **53**, 125-132.
- Hoffmann, A.A., Montgomery, B.L., Popovici, J., Iturbe-Ormaetxe, I., Johnson, P.H., Muzzi, F., . . . O'Neill, S.L. (2011) Successful establishment of *Wolbachia* in *Aedes* populations to suppress dengue transmission. *Nature*, **476**, 454-457.
- Jeffery, J.A.L., Nguyen Thi, Y., Vu Sinh, N., Le Trung, N., Hoffmann, A.A., Kay, B.H. & Ryan, P.A. (2009) Characterizing the *Aedes aegypti* population in a Vietnamese village in preparation for a *Wolbachia*-based mosquito control strategy to eliminate dengue. *PLoS Negl Trop Dis*, **3**, e552.
- Koenraadt, C.J.M., Aldstadt, J., Kijchalao, U., Sithiprasasna, R., Getis, A., Jones, J.W. & Scott, T.W. (2008) Spatial and temporal patterns in pupal and adult production of the dengue vector *Aedes aegypti* in Kamphaeng Phet, Thailand. *Am J Trop Med Hyg*, **79**, 230-238.
- LaCon, G., Morrison, A.C., Astete, H., Stoddard, S.T., Paz-Soldan, V.A., Elder, J.P., . . . Vazquez-Prokopec, G.M. (2014) Shifting Patterns of *Aedes aegypti* Fine Scale Spatial Clustering in Iquitos, Peru. *PLoS Negl Trop Dis*, **8**.
- Legros, M., Lloyd, A.L., Huang, Y.X. & Gould, F. (2009) Density-dependent intraspecific competition in the larval stage of *Aedes aegypti* (Diptera: Culicidae): Revisiting the current paradigm. *J Med Entomol*, **46**, 409-419.
- Legros, M., Otero, M., Aznar, V.R., Solari, H., Gould, F. & Lloyd, A.L. (2016) Comparison of two detailed models of *Aedes aegypti* population dynamics. *Ecosphere*, **7**.
- Maciel-De-Freitas, R., Codeco, C.T. & Lourenco-De-Oliveira, R. (2007) Daily survival rates and dispersal of *Aedes aegypti* females in Rio de Janeiro, Brazil. *Am J Trop Med Hyg*, **76**, 659-665.
- Mohammed, A. & Chadee, D.D. (2011) Effects of different temperature regimens on the development of *Aedes aegypti* (L.) (Diptera: Culicidae) mosquitoes. *Acta Tropica*, **119**, 38-43.
- Moreira, L.A., Iturbe-Ormaetxe, I., Jeffery, J.A., Lu, G.J., Pyke, A.T., Hedges, L.M., . . . O'Neill, S.L. (2009) A *Wolbachia* symbiont in *Aedes aegypti* limits infection with Dengue, Chikungunya, and *Plasmodium*. *Cell*, **139**, 1268-1278.

- Morrison, A.C., Gray, K., Getis, A., Astete, H., Sihuincha, M., Focks, D., . . . Scott, T.V. (2004) Temporal and geographic patterns of *Aedes aegypti* (Diptera : Culicidae) production in Iquitos, Peru. *J Med Ent*, **41**, 1123-1142.
- Muriu, S.M., Coulson, T., Mbogo, C.M. & Godfray, H.C.J. (2013) Larval density dependence in *Anopheles gambiae* s.s., the major African vector of malaria. *J Anim Ecol*, **82**, 166-174.
- Nasci, R.S. (1986) The size of emerging and host-seeking *Aedes-aegypti* and the relation of size to blood-feeding success in the field. *J Am Mosq Control Assoc*, **2**, 61-62.
- Nunes, M.R.T., Faria, N.R., de Vasconcelos, J.M., Golding, N., Kraemer, M.U.G., de Oliveira, L.F., . . . Vasconcelos, P.F.D. (2015) Emergence and potential for spread of Chikungunya virus in Brazil. *BMC Med*, **13**.
- O'Neill, S.L., Ryan, P.A., Turley, A.P., Wilson, G., Retski, K., Iturbe-Ormaetxe, I., . . . Simmons, C.P. (2018) Scaled deployment of *Wolbachia* to protect the community from *Aedes* transmitted arboviruses. *Gates Open Res*, **2:36**.
- Ritchie, S.A., Montgomery, B.L. & Hoffmann, A.A. (2013a) Novel Estimates of *Aedes aegypti* (Diptera: Culicidae) Population Size and Adult Survival Based on Wolbachia Releases. *J Med Ent*, **50**, 624-631.
- Ritchie, S.A., Pyke, A.T., Hall-Mendelin, S., Day, A., Mores, C.N., Christofferson, R.C., . . . van den Hurk, A.F. (2013b) An Explosive Epidemic of DENV-3 in Cairns, Australia. *Plos One*, **8**.
- Ross, P.A., Endersby, N.M. & Hoffmann, A.A. (2016) Costs of Three Wolbachia Infections on the Survival of *Aedes aegypti* Larvae under Starvation Conditions. *PLoS Negl Trop Dis*, **10**.
- Saez-Llorens, X., Tricou, V., Yu, D., Rivera, L., Tuboi, S., Garbes, P., . . . Wallace, D. (2017) Safety and immunogenicity of one versus two doses of Takeda's tetravalent dengue vaccine in children in Asia and Latin America: interim results from a phase 2, randomised, placebo-controlled study. *Lancet Infect Dis*, **17**, 615-625.
- Schmidt, T.L., Barton, N.H., Rasic, G., Turley, A.P., Montgomery, B.L., Iturbe-Ormaetxe, I., . . . Turelli, M. (2017) Local introduction and heterogeneous spatial spread of dengue-suppressing Wolbachia through an urban population of *Aedes aegypti*. *PLoS Biol*, **15**.

- 783 Schneider, J.R., Morrison, A.C., Astete, H., Scott, T.W. & Wilson, M.L. (2004) Adult
 784 size and distribution of *Aedes aegypti* (Diptera : Culicidae) associated
 785 with larval habitats in Iquitos, Peru. *J Med Ent*, **41**, 634-642.
- 786 Schofield, P. (2002) Spatially explicit models of Turelli-Hoffmann *Wolbachia*
 787 invasive wave fronts. *J Theor Biol*, **215**, 121-131.
- 788 Scott, T.W., Amerasinghe, P.H., Morrison, A.C., Lorenz, L.H., Clark, G.G., Strickman,
 789 D., ... Edman, J.D. (2000a) Longitudinal studies of *Aedes aegypti* (Diptera :
 790 Culicidae) in Thailand and Puerto Rico: Blood feeding frequency. *J Med*
 791 *Ent*, **37**, 89-101.
- 792 Scott, T.W., Morrison, A.C., Lorenz, L.H., Clark, G.G., Strickman, D., Kittayapong, P.,
 793 ... Edman, J.D. (2000b) Longitudinal studies of *Aedes aegypti* (Diptera :
 794 Culicidae) in Thailand and Puerto Rico: Population dynamics. *J Med Ent*,
 795 **37**, 77-88.
- 796 Segoli, M., Hoffmann, A.A., Lloyd, J., Omodei, G.J. & Ritchie, S.A. (2014) The Effect
 797 of Virus-Blocking *Wolbachia* on Male Competitiveness of the Dengue
 798 Vector Mosquito, *Aedes aegypti*. *PLoS Negl Trop Dis*, **8**, E3294-E3294.
- 799 Southwood, T.R.E., Murdie, G., Yasuno, M., Tonn, R.J. & Reader, P.M. (1972)
 800 Studies on the life budget of *Aedes aegypti* in Wat Samphaya, Bangkok,
 801 Thailand. *Bull World Health Org*, **46**, 211-226.
- 802 Turelli, M. (1994) Evolution of incompatibility inducing microbes and their
 803 hosts. *Evolution*, **48**, 1500-1513.
- 804 Turelli, M. & Barton, N.H. (2017) Deploying dengue-suppressing *Wolbachia*:
 805 Robust models predict slow but effective spatial spread in *Aedes aegypti*.
 806 *Theor Popul Biol*, **115**, 45-60.
- 807 Turelli, M. & Hoffmann, A.A. (1991) Rapid spread of an inherited incompatibility
 808 factor in California *Drosophila*. *Nature*, **353**, 440-442.
- 809 Walker, T., Johnson, P.H., Moreira, L.A., Iturbe-Ormaetxe, I., Frentiu, F.D.,
 810 McMeniman, C.J., ... Hoffmann, A.A. (2011) The wMel *Wolbachia* strain
 811 blocks dengue and invades caged *Aedes aegypti* populations. *Nature*, **476**,
 812 450-U101.
- 813 Williams, C.R., Johnson, P.H., Ball, T.S. & Ritchie, S.A. (2013) Productivity and
 814 population density estimates of the dengue vector mosquito *Aedes*
 815 *aegypti* (*Stegomyia aegypti*) in Australia. *Med Vet Ent*, **27**, 313-322.

Wong, J., Stoddard, S.T., Astete, H., Morrison, A.C. & Scott, T.W. (2011) Oviposition Site Selection by the Dengue Vector *Aedes aegypti* and Its Implications for Dengue Control. *PLoS Negl Trop Dis*, 5.

FIGURE AND TABLE LEGENDS

Table 1. Values of model parameters describing mosquito demographic traits and dispersal.

Table 2. Values of model parameters describing spatial structure and the sizes of mosquito releases.

Figure 1. A. Diagram of the metapopulation model showing interactions between mosquito demographic traits, larval density and habitat quality across subpopulations. The coloured triangle represents the age structure of the mosquito subpopulation in a given house, depicting the relatively long larval development stage (light blue), the short pupal development stage (dark blue) and the adult stage (pink). Houses with different habitat quality levels are arranged in a regular grid. The larval development rate and the per-capita female fecundity within a subpopulation depend on both the total larval density in the subpopulation and the house habitat quality. **B.** Model of mosquito-*Wolbachia* dynamics. Double red arrows indicate demographic rates that depend on larval density, solid arrows indicate fixed time lags, dotted arrows indicate mating interactions, and open circle terminators indicate mortality rates. Grey shading and no shading indicate *Wolbachia*-infected and uninfected life stages, respectively. Parameter symbols and values are defined in Table 1.

Figure 2. (A) The distribution of the number of pupae per house. Red bars show observations from the field surveys and black bars show the simulated distribution given by our metapopulation model. The pie chart shows the fitted proportions of houses with high (H), low (L) and empty (E) habitat quality (h_H

h_L and h_E , respectively). **(B)** Boxplots showing the distribution of per-capita female fecundity across subpopulations predicted by our model (black, left), the per-capita fecundity distribution estimated from field-caught mosquitoes (red; see text), and the distribution of the mean development times of the larval cohorts predicted by our model (black, right).

Figure 3. Comparing predicted and observed rates of spatial spread of *Wolbachia*. Black circles show the mean across all simulations of the distance of the 0.5 *Wolbachia* frequency contour from the edge of the release area and vertical black lines connect the 5% and 95% quantiles. The observed mean position of the 0.5 *Wolbachia* frequency contour from the edge of the release area following the *Wolbachia* releases conducted in Cairns is shown for two release areas, Edge Hill (red crosses) and Parramatta Park (blue crosses). Four types of temporal variation in house habitat quality are shown: Static habitat quality (Top row **(A,E,I)**); Slow stochastic temporal variation ($\bar{t}_q = 3$ months; Second row **(B,F,J)**); Fast stochastic temporal variation ($\bar{t}_q = 1$ month; Second row **(C,G,K)**); Seasonal variation (Fourth row **(D,H,L)**). Three values of the fitness cost incurred by *Wolbachia* are shown: No cost ($s=0$; First column **(A-D)**); $s=0.15$ (Second column **(E-H)**); $s=0.3$ (Third column **(I-L)**).

Figure 4. Sensitivity analysis showing the effects of variation in **A.** Daily larval mortality; **B.** Daily pupal and adult mortality, and **C.** Daily dispersal probability on predicted mosquito-*Wolbachia* dynamics. Effects on the *Wolbachia* spread distance from the release zone (red lines), the total mosquito population size (blue lines) and the average per-capita female fecundity across the metapopulation (green lines) are shown. Values of the daily pupal and adult mortality above 0.15 resulted in elimination of the mosquito population.

Table 1. Values of model parameters describing mosquito demographic traits and dispersal.

Symbol	Definition	Value		Source
T_P	Duration of the pupal stage	2 days		(Hancock <i>et al.</i> 2016)
T_G	Minimum time for females to become gravid following emergence	4 days		(Hancock <i>et al.</i> 2016)
T_H	Time between oviposition and hatching of eggs	6 days		(Hancock <i>et al.</i> 2016)
	The proportion of uninfected offspring produced by <i>Wolbachia</i> -infected females	0.01		(Walker <i>et al.</i> 2011)
s_h	The proportion of unviable offspring from an incompatible mating	0.99		(Walker <i>et al.</i> 2011)
s	The reduction in average adult lifespan caused by <i>Wolbachia</i> carriage	varies		
, ,	Parameters of functions describing mean larval development time	<i>Low quality habitats</i>	<i>High quality habitats</i>	This study
		19.6, 2.74, 0.43	1.2, 0.154, 0.68	
, ,	Parameters of functions describing the standard deviation of larval development time	-0.97, 1.0, 0.32	12, 0.24, 0.44	This study
a, b	Parameters of functions describing per-capita female fecundity	15.4, -2.03	28.5, -3.25	This study
min, max, max	Lower limit on per-capita female fecundity, and upper limits on	0.5, 60 days, 40 days		This study

	larval development time means and standard deviations			
L	Daily larval mortality	0.05		(Hancock <i>et al.</i> 2016)
A	Daily pupal and adult mortality	Field-cage	Field release simulations	(Muir & Kay 1998; Walker <i>et al.</i> 2011)
		0.03	0.1	
p_d	Daily probability that adults move to adjacent house	0.56		(Harrington <i>et al.</i> 2005)

Table 2. Values of model parameters describing spatial structure and the sizes of mosquito releases.

Symbol	Definition	Value
x_h	Side length of square grid units (houses)	25 m
x_A	Side length of square spatial region containing the target mosquito vector population	2.5 km
h_H, h_L, h_E	Proportions of houses with high, low and empty habitat quality	0.1, 0.45, 0.45
n_r	Number of releases	15
t_r	Duration of release period	15 weeks
s_r	Number of mosquitoes released per release house per release*	5*
x_r	Size of the release zone*	725 m ²
f_r	Wolbachia frequency in the release zone at the time of the final release*	0.85

*Used for model simulations where the fitness cost $s=0$ and the habitat quality landscape was static.

jpe_13423_f1.pdf

B

Infected:

This article is protected by copyright. All rights reserved.

Static:

No cost

jpe_13423_63.pdf

$s = 0.18$

$s = 0.3$

Stochastic (slow):

Stochastic (fast):

Seasonal:

This article is protected by copyright. All rights reserved.

Time after final release (months)

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Hancock, PA;Ritchie, SA;Koenraadt, CJM;Scott, TW;Hoffmann, AA;Godfray, HCJ

Title:

Predicting the spatial dynamics of Wolbachia infections in Aedes aegypti arbovirus vector populations in heterogeneous landscapes

Date:

2019-07-01

Citation:

Hancock, P. A., Ritchie, S. A., Koenraadt, C. J. M., Scott, T. W., Hoffmann, A. A. & Godfray, H. C. J. (2019). Predicting the spatial dynamics of Wolbachia infections in Aedes aegypti arbovirus vector populations in heterogeneous landscapes. JOURNAL OF APPLIED ECOLOGY, 56 (7), pp.1674-1686. <https://doi.org/10.1111/1365-2664.13423>.

Persistent Link:

<http://hdl.handle.net/11343/285971>