PLEASE READ ALL 'COMMENT' TEXT BEFORE PREPARING YOUR ARTICLE. If you do not see the Comments, select View > Print Layout. Please delete them before submitting (Review > Delete All Comments in Document) so that peer reviewers see a clean copy of the manuscript.

Remember that you are writing for an interdisciplinary audience. Please be sure to discuss interdisciplinary themes, issues, debates, etc. where appropriate. Note that the WIREs are forums for review articles, rather than primary literature describing the results of original research.

If you have any questions, contact your editorial office.

Article Title: Review of tropical cyclones in the Australian region: Climatology, variability, predictability and trends

Article Type:

- ☐ OPINION
 ☐ PRIMER
 ☐ OVERVIEW
☒ ADVANCED REVIEW
 ☐ FOCUS ARTICLE
 ☐ SOFTWARE FOCUS

Authors:

[List each person's full name, **ORCID iD**, affiliation, email address, and any conflicts of interest. Copy rows as necessary for additional authors. Please use an asterisk (*) to indicate the corresponding author.]

Savin S. Chand*

Centre for Informatics and Applied Optimization, Federation University Australia, Mt Helen, Ballarat, Victoria Australia
 Email: s.chand@federation.edu.au
 Conflict of interest: The author has declared no conflicts of interest for this article.

Andrew J. Dowdy

Bureau of Meteorology, Melbourne, Victoria, Australia
 Email: andrew.dowdy@bom.gov.au
 Conflict of interest: The author has declared no conflicts of interest for this article.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the **Version of Record**. Please cite this article as doi: [10.1002/wcc.602](https://doi.org/10.1002/wcc.602)

Hamish A. Ramsay CSIRO, Oceans and Atmospheres, Aspendale, Victoria, Australia Email: Hamish.Ramsay@csiro.au Conflict of interest: The author has declared no conflicts of interest for this article.
Kevin J. E Walsh School of Earth Science, University of Melbourne, Parkville, Australia Email: kevin.walsh@unimelb.edu.au Conflict of interest: The author has declared no conflicts of interest for this article.
Kevin J. Tory Bureau of Meteorology, Melbourne, Victoria, Australia Email: kevin.tory@bom.gov.au Conflict of interest: The author has declared no conflicts of interest for this article.
Scott B. Power Bureau of Meteorology, Melbourne, Victoria, Australia Email: scott.power@bom.gov.au Conflict of interest: The author has declared no conflicts of interest for this article.
Samuel S. Bell Centre for Informatics and Applied Optimization, Federation University Australia, Mt Helen, Ballarat, Victoria Australia Email: ss.bell@federation.edu.au Conflict of interest: The author has declared no conflicts of interest for this article.
Sally L. Lavender University of Southern Queensland, Toowoomba, QLD, Australia Email: sally.lavender@metoffice.gov.uk Conflict of interest: The author has declared no conflicts of interest for this article.
Hua Ye Bureau of Meteorology, Melbourne, Victoria, Australia Email: harvey.ye@bom.gov.au Conflict of interest: The author has declared no conflicts of interest for this article.
Yuri Kuleshov Bureau of Meteorology, Melbourne, Victoria, Australia Email: yuriy.kuleshov@bom.gov.au Conflict of interest: The author has declared no conflicts of interest for this article.

Abstract

Tropical cyclones (TCs) can have severe impacts on Australia. These include extreme rainfall and winds, and coastal hazards such as destructive waves, storm surges, estuarine flooding and coastal erosion. Various aspects of TCs in the Australian region have been documented over the past several decades. In recent years, increasing emphasis has been placed on human-induced climate change effects on TCs in the Australian region and elsewhere around the globe. However, large natural variability and the lack of consistent long-term TC observations have often complicated the detection and attribution of TC trends. Efforts have been made to improve TC records for Australia over the past decades but it is still unclear whether such records are sufficient to provide better understanding of the impacts of natural climate variability and climate change. It is important to note that the damage costs associated with tropical cyclones in Australia have increased in recent decades and will continue to increase due to growing coastal settlement and infrastructure development. Therefore, it is critical that any coastal infrastructure planning and engineering decisions, as well as disaster management decisions, strongly consider future risks from tropical cyclones. A better understanding of tropical cyclones in a changing climate will provide key insights that can help mitigate impacts of tropical cyclones on vulnerable communities. An objective assessment of the Australian TCs at regional scale and its link with climate variability and change using improved and up-to-date data records is more imperative now than before.

Graphical/Visual Abstract and Caption

Caption: Tropical cyclone frequency in the Australian region and the potential influences of monitoring methods and technologies on the best track database since 1900.

1 INTRODUCTION

It is well established that important temporal inhomogeneities exist in tropical cyclone (TC) best track data sets that limit their usefulness for climate analysis.¹⁻³ This issue largely results from the way the best track data are compiled: they are prepared at the end of each TC season using the best available observations at that time. Since the quality of observations has improved over the past few decades, there have inevitably been changes to the way that TC characteristics, especially estimates of TC intensity, are measured and incorporated in the best track data.⁴ Recognizing this issue, Kossin et al.^{5,6} constructed a global satellite-based TC intensity data set that homogenized the record by using a consistent intensity analysis routine applied to a satellite data set of consistent resolution for the entire period. Due to limited availability of relevant satellite observations, only a few decades of homogenized data are available for analysis.

In the Australian region (defined here as the region south of the equator and between 90°E and 160°E, Figure 1), efforts have been made to create more consistent observational data sets for analysis. Harper et al.⁷ employed an objective review process to create a more homogenous record for the northwest Australian region for the period 1969–2001. In recent years, a comprehensively

compiled best track data set for the entire Southern Hemisphere basin was prepared⁸ with regular revisions.^{2,9} Efforts have also been made to enhance the quality of the TC archive in the South Pacific region.^{10,11} The Australian Bureau of Meteorology also provides an Australian region TC database covering the region south of the equator between 90°E and 160°E¹². We note that the intensity estimates of TCs, as represented in these best track data sets, has steadily improved since the 1980s with increasing satellite observations and improved analysis techniques.

FIGURE 1 The Australian region and its three basins. Also shown are TC tracks over the period 1981/82 to 2016/17 seasons. Colours indicate different TC intensities for different categories of cyclones based on the Australian classification scheme defined in Table 2. TC track data are those from the Bureau of Meteorology database¹².

Over the past decades, several dedicated studies have examined various aspects of Australian region TCs including, but not limited to, climatological characteristics^{13,14}, predictability¹⁵⁻¹⁸, impact of natural climate variability such as the El Niño-Southern Oscillation and the Madden-Julian Oscillation¹⁹⁻²⁵, and climate change trends.^{8,9,26-28} However, a review that synthesises these studies with current perspectives on Australian TCs is lacking. The main emphases of the present work is to first provide a comprehensive review of Australian region TCs and then, where possible, use improved and up-to-date data records to consolidate the review with new results. We also examine the ACE index representing accumulated cyclone energy²⁹, which is an additional useful measure for examining potential climatological trends. In the next section, TC data records and homogeneity issues for Australia, as well as work being undertaken to improve the data records, are

discussed. Climatological characteristics of Australian TCs are presented in Section 3. In Section 4, we examine the impacts of natural climate variability on Australian TCs at various timescales, and in Section 5 we provide evidence of long-term trends that may be attributable to climate change. A summary is provided in Section 6.

2 AUSTRALIAN TC DATA SOURCES AND THEIR LIMITATIONS

The first definitive record of TCs over Australia, as well as in the broader South Pacific and Indian Oceans, was created for the period 1779 – 1922 by Visser.³⁰ Subsequent studies led to the compilation and reappraisal of TC data over Australia³¹, as well as in the southwest Pacific basin spanning eastern Australia^{32,33}, for the period up to the 1970s.

Holland³⁴ was the first to assess the quality and consistency of the Australian TC data for the period 1909–1979, and found that TC data prior to the 1960s were largely inconsistent and incomplete due to lack of surface observations and monitoring such as regular aircraft reconnaissance. Satellite sensing commenced in the early 1960s, and became technologically more advanced thereafter³⁵, leading to vast improvements in the way TCs were tracked and monitored. However, lack of objective methodologies for estimating TC intensities from the satellite information often placed caveats on TC intensity data prior to the 1970s.

Development of the Dvorak technique in the early 1970s,^{36,37} and the subsequent refinement of the technique,^{35,38-40} have led to significant improvements in TC intensity estimates over the last few decades (Figure 2). A review of historical TC intensities over Australia^{7,41} had cautioned against the use of TC intensity data prior to the 1980s due to instrumental biases and the evolving Dvorak technique during the 1970s. Note, unlike Atlantic and North Pacific basins, there is no *in-situ* data included in the intensity estimates for the Australian basin. Significant improvement in data quality and estimation of TC intensity occurred over the broader Southern Ocean with the availability of high-resolution satellite imagery since the mid-1980s and after the launch of the MeteoSat-5 satellite in 1998⁶. A recent global investigation⁴ recommended that TC intensity data from the 1990s are observationally more consistent and homogeneous, and therefore should form the basis of climate change and trend analysis. However, the time series since 1990 may not be sufficient to draw robust conclusions on the impact of climate variability and change.

FIGURE 2 Australian region TC frequency indicating the potential influences of monitoring methods and technologies on the best track database since 1900. (Following Harper et al.⁷)

The Australian region best track TC data set are produced by three of the six designated World Meteorological Organization Tropical Cyclone Warning Centres (WMO-TCWCs) in Brisbane, Darwin and Perth, and is comprehensively compiled and maintained by the Australian Bureau of Meteorology.¹² Other sources of the Australian best track data include the International Best Track Archive for Climate Stewardship (IBTrACS⁴²) and the Southern Hemisphere TC archive^{8,9}; both these databases use the same WMO-TCWCs in Brisbane, Darwin and Perth to source the Australian region TCs. An independent source of the TC best track data set is from the U.S Navy's Joint Typhoon Warning Center (JTWC⁴³). A study by Schreck et al.⁴⁴ compared the TC climatologies between the two databases (i.e., WMO-IBTrACS and JTWC), and found substantial discrepancies in some basic metrics such as annual number of TCs in Australia and elsewhere, potentially related to inherent differences in the TC "best-tracking" procedures where different agencies are likely to employ different operational methods, for example, to track circulations and estimate intensities throughout a storm's life cycle.⁴²

3 OBSERVED CHARACTERISTICS OF AUSTRALIAN TCS

Dare and Davidson¹⁴ provides a comprehensive climatology of the Australian region TCs for the seasons 1963/64-2002/03^a. Here we have re-examined key climatological metrics presented in that study but for the thirty-year period 1960/61-1989/90 as a standard reference and then reproduced new climatologies for the period 1980/81-2009/10 after taking into consideration the relevance of such time periods in a changing climate.⁴⁵ These metrics are shown for the entire Australian region and its three basins^b in Table 1.

TABLE 1 Climatological mean characteristics of tropical cyclones in the Australian region and its three basins for the two thirty-year climatological normals: 1960-1989 and 1980-2009. Statistically significant differences of TC characteristics between the two climatological normals at 95% and 90% levels are indicated with asterisk and plus signs respectively. Severe TCs are those that reach or exceed Category 3 (Australian Region classification).

Climatological mean field	Australian region (90°-160°E)		Western basin (90°-125°E)		Central basin (125°-142.5°E)		Eastern basin (142.5°-160°E)	
	1960-1989	1980-2009	1960-1989	1980-2009	1960-1989	1980-2009	1960-1989	1980-2009
Number of TCs per season	13.4 ⁺	10.9 ⁺	6.0 ⁺	5.2 ⁺	3.1 ⁺	2.8 ⁺	4.3 ⁺	2.9 ⁺
Number of landfalling TCs per season [% total TCs]	5.3 ⁺ [39.6]	4.4 ⁺ [40.4]	1.4 [23.0]	1.3 [25.0]	2.6 [83.9]	2.1 [75.0]	1.3 [30.2]	1.0 [34.5]
Number of severe TCs per season [% total TCs]	4.7 [34.8 ⁺]	5.4 [47.4 ⁺]	2.4 [40.0]	2.6 [48.1]	1.1 [34.0]	1.4 [43.5]	1.2 [28.5 ⁺]	1.5 [50.0 ⁺]
Number of severe landfalling TCs per season [% total landfall]	2.3 [42.3]	2.4 [53.8]	0.9 [65.1]	0.8 [62.0]	0.8 [29.1]	1.0 [46.6]	0.6 [46.2]	0.6 [56.4]
TC lifetime (days)	7.0	7.2	6.3 ⁺	7.3 ⁺	5.5	6.4	5.5 ⁺	7.3 ⁺
TC days per season	91.3	78.5	37.8	37.9	18.2	18.0	24.9	20.3
Mean latitude of genesis position	12.5	12.0	11.9	11.3	12.0	11.6	13.9	13.6
Mean latitude of Lowest Central Pressure, LCP	16.7 ⁺	15.6 ⁺	16.6 ⁺	15.8 ⁺	14.1	13.7	18.3 ⁺	16.3 ⁺
Mean latitude of dissipation	22.2 ⁺	20.3 ⁺	22.4 ⁺	20.6 ⁺	20.8	20.7	22.7 ⁺	19.3 ⁺

3.1 TC numbers, lifetime and days

^a Note that TCs in the Australian region are spread over the two calendar years. Therefore, a TC season is defined here to be the period from the appearance of the first TC from 1 July of one year to the last day of the last TC up to 30 June the following year. Thus, a season is labelled using both calendar years; for example, the season covering the period July 1963 to June 1964 is referred to as 1963/64 season. For convenience, each season in the figures is labelled using the first of the two relevant calendar years; for example, the season 1963/64 may also be referred to as 1963.

^b The Australian Bureau of Meteorology database includes all TCs that formed between 90°-160°E, as well as those that formed outside this region but eventually tracked in. To be consistent with earlier studies (e.g., Dare and Davidson¹⁴), the western basin is defined to include only those TCs that formed west of 125°E, the central basin include TCs that formed between 125°-142.5°E and the eastern basin consists of TCs that formed east of 142.5°E (Figure 1); TC formation here refers to the first track point in the database.

In the present climate (i.e., 1980/81-2009/10), about 10.9 TCs occur in the entire Australian region annually (as opposed to about 13.4 during the period 1960/61-1989/90). An average of about 5.2, 2.8 and 2.9 TCs form in the western, central and eastern basins each year, respectively. About 40% of these TCs make landfall on the Australian coasts annually, with the central basin recording the highest proportion (~75% of those formed in that basin) due to the landlocked nature of this basin and genesis locations being close to the coastline. We note a significant decline in the climatological mean frequency of TCs in the entire Australian region, as well as in the three basins, between the two climatological periods.

While most TCs in the Australian region have an average lifetime of about 7.2 days, those forming in the central basin have a shorter mean lifetime (~6.4 days) as they make landfall and dissipate early due to their close proximity to land. We note that the mean lifetime of TCs has increased in general for the Australian region, as well as for each of the three basins, with this change being statistically significant in the western and eastern basins over the two climatological periods shown. While this does not appear to be associated with a longer track into regions farther poleward, changes in observational practices between the two periods may play some role, as improvements in techniques can lead to early detection of TCs, hence the longer lifetime. Some recent studies have suggested a potential latitudinal shift in TC locations due to the influence of tropical expansion driven by climate change^{46,47}. However, the summary presented in Table 1 lists three different mean latitude measures, including genesis position, lowest central pressure and dissipation, all of which indicate an equatorward shift over the period examined. This is true for all three measures in the Australian region as a whole, as well as individually for each of its three basins, although we note that the changes are only statistically significant in some cases.

Moreover, the Australian region is exposed to about 78.5 TC days each season with the western basin recording the highest (~37.9 TC days) due to the high frequency of occurrence. The overall TC days have decreased over the two climatological periods for the entire Australian region and in the eastern basin, mainly owing to the decrease in TC number, though this decrease is not statistically significant.

3.2 TC intensity

In the entire Australian region, about 5.4 TCs reach or exceed category 3 (severe) each year over the period 1980/81-2009/10^c (Table 2), most of which occur in the western basin. While the climatological mean number of severe TCs has increased only slightly over the two climatological

^c The Bureau of Meteorology (2018) dataset contains TC intensity estimates based on the Dvorak (1984) method, and are derived using the 10-minute sustained wind and central pressure relationship (e.g., Atkinson and Holliday 1977). See Dare and Davidson (2004) for a comparison between the Australian and Saffir-Simpson categories. The main difference is that the lower intensity threshold for Saffir-Simpson category 1 is approximately the same as that for Australian category 3 (Table 2). Note that TC classification definitions vary slightly from that available at the Bureau of Meteorology website (<http://www.bom.gov.au/cyclone/about/>), which uses wind gusts as opposed to the average wind speed.

periods analysed here (i.e., 4.7 for 1960/61-1989/90 and 5.4 for 1980/81-2009/10), the proportion of severe TCs (i.e., the number of severe TCs with respect to the total frequency) has increased substantially from 34.8% to 47.4%; this marked increase is noticed mainly in the eastern basin. The number of TCs making landfall each year on Australian coasts at the severe strength category is around 2.4 per year; this accounts for about 44% of the total severe Australian TCs and 54% of the total landfalling TCs.

TABLE 2 Tropical cyclone categories and corresponding values of approximate central pressures and 10-minute averaged maximum wind speeds, modified for the Australian region from the Saffir–Simpson scale following Dare and Davidson.¹⁴

Category	Average Maximum Wind Speed (km h ⁻¹)	Central Pressure (h Pa)
1	63 – 90	> 985
2	90 – 125	985 – 970
3	125 – 165	970 – 955
4	165 – 225	955 – 930
5	> 225	< 930

3.3 Seasonality

TCs in the Australian region occur mainly between the months of November to April, which defines a typical cyclone season for the region, with the peak activity (~61% of annual numbers based on 1980/81-2009/10 climatology) occurring during January-March (Figure 3c). However, there are also cases (~7%) when TCs can occasionally develop outside this typical season. These cases are often found to be associated with early onset or late decay states of the El Niño – Southern Oscillation phenomenon, which is a major driver of TC variability in the Australian region at interannual time scale (see Section 4 for discussion). It is important to note that the overall decline in Australian TC numbers between the two climatological periods has occurred primarily during the peak season.

FIGURE 3 a) Spatial locations of TC origins (blue), location of achievement of Lowest Central Pressure (orange) and the termination locations (grey), shown for systems declared as TCs in the Australian region for at least some of their lifetime. b) Locations of TCs that achieved severe (red) and non-severe (black) status upon landfall. c) Mean seasonal distribution of TC numbers for the period 1960-1989 (grey) and 1980-2010 (black).

3.4 Spatial locations and tracks

TCs in the Australian region originate mainly between 5° S and 15°S with the mean formation near 12°S (Table 1, see also Figure 3a). Here, TCs can often form very close to the coastline, particularly in the central basin, and occasionally the point of origin can be traced back to over land.¹³ This occasional formation over land is a characteristic of Australian region TCs. A very moist monsoon environment can extend far inland, providing favourable conditions for tropical lows to intensify. Saturated land-surfaces can provide sufficient surface moisture fluxes to support lower category TC formation provided the circulation is embedded in the moist monsoonal environment, (i.e., protected from drier continental air).⁴⁸ Therefore, it is not surprising to find many TCs achieve maximum intensity near land, and only a relatively small proportion exceed Category 3 intensity upon landfall (Figure 3b). In contrast, TCs in the western and eastern basins can form far from land (up to 80° and 170°E respectively) which typically provides greater potential for intensification over the open ocean prior to landfall. As a result, a larger proportion of TCs in these basins reach Category 3 or more upon landfall.

Overall, TCs tend to reach their maximum strength between 10° S and 20°S, and even farther poleward in some cases, with the mean latitude of the lowest central pressure around 15.6°S based on the 1980/81-2009/10 climatology (e.g., Table 1 and Figure 3a). In the TC database, TC tracks often extend as far as 40°S before dissipating. However, it is important to emphasise that these data do not consistently discriminate TCs from non-TC systems as they drift poleward. It is likely that many of these storms have undergone extra-tropical transition as they move poleward or they have merged with other synoptic systems and have lost their TC characteristics. About 20% of TCs in the South Indian Ocean and the Australian region undergo extra-tropical transitions.⁸⁷

Lavender and Dowdy²⁵ produced a map showing TC track direction climatology throughout the Australian region. In contrast to previous studies that have focused on the mean direction of TC movement, the full spectrum of track directions was analysed in their study (Figure 4). This work

identified a natural split at 135°E, based on track motion. The track directions are predominantly westward between about 10°S and 20°S in the western region, while in the eastern region the distributions generally show a somewhat higher degree of variability in direction. There is an almost equal split in the direction of movement in some grid boxes, such as over the Gulf of Carpentaria (i.e., at 140°E, 10°S), noting that this provides an example of where the track information can be almost completely lost when considering the mean direction.

FIGURE 4. Polar plots for each 10°×10° boxes for the Australian region, showing the relative proportion of translational TC tracks in eight different directions (length of fans). These directional distributions are based on 6-hourly TC data. The distribution of TC translation speed is also shown for each direction, from slow (blue) to fast (red), as shown in the legend. The numbers listed for each 10°×10° box represent the number of 6 hr TC occurrences at each location during the study period, with the colour of the numbers showing regions where TCs are intensifying (red) or dissipating (blue) on average. The width of the fan also indicates the relative number of TCs in that grid box compared to the entire region. The black and white arrows represent the mean vector direction and magnitude of the TC movement. (Source: Lavender & Dowdy 2018).

Intensification of TCs in the region typically occurs in the 10°S latitude band (as shown by the red numbers in Figure 4 based on a mean decrease in central pressure), with systems tending to decay (blue numbers, Figure 4) at latitudes further to the south. Recurvature of tracks is evident from the shift in direction between 20°S and 30°S, with the red colours in the ‘fans’ showing that the translational speed tends to increase after tracks make extratropical transition south of about 30°S. Although tracks that form in the Gulf of Carpentaria are unique in some respects, including due to a large proportion making landfall, the distributions presented indicate that the direction that these tracks take in the 10°S band (i.e., prior to landfall) is more similar to those to the east than west of 135°E. At 10°S, over 40% of TCs in the Gulf of Carpentaria move in an eastward direction to the east of 135°E and less than 15% in the grid boxes to the west of 135°E.

4 INTERANNUAL VARIABILITY AND SEASONAL PREDICTABILITY

Up to this point, seasonal and climatological mean values of parameters associated with TCs in the Australian region and its three basins have been considered. In this section, we review the interannual variability and its implications for seasonal predictability.

4.1 ENSO-TC frequency relationship

The El Niño – Southern Oscillation (ENSO) phenomenon modulates TC activity in the Australian region on interannual timescales. Typically, a below average seasonal TC count occurs during El Niño events, whereas La Niña events characteristically bring above-average TC counts to the region. This statistical relationship was first uncovered by Nicholls¹⁵ using pressure anomalies over Darwin, and was developed further in the mid-1980s using sea-surface temperature (SST) data in the tropical Pacific and around northern Australia. Many other studies have since refined the ENSO-TC frequency relationship in the Australian TC region and its sub-basins over the past few decades.^{2,8,16,19-21,23,24,26,49-57} The frequency modulation by ENSO is in part due to a geographic shift in the locations where TCs typically form. For the Eastern Australian region, this migration results in a tendency for fewer storms to form between the Queensland Coast and about 160°E during El Niño years and more storms to form during La Niña years^{33,49}. For the Western Australian region, enhanced TC activity tends to occur off the northwest coast of Australia during La Niña years, as well as in the eastern Indian Ocean around 85°E-105°E,⁵⁵ as depicted in Figure 5.

FIGURE 5. a) Year-to-year variability of the seasonal number of Australian TCs (blue) and severe TCs (red), and average number of TCs per decade in $1^\circ \times 1^\circ$ grid boxes for b) all-year-climatology over the Australian region, as well as for c) El Niño, d) neutral and e) La Niña events in the Southern Hemisphere over the period 1981/82 to 2016/17 cyclone seasons. IBTrACS TC data were used to construct these figures, and the ENSO classification is based on Niño 3.4 SST anomalies $> 0.5^\circ\text{C}$ for El Niño and $< -0.5^\circ\text{C}$ for La Niña using September to February averages. Asterisks in (a) denote El Niño years while plus signs denote La Niña years; unmarked years are ENSO neutral.

4.2 Implications for seasonal predictions

The strong link between ENSO and TC frequency in the Australian region has formed the basis of several seasonal TC predictions schemes over the past few decades.^{16-18,20,26,58-60} The annual cycle of ENSO is particularly amenable for seasonal predictions in the region because the phase and potential magnitude of an ENSO event is typically evident by Austral spring, which precedes the official start of the TC season on November 1. Further, ENSO events have a strong modulating effect on large-scale environmental parameters that are themselves related to TC formation (e.g., changes to low-level

vorticity associated with the monsoon trough and mid-level relative humidity),⁶¹ and anomalies of these parameters tend to persist well into the TC season. Statistical models of seasonal TC frequency have included using simple linear regression approaches²⁶ and its variants such as Bayesian regression¹⁸ and Poisson regression^{58,59}. More recently, machine learning models have been tested in an effort to improve upon traditional linear regression schemes.^{62,63} Common predictors utilised across models include the Niño 3.4 SST, Niño 4 SST and the SOI. ENSO has also been shown to relate to the frequency of landfalling systems in the Australian region. For instance, Callaghan and Power²⁷ identified a link between pre-season SOI and the number of severe TCs making landfall over north-eastern Australia. They also noted that years in which multiple landfalls of severe TCs occurred since the SOI became available were all La Niña years.

However, despite the traditionally strong statistical link between ENSO and Australian seasonal TC frequency, a weakening of this linear relationship since the late 1990s²⁸ has made operational seasonal prediction in the region increasingly challenging. For instance, the Australian region experienced a near average TC count during the strong La Niña event of 2010/11, despite predictions of above-average seasonal activity based on expectation from historical data. While the physical mechanisms governing the non-stationarity of the ENSO-TC count relationship are currently unclear, substantial interdecadal variability in ENSO and its impact on Australian climate can arise, often from natural internally-generated variability alone⁶⁴.

4.3 Role of non-traditional ENSO regimes and the Indian Ocean

Expanding on the well-established relationship between canonical ENSO and the spatiotemporal variability of Australian TCs, Chand et al.²⁴ demonstrated that annual TC counts depend also on weak/marginal phases of ENSO (defined as 'positive-neutral' and 'negative-neutral' regimes, which traditionally fell into the ENSO-neutral category). Ramsay et al.,⁶⁵ focusing only on the Coral Sea region, showed that seasonal predictability there could be optimised using a combination of ENSO-related SST indices, including the El Niño Modoki Index (EMI) and the Trans-Niño Index (TNI), rather than just a single SST region like Niño 3.4 or Niño 4 (as are commonly used in operational seasonal prediction schemes). The EMI has also been shown to be related to TC frequency in the South Pacific region (including the eastern Australian sub-basin).⁶⁶

The Indian Ocean offers a source of potential predictability that remains relatively unexplored in the context of Australian seasonal TC activity. Liu and Chan⁶⁰ found that a negative Indian Ocean Dipole (IOD) led to increased likelihood of TCs in the eastern Australian sub-basin. Similarly, Wijnands et al.⁶³ showed that the Indian Ocean Dipole Mode Index (DMI) was the most commonly occurring predictor of the best-performing models in the region. In contrast, Werner et al.⁶⁷ found that the IOD added no additional predictive value over using traditional ENSO indices (i.e., Niño 4 SST). Finally, Ramsay et al.⁶⁸ identified a southwest-to-northeast dipole pattern of SST anomalies in the Indian Ocean and showed that a simple index based on this dipole resulted in

improved prediction of seasonal TC counts for the region compared with models that incorporated only ENSO-based predictors.

5 HISTORICAL TRENDS IN AUSTRALIAN TCS AND GLOBAL WARMING

5.1 General perspective

A number of studies have emerged in the past decades debating the link between anthropogenic global warming and historical tropical cyclone trends. In particular, much of the increased attention to this scientific debate has been sparked by the two observational studies of Webster et al.⁶⁹ and Emanuel⁷⁰ that first reported increasing trends in intense tropical cyclone activity since the 1970s. Webster et al.⁶⁹ ascribed a large and significant increase in the global frequency of Category 4 and 5 hurricanes over the period 1970-2004 to increasing sea surface temperatures in various ocean basins globally. Yet when Klotzbach and Landsea⁴ revisited that study using ten years of additional tropical cyclone data, in contrast they found a small, insignificant downward trend in the frequency of Category 4 and 5 hurricanes, particularly between 1990 and 2014. They concluded that the trends in Webster et al.⁶⁹ were artificial and due to observational challenges during the 1970s and 1980s.

On the other hand, studies that developed and utilised globally homogenised satellite-derived tropical cyclone intensity estimates to mitigate observational challenges^{6,71} have affirmed the increasing trends in intense cyclone activity for different basins. However, these studies raised concerns on whether such trends detected in a relatively short period of homogenised observational records can exceed the variability expected through natural causes²⁷, mainly from those occurring at decadal and multi-decadal time scales. Nonetheless, it is important to realise here that inconsistent evidence of historical trends in tropical cyclone intensity between studies arising from the use of different datasets or different length of data does not entail the absence of anthropogenic influences on tropical cyclone activity. Conflating “absence of evidence” with “evidence of absence” is not only inappropriate but also scientifically incorrect.⁷² There is no doubt that human activities have significantly contributed to global climate change, particularly after the mid-twentieth century when the earth has warmed rapidly, and substantial evidence now exists that large-scale environmental conditions which support tropical cyclone activity have changed significantly as a result of this anthropogenic global warming.⁷³⁻⁷⁶ In the next section, we discuss how such changes may have impacted the Australian region TCs.

5.2 Trends in Australian TC activity

Kuleshov et al.^{8,9} used their Southern Hemisphere TC data archive to examine trends in the total annual number of TCs, as well as of those that reached the severe intensity category, for Australia and the broader Southern Hemisphere regions. However, they found no apparent trends in both the total number of TCs and the number of severe TCs for the Australian region over the period 1981/82

to 2006/07, although there was some indication of a weakly positive trend in the proportion of the most intense TCs (i.e., those that reached 950-hPa or lower) in the western basin. They suggested that these lack of trends could have been affected by the shift in ENSO towards more El Niño dominance after the late-1970s.⁷⁷ Indeed, after removing the effects of ENSO, Dowdy²⁸ found a significant decreasing trend in the total annual number of TCs in the Australian region over the period 1981/82 to 2012/13. This highlights the need for considering effects of natural climate variability when examining potential trends in regional TC activity.

We have repeated the analysis of Dowdy²⁸ with an extended period of data covering the seasons 1981/82 to 2017/18 in order to examine trends in several other characteristics of Australian TCs in addition to the total annual numbers (Figure 6). Consistent with earlier studies, there are statistically significant decreasing trends in both the frequency of seasonal TCs ($\sim -0.115 \text{ year}^{-1}$) and the frequency of severe TCs ($\sim -0.112 \text{ year}^{-1}$) over Australia, with a confidence level above 95% (i.e., shown with solid trend lines in Figure 6). We note that these decreasing trends are more dominant in the western basin, and to a lesser extent in the eastern basin, while there are no apparent trends in the central Australian basin (Figure 6a). We also note that the decreasing trends in the Australian region are mainly owing to a significant decrease in the peak season (i.e., January-March) TCs over the western basin, and to some extent by a decrease in the late season (April-June) TCs over the eastern basin. There were no statistically significant trends for the early season (October-December) TC developments in all three Australian basins (Figures 6b-d).

We have also examined trends in the frequency of landfalling TCs over Australia for the period 1980/81 to 2017/18 seasons (Figure 6e). While only a slight decreasing trend in the total frequency is noted, the frequency of severe landfalling TCs has significantly decreased for the period under consideration, particularly in the western basin. However, we emphasise that the number of landfalling TCs each year is very low and care must be exercised when interpreting these trends as they are likely to have been affected by large decadal and multi-decadal variability since the mid-19th century.²⁷

A useful index for measuring basin-wide TC activity is the Accumulated Cyclone Energy (ACE) index, which combines seasonal TC numbers, lifetime and intensity into a single index.⁷⁸⁻⁸⁰ It is essentially a proxy for the integrated kinetic energy of multiple TCs throughout their lifetime. An instantaneous kinetic energy proxy, defined by the square of the estimated storm maximum wind speed, is calculated at six hourly intervals, and summed throughout the lifetime of all TCs in a specified region over a specified time period.²⁹ Since ACE is a function of the square of estimated wind speeds, errors in wind speed estimates are likely to introduce a positive bias to the results. This shouldn't affect trend results unless systematic errors have been introduced over time. Overall, ACE has significantly decreased in the Australian region and its three basins over the period 1980/81 to 2017/18 (Figure 6f). This decrease is primarily related to the decrease in the number of TCs in the Australian region. Indeed, the average kinetic energy (proxy) of TCs has not changed significantly

over the period under consideration (see the “modified-ACE” in Fig. 6f, which is ACE divided by the total number of time periods for all TCs used in the ACE calculation).

5.3 Paleotempestology of Australian TCs

While not the main focus of this work, we comment briefly on a paper by Haig et al.⁸¹ that specifically documented changes in the frequency of Australian TCs in the context of “paleotempestology”, the study of past TC activity by means of geological proxies and historical documentation. In order to circumvent short temporal resolution of TC records over Australia, they constructed a proxy of TCs – derived from carbonate layers of actively growing stalagmite records collected from regions of Queensland and Western Australia – over the last 1500 years. This proxy, referred to as a tropical cyclone activity index, described the severity of a TC season in terms of numbers, intensity, size and longevity of a storm, and was calibrated against the modern-day TC records to allow for a direct comparison of changes in TC activity between the past and present climates. They showed that the present low levels of TC activity on the northeast coasts of Australia, particularly those after the 1960s, are more pronounced than the past 550 to 1,500 years. Their work suggests that climate change cannot be ruled out as a causative factor responsible for the decrease in TC activity over Australia in the past decades, but whether the decrease will be sustained in future is not clear.

d) Late Seasonal Frequency

e) Landfall Frequency

f) Accumulated Cyclone Energy

FIGURE 6 Time series of various TC characteristics for the Australian region and its three basins over the period 1981/82 to 2016/17, and their respective linear trends. Solid (dashed) trend lines indicate statistical significance at the 95% (90%) levels; dotted trend lines are not statistically significant.

6 CONCLUSIONS

A large part of northern Australia and many thousands of kilometres of its coastal oceans are frequently exposed to threats from TC events such as storm surges, coastal flooding and wind damage. With increasing coastal infrastructure development and population growth, and growing concerns from anthropogenic climate change, these TC-induced threats are likely to be exacerbated further.⁸² The influence of anthropogenic climate change on TCs in Australia, and elsewhere around the globe, has been the focus of a growing body of literature over the past several decades. However, the ability to attribute any trend to climate change has been complicated by two main factors: the length and quality of the TC historical datasets and the presence of large interannual and decadal variability in each TC basin, which makes it difficult to separate a trend ‘signal’ from the natural variability ‘noise’. In this review, we have examined these two limitations in the context of Australian region TCs.

Key points and recommendations for future studies are summarised below.

- **Australian TC data sources:** The Australian region best track TC data set is produced by three of the six designated World Meteorological Organization Tropical Cyclone Warning Centres (WMO-TCWCs) in Brisbane, Darwin and Perth, and is comprehensively compiled and maintained by the Australian Bureau of Meteorology.¹² Other sources of the Australian best track data include the International Best Track Archive for Climate Stewardship (IBTrACS⁴²) and the Southern Hemisphere TC archive.^{8,9} An independent source of the TC best track data set is from the U.S Navy's Joint Typhoon Warning Center (JTWC⁴³). Note these data sources can have substantial discrepancies in some basic metrics of TCs.
- **Quality of observed Australian TC records:** Australian TC data for the period prior to about 1960 were largely inconsistent and incomplete.³⁴ Regular monitoring such as satellite sensing commenced after the early 1960s, with more confidence in the data from the 1980s onwards^{7,41}. A recent global investigation by Klotzbach and Landsea⁴ recommended that TC intensity data from the 1990s are observationally more consistent and homogeneous, and therefore should form the basis of climate change and trend analysis. However, the shorter time series since 1990 reduces our ability to draw robust conclusions on the impact of climate variability and change on TC intensity.
- **Climatological characteristics of Australian TCs:** Key climatological metrics of Australian TCs were examined for two thirty-year periods: a reference period (1960/61-1989/90), and a changing climate period (1980/81-2009/10). Overall, a considerable decline in the mean seasonal number of TCs between these two periods was identified for the entire Australian region (for example, about 10.9 TCs for the period 1980/81-2009/10 annually as opposed to about 13.4 during the period 1960/61-1989/90). The mean number of landfalling TCs also declined over these two periods. However, a considerable increase is noted in the proportions of severe TCs (~12%) but care needs to be taken when interpreting this increase as TC intensity data prior to the 1980s are inconsistent.
- **Interannual variability:** The interannual variability of TC frequency in the Australian region is largely related to the El Niño – Southern Oscillation (ENSO) phenomenon. Below average seasonal TC numbers typically occur during El Niño events, whereas La Niña events characteristically bring above-average TC numbers to the region. This statistical relationship has been used to develop seasonal prediction models for Australian TCs,^{18,58,60} and is the basis of operational seasonal forecasts for TC activity issued by the Bureau of Meteorology each year. However, the weakening of the linear relationship between ENSO and Australian TCs since the late 1990s²⁸ has made operational seasonal prediction in the region increasingly challenging, thus highlighting the need for additional predictors such as the Indian Ocean Dipole index to improve skill of seasonal prediction models⁶⁸.

- **TC trends:** Trends in annual Australian TC numbers were initially inconclusive.^{8,9} However, after removing the effects of ENSO variability²⁸ a significant decreasing trend in the total annual number of TCs in the Australian region over the period 1981/82 to 2012/13 was identified. Using extended period of data covering the seasons 1981/82 to 2017/18, we confirmed the decreasing trend (~ -0.115 TC year⁻¹) at an even greater statistical significance over Australia. This decreasing trend is more dominant in the western basin, due mainly to a significant decrease in the peak season (i.e., January-March) TCs as opposed to early or late season TCs. A slight decrease in landfalling TCs over Australian coasts is also noted. The overall TC activity, as measured using the ACE index, is found to have decreased significantly for the Australian region and this decrease is mainly a result of a substantial decline in TC numbers. When the effects of TC numbers and lifetime are removed so that ACE represents the contribution only due to changes in TC intensity, there are no indications of increasing strength of individual TCs over time.

In closing, it is unclear at this stage what exact mechanisms are responsible for the observed trends in TC activity over Australia. However, it is plausible that Indo-Pacific variability such as the Interdecadal Pacific Oscillation,⁸³ interdecadal variability in ENSO activity more broadly,^{27,64} weakening of the ENSO-IOD coupling arising from decadal modulation^{84,85} or climate change^{27,86} potentially have some role to play. Removing the effects of ENSO alone results in an even stronger change in TC activity over the period under consideration, particularly a decline in TC numbers. A lack of consistent long-term TC data makes it difficult to quantify Australian region changes due to decadal or multi-decadal variability, and human-induced climate change. However, as more data become available, future studies should begin to be able to make the distinction between influences from natural sources (including decadal and multi-decadal variability and other modes such as ENSO, IOD and MJO) and climate change. Additional modelling and reanalysis-based studies could also help provide guidance towards understanding the causes of long-term historical trends.

FUNDING INFORMATION

This work is supported through funding from the Earth Systems and Climate Change Hub of the Australian Government's National Environmental Science Program (NESP).

CONFLICT OF INTERESTS

The authors have declared no conflicts of interest for this article.

REFERENCES

1. Landsea CW, Harper BA, Hoarau K, Knaff JA. Can we detect trends in extreme tropical cyclones? *Science* 2006, 313:452-454.
2. Dowdy A, Kuleshov Y. An analysis of tropical cyclone occurrence in the Southern Hemisphere derived from a new satellite-era data set. *Int J Remote Sensing* 2012, 33:7382-7397.
3. Landsea CW, Franklin JL. Atlantic hurricane database uncertainty and presentation of a new database format. *Mon Weather Rev* 2013, 141:3576-3592.
4. Klotzbach PJ, Landsea CW. Extremely Intense Hurricanes: Revisiting Webster et al. (2005) after 10 Years. *J Clim* 2015, 28, 7621-7629, doi: 10.1175/JCLI-D-15-0188.1.
5. Kossin JP, Knapp KR, Vimont DJ, et al. A globally consistent reanalysis of hurricane variability and trends. *Geophys Res Lett* 2007, 34:L04815, doi:10.1029/2006GL028836.
6. Kossin JP, Olander TL, Knapp KR. Trend analysis with a new global record of tropical cyclone intensity. *J Clim* 2013, 26:9960–9976, doi:10.1175/JCLI-D-13-00262.1.
7. Harper BA, Stroud SA, McCormack M, West S. A review of historical tropical cyclone intensity in northwestern Australia and implications for climate change trend analysis. *Aust Meteor Mag* 2008, 57:121–141.
8. Kuleshov Y, Qi L, Fawcett R, Jones D. On tropical cyclone activity in the Southern Hemisphere: Trends and the ENSO connection. *Geophys Res Lett* 2008, 35:L14S08, doi:10.1029/2007GL032983.
9. Kuleshov Y, Fawcett R, Qi L, et al. Trends in tropical cyclones in the South Indian Ocean and the South Pacific Ocean. *J Geophys Res* 2010, 115:D01101, doi:10.1029/2009JD012372.
10. Diamond HJ, Lorrey AM, Knapp KR, Levinson DH. Development of an Enhanced Tropical Cyclone Tracks Database for the Southwest Pacific from 1840-2010. *Int J Climatol* 2012, 32:2240–2250.
11. Diamond HJ, Lorrey AM, Renwick JA. A southwest Pacific tropical cyclone climatology and linkages to the El Niño–Southern Oscillation. *J Clim* 2013, 26:3–25.
12. Bureau of Meteorology. Database of past tropical cyclone tracks, 2018, accessed 31 October 2018, <http://www.bom.gov.au/cyclone/history/>
13. McBride JL, Keenan TD. Climatology of tropical cyclone genesis in the Australian region. *J Climatol* 1982, 2:13–33.
14. Dare RA, Davidson NE. Characteristics of tropical cyclones in the Australian region. *Mon Weather Rev* 2004, 132:3049–3065.

15. Nicholls N. A possible method for predicting seasonal tropical cyclone activity in the Australian region. *Mon Weather Rev* 1979, 107:1221–1224.
16. Nicholls N. Predictability of interannual variations in Australian seasonal tropical cyclone activity. *Mon Weather Rev* 1985, 113:1144–1149.
17. Nicholls N. Recent performance of a method for forecasting tropical cyclone activity. *Aust Meteor Mag* 1992, 40:105–110.
18. Werner A, Holbrook NJ. A Bayesian forecast model for Australian region tropical cyclone formation. *J Clim* 2011, 24:6114–6131.
19. Nicholls N. The southern oscillation, sea-surface temperature, and internannual fluctuations in Australian tropical cyclone activity. *J Climatol* 1984, 4:661–670.
20. Solow A, Nicholls N. The relationship between the Southern Oscillation and tropical cyclone frequency in the Australian region. *J Clim* 1990, 3:1097–1101.
21. Evans JL, Allan RJ. El Niño/Southern Oscillation modification to the structure of the monsoon and tropical cyclone activity in the Australasian region. *Int J Climatol* 1992, 12:611–623.
22. Hall JD, Matthews AJ, Karoli DJ. The modulation of tropical cyclone activity in the Australian region by the Madden–Julian oscillation. *Mon Weather Rev* 2001, 129:2970–2982.
23. Ramsay HA, Leslie LM, Lamb PJ, Richman MB, Leplastrier M. Interannual variability of tropical cyclones in the Australian region: Role of large-scale environment. *J Clim* 2008, 21:1083–1103.
24. Chand SS, Tory KJ, McBride J, Wheeler M, Dare R, Walsh KJE. The different impact of positive-neutral and negative-neutral ENSO regimes on Australian tropical cyclones. *J Clim* 2013, 26:8008–8016.
25. Lavender SL, Dowdy AJ. Tropical cyclone track direction climatology and its intraseasonal variability in the Australian region. *J Geophys Res* 2016, 13:236–249, doi: 10.1002/2016JD025562.
26. Nicholls N, Landsea C, Gill J. Recent trends in Australian region tropical cyclone activity. *Meteor Atmos Phys* 1998, 65:197–205.
27. Callaghan J, Power SB. Variability and decline in the number of severe tropical cyclones making land-fall over eastern Australia since the late nineteenth century. *Clim Dyn* 2011, 37:647–662

28. Dowdy AJ. Long-term changes in Australian tropical cyclone numbers. *Atmos Sci Lett* 2014, 15:292–298.
29. Bell GD, Halpert MS, Schnell RC et al. Climate assessment for 1999. *Bull Amer Meteor Soc* 2000, 81:S1–S50.
30. Visher SS. Tropical cyclones in Australia and the South Pacific and South Indian Oceans. *Mon Weather Rev* 1922, 288-295.
31. Lourensz RS. Tropical cyclones in the Australian region July 1909 to June 1980. AGPS 1 981, Canberra.
32. Kerr IS. Tropical storms and hurricanes in the southwest Pacific. *New Zealand Meteorol Service Misc* 1976. Pub. 148.
33. Revell CG. Tropical cyclones in the southwest Pacific, Nov. 1969 to April 1979. New Zealand Meteorological Service Misc. 1981 Pub. 170, 53 pp.
34. Holland GJ. On the quality of the Australian tropical cyclone data base. *Aust Meteor Mag* 1981, 29, 169–181.
35. Velden CS, Olander TL, Zehr RM. Development of an objective scheme to estimate tropical cyclone intensity from digital geostationary satellite infrared imagery. *Weather Forecasting* 1998, 13:172–186.
36. Dvorak V F. A technique for the analysis and forecasting of tropical cyclone intensities from satellite pictures. *NOAATech Memo* 1972, NESS 36, [Available from NOAA/NESDIS, 5200 Auth Rd., Washington, DC 20333.]
37. Dvorak VF. Tropical cyclone intensity analysis and forecasting from satellite imagery. *Mon Weather Rev* 1975, 103:420–430.
38. Dvorak VF. Tropical cyclone intensity analysis using satellite data. *NOAATech Rep* 1984, 11, [Available from NOAA/NESDIS, 5200 Auth Rd., Washington, DC 20333.]
39. Veldon CS, et al. The Dvorak tropical cyclone intensity estimation technique: A satellite-based method that has endured for over 30 years. *Bull Amer Meteor Soc* 2006, 87:1195–1210.
40. Knaff JA, Brown DP, Courtney J, Gallina GM, Beven JL. An evaluation of Dvorak technique–based tropical cyclone intensity estimates. *Wea Forecasting* 2010, 25:1362–1379.
41. Harper BA, Callaghan J. On the importance of reviewing historical tropical cyclone intensities. In *27th conference on hurricanes and tropical meteorology 2006 (Vol. 2)*.

42. Knapp KR, Kruk MC, Levinson HJ, et al. The International Best Track Archive for Climate Stewardship (IBTrACS). *Bull Amer Meteor Soc* 2010, 91:363–376.
43. JTWC. Joint Typhoon Warning Centre: The U.S. Navy's Naval Oceanography Portal, Best Track Archive, accessed 4 December 2018, <http://www.metoc.navy.mil/jtwc/jtwc.html?best-tracks>
44. Schreck III CJ, Knapp KR, Kossin JP. The Impact of Best Track Discrepancies on Global Tropical Cyclone Climatologies using IBTrACS. *Mon Weather Rev* 2014, 142:3881–3899.
45. World Meteorological Organization. WMO guidelines on the calculation of climate normals. WMO/TD-No. 1203, Geneva, 2017.
46. Kossin JP, Emanuel KA, Vecchi GA. The poleward migration of the location of tropical cyclone maximum intensity. *Nature* 2014, 509:349–352.
47. Sharmila S, Walsh KJE. Recent poleward shift of tropical cyclone formation linked to Hadley cell expansion, *Nat Climate Change* 2018, 8:730–736.
48. Kilroy, G., Smith RK, Montgomery MT. Tropical low formation and intensification over land as seen in ECMWF analyses. *Quart J Roy Meteor Soc* 2016, 142:1–14.
49. Revell CG, Goulter SW. South Pacific tropical cyclones and the Southern Oscillation. *Mon Weather Rev* 1986a, 114:1138–1145.
50. Revell CG, Goulter SW. Lagged relations between the Southern Oscillation and numbers of tropical cyclones in the South Pacific region. *Mon Weather Rev* 1986b, 114:2669–2670.
51. Hastings PA. Southern Oscillation influences on tropical cyclone activity in the Australian/southwest Pacific region. *Int. J Climatol* 1990, 10:291–298.
52. Basher, RE. Zheng X. Tropical cyclones in the south- west Pacific: Spatial patterns and relationships to Southern Oscillation and sea surface temperature. *J Clim* 1995, 8:1249– 1260.
53. Broadbridge LW, Hanstrum BN. The relationship between tropical cyclones near Western Australia and the Southern Oscillation Index. *Aust Meteor Mag* 1998, 47:183–189.
54. Dong K. El Niño and tropical cyclone frequency in the Australian region and the northwest Pacific. *Aust Met Mag* 1988, 36:219–225.
55. Kuleshov Y, de Hoedt G. Tropical cyclone activity in the Southern Hemisphere. *Bull Aust Meteor Oceanogr Soc* 2003, 16:135–137.
56. Goebbert KH, Leslie LM. Interannual variability of northwest Australian tropical cyclones. *J Clim* 2010, 23:4538–4555.

57. Ramsay HA, Camargo SJ, Kim D. Cluster analysis of tropical cyclone tracks in the Southern Hemisphere. *Clim Dyn* 2012, 39:897–917.
58. McDonnell KA, Holbrook NJ. A Poisson regression model of tropical cyclogenesis for the Australian–southwest Pacific Ocean region. *Weather Forecasting* 2004a, 19:440–455.
59. McDonnell KA, Holbrook NJ. A Poisson regression model approach to predicting tropical cyclogenesis in the Australian/southwest Pacific Ocean region using the SOI and saturated equivalent potential temperature gradient as predictors. *Geophys Res Lett* 2004b, 31, L20110, doi:10.1029/2004GL020843.
60. Liu KS, Chan JCL. Interannual variation of Southern Hemisphere tropical cyclone activity and seasonal forecast of tropical cyclone number in the Australian region. *Int J Climatol* 2012, 32:190–202.
61. Gray WM. *Hurricanes: Their formation, structure and likely role in the tropical circulation*. Meteorology Over Tropical Oceans, D. B. Shaw, Ed., *Royal Meteorol Soc* 1979, 155–218.
62. Richman MB, Leslie LM. Adaptive machine learning approaches to seasonal prediction of tropical cyclones. *In: Procedia Computer Science* 12 2012, 276–281.
63. Wijnands JS, Qian G, Shelton KL, et al. Seasonal forecasting of tropical cyclone activity in the Australian and the South Pacific Ocean regions. *Math. Climate Weather Forecasting* 2015, 1:21–42.
64. Power S, Haylock M, Colman R, Wang X. The predictability of interdecadal changes in ENSO activity and ENSO teleconnections. *J Clim* 2006, 19:4755–4771.
65. Ramsay HA, Richman MB, Leslie LM. Seasonal tropical cyclone predictions using optimized combinations of ENSO regions: Application to the Coral Sea basin. *J Clim* 2014, 27:8527–8542.
66. Magee AD, Verdon-Kidd DC, Diamond HJ, Kiem AS. Influence of ENSO, ENSO Modoki and the IPO on tropical cyclogenesis: a spatial analysis of the southwest Pacific region. *Int J Climatol* 2017, 37:1118–1137.
67. Werner A, Maharaj AM, Holbrook NJ. A new method for extracting the ENSO-independent Indian Ocean dipole: Application to Australian region tropical cyclone counts. *Clim Dyn* 2012, 38:2503–2511.
68. Ramsay HA, Richman MB, Leslie LM. The modulating influence of Indian Ocean sea surface temperatures on Australian region seasonal tropical cyclone counts. *J Clim* 2017, 30:4843–4856.

69. Webster PJ, Holland GJ, Curry JA, Chang H-R. Changes in tropical cyclone number and intensity in a warming environment. *Science* 2005, 309:1844–1846, doi:10.1126/science.1116448.
70. Emanuel K. Increasing destructiveness of tropical cyclones over the past 30 years. *Nature* 2005, 436:686–688, doi:10.1038/nature03906.
71. Elsner JB, Kossin JP, Jagger TH. The increasing intensity of the strongest tropical cyclones. *Nature* 2008, 455:92–95, doi:10.1038/nature07234.
72. Sobel AH, Camargo SJ, Hall TM, et al. Human influence on tropical cyclone intensity, *Science* 2016, 353:242–246.
73. IPCC. *Climate Change 2014: Synthesis Report*. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, Pachauri RK, Meyer LA (eds.)]. Geneva, Switzerland; 2014.
74. Murakami H, Vecchi GA, Delworth TL, et al. Investigating the influence of anthropogenic forcing and natural variability on the 2014 Hawaiian hurricane season [in “Explaining Extremes of 2014 from a Climate Perspective”]. *Bull Amer Meteor Soc* 2015, 96:S115–S119, doi:10.1175/BAMS-D-15-00119.1.
75. Walsh KJE, Mcbride JL, Klotzbach PJ, et al. Tropical cyclones and climate change. *WIREs Climate Change* 2016, 7:65–89. doi: 10.1002/wcc.371.
76. Knutson TR, Camargo SJ, Chan JCL, et al. Tropical cyclones and climate change assessment: Part I, Detection and attribution. *Submitted to the Bulletin of the Amer Meteorol Soc* 2018.
77. Power SB, Smith IN. Weakening of the Walker Circulation and apparent dominance of El Niño both reach record levels, but has ENSO really changed? *Geophys Res Lett* 2007, 34:L18702, doi:10.1029/2007GL030854.
78. Camargo SJ, Sobel AH. Western North Pacific tropical cyclone intensity and ENSO. *J Clim* 2005, 18:2996–3006.
79. Chan JCL. Interannual variations of intense typhoon activity. *Tellus* 2007, 59A:455–460.
80. Chand SS, Walsh KJE. Influence of ENSO on tropical cyclone intensity in the Fiji region. *J Clim* 2011, 24:4096–4108.

81. Haig J, Nott J, Reichert GJ. Australian tropical cyclone activity lower than at any time over the past 550–1,500 years. *Nature* 2014, 505:667-773.
82. IPCC. *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation*. A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change [Field CB, Barros V, Stocker TF, et al. (eds.)]. Cambridge, UK and New York, NY: Cambridge University Press; 2012.
83. Power S, Folland C, Colman A, Mehta V. Inter-decadal modulation of the impact of ENSO on Australia. *Clim Dyn* 1999, 15, 319-324.
84. Li Z, Cai W, Lin X. Dynamics of changing impacts of tropical Indo-Pacific variability on Indian and Australian rainfall. *Scientific Reports* 2016, 6, doi: 10.1038/srep31767
85. Ham Y-G, Choi J-Y, Kug J-S. The weakening of the ENSO–Indian Ocean Dipole (IOD) coupling strength in recent decades. *Clim Dyn* 2016, 49:249–261, doi: 10.1007/s00382-016-3339-5.
86. Cai W, Sullivan A, Cowan T. Climate change contributes to more frequent consecutive positive Indian Ocean Dipole events. *Geophys Res Lett* 2009, 36:L23704, doi:10.1029/2009GL040163.
87. Bieli M, Camargo SJ, Sobel AH, Evans JL, Hall T. A global climatology of extratropical transition. Part I: Characteristics across basins. *J Clim* 2019, 32:3557-3582./JCLI-D-17-0518.1.

wcc_602_figure 1.eps

wcc_602_figure 2.eps

wcc_602_figure 3.eps

wcc_602_figure 4.eps

wcc_602_figure 6.eps

Caption: Tropical cyclone frequency in the Australian region and the potential influences of monitoring methods and technologies on the best track database since 1900.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Chand, SS;Dowdy, AJ;Ramsay, HA;Walsh, KJE;Tory, KJ;Power, SB;Bell, SS;Lavender, SL;Ye, H;Kuleshov, Y

Title:

Review of tropical cyclones in the Australian region: Climatology, variability, predictability, and trends

Date:

2019-07-01

Citation:

Chand, S. S., Dowdy, A. J., Ramsay, H. A., Walsh, K. J. E., Tory, K. J., Power, S. B., Bell, S. S., Lavender, S. L., Ye, H. & Kuleshov, Y. (2019). Review of tropical cyclones in the Australian region: Climatology, variability, predictability, and trends. WILEY INTERDISCIPLINARY REVIEWS-CLIMATE CHANGE, 10 (5), <https://doi.org/10.1002/wcc.602>.

Persistent Link:

<http://hdl.handle.net/11343/285976>