Shao Jingyi (Orcid ID: 0000-0003-0794-2810)
Zosky Graeme (Orcid ID: 0000-0001-9039-0302)
Hall Graham (Orcid ID: 0000-0002-6217-9494)
Dharmage Shyamali (Orcid ID: 0000-0001-6063-1937)
Abramson Michael (Orcid ID: 0000-0002-9954-0538)

Editorial Office Notes:

RES-18-998.R1

ORIGINAL ARTICLE

Received 6 January 2019

Invited to revise 20 March 2019

Revised 16 April 2019

Accepted 14 May 2019

Associate Editor: Stacey Peterson-Carmichael

Senior Editor: Lutz Beckert

Publication fee waiver: NO

Volume: 24

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1111/resp.13617](https://doi.org/10.1111/resp.13617)

Early life exposure to coal mine fire smoke emissions and altered lung function in young children

Jingyi Shao¹, Graeme R. Zosky^{1,2}, Graham L. Hall³, Amanda J. Wheeler^{1,4}, Shyamali Dharmage⁵, Shannon Melody¹, Marita Dalton¹, Rachel E. Foong³, Tierney O'Sullivan¹, Grant J Williamson⁶, Katherine Chappell¹, Michael J. Abramson⁷, Fay H. Johnston¹

¹Menzies Institute for Medical Research, University of Tasmania, Hobart, Tasmania 7000, Australia; ²School of Medicine, Faculty of Health, University of Tasmania, Hobart, Tasmania 7000, Australia; ³Children's Lung Health, Telethon Kids Institute and School of Physiotherapy and Exercise Science Curtin University, West Perth, Western Australia 6872, Australia; ⁴Behaviour, Environment and Cognition Research Program, Mary MacKillop Institute for Health Research, Australian Catholic University, Level 5, 215 Spring Street, Melbourne, Victoria 3000, Australia; ⁵Allergy and Lung Health Unit, Melbourne School of Population and Global Health, University of Melbourne, Carlton, Victoria 3052, Australia; ⁶School of Biological Sciences, University of Tasmania, Hobart, Tasmania 7000, Australia; ⁷Department of Epidemiology and Preventive Medicine, School of Public Health and Preventive Medicine, Monash University, Melbourne, Victoria 3004, Australia.

Correspondence: A/Prof Fay H. Johnston, Menzies Institute for Medical Research, University of Tasmania, 17 Liverpool Street, Hobart, Tasmania 7000, Australia. E-mail: Fay.Johnston@utas.edu.au

Summary at a glance

We found modest evidence for an association between infant exposure to elevated particulate matter during a six-week coal mine fire and reduced respiratory system

reactance measured three years after the fire. The magnitude of the association was small, but of potential clinical importance in the most severely exposed children.

Abbreviations

PM_{2.5}, particulate matter with an aerodynamic diameter less than 2.5 micrometers

FOT, forced oscillation technique

Rrs₅, resistance at a frequency of 5 Hz

Xrs₅, reactance at a frequency of 5 Hz

AX, the area under the reactance curve

SES, socio-economic status

CI, confidence interval

IQR, interquartile range

SD, standard deviation

Latrobe ELF Study, the Latrobe Early Life Follow-up Study

HHS, Hazelwood Health Study

Author contribution roles using the CRediT Taxonomy

Please note that not all categories are relevant to each type of research. Only select those contribution roles that are relevant to your study.

Please complete this Table on behalf of and with approval of all authors?

This information will be translated into an ‘*Author contribution statement*’ by the typesetters and included at the end of your publication.

If not done yet at submission of this study, you will also be required to complete the **Author Contribution roles in ScholarOne** at Step 4 of the First Look finalization.

<i>Role</i>	<i>Degree</i>	<i>Author initials (reflecting the full author name on the manuscript)</i>
1 -Conceptualization	Lead	FHJ
	Equal	
	Supporting	GRZ, SD, MJA, AJW
2 - Data curation	Lead	JS
	Equal	
	Supporting	MD
3 -Formal analysis	Lead	JS
	Equal	
	Supporting	FHJ, GRZ, AJW, SD, GLH, KC
4 - Funding acquisition	Lead	FHJ
	Equal	MJA
	Supporting	
5 - Investigation	Lead	JS
	Equal	
	Supporting	MD, FHJ, GJW, TO, SM, AJW
6 - Methodology	Lead	JS
	Equal	GLH, GRZ, FHJ

	Supporting	REF, KC, SD, MJA, AJW, GJW, SM
7 - Project administration	Lead	FHJ
	Equal	MD
	Supporting	
8 - Resources	Lead	FHJ
	Equal	
	Supporting	MD
9 - Software	Lead	JS
	Equal	
	Supporting	KC
10 - Supervision	Lead	FHJ
	Equal	
	Supporting	GRZ, AJW
11 - Validation	Lead	JS
	Equal	
	Supporting	FHJ, GRZ, AJW
12 - Visualization	Lead	JS
	Equal	
	Supporting	FHJ, GRZ, AJW
13 - Writing – original draft	Lead	JS
	Equal	
	Supporting	FHJ, GRZ, AJW
14 - Writing – review &	Lead	JS

editing	Equal	
	Supporting	GRZ, GLH, AJW, SD, SM, MD, REF, TO, GJW, KC, MJA, FHJ

Abstract

Background and objective: Long-term respiratory risks following exposure to relatively short periods of poor air quality early in life are unknown. We aimed to evaluate the association between exposure to a six-week episode of air pollution from a coal mine fire in children aged less than two years, and their lung function three years after the fire.

Methods: We conducted a prospective cohort study. Individual exposure to 24-hour average and peak concentrations of particulate matter less than 2.5 micrometers in diameter (PM_{2.5}) during the fire were estimated using dispersion and chemical transport modelling. Lung function was measured using the forced oscillation technique, generating standardized Z scores for resistance, reactance at a frequency of 5Hz (Rrs₅; Xrs₅) and area under the reactance curve (AX). We used linear regression models to assess the associations between PM_{2.5} exposure and lung function, adjusted for potential confounders.

Results: 84 of the 203 infants originally recruited, aged 4.3±0.5 years, completed FOT testing. Median [IQR] for average and peak PM_{2.5} were 7.9 [6.8, 16.8] and 103.4 [60.6, 150.7] µg/m³, respectively. The mean±SD Z scores were Rrs₅ 0.56±0.80, Xrs₅ -0.76±0.88 and AX 0.72±0.92. After adjustment for potential confounders including maternal smoking during pregnancy, a 10 µg/m³ increase in average PM_{2.5} was significantly associated with worsening AX (β-coefficient, 0.260; 95% CI 0.019, 0.502), while the association between a 100 µg/m³ increase in peak PM_{2.5} and AX was borderline (0.166; 95% CI -0.002, 0.334).

Conclusion: Infant exposure to coal mine fire emissions could be associated with long-term impairment of lung reactance.

Key words: long-term effects, outdoor smoke, particulate matter, preschool children, respiratory function tests

Short title: Infant smoke exposure and lung function

Introduction

Short-term health impacts associated with exposure to fine particulate matter, a major pollutant generated by fires^[1], have been well characterised. There is consistent evidence associating short-term smoke exposure with increased physician visits, emergency department presentations and hospitalizations for respiratory diseases^[2-3]. However, the long-term health risks from relatively short, that is days to weeks in duration, air pollution episodes have not been characterised, especially in children^[3]. This represents an important gap in the available evidence because severe episodic exposure to fire smoke is likely to increase with climate change^[4].

Infants and young children are more susceptible to the respiratory impacts of air pollution exposure due to their less developed airways and immune system, and faster breathing rates compared with adults^[5]. The first two years of life is a critical window for lung growth^[6]. There is a small but growing body of evidence suggesting that long-term exposure to air pollutants during infancy could result in measurable respiratory health impacts later in life. For example, exposure to traffic-related air pollution during the first year of life has been associated with impaired lung function in both children and adolescents^[7-8]. However, evidence of long-term respiratory impacts from relatively short durations of air pollution exposure during infancy is extremely limited^[9]. One study investigating the Great Smog in London suggested that exposure to the Great Smog during the first year of life could increase the risk of childhood asthma^[10].

The Hazelwood coal mine fire was ignited by embers from wildfires in February 2014 in the Latrobe Valley, Victoria, Australia and lasted for 45 days. In this mine fire episodes of moderate to extreme air pollution were experienced in several nearby towns for days to weeks. The peak 24-hour average concentration of particulate matter with an aerodynamic diameter less than $2.5\text{ }\mu\text{m}$ ($\text{PM}_{2.5}$) reached $731\text{ }\mu\text{g}/\text{m}^3$ at the closest town of Morwell, dramatically higher than the national air quality standard of $25\text{ }\mu\text{g}/\text{m}^3$ ^[11-12]. The health harms resulting from exposure to smoke from coal mine fires could be comparable to those from forest and peat fires. There is a similar spectrum of toxic components, especially in smouldering fires, and the temporal patterns of exposure are similar – often being time limited but severe^[3].

We investigated the infants aged less than two years who were exposed to smoke from the mine fire with the aim of evaluating the association between the magnitude of smoke exposure and lung function three years after the fire.

Methods

Recruitment

This study was based on a prospective cohort study of 571 children who were born between 01/03/2012 and 31/12/2015 and resided in the Latrobe Valley at the time of the fire (see Supplementary Figure S1)^[13]. At recruitment (February to September 2016), the participating parent/carer of each child completed a baseline questionnaire on

sociodemographic, health and family characteristics. Only the 203 members of the cohort who were ≤ 2 years old at the time of the fire were old enough to attempt respiratory function testing and invited to participate in this study [Figure S1 in Supplementary Information]

The Tasmanian Health and Medical Human Research Ethics Committee (reference H14875) approved this study. Additional approval was received from the Human Research Ethics Committees of Monash University, Monash Health, and the University of Melbourne. All parents or caregivers of the studied participants provided signed consent forms for both the baseline survey and the clinical testing.

Exposure estimate

Hourly PM_{2.5} concentrations during the fire period from 09/02/2014 to 31/03/2014 were estimated using meteorological, dispersion and chemical transport modelling at a spatial resolution of 1×1 km^[14]. The model quantitatively calculated PM_{2.5} emission rates at an hourly timestep based on parameters such as how much coal was burned using maps of the area and estimated emission factors. Personal exposures were then calculated for 24-hour average and peak periods based on participants' day/night locations.

Clinical testing

Children's lung function was evaluated three years after the fire, (March to July 2017), using the forced oscillation technique (FOT) (TremoFlo C-100 device, Thorasys, Montreal,

Quebec, Canada) according to ATS/ERS guidelines^[15]. We reported standardized Z scores for baseline Rrs and Xrs at a frequency of 5 Hz (Rrs₅ and Xrs₅), and the area under the reactance curve (AX)^[16]. We excluded measurements with artefacts such as mouth or tongue movement, leakage, swallowing, glottal closure or talking and those having a coherence of <90% at one or more frequencies. Three to five acceptable measurements with a coefficient of variation <10% were obtained for each child. [Supplementary material].

Covariates

All covariates were selected *a priori* according to the existing literature^[17-22] [Supplementary material].

Statistical analysis

We used Pearson's Chi-square test and Welch's t test for the comparisons of characteristics between our studied participants and the full cohort. We calculated β -coefficients and 95% confidence intervals (CIs) from multiple linear regression models to evaluate the associations between infant PM_{2.5} exposure and lung function with or without adjustment for *a priori* selected covariates. We chose the increments of 10 $\mu\text{g}/\text{m}^3$ for average PM_{2.5} and 100 $\mu\text{g}/\text{m}^3$ for peak PM_{2.5} (close to the interquartile ranges (IQRs) of average and peak PM_{2.5} in this study) to enable comparison with other air quality studies suggesting decreased lung function in children after a 10 and 71 $\mu\text{g}/\text{m}^3$ increase in average and peak PM_{2.5} exposure^[23-24]. Five participants (6.0%) had missing values for covariates. We used multiple imputation by chained equations to estimate missing data for

covariates^[25][Supplementary Table S1]. We conducted sensitivity analysis by excluding the participants with imputed data. Additionally, maternal smoking status during pregnancy was assessed as an interaction term in the multivariable regression models. We conducted stratified analysis by gender to assess the different effects of fire smoke exposure on males and females. We performed statistical analyses using R 3.5.0 (the R Foundation, Vienna, Austria). A p -value <0.050 was considered statistically significant.

Results

Participants' characteristics

Of the 203 children eligible for the study, the parents/carers of 137 provided consent for later clinical follow-up. Of these, 101 children attended the clinic and 85 (84.2%) successfully completed FOT testing. We excluded one participant from data analysis due to the poor quality of the FOT measurements (Figure 1).

Almost one fifth (17.9%) of the participants were exposed to maternal tobacco smoking while *in utero*, and two fifths (39.3%) had mothers with ≤ 12 years of education, i.e. completion of secondary education or less. Nearly a quarter (23.8%) of children lived in a house with a current smoker. The mean $\pm$ standard deviation (SD) age of the children at the time of FOT testing was 4.3 ± 0.5 years. The mean annual background $\text{PM}_{2.5}$ concentration in this area ($6.7 \mu\text{g}/\text{m}^3$) was lower than the national air quality standard of $8 \mu\text{g}/\text{m}^3$ ^[12] (Table 1).

Our participants had comparable characteristics with the entire cohort ($n=571$) in terms of gestational age, birthweight, gender, maternal alcohol or tobacco use during pregnancy, primary carer's education level and smoking status, maternal stress and breastfeeding duration (Chi-square or t test $p > 0.050$ for all comparisons; Supplementary Table S2).

$\text{PM}_{2.5}$ exposure during the fire period

The median [IQR] of the average and peak PM_{2.5} levels during the fire period were 7.9 [6.8, 16.8] and 103.4 [60.6, 150.7] µg/m³, respectively (Table 1). Children were exposed to a wide range of PM_{2.5} during the fire period as indicated by the large IQRs.

Lung function measures and risk factors

The mean±SD Z scores of baseline Rrs₅, Xrs₅ and AX were 0.56±0.80, -0.76±0.88 and 0.72±0.92, respectively. Three of the 84 children had Z scores for Rrs₅ ≥2, nine had Z scores for Xrs₅ ≤ -2 and six had Z scores for AX ≥2.

After adjustment for PM_{2.5} and other covariates, maternal smoking during pregnancy was strongly associated with impaired lung reactance, indicated by a decreased Xrs₅ (-1.152; 95%CI -1.708, -0.597; *p*=0.000) and increased AX (0.739; 95%CI 0.101, 1.377; *p*=0.026), while lower maternal education was associated with higher Xrs₅ (0.459; 95%CI 0.063, 0.855; *p*=0.026). Other covariates were not associated with any of the outcomes.

Infant fire PM_{2.5} exposure and lung function

In univariable analysis, PM_{2.5} was associated with Xrs₅ and AX, but not with Rrs₅ (Table 2).

Multivariable model suggested a linear relationship with no threshold for AX. Each 10 µg/m³ increase in average PM_{2.5} was associated with increased AX (0.260; 95%CI 0.019, 0.502; *p*=0.038). Similarly, for every 100 µg/m³ increase in peak PM_{2.5}, we observed a borderline association with AX (0.166; 95%CI -0.002, 0.334; *p*=0.058), consistent with reduced lung function. However, the association between average or peak PM_{2.5} exposure

and Xrs_5 , seen in the univariable analysis, was no longer present (Table 3; Figure 2). There was no evidence that maternal smoking modified the association between mine fire $PM_{2.5}$ and lung function (Supplementary Table S3).

Stratified analysis

Stratified analysis in girls and boys did not show statistically significant associations, although boys tended to have stronger associations than girls for measures of reactance (Supplementary Table S4).

Sensitivity analysis

After excluding participants with imputed data ($n=5$), the results were very similar, although slightly stronger (Supplementary Table S5).

Discussion

At a three-year follow-up of an infant cohort, we observed an association between elevated concentrations of PM_{2.5} during the coal mine fire and worsening peripheral lung mechanics. These results suggest that exposure to mine fire smoke in early life may have influenced lung growth and development. The measured changes in lung function associated with the fire smoke exposure were small for each incremental increase in exposure but would be likely to be of clinical relevance in the most severely exposed children. Furthermore, reductions in lung function as assessed by FOT, and measured on a single occasion, do not necessarily mean that there is a clinical problem or that one might subsequently develop. A recent study suggested that infants with low lung function during the first year could recover in later childhood^[26].

We are not aware of other published studies evaluating early life exposure to smoke from fires, or other short to moderate duration episodes of air pollution, and lung function in preschool aged children. The only comparable study, in terms of exposure, was conducted in monkeys but not in humans. The California fires of 2008 caused degraded air quality for a period of three weeks in a primate research facility soon after the birth of 50 rhesus macaque monkeys, and their lung function was evaluated in adolescence^[27]. Unlike our results, these authors found moderate reductions in airway resistance in the exposed animals compared with the unexposed indicating better lung function. Increased lung stiffness was also observed in the exposed monkeys but only in females, while our study implied stronger effects in boys than in girls after coal mine fire smoke exposure, although

the results were not statistically significant. Future studies are needed to compare the effects of early life exposure to short duration of poor air quality on lung function by gender.

Epidemiological studies have evaluated the impact of exposure to different concentrations of constant background air pollution, as distinct from short-term pollution episodes, in early life and later lung function. One study evaluated the association between exposure to traffic-related air pollution during the first year of life and adolescent lung function using a similar approach to ours^[28]. The authors reported mixed results with some associations identified between reduced lung function and exposure to oxides of nitrogen, but not particulate matter. Studies evaluating exposure to traffic-related PM_{2.5} in infancy and lung function measured with spirometry at 7-10 years of age have reported both reduced^[29], or unchanged^[30] lung function. However, these comparisons should be considered with caution because of different populations, metrics, sources and durations of air pollution exposure that were investigated.

Consistent with current literature, we found that maternal tobacco smoking during pregnancy had negative effects on children's lung function. Many epidemiological studies have indicated an adverse effect of maternal tobacco smoking on the lung health of infants and children^[19, 31-37]. Our findings further highlight the need for smoking cessation support for parents, from the pre-conception period and onwards, to improve their children's respiratory health.

The direction of associations between lower maternal education and lung function in our study was unexpected and not consistent with the weight of existing evidence regarding SES and child health^[21, 38-40]. While we do not have a good explanation for these findings, it could be chance findings in the context of multiple comparisons or inaccurate measurement of maternal education. This finding should be interpreted cautiously because of the small number of children in the subgroups (e.g. n=33 for children with mothers without post-secondary qualifications).

A strength of the study was our ability to adjust for participants' activity patterns to estimate personal PM_{2.5} exposure estimates. Furthermore, we were able to use a simple, non-invasive and objective method of evaluating lung function outcomes suitable for young children^[15]. In addition, we conducted multiple imputation to deal with missing values, which could avoid the reduction of sample size and minimize bias^[41]. Sensitivity analysis of complete cases revealed similar results to the main findings, indicating that our results were robust.

However, our study has some limitations. First, while we were able to evaluate outcomes in children exposed across a wide range of PM_{2.5}, we did not include a group of children with no smoke exposure at all because children in this group were too young to do the FOT testing. Second, the exposure estimates were drawn from modelled air quality data because the monitoring conducted during the fire across the Latrobe Valley was absent during the first week of the fire. Furthermore, our exposure estimates relied upon parental recall of their whereabouts during the fire period and there is therefore, a risk of exposure

misclassification and recall error. While most respondents reported that they were confident of their recall of events during the fire, we were unable to test this objectively. However, there is evidence suggesting a strong correlation between confidence and accuracy of recall in eyewitness studies^[42-43]. Finally, although our study population were representative of the full cohort in terms of demographic and socioeconomic characteristics, a higher proportion of children with well-educated and non-smoking parents were recruited from the local population^[13], which might influence the generalisability of the study findings.

It is possible that the results were influenced by residual confounding. We adjusted for the most important factors such as maternal tobacco smoking, secondhand smoke exposure and maternal education. Education status is a widely used proxy for SES, but in our analysis, lower educational attainment was found to have an unexpected protective association. Therefore, maternal education might not have been the most appropriate marker of SES in our participants. For this reason, our analysis also included the index of socio-economic disadvantage by region^[44] for the residential location of each participant. However, inclusion of this marker did not appreciably change the results.

It has been shown that improvements in air quality were associated with improved lung function in children^[45-46]. Given that the mine fire episode was brief and air quality in the region is generally very good, the effects observed in this study might change as children grow. Therefore, it is important to continue to monitor lung function in this group, to identify if the differences persist. Further studies with a larger sample size and wider range

of exposures by including the *in utero* and no exposure groups at follow-ups would be important for confirming these initial findings.

In conclusion, early life exposure to short-term high intensity air pollution can possibly alter lung development in children. It will be important to continue to monitor children's lung function to investigate long-term outcomes.

Acknowledgements

The Latrobe Early Life Follow-up (ELF) Study constitutes the child health and development stream of the Hazelwood Health Study. The Latrobe ELF Study forms part of the wider research program of the Hazelwood Health Study (HHS) and is run by a multidisciplinary group of researchers and administrative staff from the University of Tasmania, Monash University, the University of Melbourne, the University of Sydney and

CSIRO. We would like to acknowledge all of these staff for their important contributions. Most of all, the study team would like to acknowledge the contribution of all families and community members who have participated in the study to date.

This study is funded by the Victorian Department of Health and Human Services (Australia). The paper presents the views of the authors and does not represent the views of the Department.

Disclosure statement

This research was previously presented at the Annual Congress of the Thoracic Society of Australia and New Zealand (TSANZ) 2018 and the Joint Annual Meeting of the International Society of Exposure Science and the International Society for Environmental Epidemiology (ISES-ISEE) 2018. Outside this study, Fay Johnston received payment for expert testimony from Environment Protection Authority Victoria (Australia). Amanda Wheeler's fellowship was funded by the Centre for Air pollution, energy and health Research. Michael Abramson holds investigator initiated grants from Pfizer and Boehringer-Ingelheim for unrelated research. He has also received assistance with conference attendance from and undertaken an unrelated consultancy for Sanofi.

Author Contributions:

References

- ¹Balmes JR. Where There's Wildfire, There's Smoke. *N Engl J Med* 2018;**378**:881-3.
- ²Reid CE, Brauer M, Johnston FH, Jerrett M, Balmes JR, Elliott CT. Critical Review of Health Impacts of Wildfire Smoke Exposure. *Environ Health Perspect* 2016;**124**:1334-43.
- ³Melody SM, Johnston FH. Coal mine fires and human health: What do we know? *Int J Coal Geol* 2015;**152**:1-14.
- ⁴Liu JC, Mickley LJ, Sulprizio MP, Dominici F, Yue X, Ebisu K, Anderson GB, Khan RFA, Bravo MA, Bell ML. Particulate Air Pollution from Wildfires in the Western US under Climate Change. *Clim Change* 2016;**138**:655-66.
- ⁵Moya J, Bearer CF, Etzel RA. Children's behavior and physiology and how it affects exposure to environmental contaminants. *Pediatrics* 2004;**113**:996-1006.
- ⁶Burri PH. Structural aspects of postnatal lung development - alveolar formation and growth. *Biol Neonate* 2006;**89**:313-22.
- ⁷Schultz ES, Gruzieva O, Bellander T, Bottai M, Hallberg J, Kull I, Svartengren M, Melen E, Pershagen G. Traffic-related air pollution and lung function in children at 8 years of age: a birth cohort study. *Am J Respir Crit Care Med* 2012;**186**:1286-91.
- ⁸Schultz ES, Hallberg J, Bellander T, Bergstrom A, Bottai M, Chiesa F, Gustafsson PM, Gruzieva O, Thunqvist P, Pershagen G, Melen E. Early-Life Exposure to Traffic-related Air Pollution and Lung Function in Adolescence. *Am J Respir Crit Care Med* 2016;**193**:171-7.

⁹Black C, Tesfaigzi Y, Bassein JA, Miller LA. Wildfire smoke exposure and human health: Significant gaps in research for a growing public health issue. *Environ Toxicol Pharmacol* 2017;**55**:186-95.

¹⁰Bharadwaj P, Zivin JG, Mullins JT, Neidell M. Early-Life Exposure to the Great Smog of 1952 and the Development of Asthma. *Am J Respir Crit Care Med* 2016;**194**:1475-82.

¹¹Reisen F, Gillett R, Choi J, Fisher G, Torre P. Characteristics of an open-cut coal mine fire pollution event. *Atmos Environ* 2017;**151**:140-51.

¹²Department of the Environment and Heritage. National standards for criteria air pollutants 1 in Australia.

<http://www.environment.gov.au/protection/publications/factsheet-national-standards-criteria-air-pollutants-australia>. Accessed 17 May, 2018.

¹³Melody S, Dalton M, Dennekamp M, Wheeler A, Dharmage S, Wills K, Reeves M, Ford J, O'Sullivan T, Williamson G, Venn A, Roberts C, Johnston F. The Latrobe Early Life Follow-up (ELF) Cohort Study Volume 1 Description of the cohort and preliminary assessment of possible associations between mine fire emissions and parent-reported perinatal outcomes. http://hazelwoodhealthstudy.org.au/wp-content/uploads/2017/01/ELF_Vol-1_-Cohort_Report-v1.1.pdf. Accessed 18 September, 2018.

¹⁴Emmerson KM, Reisen F, Luhar A, Williamson G, Cope ME. Air quality modelling of smoke exposure from the Hazelwood mine fire. CSIRO Australia. <http://hazelwoodhealthstudy.org.au/wp->

Accessed 18 September, 2018.

¹⁵Beydon N, Davis SD, Lombardi E, Allen JL, Arets HG, Aurora P, Bisgaard H, Davis GM, Ducharme FM, Eigen H, Gappa M, Gaultier C, Gustafsson PM, Hall GL, Hantos Z, Healy MJ, Jones MH, Klug B, Lodrup Carlsen KC, McKenzie SA, Marchal F, Mayer OH, Merkus PJ, Morris MG, Oostveen E, Pillow JJ, Seddon PC, Silverman M, Sly PD, Stocks J, Tepper RS, Vilozni D, Wilson NM. An official American Thoracic Society/European Respiratory Society statement: pulmonary function testing in preschool children. *Am J Respir Crit Care Med* 2007;**175**:1304-45.

¹⁶Calogero C, Simpson SJ, Lombardi E, Parri N, Cuomo B, Palumbo M, de Martino M, Shackleton C, Verheggen M, Gavidia T, Franklin PJ, Kusel MM, Park J, Sly PD, Hall GL. Respiratory impedance and bronchodilator responsiveness in healthy children aged 2-13 years. *Pediatr Pulmonol* 2013;**48**:707-15.

¹⁷Lercher P, Schmitzberger R. Birth weight, education, environment, and lung function at school age: a community study in an alpine area. *Eur Respir J* 1997;**10**:2502-7.

¹⁸Dogaru CM, Strippoli MP, Spycher BD, Frey U, Beardsmore CS, Silverman M, Kuehni CE. Breastfeeding and lung function at school age: does maternal asthma modify the effect? *Am J Respir Crit Care Med* 2012;**185**:874-80.

¹⁹Balte P, Karmaus W, Roberts G, Kurukulaaratchy R, Mitchell F, Arshad H. Relationship between birth weight, maternal smoking during pregnancy and childhood and adolescent lung function: A path analysis. *Respir Med* 2016;**121**:13-20.

- ²⁰Gauthier TW, Brown LAS. In utero alcohol effects on foetal, neonatal and childhood lung disease. *Paediatr Respir Rev* 2017;**21**:34-7.
- ²¹Lee AG, Chiu YM, Rosa MJ, Cohen S, Coull BA, Wright RO, Morgan WJ, Wright RJ. Association of prenatal and early childhood stress with reduced lung function in 7-year-olds. *Ann Allergy Asthma Immunol* 2017;**119**:153-9.
- ²²Waidyatillake NT, Allen KJ, Lodge CJ, Dharmage SC, Abramson MJ, Simpson JA, Lowe AJ. The impact of breastfeeding on lung development and function: a systematic review. *Expert Rev Clin Immunol* 2013;**9**:1253-65.
- ²³Delfino RJ, Staimer N, Tjoa T, Gillen D, Kleinman MT, Sioutas C, Cooper D. Personal and Ambient Air Pollution Exposures and Lung Function Decrements in Children with Asthma. *Environ Health Perspect* 2008;**116**:550-8.
- ²⁴Yamazaki S, Shima M, Ando M, Nitta H, Watanabe H, Nishimuta T. Effect of hourly concentration of particulate matter on peak expiratory flow in hospitalized children: A panel study. *Environ Health* 2011;**10**:15-.
- ²⁵Azur MJ, Stuart EA, Frangakis C, Leaf PJ. Multiple Imputation by Chained Equations: What is it and how does it work? *Int J Methods Psychiatr Res* 2011;**20**:40-9.
- ²⁶Belgrave DCM, Granell R, Turner SW, Curtin JA, Buchan IE, Le Souef PN, Simpson A, Henderson AJ, Custovic A. Lung function trajectories from pre-school age to adulthood and their associations with early life factors: a retrospective analysis of three population-based birth cohort studies. *Lancet Respir Med* 2018;**6**:526-34.
- ²⁷Black C, Gerriets JE, Fontaine JH, Harper RW, Kenyon NJ, Tablin F, Schelegle ES, Miller LA. Early Life Wildfire Smoke Exposure Is Associated with Immune Dysregulation

and Lung Function Decrements in Adolescence. *Am J Respir Cell Mol Biol* 2017;**56**:657-66.

²⁸Schultz ES, Hallberg J, Gustafsson PM, Bottai M, Bellander T, Bergström A, Kull I, Gruzieva O, Thunqvist P, Pershagen G, Melén E. Early life exposure to traffic-related air pollution and lung function in adolescence assessed with impulse oscillometry. *J Allergy Clin Immunol* 2016;**138**:930-2.e5.

²⁹Oftedal B, Brunekreef B, Nystad W, Madsen C, Walker SE, Nafstad P. Residential outdoor air pollution and lung function in schoolchildren. *Epidemiology* 2008;**19**:129-37.

³⁰Rice MB, Rifas-Shiman SL, Litonjua AA, Oken E, Gillman MW, Kloog I, Luttmann-Gibson H, Zanobetti A, Coull BA, Schwartz J, Koutrakis P, Mittleman MA, Gold DR. Lifetime Exposure to Ambient Pollution and Lung Function in Children. *Am J Respir Crit Care Med* 2016;**193**:881-8.

³¹Chen X, Abdulhamid I, Woodcroft K. Maternal smoking during pregnancy, polymorphic CYP1A1 and GSTM1, and lung-function measures in urban family children. *Environ Res* 2011;**111**:1215-21.

³²Hu LW, Yang M, Chen S, Shah K, Hailegiorgis Y, Burgens R, Vaughn M, Huang J, Xaverius P, Paul G, Morawska L, Lu T, Lin S, Zhong SQ, Kong ML, Xie YQ, Hao YT, Zeng XW, Qian Z, Dong GH. Effects of in utero and Postnatal Exposure to Secondhand Smoke on Lung Function by Gender and Asthma Status: The Seven Northeastern Cities (SNEC) Study. *Respiration* 2017;**93**:189-97.

³³Kattan M, Bacharier LB, O'Connor GT, Cohen R, Sorkness RL, Morgan W, Gergen PJ, Jaffee KF, Visness CM, Wood RA, Bloomberg GR, Doyle S, Burton R, Gern JE.

Spirometry and Impulse Oscillometry in Preschool Children: Acceptability and Relationship to Maternal Smoking in Pregnancy. *J Allergy Clin Immunol Pract* 2018;10.1016/j.jaip.2017.12.028.

³⁴Perret JL, Walters H, Johns D, Gurrin L, Burgess J, Lowe A, Thompson B, Markos J, Morrison S, Thomas P, McDonald C, Wood-Baker R, Hopper J, Svanes C, Giles G, Abramson M, Matheson M, Dharmage S. Mother's smoking and complex lung function of offspring in middle age: A cohort study from childhood. *Respirology* 2016;**21**:911-9.

³⁵Bisgaard H, Loland L, Holst KK, Phipps CB. Prenatal determinants of neonatal lung function in high-risk newborns. *J Allergy Clin Immunol* 2009;**123**:651-7, 7.e1-4.

³⁶Gray D, Czovek D, Smith E, Willemse L, Alberts A, Gingsl Z, Hall GL, Zar HJ, Sly PD, Hantos Z. Respiratory impedance in healthy unsedated South African infants: effects of maternal smoking. *Respirology* 2015;**20**:467-73.

³⁷Tager IB, Ngo L, Hanrahan JP. Maternal smoking during pregnancy. Effects on lung function during the first 18 months of life. *Am J Respir Crit Care Med* 1995;**152**:977-83.

³⁸Gray D, Willemse L, Visagie A, Czovek D, Nduru P, Vanker A, Stein DJ, Koen N, Sly PD, Hantos Z, Hall GL, Zar HJ. Determinants of early-life lung function in African infants. *Thorax* 2017;**72**:445-50.

³⁹Slachetova H, Gehring U, Hoek G, Tomaskova H, Luttmann-Gibson H, Moshhammer H, Paldy A, Pattenden S, Slotova K, Speizer F, Zlotkowska R, Heinrich J. Parental education and lung function of children in the PATY study. *Eur J Epidemiol* 2011;**26**:45-54.

⁴⁰Martínez-Briseño D, Fernández-Plata R, Gochicoa-Rangel L, Torre-Bouscoulet L, Rojas-Martínez R, Mendoza-Alvarado L, García-Sancho C, Pérez-Padilla R. Socioeconomic

Status and Longitudinal Lung Function of Healthy Mexican Children. *PLoS One* 2015;**10**:e0136935.

⁴¹Sterne JA, White IR, Carlin JB, Spratt M, Royston P, Kenward MG, Wood AM, Carpenter JR. Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. *BMJ* 2009;**338**:b2393.

⁴²Wixted JT, Mickes L, Clark SE, Gronlund SD, Roediger HL, 3rd. Initial eyewitness confidence reliably predicts eyewitness identification accuracy. *Am Psychol* 2015;**70**:515-26.

⁴³Wixted JT, Mickes L, Dunn JC, Clark SE, Wells W. Estimating the reliability of eyewitness identifications from police lineups. *Proc Natl Acad Sci U S A* 2016;**113**:304-9.

⁴⁴ABS (Australian Bureau of Statistics). Socio-Economic Indexes for Areas (SEIFA) [http://www.ausstats.abs.gov.au/ausstats/subscriber.nsf/0/756EE3DBEFA869EFCA258259000BA746/\\$File/SEIFA%202016%20Technical%20Paper.pdf](http://www.ausstats.abs.gov.au/ausstats/subscriber.nsf/0/756EE3DBEFA869EFCA258259000BA746/$File/SEIFA%202016%20Technical%20Paper.pdf). Accessed 26 November, 2018.

⁴⁵Gauderman WJ, Urman R, Avol E, Berhane K, McConnell R, Rappaport E, Chang R, Lurmann F, Gilliland F. Association of Improved Air Quality with Lung Development in Children. *N Engl J Med* 2015;**372**:905-13.

⁴⁶Dockery DW, Ware JH. Cleaner Air, Bigger Lungs. *N Engl J Med* 2015;**372**:970-2.

Table 1. Characteristics of the participants

Characteristics	Mean±SD	Range
Birthweight* (kg)	3.4±0.6	1.4, 5.3
Gestational age (week)	39.5±1.9	33.0, 43.0
IRSD	3.3±2.8	1.0, 10.0
Background PM _{2.5} exposure (µg/m ³)	6.7±0.9	3.4, 8.3
Age at clinic visit (years)	4.3±0.5	3.4, 5.3
Height (cm)	106.8±6.2	94.5, 128.8
Weight (kg)	19.7±4.9	13.6, 48.5
	n (N) [‡]	%
Gender: Boys	41 (84)	48.8
Maternal alcohol use during pregnancy: yes	9 (83)	10.8
Maternal smoking during pregnancy: yes	15 (84)	17.9
Maternal education: secondary education or less	33 (84)	39.3
Maternal history of asthma: yes	22 (84)	26.2
Maternal stress during pregnancy: frequently stressed	12 (84)	14.3
Effect of coal mine fire on maternal stress: 'increased a lot'	31 (83)	37.3
Second hand smoke exposure: yes	20 (84)	23.8
Breastfeeding: ≤3 month	27 (83)	32.5
Respiratory medication use 24 hours before FOT testing: yes	11 (84)	13.1
Cold/flu-like illnesses in the past 3 weeks: yes	52 (84)	61.9
PM _{2.5} concentrations during the fire period (µg/m ³)	Median	IQR
24-hour average PM _{2.5}	7.9	6.8, 16.8
24-hour peak PM _{2.5}	103.4	60.6, 150.7

Abbreviations: SD, standard deviation; IRSD, Index of Relative Socio-economic Disadvantage; PM_{2.5}, particulate matter with an aerodynamic diameter less than 2.5 micrometers; IQR, interquartile range.

*Participants with complete data for birthweight (n=82).

[‡]n represents number of participants with specific characteristics, while N represents number of participants with completed data for the variable.

Table 2. Univariable analysis of infant coal mine fire PM_{2.5} exposure and other covariates on lung function

Univariable analysis (n=84)	β-coefficient (95% CI)		
	Z (Rrs ₅)	Z (Xrs ₅)	Z (AX)
Average PM _{2.5} (10 µg/m ³ increase)	0.176 (-0.004, 0.356)	-0.213 (-0.410, -0.017)*	0.241 (0.035, 0.447)*
Peak PM _{2.5} (100 µg/m ³ increase)	0.104 (-0.023, 0.231)	-0.120 (-0.260, 0.020)	0.163 (0.018, 0.308)*
Birthweight (1 kg increase)	0.058 (-0.238, 0.353)	0.034 (-0.291, 0.358)	0.104 (-0.237, 0.445)
Gestational age (per week increase)	0.042 (-0.048, 0.132)	-0.021 (-0.120, 0.079)	0.045 (-0.059, 0.149)
IRSD (per unit increase)	-0.041 (-0.103, 0.020)	0.044 (-0.023, 0.111)	-0.020 (-0.091, 0.051)
Background PM _{2.5} (1 µg/m ³ increase)	-0.030 (-0.219, 0.159)	0.098 (-0.109, 0.305)	-0.052 (-0.270, 0.166)
Maternal alcohol use during pregnancy: yes	0.163 (-0.395, 0.721)	-0.375 (-0.982, 0.231)	0.348 (-0.289, 0.986)
Maternal smoking during pregnancy: yes	0.358 (-0.084, 0.801)	-0.860 (-1.317, -0.402) [§]	0.613 (0.112, 1.115)*
Maternal education: secondary education or less	-0.010 (-0.362, 0.342)	0.265 (-0.118, 0.649)	-0.265 (-0.667, 0.138)
Maternal history of asthma: yes	-0.159 (-0.549, 0.230)	-0.117 (-0.547, 0.312)	0.096 (-0.356, 0.547)
Maternal stress during pregnancy: frequently stressed	0.141 (-0.350, 0.632)	-0.258 (-0.795, 0.280)	0.240 (-0.326, 0.805)
Effect of coalmine fire on maternal stress: 'Increased a lot'	0.092 (-0.264, 0.448)	0.042 (-0.350, 0.434)	0.009 (-0.402, 0.421)
Second hand smoke exposure: yes	0.309 (-0.090, 0.707)	-0.193 (-0.635, 0.249)	0.141 (-0.324, 0.607)
Breastfeeding duration: ≤3 months	0.229 (-0.139, 0.597)	-0.052 (-0.456, 0.353)	0.076 (-0.349, 0.502)
Recent cold/flu-like illness: yes	0.169 (-0.184, 0.521)	0.102 (-0.287, 0.491)	0.000 (-0.409, 0.410)
Respiratory medication use: yes	-0.036 (-0.546, 0.474)	0.271 (-0.287, 0.829)	-0.230 (-0.817, 0.356)

Abbreviations: CI, confidence interval; Z (Rrs₅), Z score for resistance at a frequency of 5 Hz; Z (Xrs₅), Z

score for reactance at a frequency of 5 Hz; Z (AX), Z score for the area under the reactance curve; PM_{2.5},

particulate matter with an aerodynamic diameter less than 2.5 micrometers; IRSD, Index of Relative Socio-economic Disadvantage.

* $P<0.05$.

§ $P<0.001$.

Table 3. Multivariable analysis of infant coal mine fire PM_{2.5} exposure and other covariates on lung function

Multivariable analysis (n=84)	β-coefficient [‡] (95% CI)		
	Z (Rrs ₅)	Z (Xrs ₅)	Z (AX)
Average PM _{2.5} (10 µg/m ³ increase)	0.132 (-0.088, 0.351)	-0.184 (-0.394, 0.026)	0.260 (0.019, 0.502)*
Peak PM _{2.5} (100 µg/m ³ increase)	0.071 (-0.082, 0.224)	-0.082 (-0.230, 0.065)	0.166 (-0.002, 0.334)

Abbreviations: CI, confidence interval; Z (Rrs₅), Z score for resistance at a frequency of 5 Hz; Z (Xrs₅), Z score

for reactance at a frequency of 5 Hz; Z (AX), Z score for the area under the reactance curve; PM_{2.5}, particulate matter with an aerodynamic diameter less than 2.5 micrometers.

[‡]Models adjusted for birthweight, gestational age, IRSD, background PM_{2.5} exposure, breastfeeding duration, maternal alcohol use during pregnancy, maternal smoking during pregnancy, maternal education, maternal history of asthma, maternal stress during pregnancy or during the fire, secondhand smoke exposure, cold/flu-like illnesses in the past three weeks and respiratory medication use 24 h before FOT testing. Average and peak PM_{2.5} were modelled separately, and models included all covariates listed in the table.

* $P<0.05$.

Figure Legends

Figure 1. Flow chart of the children participating in this study. DOB, date of birth; FOT, forced oscillation technique.

Figure 2. Risk differences (points) and 95% CIs (whiskers) for the associations between infant fire PM_{2.5} exposure and lung function. The risk difference presents the coefficients from multivariable linear regression analysis demonstrating the change in measures per incremental increase in exposure to fire-related PM_{2.5}. Rrs5, Resistance at a frequency of 5 Hz; Xrs5, Reactance at a frequency of 5 Hz; AX, the area under the reactance curve; PM_{2.5}, particulate matter with an aerodynamic diameter less than 2.5 micrometers.

RESP_13617_Figure 2.jpeg

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Shao, J;Zosky, GR;Hall, GL;Wheeler, AJ;Dharmage, S;Melody, S;Dalton, M;Foong, RE;O'Sullivan, T;Williamson, GJ;Chappell, K;Abramson, MJ;Johnston, FH

Title:

Early life exposure to coal mine fire smoke emissions and altered lung function in young children

Date:

2020-02-01

Citation:

Shao, J., Zosky, G. R., Hall, G. L., Wheeler, A. J., Dharmage, S., Melody, S., Dalton, M., Foong, R. E., O'Sullivan, T., Williamson, G. J., Chappell, K., Abramson, M. J. & Johnston, F. H. (2020). Early life exposure to coal mine fire smoke emissions and altered lung function in young children. RESPIROLOGY, 25 (2), pp.198-205. <https://doi.org/10.1111/resp.13617>.

Persistent Link:

<http://hdl.handle.net/11343/286028>