Article begins on page three of this document.

Title	Unintended consequences of a cautious approach to e-cigarette laws
-------	--

Authors:

	Title	First name	Mid inits	Last name	Postnom (eg, PhD) [3 only for publication]	Position1	Address1	Position2	Address2	Tel	Email
1	Dr.	Anthony	C	Catalano	MBBS (Hons), BA Arts (Journalism)	General Medicine & Clinical Pharmacology Fellow	1			03 9345 4995	christian.catalano@rch.org.au
2	Assoc. Prof.	Noel	E	Cranswick	MBBS, B Med Sc, LLB, FRACP	Director	1	Associate Professor	2	03 9345 6987	noel.cranswick@rch.org.au
3	Dr.	Joanne		Grindlay	MBBS, FACEM, FRACGP, FARGP, EMDM	Emergency Physician	1				joanne.grindlay@rch.org.au
4	Dr.	Mick		Creati	MB. BS., MPH, FRACP	Paediatrician and Consultant in Adolescent and Young Adult Medicine	1				mick.creati@rch.org.au
5	Dr.	Margie	H	Danchin	MBBS, FRACP, PhD	Paediatrician and Director	1	Co-Group Leader	3		Margie.Danchin@rch.org.au
6	Mr.	Jeff		Robinson	B. Pharm, F.S.H.P, Grad. Dip. Hosp. Pharm.	Manager	4				Jeff.ROBINSON@austin.org.au
7	Miss	Nicola		Williams	MN, PGDip Paed, BN	Nurse Consultant	1				nicola.williams@rch.org.au
8	Dr.	Amanda		Gwee	MBBS, FRACP, DTMH	General Paediatrician	1				amanda.gwee@rch.org.au
9											

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/MJA2.50278](#)

This article is protected by copyright. All rights reserved

Number of corresponding author:	1
Number of alternative corresponding author:	

Addresses:

	Institution		City	State	Post Code	
1	Royal Children's Hospital		Melbourne	VIC	3052	
2	University of Melbourne		Melbourne	VIC	3052	
3	Murdoch Children's Research Institute		Melbourne	VIC	3052	
4	Victorian Poisons Information Centre, Austin Health		Melbourne	VIC	3084	

Postal address of first corresponding author (if different from the institutional address given above)	
--	--

Primary Keywords [Office use only]	Substance-related disorders; Environment and public health; Pharmaceutical preparations
Secondary keywords [Office use only]	Legislation, drug; Overdose; Policy, drugs and alcohol; Electronic cigarettes; Prevention and control
Notes:	

Article details (press ctrl – 9 to enter details):

Article type	Letter
Blurb	

Office use

<i>Ms. Number</i>	mja19.00287. R1
<i>Medical editor</i>	Zoe Silverstone
<i>Medical editor email</i>	zsilverstone@mja.com.au
<i>Structural editor</i>	Laura Teruel
<i>Structural editor email</i>	lteruel@mja.com.au
<i>Section/Category</i>	Letters to the editor (reply)
<i>Strapheading</i>	Letter
<i>Substrap</i>	

Wiley – file data:

Filename for copyediting	cat_mja19.00287_ms.docx
Accompanying graphics	None
Stock images	None
Appendices	None

Office use – history:

Event	Date
Original submission received	28/03/2019

Event	Date
Accept	09/04/2019

Proof sent to author	
Proof returned by author	
Published (date format xx/xx/xx)	
Issue	
Vol	
DOI	
Journal	The Medical Journal of Australia
Original article DOI (for response)	

Unintended consequences of a cautious approach to e-cigarette laws

IN REPLY: We thank Mendelsohn and Wodak for their comment on our letter.¹

Although we acknowledge that there is a range of reported lethal nicotine doses in humans ranging from 0.5 mg/kg to 13 mg/kg,^{2,3} the value of 60 mg stated in our original letter is used in numerous studies and in toxicology textbooks.^{4,5} There have been documented paediatric deaths from accidental nicotine ingestion,⁶ with some articles suggesting that the lethal dose in a child may be as low as 10 mg.⁷ Of note, one case report described a lethal dose of 4.2 mg/kg in a 15-month old child. Despite vomiting immediately after the ingestion, there was rapid progression to cardiac arrest and hypoxic brain injury.⁸ Therefore, with the risk of significant morbidity and mortality, a conservative approach to an accidental ingestion is needed.

Even if a lethal dose of 500 mg is applied — equating to about 8 mg/kg per dose in an adult with a weight of 70 kg — the volume of nicotine required to produce a lethal dose in a child would still be significantly smaller. With the reported nicotine concentrations in electronic nicotine delivery systems (ENDS) ranging from 0 mg/mL to 36 mg/mL,³ the ingestion of only 5 mL (one teaspoon) of a solution with a concentration of 18 mg/mL would be needed to produce a potentially lethal dose in a child who weighs 10 kg.

We advocate for the safe labelling and childproof packaging of ENDS products in Australia to prevent deaths from accidental ingestion. The fact remains that even a small amount of ingested liquid nicotine has the potential to cause serious harm to a child, at a time when ENDS products are both widely available and, at this point, unregulated.

Competing interests: No relevant disclosures.

Author details

Anthony C Catalano¹

Noel E Cranswick^{1,2}

Joanne Grindlay¹

Mick Creati¹

Margie H Danchin^{1,3}

Jeff Robinson⁴

Nicola Williams¹

Amanda Gwee¹

¹ Royal Children's Hospital, Melbourne, VIC.

² University of Melbourne, Melbourne, VIC.

³ Murdoch Children's Research Institute, Melbourne, VIC.

⁴ Victorian Poisons Information Centre, Austin Health, Melbourne, VIC.

Christian.Catalano@rch.org.au

doi: 10.5694/mja19.00287

References

- 1 Catalano C, Cranswick NE, Robinson J, et al. Unintended consequences of a cautious approach to e-cigarette laws. *Med J Aust* 2019; 210: 143-144. <https://www.mja.com.au/journal/2019/210/3/unintended-consequences-cautious-approach-e-cigarette-laws>
- 2 Solarino B, Rosenbaum F, Riebelman B, et al. Death due to ingestion of a nicotine-containing solution: case report and review of the literature. *Forensic*

Sci Int 2010; 195: e19-e22.

- 3 American Academy of Paediatrics Policy Statement . Electronic nicotine delivery systems. *Pediatrics* 2015; 136: 1018-1026.
- 4 Cameron JM, Howell DN, White JR, et al. Variable and potentially fatal amounts of nicotine in e-cigarette nicotine solutions. *Tob Control* 2014; 23: 77-78.
- 5 Goldfrank LR, Flomenbaum NE, Lewin NA. Goldfrank's toxicologic emergencies; 6th ed. Stamford, CT: Appleton and Lange, 1998.
- 6 Kamboj A, Spiller HA, Casavant MJ, et al. Pediatric exposure to e-cigarettes, nicotine, and tobacco products in the United States. *Pediatrics* 2016; 137. pii: e20160041.
- 7 Etter JF, Bullen C, Flouris AD, et al. Electronic nicotine delivery systems: a research agenda. *Tob Control* 2011; 20: 243-248.
- 8 Seo A, Dong C, Yu H, Kang M. Accidental ingestion of e-cigarette liquid nicotine in a 15-month old child: an infant mortality case of nicotine intoxication. *Korean J Pediatr* 2016; 59: 490-493.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Catalano, AC;Cranswick, NE;Grindlay, J;Creati, M;Danchin, MH;Robinson, J;Williams, N;Gwee, A

Title:

Unintended consequences of a cautious approach to e-cigarette laws REPLY

Date:

2019-08-01

Citation:

Catalano, A. C., Cranswick, N. E., Grindlay, J., Creati, M., Danchin, M. H., Robinson, J., Williams, N. & Gwee, A. (2019). Unintended consequences of a cautious approach to e-cigarette laws REPLY. MEDICAL JOURNAL OF AUSTRALIA, 211 (4), <https://doi.org/10.5694/mja2.50278>.

Persistent Link:

<http://hdl.handle.net/11343/286146>