Article type : Letters to the editor (comment)

Article begins on page three of this document.

Title	Selecting and optimising patients for total knee arthroplasty
-------	---

Authors:

	Title	First name	Mid initials	Last name	Postnom (eg, PhD) [3 only for publication]	Position1	Address1	Position2	Address2	Tel	Email
1	Dr.	Lachlan	F	Miles	MBBS (Hons), FANZCA	Anaesthetist	1	Honorary Fellow	2		lachlan.miles@austin.org.au
2	A/Prof.	Kate	L	Burbury	MBBS(Hons) FRCAP FRCPA DPhil	Haematologist	3			613855 95000	kate.burbury@petermac.org
3	Prof.	David	A	Story	MBBS(Hons) , MD, BMedSci(Hons), FANZCA	Deputy Director	2			9035 3174	dastory@unimelb.edu.au
4											
5											

Number of corresponding author:	1
Number of alternative corresponding author:	

Addresses:

	Institution	City	State	Post Code	
1	Austin Health	Melbourne	VIC	3084	
2	Centre for Integrated Critical Care, University of Melbourne	Melbourne	VIC	3010	
3	Peter MacCallum Cancer Institute	Melbourne	VIC	3002	
4					
5					

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/MJA2.50304](#)

Postal address of first corresponding author (if different from the institutional address given above)	
--	--

Primary Keywords [Office use only]	Hematologic diseases; Surgical procedures, operative
Secondary keywords [Office use only]	Iron deficiency anemias; Perioperative care; Orthopedic procedures
Notes:	

Article details (press ctrl – 9 to enter details):

Article type	Letter
Blurb	

Office use

<i>Ms. Number</i>	mja19.00179. R1
<i>Medical editor</i>	Geoffrey Thompson/W endy Morgan
<i>Medical editor email</i>	gthompson@ mja.com.au/w morgan@mja. com.au
<i>Structural editor</i>	Laura Teruel
<i>Structural editor email</i>	lteruel@mja.c om.au
<i>Section/Category</i>	Letters to the editor (comment)
<i>Strapheading</i>	Letter
<i>Substrap</i>	

Wiley – file data:

Filename for copyediting	lac_mja19.00179_ms.docx
Accompanying graphics	None
Stock images	None
Appendices	None

Office use – history:

Event	Date
Original submission received	26/02/2019

Event	Date
Accept	14/03/2019

Proof sent to author	
Proof returned by author	
Published (date format xx/xx/xx)	
Issue	
Vol	
DOI	
Journal	
Original article DOI (for response)	

Selecting and optimising patients for total knee arthroplasty

TO THE EDITOR: We read with interest the recent review by Adie and colleagues¹ discussing the optimisation of patients planned for total knee arthroplasty (TKA) and the prevention of adverse post-operative outcomes. However, anaemia, which is a common, major and modifiable risk factor for this patient cohort,² was not addressed by the authors.

A recent observational study found pre-operative anaemia — defined as haemoglobin concentration < 130 g/L — in 32% of patients undergoing elective TKA, and, based on the results for the orthopaedic surgical cohort from this study, it is likely that most of these patients were iron depleted.³ A 2017 single-centre retrospective study similarly reported anaemia in 24% of a similar cohort of patients, and found it was associated with an increased risk of hospital stay in excess of 6 days (unadjusted odds ratio [OR], 1.97; 95% confidence interval [CI], 1.53–2.53; $P < 0.001$), which escalated further and proportionally with decreasing haemoglobin concentration.⁴ In another observational study, anaemia was associated with an increased risk of post-operative complications in aseptic revision joint arthroplasty (OR, 1.45; 95% CI, 1.24–1.70; $P < 0.001$), mortality (OR, 2.18; 95% CI, 1.09–4.36; $P = 0.028$), and increased hospital length of stay (adjusted coefficient, 1.02 days; 95% CI, 0.73–1.31; $P < 0.001$).⁵

Perhaps the most compelling argument for the recognition of anaemia in the pre-operative TKA patient is that it is a risk factor that can be addressed even as little as 2–5 days pre-operatively. A retrospective review of ultrashort term parenteral iron supplementation in major orthopaedic surgery suggested intravenous iron reduced allogeneic blood transfusion rates (8.9% v 30.1%; $P = 0.001$), which are a likely contributor to post-operative outcome. Intravenous iron was also associated with reduced length of hospital stay (8.4 days v 10.7 days; $P = 0.001$).⁶ Newer parenteral iron preparations are widely used as pre-operative rehabilitative interventions, are characterised by their ease of administration and favourable side-effect profile, and are recommended as part of best practice patient blood management guidelines.⁷

At present, large scale prospective and randomised data assessing the safety and efficacy of iron supplementation for correction of anaemia before elective TKA remain lacking. The importance of this area and the potential capacity for clinical and economic effect warrant both further prospective research and consideration by clinicians practising perioperative medicine.

Competing interests: Lachlan Miles, Kate Burbury and David Story are investigators on the NATO (non-anaemic iron deficiency and treatment outcomes in colorectal cancer surgery) study, which has received partial financial support from Vifor Pharma.

Author details

Lachlan F Miles^{1,2}

Kate L Burbury³

David A Story²

1 Austin Health, Melbourne, VIC.

2 Centre for Integrated Critical Care, University of Melbourne, Melbourne, VIC.

3 Peter MacCallum Cancer Institute, Melbourne, VIC.

lachlan.miles@austin.org.au

References

- 1 Adie S, Harris I, Chaun A, et al. Selecting and optimising patients for total knee arthroplasty. *Med J Aust* 2019; 210: 135-141. <https://www.mja.com.au/journal/2019/210/3/selecting-and-optimising-patients-total-knee-arthroplasty>
- 2 Castano-Betancourt M, Fruschein Annichino R, de Azevedo E Souza Munhoz M, et al. Identification of high-risk groups for complication after arthroplasty: predictive value of patient's related risk factors. *J Orthop Surg Res* 2018; 13: 328.
- 3 Muñoz M, Laso-Morales MJ, Gómez-Ramírez S, et al. Pre-operative haemoglobin levels and iron status in a large multicentre cohort of patients undergoing major elective surgery. *Anaesthesia* 2017; 72: 826-834.
- 4 Abdullah HR, Sim YE, Hao Y, et al. Association between preoperative anaemia with length of hospital stay among patients undergoing primary total knee arthroplasty in Singapore: a single-centre retrospective study. *BMJ Open* 2017; 7: e016403.
- 5 Lu M, Sing DC, Kuo AC, Hansen EN. Preoperative anemia independently predicts 30-day complications after aseptic and septic revision total joint arthroplasty. *J Arthroplasty* 2017; 32: S197-S201.
- 6 Muñoz M, Gómez-Ramírez S, Cuenca J, et al. Very short-term perioperative intravenous iron administration and postoperative outcome in major orthopedic surgery: a pooled analysis of observational data from 2547 patients. *Transfusion* 2014; 54: 289-299.
- 7 National Blood Authority. Patient blood management guidelines: module 2 — perioperative. Canberra: National Blood Authority, 2012. <http://www.blood.gov.au/system/files/documents/pbm-module-2.pdf> (viewed March 2019).

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Miles, LF;Burbury, KL;Story, DA

Title:

Selecting and optimising patients for total knee arthroplasty

Date:

2019-09-01

Citation:

Miles, L. F., Burbury, K. L. & Story, D. A. (2019). Selecting and optimising patients for total knee arthroplasty. MEDICAL JOURNAL OF AUSTRALIA, 211 (6), pp.285-+. <https://doi.org/10.5694/mja2.50304>.

Persistent Link:

<http://hdl.handle.net/11343/286308>