1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

Author Manuscript

P
R
O
F
.

J
U
N
Q
I
U

L
U
O

(
O
r
c
i
d

I
D
T

28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57

Author Manuscript

:
0
0
0
0
-
0
0
0
2
-
1
0
2
4
-
1
8
6
6
)
D
R
.
L
O
N

58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87

Author Manuscript

G

C
H
E
N
G

(
O
r
c
i
d

I
D

:

0
0
0
0
-
0
0
0
2
-

Author Manuscript

88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117

8
4
8
3
-
0
4
9
5
)
P
R
O
F
.

D
A
I
W
E
N

C
H
E
N

(
O

Author Manuscript

118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147

r
c
i
d

I
D

:

0
0
0
0
-
0
0
0
3
-
4
2
4
1
-
8
0
5
3
)

148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177

Author Manuscript

A
r
t
i
c
l
e

t
y
p
e

:

O
r
i
g
i
n
a
l

A
r
t
i

Corresponding author mail-id: junqluo2018@tom.com

Beta-glucan from *Agrobacterium sp.* ZX09 improves growth performance and
intestinal function in weaned piglets

Running title

Glucan improves growth and intestinal health

Full names of authors

Junqiu Luo^{1*}, Shuli Liu¹, Bing Yu¹, Jun He¹, Xiangbing Mao¹, Long Cheng², Daiwen Chen¹

Institutional affiliations

¹Institute of Animal Nutrition, Sichuan Agricultural University, and Key Laboratory of Animal
Disease Resistance Nutrition Ministry of Education, Huiming Road 211, Chengdu, Sichuan,
611130, People's Republic of China.

²Faculty of Veterinary and Agricultural Sciences, Dookie Campus 3647, The University of
Melbourne, Australia

*Corresponding author

Acknowledgments

This work was supported by the Project of International Science & Technology Cooperation,
Sichuan (grant number 2016HH0004). The authors greatly appreciate the technical assistance
provided by the staff of the laboratory and the experiment farm.

Abstract

Beta-glucan is currently under consideration as an alternative to in-feed antibiotics. The aim of the study was to investigate *Agrobacterium* sp. ZX09 beta-glucan on intestinal morphology, cytokine concentration, mucin expression and microbial populations of weaning piglets. Pigs were randomly assigned to one of five dietary treatments supplemented with 0, 25, 50, 100 and 200 mg/kg beta-glucan. Data showed an increase of ADG at the 100 mg/kg group ($P=0.03$). A significant increase in villus height and reduction in crypt depth were found in ileal tissue at the 100 mg/kg inclusion level ($P<0.05$). Dietary supplementation of 100 mg/kg beta-glucan enhanced IL-10 concentration ($P=0.04$) and gene expression of MUC1 and MUC2 ($P<0.05$) in the jejunum. Dietary supplementation of 100 mg/kg beta-glucan provoked the up-regulation of *Lactobacillus* counts and down-regulation of *Escherichia coli* counts in the cecum ($P=0.05$). Data strongly suggested that improved growth performance in response to beta-glucan supplementation at 100 mg/kg in weaned piglets may be explained by the improved intestinal function.

Keywords: β -glucan; mucin; cytokine; microbiota; piglet

1 Introduction

With widespread prohibition of in-feed antimicrobial compounds, the use of natural supplements, such as beta-glucan, is currently under consideration as an alternative to in-feed antibiotics due to its gut-modulatory function in human and farm animals (Xiong *et al.*, 2015). Beta-glucan resists hydrolysis by mammalian alimentary enzymes, modifies bacterial community, and is easily fermented by the gastrointestinal microbiota to short-chain fatty acids (SCFAs) (Murphy *et al.*, 2013). In general, SCFAs play an important role in intestinal health by affecting morphology, mucus production and mucosal gene expression and by protecting against overgrowth

with opportunistic pathogens (Yuan *et al.*, 2015).

Previously, most studies referring the stimulatory properties of glucan presenting in the cell walls of algae, fungi, yeast, cereal grains and a limited number of bacteria have been performed in rats (Belobrajdic *et al.*, 2015), chickens (Tian *et al.*, 2016), fish (Jiang *et al.*, 2016), pigs (Lee *et al.*, 2017) and cattle (Ma *et al.*, 2015). However, no substantial consistent effects of dietary glucan on the intestinal function were found. The divergent results might be caused by the quality of variant glucan used in the study. For instance, the soluble beta-glucan from *L. digitata* may be acting via a different mechanism than the insoluble beta-glucan from *L. hyperborea* and *S. cerevisiae* (Sweeney *et al.*, 2012). Apart from the solubility, beta-glucan varied in their structure, chemical composition and molecular weight, which may impact their effects on animal performance and gastrointestinal health (Zheng *et al.*, 2016). Furthermore, the composition of yeast beta-glucan is complex and not highly pure enough to rule out the complicated effects elicited by beta-glucan (Delete).

Efficacy of beta-glucan is likely to link their functional properties, such as purity, structure and molecular weight (Sweeney *et al.* 2012). In the present study, extracellular beta-glucan with high-molecular-mass was obtained from *Agrobacterium sp.* ZX09. It is a novel high purity and water-soluble polymer, which is composed of a linear chain of glucosyl residues linked through a repeat unit of seven β -(1,3) and two α -(1,3) glucosidic bonds, without any β -(1,6) glucosidic bonds. As a new beta-glucan with specific molecular structure, its safety has been demonstrated in the acute and subchronic experiment (Zhou *et al.*, 2013). The differences between *Agrobacterium sp.* beta-glucan and other sources are shown in table 1 (Zheng *et al.*, 2016). As shown in Table 1, compared to beta-glucan in cereal, yeast and fungal formulation, the composition of *Agrobacterium sp.* ZX09 β -glucan is simple and really pure. In addition, the molecular weight of *Agrobacterium sp.* ZX09 β -glucan is relatively high in comparison with other sources. The previous studies have shown that beta-glucan preparations in various sources and alteration in molecular structure enhance the growth promotion properties of the molecule by varying degrees, via eliciting gut responses through modulation of immune and

microbial systems (Khan et al. 2016). Thus, we wonder the effects of the new β -glucan source ingestion on the gut-modulating responses including intestinal barrier function, immunity and bacterial populations in piglets. In this study, we hypothesized that β -glucan derived from *Agrobacterium* sp. ZX09 may aid in gut health of piglets, and that the intestinal modulatory function may depend on the varied dosage of β -glucan in the diets. Hence, the aim of the present study was to assess the intestinal response to the dietary inclusion of *Agrobacterium* sp. ZX09 β -glucans and establish the optimum inclusion level of β -glucan derived from *Agrobacterium* sp. ZX09. However, its intestinal-modulatory activity is uncertain. Hence, the aim of the present study was to establish the optimum inclusion level of beta-glucan derived from *Agrobacterium* sp. ZX09 on growth performance, intestinal morphology, selected bacterial populations, cytokines as well as gene expression of mucins and (Delete) in the weaned piglets. To the best of our knowledge, it is the first in the literature (Delete) to determine the intestinal tissue response to beta-glucan derived from *Agrobacterium* sp. ZX09 (Delete).

2 Materials and Methods

All animal procedures described in the present experiment were approved by the Institutional Animal Care and Use Committee of Sichuan Agricultural University.

2.1 Beta-glucan

The method of beta-glucan preparation was previously described (Zhang et al. 2013). Briefly, *Agrobacterium* sp. ZX09 used in this study was isolated from a soil sample from the ocean coast, and kindly provided by Synlight Bio Co. Ltd of Sichuan, China. Cultures were maintained on HTM agar. A colony of the strain ZX09 was inoculated into a 250 mL flask containing 50 ml medium consisting of 2% sucrose and mineral salt solution. The inoculated preparation was incubated at 28 °C on a rotary shaker at 220 rpm for 24 h. A 0.5 mL portion was transferred to a 250 mL flask containing 50 mL fermentation medium. Fermentation was performed on a rotary shaker at 220 rpm for 48 h. The culture broth was diluted more than 3 times with de-ionized water and centrifuged at 12000 g for 30 min to separate cells from the supernatant. The supernatant was added to two volumes of 95% ethanol. Productivity

of beta-glucan was expressed in terms of the weight after ethanol precipitation collected by centrifugation at 6000 g for 15 min and dried under reduced pressure. The beta-glucan was further purified according to previously described methods (Chen et al. 2011). The total sugar contents of the fraction were determined by the phenol-sulfuric acid method, using glucose for the standard curve (DuBios *et al.*, 1956), and purity of the purified beta-glucan was more than 90%. The mean average molecular weight of purified beta-glucan was estimated from a calibration curve of the standard dextrans obtained by gel filtration on Sepharose CL-4B to be about 2000 kDa.

2.2 Experimental Design and Animal Diets

Transition period for feeding lasted for 3 d. After 3 d of acclimatization, a total of 180 DLY (Duroc × Landrace × Yorkshire) pigs, weaned at 25 d of age, with an initial body weight of (7.03 ± 0.03) kg, were blocked on the basis of body weight and litter of origin. Then, the pigs were randomly assigned to one of the five dietary treatments (six pens per treatment and six piglets per pen) as follows: T1, basal diet (T1); T2, basal diet supplemented with 25 mg/kg beta-glucan; T3, basal diet supplemented with 50 mg/kg beta-glucan; T4, basal diet supplemented with 100 mg/kg beta-glucan and T5, basal diet supplemented with 200 mg/kg beta-glucan. Experimental feeding period lasted for 28 d ad libitum. Purified beta-glucan from *Agrobacterium sp.* ZX09 was extracted according to the procedure described (Zhang *et al.*, 2013), and was kindly provided by Sichuan Synlight Biotech., Ltd.. The total sugar contents of the fraction were determined by the phenol-sulfuric acid method, using glucose for the standard curve (DuBios *et al.*, 1956), and purity of the purified beta-glucan was more than 90%. The mean average molecular weight of purified beta-glucan was estimated from a calibration curve of the standard dextrans obtained by gel filtration on Sepharose CL-4B to be about 2000 kDa (DELETE). Mash diets based on maize and soybean meal were formulated to approximately meet National Research Council-recommended nutrient requirements for pigs (NRC 2012) (Table 2). The basal diet contained beta-glucan, per se, so it's necessary to measure the beta-glucan contents instead of using theoretical values. Based on the analysis of beta-glucan with

Megazyme kit (Megazyme), each dietary group included 10, 32, 65, 115 and 220 mg/kg beta-glucan, respectively.

2.3 Animals and Management

Pigs were housed in pens with six animals confined to each, and had ad libitum access to feed and water. The ambient temperature within the room was approximately 30°C and decreased by 1°C each week of the experiment until reaching 24°C. At 08:00 h of d 1, 15 and 29, the body weight and feed intake of all pigs were measured to calculate average daily weight gain (ADG), average daily feed intake (ADFI) and feed conversion. Fecal consistency was scored as follows: 0, normal; 1, pasty; 2, semiliquid; and 3, liquid. The mean cumulative score of diarrhea was calculated as previously described (Zhao *et al.*, 2015). Every morning and afternoon during the experiment, the status of anal soft fecal contamination and swelling in each piglet was examined and recorded. These data were then used to calculate the incidence rate of diarrhea, which was taken as a reflection of the incidence of sick weanling pigs. The incidence rate of diarrhea was calculated according to the following formula: $[\text{total number of diarrhea pig}/(\text{total number of pigs} \times \text{days of experiment})] \times 100\%$.

2.4 Sampling and Analyses

Nutrient fecal apparent digestibility was measured over a 5 d collection period. During collections, total feces weight was recorded daily. At the end of the collection period, fecal samples were pooled and a subsample retained for laboratory analysis. On d 29, following weighing, one piglet in each pen was then anesthetized (with $\text{C}_3\text{H}_2\text{ClF}_5\text{O}$) and euthanized by intravenous administration (jugular vein) of 4% sodium pentobarbital solution (40 mg/kg BW). Then, the small intestine was removed, and the jejunum and ileum were quickly isolated, flushed with ice-cold saline. One section (about 2 cm in length) in each part was placed in 10% phosphate-buffered formalin for histologic analysis, whereas the other section (approximately 18 cm in length) in jejunal tissue was collected as mucosal sample, frozen in liquid nitrogen, and stored at -80°C until ELISA and RT-PCR analysis. Jejunum is the primary target of nutrient digestion and absorption, while cecum section accounts for microbial

fermentation and metabolite production. The mid-cecum tissues were removed and the digesta sample (approximately 10 g) of each pig was dispensed into two sterilized 5 mL centrifuge tubes for later analysis of microflora.

2.5 Chemical Analysis for Nutrient Digestibility

During the experimental period, ash insoluble in hydrochloric acid (AIA) in the diet was used as an inner indigestible marker for the determination of apparent total tract digestibility (ATTD) for nutrients. AIA analysis of the experimental diets and feces were carried out following the AOAC (2012) methods. From d15 to d 19, fecal samples were collected on a pen basis. Concentrations of dry matter (DM), crude protein (CP), ash and gross energy (GE) were determined in feed and feces according to the methods as Smith previously described (Smith et al. 2011). Briefly, following collection, feces were dried at 60°C for 72 h. The feed and dried feces samples were milled through a hammer mill provided with a 1 mm screen (Christy and Norris). The DM was assayed after drying overnight at 103°C. Ash was determined in a muffle furnace (Nabertherm) at 500°C. The nitrogen content of both feed and feces was determined using the LECO FP 528 instrument (UK Limited). The gross energy of the feed and feces was determined using a Parr 1201 oxygen bomb calorimeter (Parr). The beta-glucan content in the diets was determined by a Megazyme kit (Megazyme) (Delete). The digestibility was then calculated using the following formula:

Digestibility (%) = $[1 - (N_f \times A_d) / (N_d \times A_f)] \times 100$, in which

N_f = Nutrient concentration in feces (%DM),

N_d = Nutrient concentration in diet (%DM),

A_f = AIN concentration in feces (%DM), and

A_d = AIN concentration in diet (%DM).

2.6 Intestinal Morphology Measurements

Formalin-fixed jejunal and ileal cross-sections were embedded in paraffin wax, and 5 µm slides were cut and stained with hematoxylin and eosin. Villus height (the apex of the villus to the villus-crypt junction) and crypt depth (villus-crypt junction to the base of the crypt) were measured at 40× magnification using an image processing and

analysis system (Leica Imaging Systems Ltd., Cambridge, UK). At least 10 well-oriented intact villi and crypts were examined from each pig.

2.7 Assessment of Cytokine Concentrations in the Jejunal Mucosa

Mucosal interleukin-2 (IL-2) was measured using a commercially available swine ELISA kit (BioSource International Inc., Camarillo, CA). IL-1 β , IL-6, IL-10 and TNF- α (tumor necrosis factor- α) in the jejunal mucosa were also measured using commercially available ELISA kits by the manufacturer (R&D Systems Inc., Minneapolis, MN) for pigs, with a within-assay CV of less than 10% for all 3 assays. The assays were analyzed colorimetrically using a BioTek Synergy HT microplate reader (BioTek Instruments, Winooski, VT).

2.8 RNA Extraction and mRNA Expression of ZO-1, Occludin, Claudin-1 and Mucins in the Jejunal Mucosa

Total RNA was extracted from snap-frozen jejunal mucosa sample with TRIzol reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions. The purity of the total RNA was analysed on a NanoDrop spectrophotometer ND1000 (Thermo Scientific, Wilmington, DE, USA). The suitable RNA samples (260:280 ratio ≥ 2.0) were reverse-transcribed using high-capacity cDNA reverse transcription kit (Applied Biosystems, Foster City, CA) according to the manufacturer's instructions and then amplified by PCR. Primers used for amplification of target ZO-1, Occludin, Claudin-1, MUC1, MUC2 and housekeeping β -actin genes were shown in Table 3. Amplification was carried out in a total of 20 μ L, which contained 10 μ L Fast SYBR Green Master Mix (Applied Biosystems Inc.), 1.2 μ L forward and reverse primer mix (5 μ M) and 5 μ L cDNA by an Option DNA Engine (Bio-Rad), with PCR programme as follows: 95°C for 10 s, 40 cycles at 95°C for 5 s, 60°C for 25 s, followed by a final single extension step of 72°C for 5 min. Melt curve analysis was conducted to validate the specificity of the primers. The expression ratio of the target genes relative to the housekeeping gene (β -actin) of each sample was calculated according to the $2^{-\Delta\Delta C_t}$ method (Metzler-Zebeli *et al.*, 2012). All determinations were performed in duplicate.

2.9 DNA Extraction and Real-time PCR Analysis of Bacteria in the Caecal Digesta

The genomic DNAs were extracted from 0.1 g of digesta samples using commercially available rapid bacterial genomic DNA isolation kit by the manufacturer (Sangon Bitech, China) for pigs. The copy numbers of total bacteria, *Bifidobacterium*, *Lactobacillus* and *Escherichia coli* in the caecal samples were quantified by real-time PCR on a Bio-Rad CFX96 real-time system (Bio-Rad, USA) using SYBR Green as the fluorescent dye. The primers for each bacterial species were presented in Table 4, and programs of PCR reaction for each bacterial species were as follows: 95°C for 10 s, 40 cycles at 95°C for 5 s, 50~60°C for 25 s, followed by 95°C for 10 s. A melting curve analysis was generated after each quantitative real-time PCR assay to check and verify the specificity and purity of all PCR products: 40 cycles at 95°C for 39 s, 55°C for 1 min, and 95°C for 1 min. Bacterial counts were presented as log₁₀ copy numbers per gram of dry digesta.

2.10 Statistical Analysis

All data from the experiment were analyzed as a complete randomized design using SAS procedure the General Linear Model procedure of the Statistical Analysis Systems Institute (Delete) (SAS Institute, USA). The statistical model used for animal performance, nutrient digestibility, gut morphology, cytokine concentration and microflora (Delete) data analysis included both the linear and quadratic effects of beta-glucan inclusion levels based on regression analysis. The significance between the treatment differences was identified by Duncan's multiple comparisons test in the General Linear Model. Results were expressed as treatment means with their pooled SEM. The probability value, which denotes significance, is $P < 0.05$.

3 Results

3.1 Growth Performance and Nutrient Digestibility

During day 1-14 of the trail, linear increases in food intake were reported, as the level of beta-glucan increased in piglets (Table 5). The pigs fed 100 mg/kg beta-glucan had highly promoted ADFI ($P < 0.05$). During day 15-28 of the experiment, dietary beta-glucan inclusion led to an increase in food intake and daily gain (quadratic; $P = 0.04$), while a significant effect was observed for daily gain throughout the course of experiment (linear and quadratic; $P = 0.03$, respectively). The diarrhea

incidence was alleviated while treated with 100 mg/kg *Agrobacterium sp.* ZX09 beta-glucan compared with control group (linear; $P=0.04$). The 50 mg/kg and 100 mg/kg beta-glucan inclusion levels increased the digestibility coefficients of DM, CP, ash and gross energy ($P<0.05$) (Table 6).

3.2 Intestinal Morphology

There was no effect on jejunal morphology with increasing dietary inclusion levels of beta-glucan ($P>0.05$) (Table 7) (Delete). Effects on jejunal morphology with increasing dietary inclusion levels of beta-glucan are shown in Table 7. Increasing the level of beta-glucan from 0 to 200 mg/kg significantly improved the morphology of ileum in piglets. Increased villus height, declined crypt depth and an accelerated villus height: crypt depth ratio were detected following dietary inclusion level of beta-glucan at 100 mg/kg in the diets.

3.3 Gene Expression Related to Mucosal Barrier Function

In the study, we studied the impact of beta-glucan on mRNA expressions of tight junction proteins from jejunal mucosa (Figure 1). There was no statistical significant difference in ZO-1 (Figure 1A) and Claudin-1 (Figure 1B) expressions while comparing between dietary groups ($P>0.05$). Dietary supplementation with beta-glucan at 200 mg/kg significantly down-regulated the expression level of Occludin in the jejunum ($P<0.05$) (Figure 1C). To our surprise, 100 mg/kg beta-glucan induced the up-regulation of MUC1 (Figure 2A) and MUC2 (Figure 2B) expressions in the jejunal mucosa of piglets, whereas, the transcripts were inhibited when pigs fed with 200 mg/kg beta-glucan compared with the control animals ($P<0.05$).

3.4 Cytokine Concentrations in Jejunal Mucosa

In the present study, linear increased concentration of IL-10 (linear; $P=0.04$) and reduced concentration of TNF- α (linear; $P=0.05$) in the jejunal mucosa of weaned piglets were observed with increasing treatment of beta-glucan (Table 8). An increase in IL-10 and the decline in IL-2 and TNF- α were detected in the jejunum of pigs supplemented with 100 mg/kg beta-glucan but not at the higher dietary inclusion level.

3.5 Selected Bacterial Populations in Cecum

This study indicated that beta-glucan diet changed the selected bacterial populations in the cecum of piglets (Table 9). Increasing the level of beta-glucan from 0 to 200 mg/kg had no effect on the total bacteria in the cecum of piglets. There was no linear or quadratic effect on *Bifidobacterium* with increasing levels of beta-glucan (linear, $P=0.1$; quadratic, $P=0.44$), while the values at 25 mg/kg and 50 mg/kg dosage were significantly higher than the control group ($P<0.05$). In addition, 100 mg/kg beta-glucan further improved the number of *Lactobacillus* and decreased the copy of *Escherichia coli* ($P<0.05$).

4 Discussion

A novel high purity; water-soluble extracellular beta-glucan of specific molecular weight was used in the present study. To the best of our knowledge, it is the first in literature (Delete) It is confirmed that *Agrobacterium* sp. ZX09 beta-glucan could improve growth performance in weaned piglets and the efficacy was dependent on their inclusion levels. A complex interaction exists between growth performance and disease susceptibility in pigs fed beta-glucan (Vetvicka *et al.*, 2014). Piglets get stress and subsequently had gut misfunction in the post weaning challenge for a period of one month. Based on the improved growth performance of piglets fed by beta-glucan in our previous research (data not shown) and other study (Lee *et al.*, 2017), feeding for 28 days after weaning was considered reliable time to assess the impact of beta-glucan. The reason for differences in means of ADFI, but with no linear or quadratic effects is based on two statistical analyses used in the experiment. Linear or quadratic effects were obtained from regression analysis, while the significance difference between the means of treatment was drawn from the general linear model. No necessary connections exist between such two kinds of statistical analyses. Many studies reported that beta-glucan supplementation enhanced growth performance in pigs (Sweeney *et al.*, 2012; Lee *et al.*, 2017). But the current results are inconsistent with no effects on non-immunochallenged piglets (Hester *et al.*, 2012). An alternative explanation for this discrepancy involves in the differences of optimum concentration, purity, molecular weight, conformation, chemical modification and solubility of

beta-glucan in the diet formulation. Current report strongly demonstrates that optimal dosage of beta-glucan derived from *Agrobacterium* sp. ZX09 was 100 mg per kg of diet for weaned piglets, while the optimal dose of other tested beta-glucans were variant and their sources were mainly obtained from cell wall of *Saccharomyces cerevisiae* (Shao *et al.*, 2016; Tian *et al.*, 2016) and substantially purified from oat (Suchecka *et al.*, 2016; 2017). Beta-glucan content obtained from *Agrobacterium* sp. ZX09 is higher (>90%) than that in *Saccharomyces cerevisiae* and oat (60-80%), thereby existing more sensitive biological activities. Besides, the subjects test with beta-glucan are not the same.

In the present study, pigs fed beta-glucan had higher digestibility of DM, CP, Ash and GE. And the promoted nutrient digestibility contributes to the improved growth performance following beta-glucan ingestion. The dietary inclusion of *Agrobacterium* sp. ZX09 beta-glucan at 100 mg/kg could benefit the gut morphology, mucosal barrier function and caecal microflora, resulting in the increased digestion of nutrients during the passage through the small intestine of pigs and the decreased diarrhea incidence. The up-regulations of MUC1 and MUC2 expression are consistent with the protective role of mucins in the formation of a gut barrier after mucosal stimulation by dietary fiber (Enns *et al.*, 1994). In other studies, mixed-linked beta-glucan, supplemented either in the form of cereals or as a concentrate, was readily fermented, reduced the intestinal number of enterobacteria and increased intestinal butyrate concentrations in growing pigs (Metzler-Zebeli *et al.*, 2010). Of particular interest, the 200 mg/kg beta-glucan inclusion level inhibited the stimulation of Occludin, MUC1 and MUC2 production, which is likely to link with the relatively high diarrhea occurrence when pigs were fed with 200 mg/kg beta-glucan, which was interpreted as being deleterious for piglets, though no reduction in growth performance was detected throughout the course of the study.

The morphology is important to maintain the normal intestinal function, especially digestive and absorptive properties (Suthongsa *et al.*, 2017). In the present study, beta-glucan modified gut morphology followed by changes in an increase of villus height and a decrease of crypt depth in the ileum of weaned pigs. It is hypothesized

that dietary supplementation with beta-glucan increased the digestion and absorption of nutrients during the passage through the small intestine of pigs, which may clarify the underlined mechanisms that *Agrobacterium sp. ZX09* derived beta-glucan at dosage of 100 mg/kg in the diets improved the growth performance and reduced the diarrhea (Delete).

It is generally recognized that IL-1 β , IL-2, IL-6 and TNF- α are regarded as pro-inflammatory cytokines, which modulate immunity, regulate nutrient utilization, and depress growth performance of postnatal animals. The cardinal anti-inflammatory cytokine IL-10 can inhibit T-cell proliferation, development and function as well as the secretion of Th1- and Th2-type cytokines, furthermore, can also suppress the activity of the signal transduction of nuclear transcription factor κ B, which is a major transcription factor of pro-inflammatory cytokines (Zhang *et al.*, 2019). In this study, IL-2, IL-10 and TNF- α as three representative cytokines, had significant differences among groups. The activation of dectin-1 pathway by beta-glucan of high molecular weight, coupled with the activation of TLRs, would explain the enhanced IL-2 and TNF- α response. If feeding *Agrobacterium sp. ZX09* beta-glucan decreases secretion of pro-inflammatory cytokines and promotes secretion of anti-inflammatory cytokines, hence, less activation of the mucosal immune system would be achieved. These results concur with research groups (Wang *et al.*, 2008; Li *et al.*, 2006) who reported that dietary beta-glucan from *Saccharomyces cerevisiae* partially suppressed increase in TNF- α and IL-6 production and enhanced increase in IL-10 production following beta-glucan digestion in an LPS challenge model. On the contrary, yeast product (YP, a mixture of yeast culture, cell wall hydrolysates and yeast extracts) supplementation in the diet of weaned piglets appears to increase the incidence of diarrhea and has adverse effects on intestinal immune function (Yang *et al.*, 2016). Thus, we concluded that depending on the strain, species and preparation process, pure beta-glucan from *Agrobacterium sp. ZX09* appears to be strong modulators of intestinal inflammation in the piglets, while some beta-glucan fractions from yeasts are not.

Structure and metabolites of the gut microbial community are closely related to metagenomic function and nutrient metabolism (Zhang *et al.*, 2017; Luo *et al.* 2018).

In this study, beta-glucan had the ability to increase *Lactobacillus* and *Bifidobacterium* counts, and to reduce *Escherichia coli* number. A similar effect was demonstrated on the beneficial microbiota in pigs fed mulberry (*Morus alba* L.) leaf polysaccharides (Zhao *et al.*, 2015). However, these data are in contrast to previous studies using 0.1 g/kg beta-glucan from *Saccharomyces cerevisiae* or 300 mg/kg laminarin derived from *Laminaria digitata* (Murphy *et al.*, 2013; Zhou *et al.*, 2013), which all resulted in decreased number of *Escherichia coli* with no statistically significant effects on *Lactobacillus* and *Bifidobacterium* communities in their faecal microbiota. We hypothesize that the discrepancy may be attributed to the polymerization of beta-glucan. With a higher molecular weight, beta-glucan from *Agrobacterium* sp. ZX09 potentially remains intact for longer time period as a substrate for different bacteria phynotype, therefore exerting a prebiotic promotion of beneficial bacteria while suppression of deleterious bacteria to a greater extent than *Saccharomyces cerevisiae* and *Laminaria digitata* examined in the cecum. Additionally, alteration in microflora may also influence mucin synthesis and secretion, as adherence of beneficial bacteria to mucosal epithelia stimulates up-regulation of MUC1 and MUC2 expression in vivo and in vitro (Capaldo *et al.*, 2017). Hence, beta-glucan supplementation at 100 mg/kg may have increased MUC1 and MUC2 expression indirectly by acting as a substrate for the resident microbiota, which, in turn, upregulated mucin production.

Production of mucins and tight junction proteins in mucosa has played important roles in formation and integrity of gut barrier. In the study, dietary supplementation with *Agrobacterium* sp. ZX09 beta-glucan at 100 mg/kg decreased the diarrhea occurrence, possibly by inducing gene up-regulation of MUC1 and MUC2 in the jejunal mucosa of weaned piglets. This up-regulation is consistent with the protective role of mucins in the formation of a gut barrier after mucosal stimulation by dietary fiber (Enns *et al.*, 1994). Additionally, alteration in microflora may also influence mucin synthesis and secretion, as adherence of beneficial bacteria to mucosal epithelia stimulates up-regulation of MUC1 and MUC2 expression in vivo and in vitro (Capaldo *et al.*, 2017). Hence, beta-glucan supplementation at 100 mg/kg may have

increased MUC1 and MUC2 expression indirectly by acting as a substrate for the resident microbiota, which, in turn, upregulated mucin production. Of particular interest, the 200 mg/kg beta-glucan inclusion level inhibited the stimulation of Occludin, MUC1 and MUC2 production, which is likely to link with the relatively high diarrhea occurrence when pigs were fed with 200 mg/kg beta-glucan, as compared to the control. The decrease of mucin gene expression in the 200 mg/kg beta-glucan treatment was interpreted as being deleterious for piglets, though no reduction in growth performance was detected throughout the course of the study (Delete).

5 Conclusions

Beta-glucan derived from *Agrobacterium* sp. ZX09, at dietary inclusion level of 100 mg/kg, improved growth performance in weaned piglets through modulation of the intestinal health. Supplementation with 100 mg/kg beta-glucan positively changed the morphology in the ileum, increased IL-10 concentration, also decreased IL-2 and TNF- α concentration in the jejunal mucosa of piglets. Moreover, the dietary inclusion of 100 mg/kg beta-glucan provoked the emergence of a more improved barrier function in the jejunum, meanwhile it had the most profound effect on microbial community in the cecum of piglets. However, higher inclusion level of beta-glucan at 200 mg/kg exerted the deleterious effects on gene expression of MUC1 and MUC2 in the jejunum of weaned piglets. Therefore, we concluded that 100 mg/kg beta-glucan extracted from *Agrobacterium* sp. ZX09 was considered the dietary optimum dose for the weaned piglets in the present study.

Conflict of interest

There are not any conflicts of interest.

Acknowledgements

This work was supported by the Project of International Science & Technology Cooperation, Sichuan (grant number 2016HH0004). The authors greatly appreciate the technical assistance provided by the staff of the laboratory and the experiment

This article is protected by copyright. All rights reserved

farm.

References

- AOAC (2012). Official methods of analysis. 19th ed. Gaithersburg, MD: Association of Official Analytical Chemists.
- Belobrajdic, D.P., Jobling, S.A., Morell, M.K., Taketa, S. & Bird, A.R. (2015). Wholegrain barley β -glucan fermentation dose not improve glucose tolerance in rats fed a high-fat diet. *Nutrition Research*, **35**, 162-168.
- Capaldo, C.T., Powell, D.N. & Kalman, D. (2017). Layered defense: how mucus and tight junctions seal the intestinal barrier. *Journal of Molecular Medicine*, **95**, 927-934.
- Chen, P., Wang, Z., Zeng, L., Yang, X. & Wang, S. (2011). A novel soluble beta-glucan salean protects against acute alcohol-induced hepatotoxicity in mice. *Bioscience Biotechnology and Biochemistry*, **75**, 1990-1993.
- DuBios, M., Gilles, K.A., Hamilton, J.K., Rebers, P.A. & Smith, F. (1956). Colorimetric Method for Determination of Sugars and Related Substances. *Analytical Chemistry*, **28**, 350-356.
- Enns, M.L., Schmidt-Wittig, U., Honer, K., Kownatzki, R. & Gartner, K. (1994). Mechanical challenge causes alterations of rat colonic mucosa and released mucins. Alterations of mucosa and mucins. *Journal of Experimental Animal Science*, **36**, 128-140.
- Hester, S.N., Comstock, S.S., Thorum, S.C., Monaco, M.H., Pence, B.D., Woods, J.A. & Donovan, S.M. (2012). Intestinal and systemic immune development and response to vaccination are unaffected by dietary (1,3/1,6)- β -D-glucan supplementation in neonatal piglets. *Clinical and Vaccine Immunology*, **19**, 1499-1508.
- Jiang, C., Wang, P., Li, M., Liu, S. & Zhang, S. (2016). Dietary β -glucan enhances the contents of complement component 3 and factor B in eggs of zebrafish. *Developmental and Comparative Immunology*, **65**, 107-113.
- Khan, A.A., Gani, A., Masoodi, F. A., Amin, F., Wani, I. A., Khanday, F. A. & Gani A. (2016). Structural thermal, functional, antioxidant & antimicrobial properties of

656 β -d-glucan extracted from baker's yeast (*Saccharomyces cerevisiae*)—Effect of
657 irradiation. *Carbohydrate Polymers*, **140**, 442–450.

658 Lee, S.I., Kim, J.K., Hancock, J.D. & Kim, I.H. (2017). β -glucan from mulberry
659 leaves and curcuma can improve growth performance and nutrient digestibility in
660 early weaned pigs. *Journal of Applied Animal Research*, **45**, 209-214.

661 Li, J., Li, D.F., Xing, J.J., Cheng, Z.B. & Lai, C.H. (2006). Effects of β -glucan
662 extracted from *Saccharomyces cerevisiae* on growth performance, and immunological
663 and somatotropic responses of pigs challenged with *Escherichia coli*
664 lipopolysaccharide. *Journal of Animal Science*, **84**, 2374-2381.

665 Luo, Y., Chen, H., Yu, B., He, J., Zheng, P., Mao, X., Yu, J., Luo, J., Huang, Z. &
666 Chen, D. (2018). Dietary pea fibre alters the microbial community and fermentation
667 with increase in fibre degradation-associated bacterial groups in the colon of pigs.
668 *Journal of Animal Physiology and Animal Nutrition*, **102**, e254–e261.

669 Ma, T., Tu, Y., Zhang, N., Guo, J., Deng, K., Zhou, Y., Yun, Q. & Diao, Q. (2015).
670 Effects of dietary yeast β -glucan on nutrient digestibility and serum profiles in
671 pre-reminant Holstein calves. *Journal of Integrative Agriculture*, **14**, 749-757.

672 Metzler-Zebeli, B.U., Ganzle, M.G., Mosenthin, R. & Zijlstra, R.T. (2012). Oat
673 β -glucan and dietary Calcium and phosphorus differentially modify intestinal
674 expression of proinflammatory cytokines and monocarboxylate transporter 1 and
675 cecal morphology in weaned pigs. *Journal of Nutrition*, **142**, 668-674.

676 Murphy, P., Bello, F.D., O'Doherty, J., Arendt, E.K., Sweeney, T. & Coffey, A.
677 (2013). Analysis of bacterial community shifts in the gastrointestinal tract of pigs fed
678 diets supplemented with β -glucan from *Laminaria digitata*, *Laminaria hyperborea* and
679 *Saccharomyces cerevisiae*. *Animal*, **7**, 1079-1087.

680 Shao, Y., Wang, Z., Tian, X., Guo, Y. & Zhang, H. (2016). Yeast β -D-glucans
681 induced antimicrobial peptide expressions against *Salmonella* infection in broiler
682 chickens. *International Journal of Biological Macromolecules*, **85**, 573-584.

683 Smith, A.G., O'Doherty, J.V., Reilly, P., Ryan, M.T., Bahar, B. & Sweeney, T. (2011).
684 The effects of laminarin derived from *Laminaria digitat* on measurements of gut
685 health: selected bacterial populations, intestinal fermentation, mucin gene expression

and cytokine gene expression in the pig. *British Journal of Nutrition*, **105**, 669-677.

Suchecka, D., Harasym, J., Wilczak, J. & Gromadzka-Ostrowska, J. (2016). Hepato- and gastro- protective activity of purified oat 1-3,1-4- β -D-glucans of different molecular weight. *International Journal of Biological Macromolecules*, **91**, 1177-1185.

Suchecka, D., Blaszczyk, K., Harasym, J., Gudej, S., Wilczak, J. & Gromadzka-Ostrowska, J. (2017). Impact of purified oat 1-3,1-4- β -d-glucan of different molecular weight on alleviation of inflammation parameters during gastritis. *Journal of Functional Foods*, **28**, 11-18.

Suthongsa, S., Pichyangkura, R., Kalandakanond-Thongsong, S. & Thongsong, B. (2017). Effects of dietary levels of chito-oligosaccharide on ileal digestibility of nutrients, small intestinal morphology and crypt cell proliferation in weaned pigs. *Livestock Science*, **198**, 37-44. (Delete)

Sweeney, T., Collins, C.B., Reilly, P., Pierce, K.M., Ryan, M. & O'Doherty, J.V. (2012). Effect of purified β -glucans derived from *Laminaria digitata*, *Laminaria hyperborea* and *Saccharomyces cerevisiae* on piglet performance, selected bacterial populations, volatile fatty acids and pro-inflammatory cytokines in the gastrointestinal tract of pigs. *British Journal of Nutrition*, **108**, 1226-1234.

Tian, X., Shao, Y., Wang, Z. & Guo, Y. (2016). Effects of dietary yeast β -glucans supplementation on growth performance, gut morphology, intestinal *Clostridium perfringens* population and immune response of broiler chickens challenged with necrotic enteritis. *Animal Feed Science and Technology*, **215**, 144-155.

Vetvicka, V. & Oliveira, C. (2014). B(1-3)(1-6)-D-glucans modulate immune status in pigs: potential importance for efficiency of commercial farming. *Annals of Translational Medicine*, **2**, 16.

Wang, Z., Guo, Y., Yuan, J. & Zhang, B. (2008). Effect of β -1,3/1,6-glucan supplementation on growth performance, immune response and plasma prostaglandin E₂, growth hormone and ghrelin in weanling piglets. *Asian-Australasian Journal of Animal Sciences*, **21**, 707-714.

Xiong, X., Yang, H., Li, B., Liu, G., Huang, R., Li F., Liao, P., Zhang, Y., Nyachoti, C.M. & Deng, D. (2015). Dietary supplementation with yeast product improves intestinal function, and serum and ileal amino acid contents in weaned piglets. *Livestock Science*, **171**, 20-27.

Yang, H., Wu, F., Long, L., Li, T. & Xiong, X. (2016). Effects of yeast products on the intestinal morphology, barrier function, cytokine expression, and antioxidant system of weaned piglets. *Journal of Zhejiang University-Science B (Biomedicine & Biotechnology)*, **17**, 752-762.

Yuan, Y., Wang, S., Liu, Y., Li, B., Wang, B. & Peng, Y. (2015). Long-term effect of pH on short-chain fatty acids accumulation and microbial community in sludge fermentation systems. *Bioresource Technology*, **197**, 56-63.

Zhang, C., Zhao, X.H., Yang, L., Chen, X.Y., Jiang, R.S., Jin, S.H. & Geng, Z.Y. (2017). Resveratrol alleviates heat stress-induced impairment of intestinal morphology, microflora, and barrier integrity in broilers. *Poultry Science*, **96**, 4325-4332.

Zhang, C., Chen, K.K., Zhao, X.H., Wang, C. & Geng, Z.Y. (2019). Effect of L-theanine on the growth performance, immune function, and jejunum morphology and antioxidant status of ducks. *Animal*, **13**, 1145-1153.

Zhang, Y., Xia, L., Pang, W., Wang, T., Chen, P., Zhu, B. & Zhang, J. (2013). A novel soluble β -1,3-D-glucan salean reduces adiposity and improves glucose tolerance in high-fat diet-fed mice. *British Journal Nutrition*, **109**, 254-262.

Zhao, X., Li, L., Luo, Q., Ye, M., Luo, G. & Kuang, Z. (2015). Effects of mulberry (*Morus alba* L.) leaf polysaccharides on growth performance, diarrhea, blood parameters, and gut microbiota of early-weanling pigs. *Livestock Science*, **177**, 88-94.

Zheng, X., Zou, S., Xu, H., Liu, Q., Song, J. & Xu, M. (2016). The linear structure of β -glucan from baker's yeast and its activation of macrophage-like RAW264.7 cells. *Carbohydrate Polymers*, **148**, 61-68.

Zhou, T.X., Jung, J.H., Zhang, Z.F. & Kim, I.H. (2013). Effect of dietary β -glucan on growth performance, fecal microbial shedding and immunological responses after

lipopolysaccharide challenge in weaned pigs. *Animal Feed Science and Technology*,
179, 85-92.

Table 1. Structure and content of beta-glucan preparations

Beta-glucan source	Molecular Weight (kDa)	Content (%)	Structure
Yeast	5-80	≤30%	β-1,3/1,6; β-1,3/1,4
Oat	5-250	≤85%	β-1,3/1,4
Algal	<5	≤80%	β-1,3/1,6
<i>Agrobacterium</i> sp. ZX09	200-3000	≥90%	β-1,3

Table 2. The composition and nutrient content of basal diet

Ingredients	Composition, g/kg
Corn	306.2
Extruded corn	277.0
Fish meal	47.0
Whey powder	80.0
Soybean meal	100.0
Extruded full-fat soybean	60.0
Soy protein concentrate	70.0
Sucrose	25.0
L-Lysine·HCL(78%)	4.3
L-Threonine(98.5%)	1.0
DL-Methionine(99%)	1.9
L-Tryptophan	0.2
Choline chloride	1.0
Sodium chloride	2.0
Calcium carbonate	8.8
Dicalcium phosphate	3.1
Soybean oil	10.0
Vitamin premix ¹	0.5
Mineral premix ²	2.0
Nutrient composition, g/kg	
Digestible energy ³ , MJ/kg	14.52
Crude protein ⁴	193..9
Total lysine ⁴	13.6
Total methionine and cystine ³	7.4
Total tryptophan ³	2.2
Total threonine ³	8.0
Calcium ³	8.2
Phosphorus available ³	4.0

¹ Provided the following per kg of diet: Vitamin A, 8000 IU; Vitamin D₃, 1500 IU; Vitamin E, 25 IU; Vitamin K₃, 2.0 mg; Vitamin B₁, 2.0 mg; Vitamin B₂, 5.0 mg; Vitamin B₆, 4.0 mg; Vitamin B₁₂, 0.1 mg; Nicotonic, 25 mg; Pantothenic, 12 mg; Folic acid, 0.75 mg; Biotin, 0.2 mg.

² Provided the following per kg of diet: Fe (FeSO₄·7H₂O), 100 mg; Cu (CuSO₄·5H₂O), 6 mg; Zn (ZnSO₄·7H₂O), 100 mg; Mn (MnSO₄·H₂O), 4 mg; Se (Na₂SeO₃·5H₂O), 0.35 mg; I (KI), 0.14 mg.

³ Calculated values: nutrient level in each ingredient × ingredient content in the diet.

⁴ Measured values.

Table 3. Primers used for quantitative RT-PCR of mucosal barrier function

Gene	Primers and sequence (5' to 3')	Accession number	Product length
ZO-1	F: CAGCCCCCGTACATGGAGA R: GCGCAGACGGTGTTCATAGTT	XM_005659811	114bp
Occludin	F: CTACTCGCTCAACGGGAAAG R: ACGCCTCCAAGTTACCZCTG	NM_001163647.2	158bp
Claudin-1	F: TCTTAGTTGCCACAGCATGG R: CCAGTGAAGAGAGCCTGACC	NM001244539	106bp
MUC1	F: GTGCCGCTGCCCACAACCTG	XM_001926883.4	141bp

	R: AGCCGGGTACCCCAGACCCA		
MUC2	F:GGTCATGCTGGAGCTGGACAGT	XM_003122394.1	181bp
	R: TGCCTCCTCGGGGTCGTCAC		
β -actin	F: TCTGGCACCACACCTTCT	DQ178122	114bp
	R: TGATCTGGGTCATCTTCTCAC		

Table 4. Primers for real-time PCR of bacteria

Items	Primers and sequence (5'-3')	Product length
<i>Total bacteria</i>	F: ACTCCTACGGGAGGCAGCAG R: ATTACCGCGGCTGCTGG	200 bp
<i>Escherichia coli</i>	F: CATGCCGCGTGTATGAAGAA R:CGGGTAACGTCAATGAGCAAA	96 bp
<i>Bifidobacterium</i>	F: CGCGTCCGGTGTGAAAG R: CTTCCCGATATCTACACATTCCA	121 bp
<i>Lactobacillus</i>	F: GAGGCAGCAGTAGGGAATCTTC R: CAACAGTTACTCTGACACCCGTTCTTC	126 bp

Table 5. Effects of *Agrobacterium sp.* ZX09 beta-glucan supplementation on growth performance of weaned piglets

Item	Beta-glucan inclusion level (mg/kg)					Significance		
	0	25	50	100	200	SEM	Linear	Quadratic
1-14d								
ADFI (g) ¹	364.65 ^b	370.40 ^b	377.70 ^{ab}	416.58 ^a	348.95 ^b	6.44	0.05	0.42
ADG (g) ²	225.60	227.18	256.83	259.21	220.83	8.77	0.49	0.22
F/G ³	1.61	1.62	1.48	1.60	1.59	0.06	0.56	0.29
15-28d								

ADFI (g)	615.40 ^{ab}	617.93 ^{ab}	664.03 ^a	660.22 ^{ab}	609.41 ^b	8.69	0.11	0.04
ADG (g)	336.39	348.82	368.54	366.50	336.23	5.38	0.14	0.04
F/G	1.83	1.78	1.80	1.81	1.81	0.02	0.90	0.99
1-28d								
ADFI (g)	480.47 ^{ab}	494.16 ^{ab}	518.89 ^{ab}	529.66 ^a	485.16 ^b	7.20	0.16	0.12
ADG (g)	280.99 ^b	288.00 ^b	310.20 ^{ab}	325.12 ^a	278.53 ^b	5.49	0.03	0.03
F/G	1.71	1.72	1.67	1.63	1.75	0.02	0.38	0.24
Diarrhea incidence	6.83 ^{ab}	4.17 ^b	5.00 ^b	2.67 ^c	9.83 ^a	0.72	0.04	0.08

795 ¹ADFI: Average daily feed intake

796 ²ADG: Average daily gain

797 ³F/G: Feed intake/daily gain

798 ^{a,b,c} Mean values within a row with unlike superscript letters were significantly different (P<0.05).

799

800

801

802

803

804

805

806

Table 6. Effects of *Agrobacterium* sp. ZX09 beta-glucan supplementation on

nutrient digestibility

Item	Beta-glucan inclusion level (mg/kg)					SEM	Significance	
	0	25	50	100	200		Linear	Quadratic
GE (%) ¹	79.78 ^{bc}	78.72 ^c	83.65 ^a	82.45 ^a	82.16 ^{ab}	0.50	<0.01	0.03
Ash (%)	37.74 ^b	38.21 ^b	49.12 ^a	45.83 ^a	45.38 ^a	1.21	<0.01	<0.01
DM (%) ²	80.15 ^{bc}	79.57 ^c	83.87 ^a	82.67 ^a	82.24 ^{ab}	0.44	<0.01	0.03
CP (%) ³	72.22 ^{bc}	70.20 ^c	75.94 ^a	76.52 ^a	75.45 ^{ab}	0.70	<0.01	0.02

809 ¹GE: Gross energy

810 ²DM: Dry matter

811 ³CP: Crude protein

812 ^{a,b,c} Mean values within a row with unlike superscript letters were significantly different (P<0.05).

813

814 **Table 7. Effects of *Agrobacterium* sp. ZX09 beta-glucan supplementation on**
815 **intestinal morphology of weaned piglets**

Item	Beta-glucan inclusion level (mg/kg)					SEM	Significance	
	0	25	50	100	200		Linear	Quadratic
Jejunum								
Villus height (μm)	406.62	394.71	455.98	466.91	408.72	17.83	0.45	0.40
Crypt depth (μm)	201.29	215.50	200.22	203.11	188.18	8.25	0.90	0.55
VH:CD ¹	2.09	1.84	2.31	2.20	2.18	0.11	0.62	0.86
Ileum								
Villus height (μm)	409.48 ^b	400.23 ^b	426.13 ^{ab}	502.89 ^a	447.93 ^{ab}	14.25	0.07	0.09
Crypt depth (μm)	258.65 ^a	188.68 ^b	211.29 ^b	184.02 ^b	203.20 ^b	9.02	0.03	0.05
VH:CD	1.62 ^c	2.12 ^{bc}	2.02 ^{bc}	2.75 ^a	2.22 ^b	0.12	0.09	<0.01

816 ¹VH: CD: Ratio of villus height to crypt depth

817 ^{a,b,c} Mean values within a row with unlike superscript letters were significantly different (P<0.05).

818

819

820 **Table 8. Effects of *Agrobacterium* sp. ZX09 beta-glucan supplementation on**
821 **cytokine concentrations in the jejunum of weaned piglets**

Item	Beta-glucan inclusion level (mg/kg)					SEM	Significance	
	0	25	50	100	200		Linear	Quadratic
IL-1 β (ng/ mg protein)	116.72	109.25	112.67	110.93	118.49	2.12	0.19	0.56
IL-2 (pg/mg protein)	79.13 ^a	79.71 ^a	78.98 ^a	76.81 ^b	77.09 ^{ab}	2.04	0.07	0.21
IL-6 (pg/ mg protein)	540.33	556.39	545.98	542.19	545.04	1.96	0.13	0.98
IL-10 (ng/ mg protein)	18.66 ^a	20.14 ^a	22.82 ^{ab}	25.17 ^b	18.58 ^a	1.25	0.04	0.15

TNF- α (pg/ mg protein)

150^a

140^a

110^b

110^b

130^{ab}

0.03

0.05

0.36

822

^{a,b} Mean values within a row with unlike superscript letters were significantly different (P<0.05).

823

824

Table 9. Effects of *Agrobacterium* sp. ZX09 beta-glucan supplementation on cecal

825

microflora of weaned piglets (Unit: lg copies/gram of dry digests)

Item	Beta-glucan inclusion level (mg/kg)					Significance		
	0	25	50	100	200	SEM	Linear	Quadratic
Lactobacillus	8.20 ^b	8.29 ^b	8.73 ^{ab}	9.03 ^a	8.77 ^{ab}	0.11	0.08	0.49
Bifidobacterium	8.54 ^b	10.26 ^a	10.41 ^a	9.57 ^{ab}	9.24 ^{ab}	0.24	0.10	0.44
Escherichia coli	8.80 ^a	7.73 ^{ab}	7.87 ^{ab}	7.68 ^b	7.37 ^b	0.24	0.05	0.41
Total bacteria	10.51	11.49	11.58	11.46	11.53	0.21	0.50	0.209

826

^{a,b} Mean values within a row with unlike superscript letters were significantly different (P<0.05).

Fig.1 mRNA expression level of tight junction proteins in the jejunal mucosa of piglets fed diets with different inclusion levels of beta-glucan. Data are shown as means $\pm$ SEM. Fig. 1A: ZO-1 mRNA expression level. Fig. 1B: Claudin-1 mRNA expression level. Fig. 1C: Occludin mRNA expression level. Difference letters indicate statistically significant differences between groups ($P < 0.05$). T1-control group fed with basal diet; T2-experimental group fed with 25 mg/kg beta-glucan; T3-experimental group fed with 50 mg/kg beta-glucan; T3-experimental group fed with 50 mg/kg beta-glucan; T4-experimental group fed with 100 mg/kg beta-glucan; T5-experimental group fed with 200 mg/kg beta-glucan

Fig.2 mRNA expression level of MUC1 and MUC2 in the jejunal mucosa of piglets fed diets with different molecular weight beta-glucan. Data are shown as means±SEM. Fig. 2A: MUC1 mRNA expression level. Fig. 2B: MUC2 mRNA expression level. Difference letters indicate statistically significant differences between groups ($P<0.05$). T1-control group fed with basal diet; T2-experimental group fed with 25 mg/kg beta-glucan; T3-experimental group fed with 50 mg/kg beta-glucan; T4-experimental group fed with 100 mg/kg beta-glucan; T5-experimental group fed with 200 mg/kg beta-glucan

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Luo, J;Liu, S;Yu, B;He, J;Mao, X;Cheng, L;Chen, D

Title:

Beta-glucan from *Agrobacterium* sp. ZX09 improves growth performance and intestinal function in weaned piglets

Date:

2019-11-01

Citation:

Luo, J., Liu, S., Yu, B., He, J., Mao, X., Cheng, L. & Chen, D. (2019). Beta-glucan from *Agrobacterium* sp. ZX09 improves growth performance and intestinal function in weaned piglets. *Journal of Animal Physiology and Animal Nutrition*, 103 (6), pp.1818-1827. <https://doi.org/10.1111/jpn.13163>.

Persistent Link:

<http://hdl.handle.net/11343/286321>