PROF. ANDREA BERNASCONI (Orcid ID : 0000-0001-9358-5703)

DR. FERNANDO CENDES (Orcid ID : 0000-0001-9336-9568)

DR. WILLIAM H THEODORE (Orcid ID : 0000-0002-4669-5747)

PROF. ANGELO LABATE (Orcid ID : 0000-0002-8827-7324)

PROF. PHILIPPE RYVLIN (Orcid ID : 0000-0001-7775-6576)

Article type : Letter

Response to commentary on Recommendations for the use of structural MRI in the care of patients with epilepsy: A consensus report from the ILAE Neuroimaging Task Force

Andrea Bernasconi ^{1/*}, Fernando Cendes ², William Theodore ³, Ravnoor S Gill ¹, Matthias Koepp ⁴, R. Edward Hogan ⁵, Graeme Jackson ⁶, Paolo Federico ⁷, Angelo Labate ⁸, Anna Elisabetta Vaudano ⁹, Ingmar Blümcke ¹⁰, Philippe Ryvlin ¹¹, Neda Bernasconi ^{1/*}

- 1) *Neuroimaging of Epilepsy Laboratory, McConnell Brain Imaging Centre and Montreal Neurological Institute and Hospital, McGill University, Montreal, Quebec, Canada*
- 2) *Department of Neurology, University of Campinas – UNICAMP, Campinas, SP, Brazil*
- 3) *Clinical Epilepsy Section, NIH Bethesda Maryland USA, NIH*
- 4) *Institute for Neurology, University College London, UK*
- 5) *Department of Neurology, Washington University Scholl of Medicine, St. Louis, MO, USA*
- 6) *The Florey Institute of Neuroscience and Mental Health and The University of Melbourne, Austin Campus, Heidelberg, Victoria, Australia*
- 7) *Hotchkiss Brain Institute, University of Calgary, Calgary, Alberta, Canada*
- 8) *Institute of Neurology, University of Catanzaro, Italy*
- 9) *Neurology Unit, OCASE Hospital, AOU Modena, University of Modena and Reggio-Emilia, Modena, Italy*
- 10) *Department of Neuropathology, University Hospital Erlangen, Germany*
- 11) *Clinical Neurosciences, Centre Hospitalier Universitaire Vaudois (CHUV), Lausanne, Switzerland*

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/EPI.16324](https://doi.org/10.1111/EPI.16324)

This article is protected by copyright. All rights reserved

** These authors contributed equally to this work*

KEY WORDS: Epilepsy, structural magnetic resonance imaging, adults, pediatrics

Address correspondence to

Andrea Bernasconi, MD

Neuroimaging of Epilepsy Laboratory

McConnell Brain Imaging Centre and Montreal Neurological Institute and Hospital

McGill University

3801 University Street

Montreal, Quebec, Canada H3A 2B4

Telephone: (514) 398-3361

E-mail: andrea.bernasconi@mcgill.ca

We wish to thank Rosenow and colleagues for their interest and comments on our ILAE recommendations on the use of MRI in the care of people with epilepsy ¹.

The authors point out that gradient recall echo (GRE-T2*) or susceptibility weighted imaging (SWI) should be integrated in a standard work-up of epilepsy. Although we agree in principle with the utility of these imaging contrasts in the evaluation of patients with cavernoma-related epilepsy, we would like to counterargue.

Cerebral cavernous malformations (CCM), also called cavernous hemangiomas, comprise 10-15% of all CNS vascular lesions ². Their annual detection rate incidence has been estimated at 0.56 per 100,000 per year for adults >16 years of age ³. The most common clinical symptoms of CCM include seizures (50%), intracranial hemorrhage (25%), and focal neurological deficits without evidence of recent hemorrhage (25%) ⁴. A significant fraction of cases (20%-50%) have no symptoms and are discovered incidentally due to widespread availability and utilization of MRI ⁵.

CCM are generally sporadic and characterized by a single lesion. Conversely, the autosomal dominant familial form, which accounts for 10–30% of all cases ⁶, is associated with multiple lesions. In these cases, the probability that new lesions develop during lifetime makes it necessary to have imaging surveillance of family members.

In clinically symptomatic patients, conventional MRI sequences, including high-resolution T1 magnetization prepared rapid gradient echo (MPRAGE), T2-weighted and FLAIR sequences as those proposed in the harmonized neuroimaging of epilepsy structural sequences (HARNESS) MRI protocol, identify CCM with a specificity and sensitivity nearing 100%. Their image appearance include a central reticulated core containing blood products (with intensity variations depending on the time of bleeding), as well as a surrounding hemosiderin ring ^{3; 7; 8}. GRE-T2* and SWI are the mainstay sequences to identify familial CMM, as these sequences are most sensitive to small lesions undetected on conventional sequences ^{9; 10}.

While we agree that electrophysiology cannot disentangle the specific effects of hemosiderin or blood products on electrical activity, clinicians use clues such as family history to estimate the risk for CCM in a given patient. Thus, as specified in the current ILAE recommendation ¹, the Committee is of the opinion that dedicated sequences could be added in appropriate clinical circumstances. In other words, an SWI sequence would be relevant only in patients in whom the HARNESS-MRI protocol is non-informative.

We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this report is consistent with those guidelines.'

Disclosure

Neither of the authors has any conflict of interest to disclose.

References

1. Bernasconi A, Cendes F, Theodore WH, et al. Recommendations for the use of structural magnetic resonance imaging in the care of patients with epilepsy: A consensus report from the International League Against Epilepsy Neuroimaging Task Force. *Epilepsia* 2019;60:1054-1068.
2. Barrow DL. Classification and natural history of cerebral vascular malformations: arteriovenous, cavernous, and venous. *J Stroke Cerebrovasc Dis* 1997;6:264-267.
3. Akers A, Al-Shahi Salman R, I AA, et al. Synopsis of Guidelines for the Clinical Management of Cerebral Cavernous Malformations: Consensus Recommendations Based on Systematic Literature Review by the Angioma Alliance Scientific Advisory Board Clinical Experts Panel. *Neurosurgery* 2017;80:665-680.
4. Al-Shahi Salman R, Hall JM, Horne MA, et al. Untreated clinical course of cerebral cavernous malformations: a prospective, population-based cohort study. *Lancet Neurol* 2012;11:217-224.
5. Morris Z, Whiteley WN, Longstreth WT, Jr., et al. Incidental findings on brain magnetic resonance imaging: systematic review and meta-analysis. *BMJ* 2009;339:b3016.
6. Goldstein HE, Solomon RA. Epidemiology of cavernous malformations. *Handbook of clinical neurology* 2017;143:241-247.
7. Rigamonti D, Johnson PC, Spetzler RF, et al. Cavernous malformations and capillary telangiectasia: a spectrum within a single pathological entity. *Neurosurgery* 1991;28:60-64.
8. Rosenow F, Alonso-Vanegas MA, Baumgartner C, et al. Cavernoma-related epilepsy: review and recommendations for management--report of the Surgical Task Force of the ILAE Commission on Therapeutic Strategies. *Epilepsia* 2013;54:2025-2035.
9. Wang KY, Idowu OR, Lin DDM. Radiology and imaging for cavernous malformations. *Handbook of clinical neurology* 2017;143:249-266.
10. Sparacia G, Speciale C, Banco A, et al. Accuracy of SWI sequences compared to T2*-weighted gradient echo sequences in the detection of cerebral cavernous malformations in the familial form. *Neuroradiol J* 2016;29:326-335.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Bernasconi, A;Cendes, F;Theodore, W;Gill, RS;Koepp, M;Hogan, RE;Jackson, G;Federico, P;Labate, A;Vaudano, AE;Bluemcke, I;Ryvlin, P;Bernasconi, N

Title:

Response to commentary on recommendations for the use of structural MRI in the care of patients with epilepsy: A consensus report from the ILAE Neuroimaging Task Force

Date:

2019-08-29

Citation:

Bernasconi, A., Cendes, F., Theodore, W., Gill, R. S., Koepp, M., Hogan, R. E., Jackson, G., Federico, P., Labate, A., Vaudano, A. E., Bluemcke, I., Ryvlin, P. & Bernasconi, N. (2019). Response to commentary on recommendations for the use of structural MRI in the care of patients with epilepsy: A consensus report from the ILAE Neuroimaging Task Force. *EPILEPSIA*, 60 (10), pp.2143-2144. <https://doi.org/10.1111/epi.16324>.

Persistent Link:

<http://hdl.handle.net/11343/286343>