LPS induces steroid-resistant exacerbations in a mouse model of allergic airway disease collectively through IL-13 and pulmonary macrophage activation

Sara Hadjigol^{1, *}, Keilah, G. Netto^{1, *}, Steven Maltby¹, Hock L. Tay¹, Thi H. Nguyen¹, Nicole G. Hansbro¹, Fiona Evers¹, Philip M. Hansbro¹, Ming Yang^{1, ¶} & Paul S. Foster^{1, ¶}

¹ Priority Research Centre for Healthy Lungs, Department of Microbiology and Immunology, School of Biomedical Sciences & Pharmacy, Faculty of Health and Priority Research Centre for Healthy Lungs, Hunter Medical Research Institute, University of Newcastle, Callaghan, NSW 2300, Australia

²Centre for Inflammation, Centenary Institute, Sydney, NSW 2050, and University of Technology Sydney, Faculty of Science, Ultimo NSW 2007, Australia

* Hadjigol S. and Netto K. G. contributed equally; ¶Yang M. and Foster P. S. contributed equally

Running Title: LPS-induced exacerbations require IL-13 and lung macrophages

Key Words: asthma exacerbation, LPS, macrophages, IL-13, steroid-resistance, AHR.

ABSTRACT

Background: Acute exacerbations of asthma represent a major burden of disease and are often caused by respiratory infections. Viral infections are recognised as significant triggers of exacerbations, however less is understood about the how microbial bioproducts such as the endotoxin (lipopolysaccharide (LPS)) trigger episodes. Indeed, increased levels of LPS have been linked to asthma onset, severity and steroid-resistance

Objective: The goal of this study was to identify mechanisms underlying bacterial-induced exacerbations by employing LPS as a surrogate for infection.

Methods: we developed a mouse model of LPS induced exacerbation on the background of pre-existing Type-2 allergic airways disease (AAD).

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/CEA.13505](https://doi.org/10.1111/CEA.13505)

Results: LPS induced exacerbation was characterised by steroid resistant airways hyper-responsiveness (AHR) and an exaggerated inflammatory response distinguished by increased numbers of infiltrating neutrophils/macrophages and elevated production of lung inflammatory cytokines, including TNF α , IFN γ , IL-27 and MCP-1. Expression of the Type-2 associated inflammatory factors such as IL-5 and IL-13 were elevated in AAD but not altered by LPS exposure. Furthermore, AHR and airways inflammation were no longer suppressed by corticosteroid (dexamethasone) treatment after LPS exposure. Depletion of pulmonary macrophages by administration of 2-chloroadenosine into the lungs suppressed AHR and reduced IL-13, TNF α and IFN γ expression. Blocking IL-13 function, through either IL-13-deficiency or administration of specific blocking antibodies, also suppressed AHR and airway inflammation.

Conclusions & Clinical Relevance: We present evidence that IL-13 and innate immune pathways (in particular pulmonary macrophages) contribute to LPS-induced exacerbation of pre-existing AAD and provide insight into the complex molecular processes potentially underlying microbial-induced exacerbations.

INTRODUCTION

Asthma is a complex and heterogeneous chronic airway disease, which is characterised by airways inflammation, mucus hyperplasia and airway hyper-responsiveness (AHR)¹. Acute asthma exacerbations are a significant clinical issue as they worsen disease symptoms, often fail to respond to corticosteroid treatment and contribute significantly to healthcare costs². Respiratory infections are associated with the induction, progression and exacerbation of asthma and have been associated with up to 80% of acute asthma exacerbations in children and 50% in adults³. Bacterial infections including *Haemophilus influenzae*, *Mycoplasma pneumoniae* and *Chlamydia pneumoniae* are associated with exacerbations⁴.

Further, exposure to the bacterial component lipopolysaccharide (LPS; endotoxin) has independently been linked to asthma onset, severity and steroid-resistance^{5,6}. Microbiome studies have also highlighted potential pathogenic roles for microbiota and microbial products on disease pathogenesis, including asthma onset, severity and disease phenotype^{7,8}. However, the mechanisms linking endotoxin exposure to asthma exacerbations remain poorly understood.

A number of lines of evidence link lung bacterial exposure with asthma outcomes. Individuals with asthma have altered bacterial profiles in their lungs as determined by 16S rRNA sequencing, compared to healthy controls⁹. Differences in bacterial colonisation of the lungs are associated with altered bronchial hyperreactivity¹⁰ and clinical phenotypes of

asthma¹¹. Pathogenic bacterial infection is associated with more severe airway obstruction and neutrophil recruitment to the airways, which fails to respond to corticosteroid therapy¹². A number of epidemiological studies have also linked high exposure levels to the bacterial cell-wall component LPS with severity of asthma^{5,6}. Low-dose LPS inhalation in healthy controls can induce sputum neutrophilia¹³, but limited data is available on the effect of LPS challenge in individuals with pre-existing asthma.

A number of mouse models have been developed to assess the effects of bacterial infection or LPS exposure on the development of allergic airways disease (AAD)⁴. AAD models of bacterial infection have highlighted a range of mechanisms contributing to pathogenesis, including induction of neutrophilic inflammation, potentiation of T_H2 responses and induction of T_H1 and/or T_H17 responses¹⁴ in infected airways. Most studies assessing the pathogenic role of LPS have focussed on the effects of pre-exposure on subsequent allergen induced induction of AAD¹⁵ or co-exposure during allergen sensitisation and/or aerochallenge¹⁶⁻¹⁸. LPS co-exposure has been demonstrated to modulate the local immune response, through activation of T_H1 and/or T_H2 responses and can induce steroid-resistant inflammation. However, there is only limited data on the effects of LPS exposure in the lung after allergen challenge and establishment of Type-2 AAD, which more appropriately models disease exacerbations in individuals with existing asthma.

Asthma exacerbations often occur on the background of an underlying Type-2 inflammatory response. In the current study, we have generated a model of LPS-induced exacerbation on the background of pre-existing AAD. This builds on our previous work demonstrating that prolonged AHR persists for at least 1 week after allergen challenge in models of ovalbumin (OVA) induced AAD¹⁹. Importantly, this prolonged AHR is associated with AHR originating specifically from the airways (like asthma) and activation of pulmonary macrophages¹⁹, which we hypothesised may be modulated by allergen-associated LPS exposure. We now demonstrate that LPS administration to mice with pre-existing type-2 AAD led to an exacerbation characterised by increased AHR and dampened responsiveness to dexamethasone (DEX) treatment. Exacerbation was accompanied by significant infiltration of neutrophils (24hrs) and macrophages into the lung, increased inflammatory cytokine production and persistence of Type-2-inflammation (eosinophils, IL-5, IL-13 and eotaxin). Macrophage depletion or neutralisation of IL-13 (or genetic IL-13-deficiency) after allergen challenge suppressed AHR and the severity of LPS induced inflammatory response and background Type-2 inflammation. These investigations provide a model of bacterial-induced exacerbations of pre-existing Type-2 AAD and demonstrate that both activated pulmonary

macrophages and IL-13 are required for LPS-induced exacerbation. Thus, Type-2 inflammatory factors and macrophages may cooperate to regulate endotoxin-induced exacerbations of asthma. Characterising the interplay between underlying chronic Type-2 responses in asthma and activation of innate immune responses by infection may improve our understanding of the mechanisms of exacerbations and identify new approaches to treatment.

METHODS

Mice

Adult male wild type mice (6-8 weeks old) and IL-13 gene-deficient mice (IL-13^{-/-} mice) on a BALB/c background were obtained from the University of Newcastle Animal Services Unit and experiments performed in the Hunter Medical Research Institute (HMRI) animal facility, under specific pathogen-free conditions. IL-13^{-/-} mice have been backcrossed onto a BALB/c background for greater than 10 generations. All protocols were approved by the local Animal Care and Ethics Committee (The University of Newcastle; Approval Number A-2013-317).

Experimental procedures

As previously described²⁰, mice were systemically sensitized by intra-peritoneal (i.p. day 0) injection of chicken egg ovalbumin (OVA, 50 µg; Sigma) or saline control Alhydrogel (1 mg; Sigma) in 200µl sterile saline. Mice were then aerosol-challenged with OVA (10 mg/ml in 0.9% saline) for 30 min/day on days 13-16. Where indicated, mice were instilled intratracheally (i.t.) with LPS (50 ng; Sigma-Aldrich) under Aflaxan anaesthetic (1:4; Jurox) in PBS, on days 19 and 21. Endpoints were assessed on day 24.

To assess responsiveness to corticosteroid treatment, mice were treated with DEX (1mg/kg i.p.; Sigma-Aldrich) on days 19 and 21. To deplete macrophages, mice were treated with 2-Chloroadenosine (2-CA, 50µl of 1mM, i.t.; Sigma-Aldrich) or PBS control on days 22 and 23, as previously described¹⁹. To neutralize IL-13, mice were injected with anti-IL-13 antibody (150µg, i.p, clone eBio1316H; eBioscience) or isotype control (rat IgG1_K, clone eBRG1, eBioscience) on days 19 and 21.

Lung function

Airway resistance (Raw) in response to methacholine (MCh, Sigma-Aldrich) was measured using a Flexivent apparatus (FX1 system; Scireq)²⁰. Briefly, mice were anesthetized by injection of xylazine (2 mg/ml, i.p.; Troy Laboratories), ketamine (40 mg/ml; Parnell) and PBS (5:4:1). A cannula was inserted into the trachea and mice were ventilated with a tidal volume of 8 ml/kg at a frequency of 450 breaths/minute. Mice were challenged with saline aerosol followed by increasing concentrations of Methacholine (0.1, 0.3, 1, 3, 10 and 30

mg/ml). Measurements were excluded if the coefficient of determination was lower than 95%. Airway resistance was recorded and presented as percentage increase over baseline.

Bronchoalveolar lavage fluid (BALF) differential cell counts

BALF was collected immediately following lung function measurements, as previously described²¹. The left lobe of the lung was tied off and the right lung was flushed twice with 700µl Hanks Buffered Saline Solution (HBSS; Invitrogen). BALF samples were centrifuged to pellet cells and supernatant stored at -80°C for protein analysis. Red blood cells were lysed in hypotonic red cell lysis buffer and remaining cells cytospun onto glass slides. Differential leukocyte counts were determined based on morphological criteria by light microscopy (×100) on May-Grunwald and Giemsa-stained slides, counting of 300 cells/slide/sample.

Macrophage isolation and treatment

Mouse pulmonary macrophages, which include alveolar macrophages, interstitial macrophages and monocytes, were isolated from mouse lungs as previously described²². Briefly, lung tissues were minced and forced through a 70µm cell strainer. Macrophages were purified by density gradient centrifugation (Histopaque-1083; Sigma-Aldrich) and seeded onto tissue culture-treated 6 well plates at 37°C at a concentration of 1×10^6 cells/ml in DMEM containing 20% fetal calf serum (FCS). After 3 h, >95% of adherent cells were macrophages. Where indicated, isolated primary macrophages were collected into Trizol and stored at -80°C for RNA analysis.

In specified experiments, purified macrophages were isolated from naïve mice and cultured *in vitro*. Isolated macrophages were cultured in fresh complete Dulbecco's Modified Eagle Medium (DMEM) supplemented with 20% FCS, 2 mM L-glutamine, 0.1 mM sodium pyruvate, 20 mM HEPES and 2-mercaptoethanol (β-ME). Where indicated, macrophages were incubated with recombinant murine IL-13 (100 ng/ml; Lonza) for 16 hours, then exposed to LPS (100 ng/ml; Sigma) and incubated for a further 12 hours. Cells were collected into Trizol and stored at -80°C for RNA analysis and supernatant were collected and stored at -80°C for protein analysis.

RNA isolation, reverse-transcription (RT)-PCR and quantitative PCR (qPCR) assessment

As previously described²³, lung tissue was immediately placed in RNAlater (Ambion, Austin, TX) and stored at -80°C for further analysis. RNA was isolated using Trizol reagent, following manufacturer instructions (Invitrogen Life Technologies), followed by phenol-chloroform separation and isopropanol precipitation. RNA samples were quantified on a

NanoDrop 1000 spectrophotometer (ND-1000; Thermo Scientific). cDNA was synthesised using random hexamer primers and MMLV reverse transcriptase (Invitrogen) on a T100 thermal cycler (Bio-Rad). Relative qRT-PCR quantification was performed on a Viia7 real-time PCR machine (Life Technologies), using SYBR reagents with following primer sequences: 5'-AGGCCAGACTTTGTTGGATTGAA-3' (forward) and 5'-CAACTTGCGCTCATCTTAGGCTTT-3' (reverse) for HPRT; 5'-CTGTTGCTGCTACCCTTGCTT-3' (forward) and 5'-CACTCCTGGCAATCGAGATTC-3' (reverse) for IL-27 (p28); 5'-GTCTACTGAACTTCGGGGTGATCG-3' (forward) and 5'-AGCCTTGTCCTTGAAGAGAACC-3' (reverse) for TNF α ; 5'-TCTTGAAAGACAATCAGGCCATCA-3' (forward) and 5'-GAATCAGCAGCGACTCCTTTTCC-3' (reverse) for IFN γ ; 5'-AGAGCCAGACGGGAGGAAG-3' (forward) and 5'-CCAGCCTACTCATTGGGATC-3' (reverse) for MCP-1; 5'-AGCTGAGCAACATCACACAAGACC-3' (forward) and 5'-TGGGCTACTTCGATTTTGGTATCG-3' (reverse) for IL-13; 5'-CGGCCGGAGAAAGTTGGTCCC-3' (forward) and 5'-GCACACCCGCCTGGTATGTCC-3' (reverse) for Muc5ac. Thermal cycling conditions consisted of an initial denaturing step (95°C: 3 min) followed by 40 cycles of denaturing (95°C, 5 s) and annealing (60°C, 30 s). All mRNA levels were normalized to hypoxanthine-guanine phosphoribosyl transferase (HPRT).

Flow cytometry analysis

Single-cell suspensions from lungs were stimulated with PMA (50 ng/ml) and ionomycin (1 μ g/ml) followed by Brefeldin A (5 μ g/ml) for a total of 5 hours of incubation. Cells were then washed in PBS and resuspended in ZombieYellow Fixable Viability Kit (BioLegend) for exclusion of dead cells. Fc receptor block (CD16/32) was added followed by anti-mouse CD3, CD4, CD45, CD90.2, ICOS and ST2 antibodies (BD Pharmingen, BioLegend) and washed with FACS buffer (PBS + 1% FCS + 2 mM EDTA) twice. Cells were then incubated on ice in 4% paraformaldehyde, permeabilized using BD Perm/Wash buffer (BD Pharmingen) and stained for intracellular marker (anti - IL-13). Analysis was performed using FACS Fortessa (BD Pharmingen). IL-13-producing ILC2 cells are defined as CD45⁺Lineage⁻ICOS⁺CD90.2⁺ST2⁺IL-13⁺.

ELISA

Lung tissue was disrupted in radioimmunoprecipitation assay buffer (RIPA; Sigma-Aldrich) with protease/phosphatase inhibitor cocktail (Cell Signaling Technology), on a Tissuelyser LT tissue disruptor (Qiagen) at 50 Hz for 5 minutes and stored at -80°C. Levels of TNF α , IFN γ , IL-5, IL-13, IL-27 (p28), eotaxin-1 and MCP-1 were assessed using ELISA kits

(DuoSet, R&D Systems) according to manufacturer's instructions. Total protein concentrations were determined using a BCA Protein Assay kit (Thermo Scientific). Individual cytokine levels were normalised to total protein.

Lung histology

Lung sections presenting central and peripheral airways were fixed in 10% formalin buffer for 24h, then transferred to 70% ethanol. The fixed tissue was embedded in paraffin and cut into 8µm sections. After heat-affixing the section to the glass slide, the slides were immersed in xylene to remove the paraffin from the section and stained with AB-PAS (Alcian Blue/Periodic Acid-Schiff) for recognition of mucus secreting cells (MSC) or Carbol's Chromotrope-hematoxylin to detect eosinophils. Peribronchial eosinophil numbers were determined by counting the number of eosinophils surrounding the airways in 20 high power fields (HPF×1000) using a green lens filter (Olympus, Japan). To count mucus-secreting cells, we identified intact airways and quantified cells at 100x magnification under oil immersion. A gradicule was used along the edge of the airway. Pink-staining MSCs were counted in 10 gradicule areas and data were then averaged to provide results in cells/100 µm.

Statistical analysis

Statistical analysis was performed using Prism version 6.0 (GraphPad Software, USA). Two-way ANOVA was used to identify differences between two or more experimental groups, and Student's unpaired t-tests was used where comparison were made between two treatment groups. All results are presented as means ± SEM and *P* values < 0.05 were considered statistically significant.

RESULTS

LPS administration exacerbates AHR, increases lung macrophage numbers and attenuates steroid-responsiveness.

To establish a mouse model of LPS-induced exacerbation of AHR and airway inflammation, BALB/c WT mice were sensitized with OVA (or PBS control) in Alum and exposed to aerosolised OVA to induce AAD. Three days after the final OVA aerosol challenge, some groups were treated with LPS (OVA/LPS) or LPS + DEX (OVA/LPS/DEX) on days 19 and 21 and lung function and inflammatory responses were assessed on day 24 (Fig. 1).

OVA-sensitized/challenged (OVA) mice exhibited significantly increased AHR in response to increasing doses of methacholine, compared with control PBS-sensitized/OVA-challenged (PBS) mice (Fig. 1A), as previously described¹⁹. This response is reduced in comparison to day 17, where allergic airway inflammation (type 2 cytokine levels, eosinophil and Th2 cell numbers) is most prominent after allergen challenge (data not shown). LPS

administration exacerbated AHR in OVA-treated mice (OVA/LPS), as compared to that in OVA-treated mice (OVA) (Fig. 1A). We previously demonstrated that DEX treatment of OVA-treated mice in this model effectively suppressed AAD and decreased AHR to baseline (PBS) levels on day 17 and at this time point²⁰. Conversely, following LPS administration, DEX treatment (OVA/LPS/DEX) failed to suppress AHR (Fig. 1A). These findings indicate that LPS exposure exacerbates AHR in mice with pre-existing Type-2 AAD, which is resistant to systemic corticosteroid treatment.

Differential BALF immune cell numbers were quantified to characterize the extent and type of inflammatory infiltration (Fig. 1B). OVA-treated mice had increased neutrophil, macrophage, eosinophil and lymphocyte numbers in BALF, compared to the PBS control group (Fig. 1B). LPS exposure induced a transient increase in neutrophils at day 22 (data not shown) and increased macrophage numbers at day 24 (Fig. 1B), compared to OVA-treated mice. Eosinophil numbers were decreased following LPS exposure (Fig. 1B). We previously demonstrated that DEX treatment of OVA-treated mice decreased total inflammatory cell numbers in BALF, by reducing eosinophil and lymphocyte numbers at the same timepoint²⁰. DEX treatment following LPS exposure further reduced eosinophil numbers but had no effect on macrophage numbers (Fig. 1B), which remained elevated compared to OVA-treated mice. These findings indicate that LPS exacerbation of pre-existing AAD is accompanied by a corticosteroid-resistant increase in macrophage numbers in BALF.

LPS administration to OVA-treated mice induces steroid-resistant inflammatory cytokine production but has no effect on T_H2-associated cytokines.

We next assessed the effect of LPS exposure of OVA-treated mice on cytokine production within the lung. As previously described²⁴, OVA-treated mice had increased protein levels of IL-5, IL-13, eotaxin-1, IL-27(p28), IFN γ and MCP-1 in the lung (Fig. 1D). DEX treatment significantly decreased IL-5 and IL-13 production in OVA-treated mice. Following LPS administration, levels of IL-5 and IL-13 remained elevated, but were not changed compared with the OVA group (Fig. 1D). Of note, increased levels of IL-13 were not altered by DEX treatment in OVA/LPS/DEX-treated mice. The same was observed for muc5ac expression in lung (Fig. 1C), in the OVA/LPS/DEX treated mice.

We also quantified the production of inflammatory cytokines, which have previously been implicated in asthma exacerbations. TNF α levels were unchanged in OVA-treated mice, compared to PBS controls (Fig. 1D). We previously demonstrated that DEX treatment has no effect on the levels of eotaxin, IL-27p28, TNF α and IFN γ in OVA-treated mice²⁰⁻²⁵. Eotaxin-1, IL-27 (p28), TNF α , IFN γ and MCP-1 protein levels were all increased in OVA/LPS-

treated mice compared to those in OVA-treated mice (Fig. 1D). DEX treatment had no effect on the protein levels of IL-27 (p28), TNF α or IFN γ induced by LPS administration (Fig. 1D). Thus, LPS exposure increased the production of multiple inflammatory cytokines, without altering levels of IL-5 and IL-13, and this increase in cytokine (IL-27 (p28), TNF α and IFN γ) production was steroid-resistant.

Pulmonary macrophages from OVA/LPS-treated mice upregulate expression of inflammatory cytokines.

Based on the proposed role of pulmonary macrophages in steroid-resistant asthma exacerbations²⁶ and the increase in BALF macrophage numbers following LPS exposure, we also assessed expression of innate cytokines elevated in the lung of purified primary lung macrophages isolated at day 24 (as it is not experimentally practical to measure protein levels in these cells). TNF α expression was only significantly increased in macrophages isolated from OVA-treated mice (Fig. 2). By contrast, in macrophages isolated from OVA/LPS-treated mice, the levels of IL-27 (p28), TNF α , IFN γ and MCP-1 expression were all increased (Fig. 2), compared to the OVA-treated group. DEX treatment partially reduced expression of IL-27 (p28), TNF α and MCP-1 and completely abolished expression of IFN γ , compared with macrophages from OVA/LPS-treated mice (Fig. 2). These data provide evidence that pulmonary macrophages upregulate expression of inflammatory cytokines following OVA/LPS stimulation and expression is resistant to steroid treatment.

Macrophage depletion suppresses AHR and reduces pro-inflammatory cytokine levels during LPS-induced exacerbation of AAD.

To determine whether macrophages have a functional role in disease exacerbation in our model, we depleted macrophages using the established method of 2-CA administration [a purine analogue that specifically depletes pulmonary macrophages^{19,27}]. 2-CA administration reduced OVA/LPS-induced AHR, compared with vehicle-treated controls, to the same level observed in OVA-treated mice (Fig. 3A). 2-CA administration decreased macrophage and lymphocyte numbers in the OVA/LPS group (Fig. 3B). BALF neutrophil and eosinophil numbers were not affected by 2-CA administration (Fig. 3B). 2-CA administration decreased the expression of muc5ac in OVA and OVA/LPS treated mice (Fig. 3C). Furthermore, 2-CA significantly decreased the levels of IL-13, IL-27 (p28), TNF α and IFN γ in OVA/LPS group, compared to those in OVA treated group, as assessed by ELISA (Fig. 3D). In contrast, 2-CA administration had no effect on IL-5, eotaxin-1 or MCP-1 levels (Fig. 3D). Together, these findings indicate that macrophages are required for LPS-induced exacerbation of AHR and

that macrophage ablation results in a reduction in the levels of a subset of cytokines (IL-13, IL-27 (p28), TNF α and IFN γ) in the lung.

IL-13 blockade suppresses AHR, inflammatory cell recruitment and pro-inflammatory cytokine levels during LPS-induced exacerbation.

IL-13 plays a key role in contributing to induction of AHR and goblet cell hyperplasia²⁸. While IL-13 expression was not increased following LPS exposure, we did note that IL-13 levels were significantly elevated in all groups (as compared to that in PBS group) and that steroid treatments did not suppress IL-13 levels after LPS exposure (by contrast to OVA treated group) (Fig. 1). We next assessed the potential role of IL-13 crosstalk with LPS activation of the innate macrophage driven immune response by employing both IL-13^{-/-} mice and treatment with neutralising anti-IL-13 antibodies just prior to induction of the exacerbation by LPS.

Induction of AHR was suppressed in IL-13^{-/-} mice, compared to that in WT controls, following either OVA sensitisation/challenge alone or subsequent LPS exposure (Fig. 4A). In differential counts, all inflammatory cell types were reduced in IL-13^{-/-} mice following OVA- and OVA/LPS- treatments, compared to the respective WT responses to these treatments (Fig. 4B). The expression of muc5ac was not altered in OVA or OVA/LPS groups in IL-13^{-/-} mice when compared to PBS treated controls (Fig. 4C). Protein levels of IL-13 were not detected after OVA or OVA/LPS exposure in IL-13^{-/-} mice (Fig. 4D). Further, eotaxin-1, IL-27 (p28), IFN γ and MCP-1 levels were lower in IL-13^{-/-} mice following OVA/LPS treatment, compared to the respective WT responses to these treatments (Fig. 4D). IL-5 and TNF α levels were not increased in IL-13^{-/-} mice (Fig. 4D). These findings provide evidence that IL-13 production is required for OVA and subsequently LPS-induced exacerbation of AHR, and severity of inflammation and cytokine production. However, these experiments do not differentiate between the role of IL-13 during the initial induction of AAD by allergen exposure and the LPS-induced exacerbation phase.

To assess the functional role of IL-13 during LPS induced exacerbation, we administered neutralising antibody against IL-13 after AAD was established (Fig. 5). IL-13 neutralization abolished OVA/LPS-induced AHR, compared with isotype control (Fig. 5A). Further, anti-IL-13 treatment reduced neutrophil and eosinophil numbers and to a lesser extent macrophage and lymphocyte numbers (Fig. 5B). The expression of muc5ac was significantly reduced in the OVA/LPS group that was treated with anti-IL-13 when compared to OVA/LPS group (Fig. 5C). Blocking IL-13 decreased IL-5, IL-13 and MCP-1 production in lung tissue (Fig. 5D). In contrast, TNF α , IL-27 (p28) and eotaxin-1 levels were unaffected

by anti-IL-13 administration (Fig. 5D). These results indicate that IL-13 is required during the LPS induced exacerbation phase, to maintain the Type-2 response and that this cytokine plays a cooperative role with macrophages (Fig. 3) in the induction of AHR and airway inflammation.

IL-13 and LPS stimulation of isolated naïve alveolar macrophages modulates inflammatory cytokine production.

Our intervention studies indicate roles for both pulmonary macrophages and IL-13 in LPS-induced exacerbations. Next, we assessed the direct response of naïve pulmonary macrophages to LPS and/or IL-13 stimulation *in vitro*. LPS stimulation alone induced marked expression of IL-27 (p28), TNF α , IFN γ and MCP-1 in primary macrophages (Fig. 6). IL-13 stimulation alone had limited effect on cytokine expression when compared to PBS. Interestingly, IL-13/LPS co-stimulation further increased MCP-1 expression, compared to LPS or IL-13 stimulation alone (Fig. 6), but had no further effect on IL-27, TNF α or IFN γ expression. These findings provide evidence that IL-13 and LPS modulate inflammatory cytokine expression by pulmonary macrophages and indicate additive effects of IL-13/LPS co-stimulation on chemokine expression.

Steroid insensitivity in Group 2 ILC cells in the lungs following LPS exposure.

To further understand the pathways involved in DEX resistance to LPS induced exacerbation (OVA/LPS group) we stain lung cells with antibody markers for ILC2s and enumerated their numbers by flow cytometry (Fig. 7A). We observed a significantly increased amount of ILC2s in the OVA treated group but numbers were not increased by LPS exposure (OVA/LPS group), and numbers were not suppressed by DEX treatment (Fig. 7A). The protein levels of IL-25, IL-33 and TSLP were also increased by OVA treatment group by not further by LPS exposure and levels were not reduced by DEX treatment (Fig. 7B).

Effect of DEX, 2-CA and anti-IL-13 treatments on histopathological changes in lung tissues during OVA/LPS-induced allergic asthma.

To characterize the changes in lung histopathology caused by OVA and OVA/LPS exposure, and the effects of different treatments (DEX, 2-CA and anti-13), we stained the lung samples with AB-PAS and CR (Fig 8A). Mucus producing cells and eosinophils numbers were then quantified (Fig. 8B). Goblet cell hyperplasia, mucus hypersecretion and increased numbers of eosinophils were observed after OVA treatment but not further increased by LPS exposure (OVA/LPS). DEX treatment of the OVA/LPS group significantly reduced eosinophil numbers but not the levels of PAS positive cells. By contrast the OVA/LPS group treated with 2-CA or anti-IL-13 had diminished levels of PAS-positive cells. Evidence of

inflammatory cell infiltration was observed in CR stained samples. OVA, OVA/LPS and DEX treated groups showed an extensive cell infiltration around the airways and blood vessels when compared to control group. Mice treated with 2-CA and anti-IL-13 after OVA or OVA/LPS exposure had significantly reduced inflammatory infiltrates.

DISCUSSION

Respiratory infections are clinically associated with asthma exacerbations, which negatively contribute to disease burden, healthcare utilisation and associated healthcare costs for asthma³. Further, epidemiological studies have linked endotoxin (LPS) exposure to worsened disease outcomes and exacerbations^{5,6}. There is also growing interest in the effects of the microbiota and microbial products on disease pathogenesis in asthma^{7,8}. We provide evidence that LPS exposure exacerbates AHR in the context of pre-established AAD, which models the effect of microbial exposure in patients with pre-existing allergic asthma. LPS-induced exacerbations were steroid-resistant, which is an important clinical feature of severe asthma and disease exacerbations²⁹. In functional studies, we provide evidence that pulmonary macrophages and IL-13 are required for disease exacerbation. These data highlight that integration of IL-13 and innate immune responses predispose to steroid-resistant inflammatory pathways and AHR associated with LPS induced exacerbation.

The effects of LPS or bacterial exposure on AAD have previously been assessed in a range of protocols^{4, 30, 31}. Repeated administration of high-dose LPS alone (4 consecutive days; 100 µg) in the absence of pre-existing AAD induced AHR³². This may be linked to neutrophil recruitment and/or macrophage activation. However, at the dose used in the current study (0.05 µg per exposure to naïve WT mice), LPS alone transiently increases neutrophil numbers (not shown) but does not induce AHR²¹. Early-life LPS exposure has been shown to protect from development of AAD, by dampening T_H2 responses¹⁵. However, LPS exposure during allergen sensitisation or challenge was shown to exacerbate disease, through increased T_H1 activation and/or increased mast cell activation and T_H2 immune responses^{16,17}. Similarly, *M. pneumoniae* bacterial lung infection prior to allergen sensitisation protected from AAD induction, while infection after sensitisation promoted Type-2 inflammation and severity of inflammation³³. In contrast, *Chlamydia muridarum* lung infection before induction of AAD resulted in an allergen driven mixed T_H1/T_H2 response and increased IL-13 production³⁴ or increased neutrophil accumulation and T_H1/T_H17-associated gene expression³⁵. *H. influenzae* infection prior to allergen sensitisation led to chronic bacterial infection and persistent steroid-resistant, neutrophilic airways disease¹⁴.

Furthermore, *H. influenzae* infection worsened disease through increased T_H17 responses in an exacerbation model³⁶. LPS co-exposure during allergen challenge has also been shown to dampen steroid-responsiveness¹⁸. We now provide evidence that LPS exposure after establishment of AAD activates innate proinflammatory pathways and activates macrophage to induce AHR in concert with IL-13 regulated Type-2 responses.

In our model, LPS-induced exacerbation was associated with increased expression of a number of inflammatory cytokines that have been implicated in asthma severity and/or exacerbations. Notably, IL-27, TNF α , IFN γ and IL-13 were increased in lung tissue, and expression persisted despite DEX treatment (Fig. 1C). IL-27 is a heterodimeric cytokine consisting of two subunits (EBI3 and p28), which promotes T_H1 polarization and IFN γ production³⁷. IFN γ is associated with Type-1 anti-viral responses and bacterial infections, and has been suggested as a potential target in chronic asthma³⁸. IFN γ expression is increased in individuals with severe asthma, compared to moderate disease³⁹. Increased IL-27 and IFN γ production may also reflect a shift towards T_H1 polarisation, from a T_H2 allergic background. We previously demonstrated that administration of LPS and recombinant IFN γ into the lungs is sufficient to induce steroid-resistant AHR in a short-term model²¹. Further, administration of recombinant IL-27 and IFN γ to mice also induced steroid-resistant AHR, through activation of pulmonary macrophages⁴⁰. IFN γ and IL-27 expression was also found to be increased in sputum samples from patients with neutrophilic asthma, a patient subset that are typically less responsive to corticosteroid treatment⁴⁰. TNF α promotes AHR induction and neutrophil recruitment (as shown in animal and clinical studies) in mouse models and control human subjects⁴¹⁻⁴³. TNF α is significantly increased in samples from individuals with severe asthma^{44,45}, individuals with neutrophilic asthma²⁰ and following exacerbation, where levels were associated with exacerbation severity⁴⁶. Clinical trials targeting TNF α in asthma have yielded some positive results, with reduced exacerbation rates^{47,48}, improved quality of life, lung function⁴⁴ and asthma control⁴⁹. However, trial outcomes have been variable and there are concerns about increased risk of infections following treatment. MCP-1 is produced by activated macrophages and promotes macrophage migration⁵⁰. We have detected heightened expression of MCP-1 in purified pulmonary macrophages from OVA/LPS treated mice and LPS/IL-13 stimulated naive macrophages (Fig 2 & 6). MCP-1 expression is increased by virus infection in individuals with asthma⁵¹ and in sputum from individuals with neutrophilic asthma²⁰. Interestingly, MCP-1 may not only regulate macrophage activation but also attract either Th2 cells or ILC2s into inflamed tissues, as these cells express receptor specific to this chemokine⁵². Of note, the latter two cells are the predominant cellular source of IL-13⁵³. We

identified increased expression of each of these molecules in our LPS-induced exacerbation model. However, the specific role each of these cytokines play in asthma exacerbation remains unclear.

A growing body of literature has highlighted functional roles for macrophages in steroid-resistant asthma^{16,54}. In line with a role for pulmonary macrophages in our model, we observed increased production of IL-27, IFN γ , TNF α and MCP-1 in primary pulmonary macrophages isolated from OVA/LPS-exposed mice (Fig. 2). By contrast to the Type 2 factors (IL-5, eosinophil and eotaxin, Fig. 1), DEX treatment did not affect the levels of IL-13, IL-27, IFN γ or TNF α production in mice treated with LPS (LPS/DEX group, Fig. 1). In intervention studies, macrophage depletion markedly reduced LPS induced AHR, which was associated with reduced IL-13, TNF α and IFN γ production in lung tissue. We have previously demonstrated that macrophage depletion also reduces AHR induced by OVA treatment alone at the examined time point, with no effect on the numbers of eosinophils or mucus-secreting cells¹⁹. Further, we have observed similar role for macrophages in regulating AHR induced by exposure of the lung to both IFN γ /LPS- or RSV- induced exacerbation of pre-existing AAD^{19,20}. In the current study we demonstrate that macrophages play a critical role in LPS-induced exacerbations of AHR and inflammation in concert with IL-13 and possibly downstream of these factors through processes involving TNF α and/or IFN γ .

Key roles for Type-2 inflammation (including eosinophils, IL-5, IL-13 and eotaxin) are well-recognised in the pathogenesis of AAD⁵⁵. However, the role of Type-2 inflammation in exacerbations is less clear. LPS co-exposure during allergen challenge has been shown to increase¹⁷ or decrease¹⁸ Type-2 cytokine levels. IL-13 in particular plays a key role in allergic asthma pathogenesis by inducing AHR and promotes eosinophil recruitment and goblet cell hyperplasia²⁸. IL-13 administration to the lung alone is sufficient to induce AHR⁵⁶ and viral delivery of IL-13 to the lung induces mucus hypersecretion, neutrophilia and AHR, which is resistant to DEX treatment⁵⁷. Further, IL-13 over-expressing transgenic mice develop spontaneous AHR and mixed granulocytic lung inflammation⁵⁸. In our model, LPS exposure did not alter levels of IL-5 or IL-13. However, treatment with DEX (LPS/DEX group) reduced IL-5 (and eotaxin) but not IL-13 levels and the reduction in IL-5 and eotaxin correlated with decreased numbers of eosinophils in BALF. Notably, IL-13 levels remain elevated after corticosteroid treatment in individuals with steroid-insensitive asthma⁵⁹. By contrast, DEX treatment reduced Type-2 cytokine expression, including IL-13 production, in the absence of LPS exposure²⁰, and as demonstrated in our Fig. 1. Thus, persistent and steroid-resistant IL-13 expression following LPS exposure may contribute to the

pathogenesis. It should be noted that steroid resistance in asthma is heterogenous⁶⁰. For instance, the expression of defective glucocorticoid receptors in non-hematopoietic lung cells is closely associated with steroid resistance in a mouse model of allergic asthma⁶¹. Persistent eosinophilic inflammation is also considered as one hallmark of a subgroup of severe asthma⁶².

IL-13 production was required for LPS-induced exacerbations in our model, as IL-13^{-/-} mice failed to develop AHR and airway inflammation was significantly attenuated. These findings were consistent with previous observations that IL-13-deficient mice fail to develop AAD, including in the context of *Chlamydia*-induced persistent AHR⁶³. To confirm a role for IL-13 in the exacerbation phase, we administered IL-13 neutralising antibodies after the establishment of AAD (on days 19 and 21). Neutralization of IL-13 inhibited AHR, reduced infiltrating inflammatory cell numbers (neutrophils, eosinophils and lymphocytes) and decreased IL-5, IFN γ , MCP-1 and IL-13 protein levels. Numerous studies have demonstrated that genetic or pharmacologic blockade of IL-13 protects from induction of characteristic features of AAD, including AHR, inflammation and airway remodelling²⁸. In clinical trials, treatment with lebrikizumab (anti-IL-13) significantly reduced asthma exacerbations⁶⁴. However, targeting IL-13 has also had only limited effects on exacerbations⁶⁵. Further stratification of patients and identification of the exacerbation trigger (e.g. viral and/or bacterial) may demonstrate greater therapeutic benefit. Our findings provide evidence that blocking IL-13 and/or macrophage function can also inhibit LPS-induced steroid-resistant exacerbations of AAD, suggesting a potential role for this cytokine in bacterial regulated components of exacerbations.

Understanding how microbial bioproducts such as LPS/endotoxin contribute to the progression and exacerbations of asthma will provide insights in new approaches to treat disease. Here we demonstrate that LPS induces an exacerbation of AAD characterised by steroid resistant AHR and increased numbers of macrophages in the lung activation of innate inflammatory pathways and elevated levels of IL-13. Depletion of macrophages suppressed AHR and reduced IL-13, IL-27, TNF α and IFN γ levels in the lung. Further, IL-13 was required for LPS-induced exacerbations and blocking IL-13 function reduced IL-5, eotaxin, MCP-1 and IFN γ production. Our findings provide support for a mechanism whereby LPS exposure in the context of Type-2 AAD exacerbates disease through parallel activation of innate inflammatory pathways in macrophages and the induction of a steroid-resistant IL-13 pathway. These findings provide further insight into the mechanisms underlying LPS and

bacterial-induced exacerbations, which are relevant to existing therapies (e.g. anti-IL-13) and identify further potential targets for future intervention.

FOOTNOTES

§Corresponding Author Information: Paul S. Foster & Ming Yang, Hunter Medical Research Institute, Lot 1 Kookaburra Circuit, New Lambton Heights, NSW 2305, Australia. Telephone (+61-2) 4042-0163 or (+61-2) 4042-0183, FAX: (+61-2) 4042-0024, Email Address: Paul.Foster@newcastle.edu.au or Ming.Yang@newcastle.edu.au

Grant Support: This work was supported by project grants from the National Health and Medical Research Council (NHMRC) of Australia. SM was supported by fellowships from the Canadian Institutes of Health Research (CIHR) and the University of Newcastle. PMH is funded by a Fellowship and grants from the NHMRC (1079187, 1175134).

REFERENCES

1. Agache ICA, Jutel M, Virchow JC. Untangling asthma phenotypes and endotypes. *Allergy* 2012;67:835-846.
2. Graham LM, Eid N. The impact of asthma exacerbations and preventive strategies. *Curr Med Res Opin* 2015;31:825-835.
3. Busse WW, Lemanske RF Jr., Gern JE. Role of viral respiratory infections in asthma and asthma exacerbations. *Lancet* 2010;376:826-834.
4. Maltby S, Tay HL, Yang M, Foster PS. Mouse models of severe asthma: Understanding the mechanisms of steroid resistance, tissue remodelling and disease exacerbation. *Respirology*. 2017; 22(5):874-885.
5. Michel O, Kips J, Duchateau J, Vertongen F, Robert L, Collet H, Pauwels R, Sergysels R. Severity of asthma is related to endotoxin in house dust. *Am J Respir Crit Care Med* 1996;154: 1641-1646.
6. Thorne PS, Kulhankova K, Yin M, Cohn R, Arbes SJ Jr., Zeldin DC. Endotoxin exposure is a risk factor for asthma: the national survey of endotoxin in United States housing. *Am J Respir Crit Care Med* 2005;172: 1371-1377.
7. Maslowski KM, Mackay CR. Diet, gut microbiota and immune responses. *Nat Immunol* 2011;12: 5-9.
8. Singanayagam A, Ritchie AI, Johnston SL. Role of microbiome in the pathophysiology and disease course of asthma. *Curr Opin Pulm Med* 2017;23: 41-47.

- 534 9. Hilty M, Burke C, Pedro H, Cardenas P, Bush A, Bossley C, Davies J, Ervine A, Poulter L,
535 Pachter L, Moffatt MF, Cookson WO. Disordered microbial communities in asthmatic
536 airways. *PLoS One* 2010;5: e8578.
- 537 10. Huang YJ, Nelson CE, Brodie EL, Desantis TZ, Baek MS, Liu J, Woyke T, Allgaier M,
538 Bristow J, Wiener-Kronish JP, Sutherland ER, King TS, Icitovic N, Martin RJ, Calhoun WJ,
539 Castro M, Denlinger LC, Dimango E, Kraft M, Peters SP, Wasserman SI, Wechsler ME,
540 Boushey HA, Lynch SV. Airway microbiota and bronchial hyperresponsiveness in patients
541 with suboptimally controlled asthma. *J Allergy Clin Immunol* 2011;127: 372-381 e371-373.
- 542 11. Taylor SL, Leong LEX, Choo JM, Wesselingh S, Yang IA, Upham JW, Reynolds PN,
543 Hodge S, James AL, Jenkins C, Peters MJ, Baraket M, Marks GB, Gibson PG, Simpson JL,
544 Rogers GB. Inflammatory phenotypes in patients with severe asthma are associated with
545 distinct airway microbiology. *J Allergy Clin Immunol*. 2017; 141(1):94-103
- 546 12. Green BJ, Wiriyaichaiorn S, Grainge C, Rogers GB, Kehagia V, Lau L, Carroll MP,
547 Bruce KD, Howarth PH. Potentially pathogenic airway bacteria and neutrophilic
548 inflammation in treatment resistant severe asthma. *PLoS One* 2014;9: e100645.
- 549 13. Janssen O, Schaumann F, Holz O, Lavae-Mokhtari B, Welker L, Winkler C, Biller H,
550 Krug N, Hohlfeld JM. Low-dose endotoxin inhalation in healthy volunteers--a challenge
551 model for early clinical drug development. *BMC Pulm Med* 2013;13: 19.
- 552 14. Essilfie AT, Simpson JL, Dunkley ML, Morgan LC, Oliver BG, Gibson PG, Foster PS,
553 Hansbro PM. Combined *Haemophilus influenzae* respiratory infection and allergic airways
554 disease drives chronic infection and features of neutrophilic asthma. *Thorax* 2012; 67: 588-
555 599.
- 556 15. Kuipers H, Hijdra D, De Vries VC, Hammad H, Prins JB, Coyle AJ, Hoogsteden HC,
557 Lambrecht BN. Lipopolysaccharide-induced suppression of airway Th2 responses does not
558 require IL-12 production by dendritic cells. *J Immunol* 2003;171: 3645-3654.
- 559 16. Delayre-Orthez C, Becker J, de Blay F, Frossard N, Pons F. Exposure to endotoxins
560 during sensitization prevents further endotoxin-induced exacerbation of airway inflammation
561 in a mouse model of allergic asthma. *Int Arch Allergy Immunol* 2005; 138: 298-304.
- 562 17. Murakami D, Yamada H, Yajima T, Masuda A, Komune S, Yoshikai Y.
563 Lipopolysaccharide inhalation exacerbates allergic airway inflammation by activating mast
564 cells and promoting Th2 responses. *Clin Exp Allergy* 2007;37: 339-347.
- 565 18. Komlosi ZI, Pozsonyi E, Tabi T, Szoko E, Nagy A, Bartos B, Kozma GT, Tamasi L,
566 Orosz M, Magyar P, Losonczy G. Lipopolysaccharide exposure makes allergic airway

inflammation and hyper-responsiveness less responsive to dexamethasone and inhibition of iNOS. *Clin Exp Allergy* 2006;36: 951-959.

19. Yang M, Kumar RK, Foster PS. Interferon-gamma and pulmonary macrophages contribute to the mechanisms underlying prolonged airway hyperresponsiveness. *Clin Exp Allergy* 2010; 40: 163-173.

20. Nguyen TH, Maltby S, Simpson JL, Eysers F, Baines KJ, Gibson PG, Foster PS, Yang M. TNF-alpha and Macrophages Are Critical for Respiratory Syncytial Virus-Induced Exacerbations in a Mouse Model of Allergic Airways Disease. *J Immunol* 2016; 196: 3547-3558.

21. Yang M, Kumar RK, Foster PS. Pathogenesis of steroid-resistant airway hyperresponsiveness: interaction between IFN-gamma and TLR4/MyD88 pathways. *J Immunol* 2009; 182: 5107-5115.

22. Li JJ, Tay HL, Maltby S, Xiang Y, Eysers F, Hatchwell L, Zhou H, Toop HD, Morris JC, Nair P, Mattes J, Foster PS, Yang M. MicroRNA-9 regulates steroid-resistant airway hyperresponsiveness by reducing protein phosphatase 2A activity. *J Allergy Clin Immunol*. 2015; 136(2):462-73

23. Maltby S, Hansbro NG, Tay HL, Stewart J, Plank M, Donges B, Rosenberg HF, Foster PS. Production and differentiation of myeloid cells driven by proinflammatory cytokines in response to acute pneumovirus infection in mice. *J Immunol* 2014; 193: 4072-4082.

24. Chung KF Targeting the interleukin pathway in the treatment of asthma. *Lancet* 2015; 386: 1086-1096.

25. Nguyen TH, Maltby S, Tay HL, Eysers F, Foster PS, Yang M. Identification of IFN-γ and IL-27 as Critical Regulators of Respiratory Syncytial Virus-Induced Exacerbation of Allergic Airways Disease in a Mouse Model. *The Journal of Immunology*. 2017; 200(1):237-247

26. Yang M, Kumar RK, Hansbro PM, Foster PS. Emerging roles of pulmonary macrophages in driving the development of severe asthma. *J Leukoc Biol* 2012; 91: 557-569.

27. Kubota Y, Iwasaki Y, Harada H, Yokomura I, Ueda M, Hashimoto S, Nakagawa M. Depletion of alveolar macrophages by treatment with 2-chloroadenosine aerosol. *Clin Diagn Lab Immunol* 1999; 6: 452-456.

28. Grünig G, Corry DB, Reibman J, Wills-Karp M. Interleukin 13 and the evolution of asthma therapy. *American journal of clinical and experimental immunology* 2012; 1: 20-27.

29. Singh AM, Busse WW. Asthma exacerbations. 2: aetiology. *Thorax* 2006; 61: 809-816.

30. Zhu Z, Oh SY, Zheng T, Kim YK. Immunomodulating effects of endotoxin in mouse models of allergic asthma. *Clin Exp Allergy* 2010; 40: 536-546.

31. Kumar RK, Herbert C, Foster PS. Mouse models of acute exacerbations of allergic asthma. *Respirology* 2016; 21: 842-849.
32. Starkhammar M, Kumlien Georen S, Swedin L, Dahlen SE, Adner M, Cardell LO. Intranasal administration of poly(I:C) and LPS in BALB/c mice induces airway hyperresponsiveness and inflammation via different pathways. *PLoS One* 2012; 7: e32110.
33. Chu HW, Honour JM, Rawlinson CA, Harbeck RJ, Martin RJ. Effects of respiratory Mycoplasma pneumoniae infection on allergen-induced bronchial hyperresponsiveness and lung inflammation in mice. *Infect Immun* 2003; 71: 1520-1526.
34. Horvat JC, Beagley KW, Wade MA, Preston JA, Hansbro NG, Hickey DK, Kaiko GE, Gibson PG, Foster PS, Hansbro PM. Neonatal chlamydial infection induces mixed T-cell responses that drive allergic airway disease. *Am J Respir Crit Care Med* 2007; 176: 556-564.
35. Kim RY, Horvat JC, Pinkerton JW, Starkey MR, Essilfie AT, Mayall JR, Nair PM, Hansbro NG, Jones B, Haw TJ, Sunkara KP, Nguyen TH, Jarnicki AG, Keely S, Mattes J, Adcock IM, Foster PS, Hansbro PM. MicroRNA-21 drives severe, steroid-insensitive experimental asthma by amplifying phosphoinositide 3-kinase-mediated suppression of histone deacetylase 2. *J Allergy Clin Immunol.* 2016; 139(2):519-532
36. Essilfie AT, Simpson JL, Horvat JC, Preston JA, Dunkley ML, Foster PS, Gibson PG, Hansbro PM. Haemophilus influenzae infection drives IL-17-mediated neutrophilic allergic airways disease. *PLoS Pathog* 2011; 7: e1002244.
37. Pflanz S, Timans JC, Cheung J, Rosales R, Kanzler H, Gilbert J, Hibbert L, Churakova T, Travis M, Vaisberg E, Blumenschein WM, Mattson JD, Wagner JL, To W, Zurawski S, McClanahan TK, Gorman DM, Bazan JF, de Waal Malefyt R, Rennick D, Kastelein RA.. IL-27, a heterodimeric cytokine composed of EBI3 and p28 protein, induces proliferation of naive CD4+ T cells. *Immunity* 2002; 16: 779-790.
38. Kumar RK, Webb DC, Herbert C, Foster PS. Interferon-gamma as a possible target in chronic asthma. *Inflamm Allergy Drug Targets* 2006; 5: 253-256.
39. Shannon J, Ernst P, Yamauchi Y, Olivenstein R, Lemiere C, Foley S, Cicora L, Ludwig M, Hamid Q, Martin JG. Differences in airway cytokine profile in severe asthma compared to moderate asthma. *Chest* 2008.133: 420-426.
40. Li JJ, Wang W, Baines KJ, Bowden NA, Hansbro PM, Gibson PG, Kumar RK, Foster PS, Yang M. IL-27/IFN-gamma induce MyD88-dependent steroid-resistant airway hyperresponsiveness by inhibiting glucocorticoid signaling in macrophages. *J Immunol* 2010; 185: 4401-4409.

- 634 41. Nakae S, Lunderius C, Ho LH, Schafer B, Tsai M, Galli SJ. TNF can contribute to
635 multiple features of ovalbumin-induced allergic inflammation of the airways in mice. *J*
636 *Allergy Clin Immunol* 2007; 119: 680-686.
- 637 42. Lukacs NW, Strieter RM, Chensue SW, Widmer M, Kunkel SL. TNF-alpha mediates
638 recruitment of neutrophils and eosinophils during airway inflammation. *J Immunol* 1995;
639 154: 5411-5417.
- 640 43. Thomas PS, Yates DH, Barnes PJ. Tumor necrosis factor-alpha increases airway
641 responsiveness and sputum neutrophilia in normal human subjects. *Am J Respir Crit Care*
642 *Med* 1995; 152: 76-80.
- 643 44. Berry MA, Hargadon B, Shelley M, Parker D, Shaw DE, Green RH, Bradding P,
644 Brightling CE, Wardlaw AJ, Pavord ID. Evidence of a role of tumor necrosis factor alpha in
645 refractory asthma. *N Engl J Med* 2006; 354: 697-708.
- 646 45. Howarth PH, Babu KS, Arshad HS, Lau L, Buckley M, McConnell W, Beckett P, Al Ali
647 M, Chauhan A, Wilson SJ, Reynolds A, Davies DE, Holgate ST. Tumour necrosis factor
648 (TNFalpha) as a novel therapeutic target in symptomatic corticosteroid dependent asthma.
649 *Thorax* 2005; 60: 1012-1018.
- 650 46. Manthei DM, Schwantes EA, Mathur SK, Guadarrama AG, Kelly EA, Gern JE, Jarjour
651 NN, Denlinger LC. Nasal lavage VEGF and TNF-alpha levels during a natural cold predict
652 asthma exacerbations. *Clin Exp Allergy* 2014; 44: 1484-1493.
- 653 47. Erin EM, Leaker BR, Nicholson GC, Tan AJ, Green LM, Neighbour H, Zacharasiewicz
654 AS, Turner J, Barnathan ES, Kon OM, Barnes PJ, Hansel TT. The effects of a monoclonal
655 antibody directed against tumor necrosis factor-alpha in asthma. *Am J Respir Crit Care Med*
656 2006.174: 753-762.
- 657 48. Edwards CJ, Polosa R. Study of infliximab treatment in asthma. *Am J Respir Crit Care*
658 *Med* 2007.175: 196; author reply 196-197.
- 659 49. Morjaria JB, Chauhan AJ, Babu KS, Polosa R, Davies DE, Holgate ST. The role of a
660 soluble TNFalpha receptor fusion protein (etanercept) in corticosteroid refractory asthma: a
661 double blind, randomised, placebo controlled trial. *Thorax* 2008. 63: 584-591.
- 662 50. Deshmane SL, Kremlev S, Amini S, Sawaya BE. Monocyte chemoattractant protein-1
663 (MCP-1): an overview. *J Interferon Cytokine Res* 2009; 29: 313-326.
- 664 51. Puthothu B, Bierbaum S, Kopp MV, Forster J, Heinze J, Weckmann M, Krueger M,
665 Heinzmann A. Association of TNF-alpha with severe respiratory syncytial virus infection and
666 bronchial asthma. *Pediatr Allergy Immunol* 2009; 20: 157-163.

667 52. Gu L., Tseng S., Horner R.M., Tam C., Loda M., Rollins B.J. Control of TH2 polarization
668 by the chemokine monocyte chemoattractant protein-1. *Nature*. 2000;404(6776):407-11.

669 53. Zhu J. T helper 2 (Th2) cell differentiation, type 2 innate lymphoid cell (ILC2)
670 development and regulation of interleukin-4 (IL-4) and IL-13 production. *Cytokine*.
671 2015;75(1):14-24.

672 54. Byrne AJ, Mathie SA, Gregory LG, Lloyd CM. Pulmonary macrophages: key players in
673 the innate defence of the airways. *Thorax* 2015; 70: 1189-1196.

674 55. Foster PS, Maltby S, Rosenberg HF, Tay HL, Hogan SP, Collison AM, Yang M, Kaiko
675 GE, Hansbro PM, Kumar RK, Mattes J. Modeling TH 2 responses and airway inflammation
676 to understand fundamental mechanisms regulating the pathogenesis of asthma. *Immunol Rev*
677 2017; 278: 20-40.

678 56. Walter DM, McIntire JJ, Berry G, McKenzie ANJ, Donaldson DD, DeKruyff RH,
679 Umetsu DT. Critical role for IL-13 in the development of allergen-induced airway
680 hyperreactivity. *Journal of Immunology* 2001; 167: 4668-4675.

681 57. Therien AG, Bernier V, Weicker S, Tawa P, Falgout JP, Mathieu MC, Honsberger J,
682 Pomerleau V, Robichaud A, Stocco R, Dufresne L, Houshyar H, Lafleur J, Ramachandran C,
683 O'Neill GP, Slipetz D, Tan CM. Adenovirus IL-13-induced airway disease in mice: a
684 corticosteroid-resistant model of severe asthma. *Am J Respir Cell Mol Biol* 2008; 39: 26-35.

685 58. Fulkerson PC, Fischetti CA, Rothenberg ME. Eosinophils and CCR3 Regulate
686 Interleukin-13 Transgene-Induced Pulmonary Remodeling. *The American Journal of*
687 *Pathology* 2006; 169: 2117-2126.

688 59. Naseer T, Minshall EM, Leung DY, Laberge S, Ernst P, Martin RJ, Hamid Q. Expression
689 of IL-12 and IL-13 mRNA in asthma and their modulation in response to steroid therapy. *Am*
690 *J Respir Crit Care Med* 1997;155: 845-851.

691 60. Sasse SK, Kadiyala V, Danhorn T, Panettieri RA, Phang TL, Gerber AN. Glucocorticoid
692 Receptor ChIP-Seq Identifies PLCD1 as a KLF15 Target that Represses Airway Smooth
693 Muscle Hypertrophy. *American journal of respiratory cell and molecular biology*.
694 2017;57(2):226-37.

695 61. Corren J. Cytokine inhibition in severe asthma: current knowledge and future directions.
696 *Current opinion in pulmonary medicine*. 2011;17(1):29-33.

697 62. Wenzel SE, Busse WW. Severe asthma: lessons from the Severe Asthma Research
698 Program. *J Allergy Clin Immunol*. 2007;119(1):14-21; quiz 2-3.

63. Starkey MR, Essilfie AT, Horvat JC, Kim RY, Nguyen DH, Beagley KW, Mattes J, Foster PS, Hansbro PM. Constitutive production of IL-13 promotes early-life Chlamydia respiratory infection and allergic airway disease. *Mucosal Immunol* 2013; 6: 569-579.

64. Hanania NA, Noonan M, Corren J, Korenblat P, Zheng Y, Fischer SK, Cheu M, Putnam WS, Murray E, Scheerens H, Holweg CT, Maciucă R, Gray S, Doyle R, McClintock D, Olsson J, Matthews JG, Yen K. Lebrikizumab in moderate-to-severe asthma: pooled data from two randomised placebo-controlled studies. *Thorax* 2015; 70: 748-756.

65. Li H, Wang K, Huang H, Cheng W, Liu X. A meta-analysis of anti-interleukin-13 monoclonal antibodies for uncontrolled asthma. *PLoS ONE*. 2019;14(1):e0211790.

FIGURE LEGENDS

FIGURE 1. LPS administration induces clinical features of asthma exacerbation characterised by steroid-resistant AHR and an exaggerated inflammation response in mice with pre-existing AAD. BALB/c mice were sensitized with OVA or PBS plus alum on day 0 and exposed to aerosol OVA on days 13-16. LPS or PBS was administered i.t. on days 19 and 21 and endpoint assessed on day 24. DEX or PBS was administered i.p., three hours before LPS administration. (A) Airway resistance to increased doses of methacholine (mg/ml), presented as percentage change over baseline (saline). (B) Differential BALF cell counts of neutrophils, macrophages, eosinophils and lymphocytes presented as $1 \times 10^4/\text{ml}$ of BALF. (C) Mucus levels were assessed by qPCR, normalised to HPRT. (D) Cytokine levels in lung tissue were assessed by ELISA, normalised to total lung protein (pg/ml). Presented as mean $\pm$ SEM (n=8-10 mice per group); *p<0.05, **p<0.01, ***p<0.001, #p<0.0001. # symbol on top of the bar column shows significant difference (p<0.0001) compared to all groups.

FIGURE 2. LPS administration to OVA-sensitised and challenged mice increased innate inflammatory cytokine expression in primary lung macrophages. Primary pulmonary macrophages were isolated from treated mice on day 24. Cytokine levels were assessed by qPCR, normalised to HPRT. Presented as mean $\pm$ SEM (n=8-10 mice per group); *p<0.05, **p<0.01, ***p<0.001, #p<0.0001.

FIGURE 3. 2-CA-mediated macrophage depletion suppresses AHR and inflammatory cytokine levels in OVA/LPS-treated mice. Mice were treated i.t. with 2-CA or PBS (vehicle control) after LPS challenge in OVA-treated mice on days 22/23 and endpoints assessed on day 24. (A) Airway resistance to increased doses of methacholine (mg/ml), presented as

percentage change over baseline (saline). (B) Differential BALF cell counts of neutrophils, macrophages, eosinophils and lymphocytes presented as $1 \times 10^4/\text{ml}$ of BALF. (C) Mucus levels were assessed by qPCR, normalised to HPRT. (D) Cytokine levels in lung tissue assessed by ELISA, normalised to total lung protein, showed as pg/ml. Presented as mean $\pm$ SEM (n=8-10 mice per group); *p<0.05, **p<0.01, ***p<0.001, # p<0.0001. # symbol on top of the bar column shows significant difference (p<0.0001) compared to all groups.

FIGURE 4. IL-13-deficiency suppresses AHR, inflammatory cell infiltrates and cytokine levels in OVA/LPS-treated mice. (A) Airway resistance presented as percentage change over baseline (saline) on day 24 in wild-type (WT) and IL-13^{-/-} mice. (B) Differential BALF cell counts of neutrophils, macrophages, eosinophils and lymphocytes presented as $1 \times 10^4/\text{ml}$ of BALF. (C) Mucus levels were assessed by qPCR, normalised to HPRT. (D) Cytokine levels in lung tissue assessed by ELISA, normalised to total lung protein (pg/ml). Presented as mean $\pm$ SEM (n=5-6 mice per group); *p<0.05, **p<0.01, ***p<0.001, # p<0.0001. # symbol on top of the bar column shows significant difference (p<0.0001) compared to all groups.

FIGURE 5. IL-13 neutralization suppresses AHR and inflammatory cell numbers in OVA/LPS-treated mice. Anti-IL-13 antibody or isotype control was administered i.p. to OVA/LPS-treated mice, 4 h before LPS administration on day 19 and 21 and endpoints assessed on day 24. (A) Airway resistance to increased doses of methacholine (mg/ml), presented as percentage change over baseline (saline). (B) Differential BALF cell counts of neutrophils, macrophages, eosinophils and lymphocytes presented as $1 \times 10^4/\text{ml}$ of BALF. (C) Mucus levels were assessed by qPCR, normalised to HPRT. (D) Cytokine levels in lung tissue assessed by ELISA, normalised to total lung protein (pg/ml). Presented as mean $\pm$ SEM (n=8-10 mice per group); *p<0.05, **p<0.01, # p<0.0001. # symbol on top of the bar column shows significant difference (p<0.0001) compared to all groups.

FIGURE 6. Cytokine expression is induced in naïve pulmonary macrophages following LPS and/or IL-13 exposure *in vitro*. Primary lung macrophages were isolated from naïve mice and stimulated with IL-13 (100 ng/ml) for 16 hours, before addition of LPS (100 ng/ml) for 12 hours. Cytokine or chemokine levels (as indicated) were assessed by qPCR, normalised to HPRT. Presented as mean $\pm$ SEM (n=6 samples per group); ***p<0.001, # p<0.0001.

FIGURE 7. Steroid insensitivity in Group 2 ILC cells in the lungs following LPS exposure. Mice were treated i.p. with DEX after LPS challenge in OVA-treated mice on days

22/23 and endpoints assessed on day 24. (A) Levels of IL-13-producing ILC2 cells (CD45⁺Lineage⁻ICOS⁺CD90.2⁺ST2⁺IL-13⁺) in lung tissue were assessed by flow cytometry analysis. (B) Type 2 cytokine levels in lung tissue were assessed by ELISA, normalised to total lung protein. Presented as mean±SEM (n=6 mice per group); *p<0.05, **p<0.01, ***p<0.001, # p<0.0001.

FIGURE 8. Effect of DEX, 2-CA and anti-IL-13 treatments on histopathological changes in lung tissues during OVA/LPS-induced allergic asthma.

Lungs samples were collected on day 24, fixed in 10% formalin, paraffinized and cut into slices. The slices were stained with Alcian Blue/Periodic Acid-Schiff (AB-PAS) to examine mucus production or Carbol's Chromotrope-hematoxylin (CR) to examine cell infiltration and hypertrophy of structural cells. (A) shows histopatology of the lung samples from different groups and treatments, all the histological images were randomly selected using 20X magnification lens. (B) Positive mucosal goblet cells and eosinophils around the bronchial airway were counted as described in the Methods section. Presented as mean±SEM (n=8-10 mice per group); *p<0.05, #p<0.0001.

Figure 1.

cea_13505_f1.tif

cea_13505_f2.tif

Figure 3.

cea_13505_f3.tif

Figure 4.

cea_13505_f4.tif

A AHR

B BALF

C Mucus

D Protein level

Figure 5.

cea_13505_f5.tif

Figure 6.

cea_13505_f6.tif

Figure 7.

cea_13505_f7.tif

Figure 8.

cea_13505_f8.tif

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Hadjigol, S;Netto, KG;Maltby, S;Tay, HL;Nguyen, TH;Hansbro, NG;Eyers, F;Hansbro, PM;Yang, M;Foster, PS

Title:

Lipopolysaccharide induces steroid-resistant exacerbations in a mouse model of allergic airway disease collectively through IL-13 and pulmonary macrophage activation

Date:

2019-10-18

Citation:

Hadjigol, S., Netto, K. G., Maltby, S., Tay, H. L., Nguyen, T. H., Hansbro, N. G., Eyers, F., Hansbro, P. M., Yang, M. & Foster, P. S. (2019). Lipopolysaccharide induces steroid-resistant exacerbations in a mouse model of allergic airway disease collectively through IL-13 and pulmonary macrophage activation. CLINICAL AND EXPERIMENTAL ALLERGY, 50 (1), pp.82-94. <https://doi.org/10.1111/cea.13505>.

Persistent Link:

<http://hdl.handle.net/11343/286514>