Richmond Christopher (Orcid ID: 0000-0002-9778-4344)

Title

Cerebellar ataxia with normal intellect associated with a homozygous truncating variant in *CA8*

Running Title

Truncating mechanism for *CA8* cerebellar ataxia

Bylines

Christopher. M. Richmond¹, Richard Leventer^{1,2,3}, Monique M. Ryan^{1,2,3}, Martin. B. Delatycki^{1,2}

Affiliation lines

1. Murdoch Children's Research Institute, Parkville, Victoria, Australia
2. Department of Paediatrics, University of Melbourne, Parkville, Victoria, Australia
3. Department of Neurology, Royal Children's Hospital, Parkville, Victoria, Australia

Corresponding author

Prof. Martin Delatycki. *Address:* Victorian Clinical Genetics Services, Murdoch Children's Research Centre, Flemington Rd, Parkville, VIC, Australia 3052.

Email: martin.delatycki@vcgs.org.au. *Phone:* +61 3 8341 6290. *Fax:* +61 3 8341 6385.

Acknowledgements

The authors wish to acknowledge and value the contributions of clinical and laboratory personnel involved in the care and diagnosis of this patient. We would also like to thank the family of the proband for their time and participation.

Conflict of Interest

The authors declare no competing financial interests.

Data Availability Statement

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/cge.13666](https://doi.org/10.1111/cge.13666)

The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy restrictions.

Abstract

Biallelic pathogenic variants in *CA8* cause cerebellar ataxia, mental retardation and dysequilibrium syndrome 3 (CAMRQ3), a rare form of hereditary ataxia characterised by cerebellar hypoplasia/atrophy, variable intellectual disability and often quadrupedal gait. The few cases reported in the medical literature are all caused by pathogenic homozygous or compound heterozygous missense variants in *CA8*. We report a 9 year-old boy with marked gross motor delay, ataxia and progressive cerebellar atrophy with limited bipedal gait, but without intellectual disability. Singleton whole exome sequencing was performed. A novel homozygous truncating variant in *CA8* (c.232C>T) with a predicted premature termination codon at position 78 (p.Arg78*) was identified. Both parents and the proband's healthy sister are heterozygous for the variant. This variant is likely pathogenic and the cause of the condition in this child. Functional evidence in the form of a spontaneous mouse model involving homozygous intragenic deletion of the mouse analogue of *CA8* with nonsense-mediated decay and similar clinical features to the proband support pathogenicity. Identification of this truncating variant broadens the genotypic and phenotypic spectrum of *CA8*-related cerebellar ataxia.

Keywords

Cerebellar ataxia, hereditary ataxia, cerebellar hypoplasia, neurogenetics, medical genomics

Introduction

The hereditary ataxias, a heterogeneous group of conditions characterised by impaired balance and coordination, are collectively common and confer considerable morbidity across the lifespan¹. Increased availability of next generation sequencing (NGS) has permitted molecular characterisation of many of these conditions and improved understanding of their neuropathophysiological underpinnings. Considerable locus and clinical heterogeneity is now recognised. The existence of an autosomal recessive form of nonprogressive cerebellar ataxia with variable intellectual disability, disturbed equilibrium, cerebro-cerebellar hypoplasia and loss of bipedal locomotion was described as early as 1940² and further analysed in a large consanguineous Hutterite family in 1981³. Diagnostic labels of ‘dysequilibrium’ syndrome⁴ and Unertan syndrome⁵ have been historically applied, with ‘cerebellar ataxia, mental retardation and dysequilibrium syndrome’ (CAMRQ)⁶ now in use. Biallelic mutations in four genes have been implicated in CAMRQ, giving rise to four subtypes (CAMRQ1-4) caused by pathogenic variants in *VLDLR*⁷, *WDR81*⁶, *CA8*⁸ and *ATP8A2*⁹ respectively.

CAMRQ1 (MIM 224050), or *VLDLR*-associated cerebellar hypoplasia (*VLDLR*-CH), is the most well-described subtype, first described in the Hutterite population due to a founder mutation^{10, 11}. In addition to the core features, *VLDLR*-CH causes moderate-to-profound intellectual disability, dysarthria, inferior cerebellar and pontine hypoplasia and gyral simplification, frequently with quadrupedal gait¹¹. Confirmed mutations in *WDR81* and *ATP8A2* (causing CAMRQ2 and CAMRQ4) have only been described in a small number of families^{6, 9, 12}. There are no established genotype-phenotype correlations.

CAMRQ3 (MIM 613227) is caused by biallelic mutations in *CA8*, a gene that encodes carbonic anhydrase-related peptide 8 (CARP8)⁸. There are only three published cases, all caused by biallelic missense mutations^{8, 13, 14} (Table 1). We describe a 9 year-old boy with ataxia and normal intellect who has a homozygous novel truncating variant in *CA8*, to our knowledge the first published report of this mutational mechanism causing the condition. Our findings expand the phenotypic and genotypic spectrum of *CA8*-related cerebellar ataxia.

Materials and Methods

Case Report

The 9 year-old male proband is the first child born to healthy non-consanguineous Caucasian parents. Labour was induced at 35 weeks for spontaneous rupture of membranes after an otherwise uncomplicated pregnancy. He required brief support in the special care nursery for temperature regulation and feeding difficulties. Moderate generalised hypotonia was apparent in infancy, particularly involving the trunk and lower limbs, with severe head-lag that continued until age 5 years. He sat at 14 months and crawled at 18 months. First independent steps were achieved at 6 years with significant cerebellar ataxia. At 9 years, gait is wide-based, slow and limited to 10-20 steps, requiring assistance for longer distances. The proband has poor fine motor skills and incoordination with dysmetria. Speech was delayed with his first word at 12 months but no additional language until 24 months. At 9 years speech content and vocabulary is age-appropriate but with reduced velocity and significant dysarthria. He has poor oromotor coordination without bulbar or respiratory compromise. Visual acuity is normal although some tracking and accommodative difficulties are apparent.

He attends a mainstream school and does not have an intellectual disability. At no stage was there developmental plateau or regression.

On last examination at age nine years there were no significant dysmorphic features. His weight and height were on the 5th percentile and head circumference on the 30th percentile. Although nystagmus was not present, he had difficulty with visual tracking and smooth pursuit. He had marked cerebellar dysarthria. His gait was wide based and unsteady, and he showed truncal instability. There was normal tone in his limbs and normal strength. Reflexes in the upper limbs were normal but reflexes in the lower limbs were increased with subtle crossing and an extensor right plantar response. The Romberg test was normal. He had an intention tremor and difficulties with rapid alternating movements.

Magnetic resonance imaging (MRI) of the brain was normal at 2 years, but repeat MRI at age 7 years showed cerebellar vermian and hemispheric cerebellar atrophy superiorly with thinning of the folia (Figure 1). Normal investigations included chromosomal microarray, urine and plasma metabolic screens, transferrin isoforms, lysosomal enzymes, mitochondrial and *POLG* gene panel, mitochondrial deletion/duplication panel and testing for the *FXN* intron 1 GAA expansion that underlies most Friedreich ataxia.

Whole exome sequencing

Informed consent was obtained for genomic testing and case report publication, in accordance with institutional requirements. Singleton whole exome sequencing with deletion/duplication analysis was undertaken on the proband. DNA was extracted from peripheral blood using QIAamp DNA blood mini kits (Qiagen, Düsseldorf, Germany) and sample quality confirmed using the Agilent Tape Station Genomic DNA ScreenTape (Agilent

Technologies, Santa Clara, CA, USA). Sequencing was performed at PerkinElmer Genomics Laboratory (Branford, CT, USA) by massively-parallel sequencing using a targeted sequence capture method (Agilent v6CREv2) followed by NGS of the amplified captured regions (Illumina, San Diego, CA, USA). Alignment to reference genome GRCh37 was performed. The DRAGEN (Dynamic Read Analysis for GENomics) Bio-IT Platform (Illumina) generated annotated variant calls within the target region and copy number variation (CNV) analysis was performed using the DRAGEN CNV platform (Illumina). Variants were classified according to the American College of Medical Genetics and Genomics (ACMG) variant classification criteria¹⁵.

Results

A novel homozygous nonsense variant in exon 2 of *CA8* (NM_004056.4[CA8]: c.232C>T) was identified, with predicted substitution of arginine for a premature termination codon (PTC) at position of 78 of the protein (p.Arg78*). This variant has not previously been reported in clinical cases and is absent in population databases (gnomAD, 1000 Genomes). Both healthy parents were heterozygous for the variant by polymerase chain reaction (PCR) amplification and Sanger sequencing, as was the proband's healthy younger sister. There were no additional variants of interest considered compatible with the phenotype of the proband.

Discussion

CAMRQ3 is a rare hereditary ataxia reported in only three published families to date. Core features of (usually superior) cerebellar atrophy, ataxia and dysarthria appear to be fully penetrant, while intellectual disability, tremor and loss of bipedal gait are variably reported. This is similar to *VLDLR*-associated cerebellar hypoplasia. Possible progression of the phenotype has been reported in one individual with CAMRQ3 (Table 1). The proband reported here has not experienced clinical regression but the differing MRI findings between age 2 and age 7 may suggest some progression the phenotype. The small number of individuals reported precludes meaningful conclusions around genotype-phenotype correlation. It would appear, however, that intellectual disability and quadrupedal gait are not necessary in making the diagnosis, and a diagnostic label of 'CA8-related cerebellar ataxia' is proposed in place of CAMRQ3 in this child.

There has been some debate regarding the biological mechanisms driving the quadrupedal gait tendency which has historically defined the condition. Quadrupedal

locomotion may represent an adaptive compensation for impaired balance and coordination rather than a devolution of bipedality^{16, 17} although it is generally not seen in other early-onset ataxias. The proband has achieved limited bipedal gait with intensive support and physical therapy, possibly representing a modified phenotype and extending the phenotypic spectrum of the condition.

Given the paucity of truncating *CA8* variants in the medical literature and disease databases, additional functional evidence was considered in assigning pathogenicity to the p.(Arg78*) variant. This variant is located in exon 2, predicting nonsense-mediated decay (NMD). Should the transcript escape NMD, there would be loss of the majority of the carbonic anhydrase domain, including a pfam-predicted active site at position 87. There is a spontaneous mouse model in which a biallelic 19 bp deletion in the murine homolog of *CA8* that results in ataxia and truncal dystonia, providing persuasive evidence for a loss of function mechanism¹⁸. Homozygous *wdl* mutant mice exhibit reduced *CA8* mRNA transcript and protein levels compared to wildtype, indicating NMD and supporting a loss of function mechanism¹⁸.

The ataxic phenotype in *wdl* mice is mediated through disordered granule cell proliferation and circuit patterning of cerebellar Purkinje cells, providing insight into the pathogenesis of the condition¹⁹. Development of paraneoplastic cerebellar degeneration due to anti-CARP8 antibodies targeting Purkinje cells in an individual with malignant melanoma provides additional *in vivo* evidence that loss of function of this protein results in cerebellar dysfunction²⁰.

We conclude that the p.(Arg78*) variant is likely pathogenic and is the cause of this child's condition. A likely pathogenic classification was applied, based on ACMG criteria¹⁵, in

consideration of the variant's absence in population databases (PM2), compelling functional evidence for a loss of function mechanism (PS3) and close concordance of clinical features with those reported in an animal model. In this family, molecular characterisation has permitted diagnostic confidence and informed reproductive planning. Identification of this truncating variant broadens the mutational spectrum of this rare condition and highlights the value of whole exome sequencing in diagnosis of rare forms of hereditary ataxia.

References

1. Musselman KE, Stoyanov CT, Marasigan R, Jenkins ME, Konczak J, Morton SM, et al. Prevalence of ataxia in children: a systematic review. *Neurology*. 2014; 82(1):80-9.
2. Norman RM. Primary Degeneration of the Granular Layer of the Cerebellum: An Unusual Form of Familial Cerebellar Atrophy Occurring in Early Life. *Brain*. 1940; 63(4):365-379.
3. Schurig V, Orman AV, Bowen P. Nonprogressive cerebellar disorder with mental retardation and autosomal recessive inheritance in Hutterites. *Am J Med Genet*. 1981; 9(1):43-53.
4. Hagberg B, Sanner G, Steen M. The dysequilibrium syndrome in cerebral palsy. Clinical aspects and treatment. *Acta Paediatr Scand Suppl*. 1972; 226:1-63.
5. Tan U, Karaca S, Tan M, Yilmaz B, Bagci NK, Ozkur A, et al. Unertan syndrome: a case series demonstrating human devolution. *Int J Neurosci*. 2008; 118(1):1-25.
6. Gulsuner S, Tekinay AB, Doerschner K, Boyaci H, Bilguvar K, Unal H, et al. Homozygosity mapping and targeted genomic sequencing reveal the gene responsible for cerebellar hypoplasia and quadrupedal locomotion in a consanguineous kindred. *Genome Res*. 2011; 21(12):1995-2003.
7. Ozcelik T, Akarsu N, Uz E, Caglayan S, Gulsuner S, Onat OE, et al. Mutations in the very low-density lipoprotein receptor VLDLR cause cerebellar hypoplasia and quadrupedal locomotion in humans. *Proc Natl Acad Sci U S A*. 2008; 105(11):4232-6.

8. Türkmen S, Guo G, Garshasbi M, Hoffmann K, Alshalah AJ, Mischung C, et al. CA8 mutations cause a novel syndrome characterized by ataxia and mild mental retardation with predisposition to quadrupedal gait. *PLoS Genet*. 2009; 5(5):e1000487.
9. Onat OE, Gulsuner S, Bilguvar K, Nazli Basak A, Topaloglu H, Tan M, et al. Missense mutation in the ATPase, aminophospholipid transporter protein ATP8A2 is associated with cerebellar atrophy and quadrupedal locomotion. *Eur J Hum Genet*. 2013; 21(3):281-5.
10. Ali BR, Silhavy JL, Gleeson MJ, Gleeson JG, Al-Gazali L. A missense founder mutation in VLDLR is associated with Dysequilibrium Syndrome without quadrupedal locomotion. *BMC Med Genet*. 2012; 13:80.
11. Boycott KM, Parboosingh JS. VLDLR-Associated Cerebellar Hypoplasia. In: Adam MP, Ardinger HH, Pagon RA, Wallace SE, Bean LJH, Stephens K, et al., editors. *GeneReviews*. Seattle (WA): University of Washington, Seattle; 2008.
12. Komara M, John A, Suleiman J, Ali BR, Al-Gazali L. Clinical and molecular delineation of dysequilibrium syndrome type 2 and profound sensorineural hearing loss in an inbred Arab family. *Am J Med Genet A*. 2016; 170a(2):540-543.
13. Kaya N, Aldhalaan H, Al-Younes B, Colak D, Shuaib T, Al-Mohaileb F, et al. Phenotypical spectrum of cerebellar ataxia associated with a novel mutation in the CA8 gene, encoding carbonic anhydrase (CA) VIII. *Am J Med Genet B Neuropsychiatr Genet*. 2011; 156b(7):826-34.
14. Najmabadi H, Hu H, Garshasbi M, Zemojtel T, Abedini SS, Chen W, et al. Deep sequencing reveals 50 novel genes for recessive cognitive disorders. *Nature*. 2011; 478(7367):57-63.
15. Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genetics in Medicine*. 2015; 17(5):405-24.
16. Humphrey N, Mundlos S, Turkmen S. Genes and quadrupedal locomotion in humans. *Proc Natl Acad Sci U S A*. 2008; 105(21):E26.
17. Herz J, Boycott KM, Parboosingh JS. "Devolution" of bipedality. *Proc Natl Acad Sci U S A*. 2008; 105(21):E25.

18. Jiao Y, Yan J, Zhao Y, Donahue LR, Beamer WG, Li X, et al. Carbonic anhydrase-related protein VIII deficiency is associated with a distinctive lifelong gait disorder in waddles mice. *Genetics*. 2005; 171(3):1239-46.
19. Miterko LN, White JJ, Lin T, Brown AM, O'Donovan KJ, Sillitoe RV. Persistent motor dysfunction despite homeostatic rescue of cerebellar morphogenesis in the Car8 waddles mutant mouse. *Neural Dev*. 2019; 14(1):6.
20. Bataller L, Sabater L, Saiz A, Serra C, Claramonte B, Graus F. Carbonic anhydrase-related protein VIII: autoantigen in paraneoplastic cerebellar degeneration. *Ann Neurol*. 2004; 56(4):575-9.

Figure Legends

Table 1. Previously reported cases of CAMRQ3 with confirmed pathogenic or likely pathogenic variants in CA8

Figure 1. Brain MRI showing progressive cerebellar atrophy. Midline sagittal and coronal T1-weighted images at age two years (left column) and seven years (right column) showing progressive cerebellar atrophy affecting both the vermis and superior cerebellar hemispheres.

Clinical and Research Publication Consent Form

Genetic File Number: 55776

DETAILS OF PATIENT

First Name: [REDACTED] Surname: [REDACTED]

Date of Birth: [REDACTED]

I, [REDACTED], being the patient / legal capacity appointee, or, being the parent / legal guardian / next of kin
[Full name of the person giving consent]

of [REDACTED]
[Full name of the patient / child / family member/s]

consent for medical research or scientific publication of clinical information.

VCGS Clinician to request person giving consent to please circle *agree* or *do not agree* option

☒ **I agree** ☐ **do not agree** (please select) that clinical details, including information about medical conditions, the family history, and results of investigations (including images such as X-rays and pictures taken through a microscope) may be included.

☒ **I agree** ☐ **do not agree** (please select) that a drawn family tree (pedigree) may be included (if applicable).

☒ **I agree** ☐ **do not agree** (please select) that clinical photographs may be included (if applicable).

I understand that the clinical information will be published without names attached, however I understand that complete anonymity cannot be guaranteed. It is possible that someone may recognise the information provided as related to me or my family.

I understand that the clinical information may include details of medical conditions and their prognosis, and treatment that has been provided or may be required in the future.

I understand that the clinical information may be published in a journal that is distributed worldwide, which go mainly to doctors and other health professionals, but may also be seen by others including academics, students and journalists. It may be placed on the journal's website and made available on other medical / scientific websites or linked to in social media.

I understand that in some cases, the journal may require an additional consent form to be completed at a later date.

I understand that I may revoke my consent up until the information is committed to publication in a medical journal ("gone to press"). I understand that after the article has "gone to press", it is not possible to revoke my consent.

I understand that VCGS staff will not publish images on non medical / scientific websites such as Google, but others may take images published in medical / scientific publications and place those images on non medical / scientific websites. VCGS has no control over such actions.

VCGS handles all medical information in accordance with the Privacy Act 1988 (Cth) and the Health Records Act 2001 (Vic)

I have been given the opportunity to ask questions and I am satisfied with the explanation and the answers to my questions. I hereby give my consent.

Signature: (Patient/appointee/parent/guardian/NoK): [Signature] Date: 23/05/2019

VCGS CLINICIAN: I have a good understanding of the service and I have explained the process to the above named and answered any questions. In my opinion, she/he/they has understood this explanation and has signed and consents without compulsion.

Clinician's Name: Dr. Chris Ridwood Signature: [Signature] Date: 21/5/2019

If interpreter service used:

Name of Interpreter: _____ Signature: _____ Date: _____

CGE_13666_Figure 1.tiff

CA8-related cerebellar ataxia: novel truncating mechanism

CAMRQ3

cerebellar ataxia
mental retardation
dysequilibrium

Cause: biallelic
pathogenic variants
in **CA8**...

9 year-old male

Congenital ataxia
Limited bipedal gait
Cerebellar atrophy
Normal intellect

Previously published
variants all **missense**

whole exome
sequencing

p.Ser100Pro

p.Gly162Arg

p.Arg237Gln

c.232C>T

p.(Arg78*)

Mouse model
supports loss
of function

CGE_13666_Graphical Abstract.PNG

Table 1. Previously reported cases of CAMRQ3 with confirmed pathogenic or likely pathogenic variants in CA8

Family	Nucleotide Change [†]	Predicted Protein Change	Variant Type	Clinical Presentation	MRI Findings	Source, Evidence
1	c.298T>C	p.Ser100Pro	Missense	CAMRQ3 Mild ID Cerebellar ataxia, dysarthria Tremor Quadrupedal gait	n/a	Türkmen et al ⁸ : 4 affected siblings
2	c.710G>A	p.Arg237Gln	Missense	CAMRQ [‡] Ataxia	Cerebellar hypoplasia	Najmabadi et al ¹⁴ : single proband
3	c.484G>A	p.Gly162Arg	Missense	CAMRQ3 Mild ID Cerebellar ataxia, dysarthria Delayed/absent walking No quadrupedal gait	Cerebellar volume loss (possibly progressive) Peritrigonal white matter irregularities PET: Hypometabolic cerebellar hemispheres, temporal lobes, mesial cortex	Kaya et al ¹³ : 7 affected individuals, 3 related families
4	c.232C>T	p.Arg78*	Nonsense	CAMRQ3 Cerebellar ataxia Significant gross motor delay Generalised hypotonia No quadrupedal gait, walking limited to 10-20 steps Normal intellect	Cerebellar atrophy (vermis, superior cerebellar hemispheres), progressive	This report

[†]Transcript NM_004056.4

[‡]Limited clinical information available

ACMG denotes American College of Medical Genetics and Genomics¹⁵; CAMRQ3, ‘cerebellar ataxia, mental retardation and dysequilibrium syndrome 3’; ID, intellectual disability; n/a, not available; PET, positron emission tomography.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Richmond, CM;Leventer, R;Ryan, MM;Delatycki, MB

Title:

Cerebellar ataxia with normal intellect associated with a homozygous truncating variant in CA8

Date:

2019-11-14

Citation:

Richmond, C. M., Leventer, R., Ryan, M. M. & Delatycki, M. B. (2019). Cerebellar ataxia with normal intellect associated with a homozygous truncating variant in CA8. CLINICAL GENETICS, 97 (3), pp.516-520. <https://doi.org/10.1111/cge.13666>.

Persistent Link:

<http://hdl.handle.net/11343/286619>