Article begins on page three of this document.

Title	Diagnosis of West Nile Virus encephalitis in a returned traveller
-------	---

Authors:

Author Manuscript

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/MJA2.50416](https://doi.org/10.1002/MJA2.50416)

This article is protected by copyright. All rights reserved

	Title	First name	Mid initials	Last name	Postnom (eg, PhD)	Position1	Address1	Position2	Address2	Tel	Email
1	Dr.	Naomi	C A	Whyler	MBChB, MRCP, MSc, BMedSci	ICU Registrar	1	General Medicine Registrar	2		naomicawhyler@doctors.org.uk
2	Dr.	Jasmine	C	Teng		Infectious Diseases & Microbiology Registrar	3				jasminecteng@gmail.com
3	Assoc. Prof.	David	J	Brewster	MBBS, FCICM, FANZCA, GCHPE	Associate Professor and Clinical Dean	4	Deputy Director of Intensive Care	1		dbrewster@cabrini.com.au
4	Dr.	Ruth		Chin		Consultant in General Medicine & Infectious Diseases	1	Consultant in Infectious Diseases	5		Ruth.Chin@mh.org.au
5	Dr.	Ian		Cox		Consultant Radiologist	1				Ian.Cox@i-med.com.au
6	Dr.	Julian		Druce	MSc, PhD	Senior Scientist	6			03 9342 9623	julian.druce@mh.org.au
7	Prof.	Henry	M	Prince	MBBS, MD, FRACP, FRCPA, MAICD	Director of Blood Cell Therapies	7	Consultant Haematologist	1	+61 (0)3 965617 00	miles.prince@petermac.org
8	Dr.	David	A	Sheffield		Consultant in Infectious Diseases	1	Consultant in Infectious Diseases	8		sheffield010@gmail.com
9	Dr.	Eugene		Teh		Consultant in General Medicine	1	Consultant in General Medicine	9		eugene.tehjukee@gmail.com
10	Assoc. Prof.	Vineet		Sarode	MBBS, MD, MPH, FCICM, FRACMA	Director of Intensive Care	1		4	03 9508 1712	vsarode@cabrini.com.au

Number of corresponding author:	1
Number of alternative corresponding author:	

Addresses:

	Institution	Postal address	City	State	Post Code	Nation
1	Cabrini Malvern	181-183 Wattletree Road	Melbourne	VIC	3144	
2	Alfred Health	The Alfred	Melbourne	VIC	3004	
3	Melbourne Pathology	103 Victoria Parade	Melbourne	VIC	3066	
4	Cabrini Clinical School, Monash University	99 Commercial Road	Melbourne	VIC	3004	
5	Royal Melbourne Hospital	300 Grattan Street	Melbourne	VIC	3050	
6	Victorian Infectious Diseases Reference Laboratory	792 Elizabeth Street	Melbourne	VIC	3000	
7	Peter MacCallum Centre	Locked Bag 1	Melbourne	VIC	8006	
8	Monash Health	246 Clayton Road	Melbourne	VIC	3168	
9	Eastern Health	5 Arnold Street	Melbourne	VIC	3128	

Postal address of first corresponding author (if different from the institutional address given above)	
--	--

Primary Keywords [Office use only]	Infectious diseases; Health occupations; Diagnostic techniques and procedures; General medicine; Environment and public health
Secondary keywords [Office use only]	Encephalitis, viral; Travel medicine; Neuroimaging; Pathology, molecular; Diagnostic tests and procedures; Arbovirus infections; Central nervous system viral diseases; Climate change
Notes:	

Article details (press ctrl – 9 to enter details):

Office use

Article type	Lessons from practice
Blurb	First case of imported West Nile virus infection in Australia causing fatal neuroinvasive disease
	uscript

<i>Ms. Number</i>	mja19.00408.R2
<i>Medical editor</i>	Francis Geronimo
<i>Medical editor email</i>	fgeronimo@mja.com.au
<i>Structural editor</i>	Graeme Prince
<i>Structural editor email</i>	gprince@mja.com.au
<i>Section/Category</i>	Medical education/Lessons from practice
<i>Strapheading</i>	Medical education
<i>Substrap</i>	Lessons from practice

Wiley – file data:

Filename for copyediting	why_mja19.00408_ms
Accompanying graphics	why_mja19.00408_gr
Stock images	
Appendices	

Office use – history:

Event	Date
Original submission received	05/05/2019

Event	Date
Accept	27/09/2019

Proof sent to author	
Proof returned by author	
Published (date format xx/xx/xx)	09/12/19
Issue	11
Vol	211
DOI	10.5694/mja19.00408
Journal	The Medical Journal of Australia
Original article DOI (for response)	

Diagnosis of West Nile virus encephalitis in a returned traveller

Clinical record

A 63-year-old woman developed chills, vomiting and diarrhoea the day after returning from 2 months' travel in South East Europe. Travel included a Danube River cruise trip and time in Serbia, Croatia, Germany, Austria, Slovakia and Hungary. She visited rural areas and sustained mosquito bites, but reported no unpasteurised food consumption, animal contact or bites. Pre-travel consultation was not undertaken. Past medical history included follicular lymphoma in remission, with rituximab-induced hypogammaglobulinaemia.

She was hospitalised on day 3 of her illness. On examination, she was febrile (39.1°C), dehydrated but cognitively intact, without meningism, rash or arthritis. Blood pressure of 80/60 mmHg responded to intravenous hydration. Investigations including full blood examination, biochemistry, liver function, C-reactive protein and stool cultures were unremarkable. Initial presumptive diagnosis was infectious gastroenteritis, and empirical ciprofloxacin was administered.

On day 5, she developed confusion, nominal aphasia and slow affect. Brain computed tomography and magnetic resonance imaging (MRI) were unremarkable. Cerebrospinal fluid (CSF) analysis revealed mild lymphocytic pleocytosis ($24 \times 10^6/L$; reference interval [RI], $0-5 \times 10^6/L$), raised erythrocytes ($224 \times 10^6/L$; RI, $< 1 \times 10^6/L$) and a mildly raised protein level (0.54 g/L; RI, 0.15–0.45 g/L), consistent with infection rather than traumatic tap. Aciclovir, ampicillin and piperacillin–tazobactam were administered to cover gram-positive and gram-negative bacterial and viral meningoencephalitis, including listeria and nosocomial pathogens given deterioration 48 hours after admission. Intravenous immunoglobulin, the patient's longstanding treatment for hypogammaglobulinaemia, was administered.

Intubation for mechanical ventilation was performed on day 10, due to worsening symptoms of fluctuating conscious state and uncoordinated motor responses. MRI identified new symmetrical bilateral thalamic, mesial temporal and brainstem signal abnormality (Box 1). Electroencephalography showed subclinical seizure activity.

Initial laboratory investigations were negative: CSF bacterial cultures, viral polymerase chain reaction (PCR) for enterovirus and herpes viruses (herpes simplex virus types 1 and 2, varicella virus and cytomegalovirus), cryptococcal antigen, fungal cultures, and treponemal and human immunodeficiency virus serology.

Imported flavivirus meningoencephalitis including West Nile virus and tick-borne encephalitis virus were considered. Negative flavivirus serology was difficult to interpret

due to hypogammaglobulinaemia. Flavivirus PCR testing was performed on CSF (day 14), whole blood (days 9 and 11) and urine (days 12 and 24). All results were negative except day 12 urine flavivirus PCR; DNA sequencing of this sample identified West Nile virus imported type with the closest match being West Nile virus Bulgaria 2015, confirmed by positive urine viral culture.

Treatment was supportive, with phenytoin, levetiracetam, intravenous immunoglobulin (total 1 g/kg), ventilatory support and tracheostomy.

Despite oedema improving on day 14 MRI, neurological status plateaued: the patient continued to exhibit persistent central apnoeas, decerebrate posturing and reduced consciousness level. Day 28 MRI demonstrated improvement of thalamic and midbrain oedema, but showed new bilateral, punctate foci of marked T2 hyperintensity in midbrain corticospinal tracts, consistent with a diagnosis of severe acquired brain injury secondary to encephalitis. Given limited prognosis, the patient was palliated and died on day 32.

Discussion

This is one of two reported cases of imported West Nile virus infection into Australia causing neuroinvasive disease. In 2009, Rogers and colleagues described an Australian laboratory-confirmed imported West Nile virus infection in a visiting Israeli tourist.¹

West Nile virus is a mosquito-borne neurotropic flavivirus endemic to North America, Africa, Asia, Europe, and Northern Australia (Kunjin subtype).^{2,3} Its transmission cycle includes bird species as amplifier hosts, mosquito vectors, and mammals including humans as dead-end hosts.² Most human infections are asymptomatic, but 20% develop influenza-like symptoms including malaise and fever, and < 1% develop West Nile virus neuroinvasive disease with encephalitis, meningitis or acute flaccid paralysis.² West Nile virus Kunjin subtype causes milder disease, lower rates of West Nile virus neuroinvasive disease, and no reported deaths.^{2,3} Exposure is hypothesised to protect against other West Nile virus subtypes.³ Risk factors for severe disease include advanced age, chronic conditions and immunosuppression, with mortality rates of 3–19%.²

Travel history is essential for diagnosis, and real time monitoring resources describing local West Nile virus outbreaks in endemic areas are useful. The European Centre for Disease Prevention and Control reported a 7.2-fold increase in West Nile virus cases from 2017 to 2018⁴ and aided diagnosis in our case.

Rogers and colleagues relied on paired flavivirus serology for diagnosis,¹ whereas our case relied on urine flavivirus PCR for diagnosis. Flavivirus diagnosis can be difficult due to low viraemia levels, potential for serological cross-reactivity,⁵ or false negative results in the presence of hypogammaglobulinaemia. Emerging evidence suggests flavivirus PCR may be detectable for a number of weeks in urine and whole blood,⁵ particularly in neuroinvasive disease,⁶ compared with serum, which may remain positive for up to a week after symptom onset.⁵

Radiological changes can help narrow down the differential diagnoses of encephalitis. In our case, thalamic MRI changes supported the diagnosis of West Nile virus. Bilateral

thalamic and basal ganglia involvement is reported with specific encephalitis aetiologies, particularly respiratory viruses, *Mycobacterium* species, and arboviruses including West Nile virus.⁷

West Nile virus management is predominantly supportive, although intravenous immunoglobulin with high anti-West Nile virus antibody titres may provide benefit.⁸ Box 2 summarises current avenues of pharmacotherapy.

Our case highlights the importance of travel history, knowledge of endemic disease outbreaks, appropriate sampling to increase diagnostic yield, and radiology to assist diagnosis of undifferentiated infection in returned travellers.

Lessons from practice

- The evolving spectrum of disease in an era of increasing international travel and climate change presents new challenges to local clinicians. Use of health monitoring internet resources can help identify outbreaks at travel destinations to aid diagnosis and management of undifferentiated infection in returned travellers.
- Use of flavivirus serology for diagnosis is challenging in the presence of hypogammaglobulinaemia, and may produce false negatives. Close liaison with a laboratory for appropriate tests is essential.
- Consider flavivirus polymerase chain reaction testing of urine, which remains positive for longer than serum/plasma. Urine and whole blood polymerase chain reaction specimens are increasingly recognised as the best basis for molecular flavivirus diagnosis.
- MRI imaging demonstrating typical features of bilateral thalamic and basal ganglia changes supports a diagnosis of West Nile virus encephalitis and can assist with diagnosis, management and prognostication.

Competing interests: No relevant disclosures.

Provenance: Not commissioned; externally peer reviewed.

Author details

Naomi CA Whyler^{1,2}

Jasmine C Teng³

David J Brewster^{1,4}

Ruth Chin^{1,5}

Ian Cox¹

Julian Druce⁶

Henry M Prince^{1,7}

David A Sheffield^{1,8}

Eugene Teh^{1,9}

Vineet Sarode^{1,4}

1 Cabrini Malvern, Melbourne, VIC.
2 Alfred Health, Melbourne, VIC.
3 Melbourne Pathology, Melbourne, VIC.
4 Cabrini Clinical School, Monash University, Melbourne, VIC.
5 Royal Melbourne Hospital, Melbourne, VIC.
6 Victorian Infectious Diseases Reference Laboratory, Melbourne, VIC.
7 Peter MacCallum Centre, Melbourne, VIC.
8 Monash Health, Melbourne, VIC.
9 Eastern Health, Melbourne, VIC.
naomicawhyler@doctors.org.uk
doi: 10.5694/mja19.00408

References

- 1 Rogers BA, Hueston L, Ratnam I. Imported West Nile virus encephalitis in an Israeli tourist. *Med J Aust* 2009; 191: 232-234. <https://www.mja.com.au/journal/2009/191/4/imported-west-nile-virus-encephalitis-israeli-tourist>
- 2 Chancey C, Grinev A, Volkova E, Rios M. The global ecology and epidemiology of West Nile virus. *BioMed Res Int* 2015; 376230: 1-20.
- 3 Mackenzie JS, Hall RA, Smith DW. West Nile virus: is there a message for Australia? *Med J Aust* 2003; 178: 5-6. <https://www.mja.com.au/journal/2003/178/1/west-nile-virus-there-message-australia>
- 4 European Centre for Disease Prevention and Control. Epidemiological update: West Nile virus transmission season in Europe, 2018. <https://ecdc.europa.eu/en/news-events/epidemiological-update-west-nile-virus-transmission-season-europe-2018> (viewed July 2019).
- 5 Lustig Y, Sofer D, Bucris ED, Mendelson E. Surveillance and diagnosis of West Nile virus in the face of flavivirus cross-reactivity. *Front Microbiol* 2018; 9: 2421.
- 6 Barzon L, Pacenti M, Franchin E et al. Excretion of West Nile virus in urine during acute infection. *J Infect Dis* 2013; 208: 1086-1092.
- 7 Beattie GC, Glaser CA, Sheriff H, et al. Encephalitis with thalamic and basal ganglia abnormalities: etiologies, neuroimaging, and potential role of respiratory viruses. *Clin Infect Dis* 2013; 56: 825-832.
- 8 Shimoni Z, Bin, H, Bulvik S, et al. The clinical response of West Nile virus neuroinvasive disease to intravenous immunoglobulin therapy. *Clin Pract* 2012; 2: e18.
- 9 Acharya D, Bai F. An overview of current approaches toward the treatment and prevention of West Nile virus infection. *Methods Mol Biol* 2016; 1435: 249-291.
- 10 David S, Abraham AM. Epidemiological and clinical aspects on West Nile virus, a globally emerging pathogen. *Infect Dis* 2016; 48: 571-585.

[Boxes]

1 T2-weighted magnetic resonance imaging (day 10), showing bilateral thalamic hyperintensity (arrows)

[why_mja19.00408_gr]

2 Current avenues of pharmacotherapy for West Nile virus

Pharmacotherapy	Current research avenues
Vaccines	Several equine vaccines are available; with some human vaccines in clinical trial ⁹
Ribavirin	In vitro activity has been demonstrated, but ribavirin has been associated with higher mortality when used in human West Nile virus outbreaks ¹⁰
Mycophenolic acid	In vitro activity has been demonstrated, but immunosuppressive effects seen with human use ¹⁰
Intravenous immunoglobulin	Case reports suggest improved outcomes particularly for West Nile neuroinvasive disease requiring respiratory support; ⁸ benefit may be due to the presence of high titres of West Nile virus antibodies in the sourced intravenous immunoglobulin ⁸
Targeted antiviral therapy	Candidates for viral entry/fusion inhibitors, protease inhibitors and other viral replication inhibitors are in early stages of research ⁹
Monoclonal antibodies	Humanised anti-West Nile virus monoclonal antibodies appear promising for post-exposure therapy, and are currently undergoing clinical trial evaluation ⁹

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Whyler, NCA;Teng, JC;Brewster, DJ;Chin, R;Cox, I;Druce, J;Prince, HM;Sheffield, DA;Teh, E;Sarode, V

Title:

Diagnosis of West Nile virus encephalitis in a returned traveller

Date:

2019-12-09

Citation:

Whyler, N. C. A., Teng, J. C., Brewster, D. J., Chin, R., Cox, I., Druce, J., Prince, H. M., Sheffield, D. A., Teh, E. & Sarode, V. (2019). Diagnosis of West Nile virus encephalitis in a returned traveller. MEDICAL JOURNAL OF AUSTRALIA, 211 (11), pp.501-502. <https://doi.org/10.5694/mja2.50416>.

Persistent Link:

<http://hdl.handle.net/11343/286626>