Article begins on page three of this document.

Title	Sarcopenia: a deserving recipient of an Australian ICD-10-AM code
-------	---

Authors:

	Title	First name	Mid inits	Last name	Postnom (eg, PhD) [3 only for publication]	Position1	Address1	Position2	Address2	Tel	Email
1	Dr.	Jesse		Zanker	MBBS, MPHTM, FRACP	Research Fellow	1	Geriatrician	2	038345 6333	jesse.zanker@wh.org.au
2	Dr.	David		Scott	PhD	Research Fellow	2	Research Fellow	3		d.scott@unimelb.edu.au
3	Dr.	Sharon	L	Brennan-Olsen	PhD, BA(Hons), GCALL	Senior Research Fellow	1		2		sbrennan@unimelb.edu.au
4	Prof.	Gustavo		Duque	MD, PhD, FRACP	Professor and Chair of Medicine	1	Professor and Chair of Medicine	2		gustavo.duque@unimelb.edu.au
5											

Number of corresponding author:	1
Number of alternative corresponding author:	

Addresses:

	Institution		City	State	Post Code	
1	Australian Institute of Musculoskeletal Science, University of Melbourne		Melbourne	VIC	3021	
2	Western Health		Melbourne	VIC	3021	
3	Monash University		Melbourne	VIC	3168	
4						
5						

Postal address of first corresponding author (if different from the institutional address given above)	
--	--

Primary Keywords [Office use only]	Gerontology; Musculoskeletal diseases; Health occupations
Secondary keywords [Office use only]	Aged; Geriatrics; Health services for the aged
Notes:	

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/MJA2.50432](#)

This article is protected by copyright. All rights reserved

Article details (press ctrl – 9 to enter details):

Article type	Letter
Blurb	

Office use

<i>Ms. Number</i>	mja19.00552. R1
<i>Medical editor</i>	Zoe Silverstone
<i>Medical editor email</i>	zsilverstone@ mja.com.au
<i>Structural editor</i>	Laura Teruel
<i>Structural editor email</i>	lteruel@mja.c om.au
<i>Section/Category</i>	Letters to the editor (new)
<i>Strapheading</i>	Letter
<i>Substrap</i>	

Wiley – file data:

Filename for copyediting	zan_mja19.00552_ms.docx
Accompanying graphics	None
Stock images	None
Appendices	None

Office use – history:

Event	Date
Original submission received	14/06/2019

Event	Date
Accept	01/07/2019

Proof sent to author	
Proof returned by author	
Published (date format xx/xx/xx)	13/01/20
Issue	1
Vol	212
DOI	10.5694/mja1 9.00552
Journal	The Medical Journal of Australia
Original article DOI (for response)	

Sarcopenia: a deserving recipient of an Australian ICD-10-AM code

TO THE EDITOR: In July 2019, sarcopenia — a progressive and generalised skeletal muscle condition involving loss of skeletal muscle mass and function¹ — was awarded a code in the International Classification of Diseases, tenth revision, Australian modification (ICD-10-AM). This recognition has arrived 30 years after Irwin Rosenberg first described the condition in 1989.²

Sarcopenia is independently associated with poor quality of life, falls, fractures, institutionalisation and mortality.¹ About 13–19% of community-dwelling older adults may have this condition, and prevalence is highest among those living in residential care.¹

All individuals experience declines in muscle mass and function during ageing, but only those who meet the criteria described in the Box are considered to have sarcopenia. The definition currently promoted by the Australian and New Zealand Society for Sarcopenia and Frailty Research is the initial European Working Group on Sarcopenia in Older People definition,³ which was adopted after a Delphi consensus.⁵ Measures of muscle strength and physical performance such as grip strength, chair stands and gait speed are cost-effective and easy to perform in clinical practice. Obtaining measures of muscle and lean mass may be challenging outside of the research setting. Therefore, in individuals with low muscle strength or physical performance, in the absence of other potential causes (eg, osteoarthritis), sarcopenia should be suspected and safe and effective interventions can be offered.

Patients with, or at risk of, sarcopenia should be recommended exercise therapy, in particular, progressive resistance training.¹ This type of training prescribed by treating clinicians can be implemented by allied health professionals, including exercise physiologists and physiotherapists. Protein supplementation can prevent loss of muscle, but this is most beneficial when combined with progressive resistance training.¹ A number of randomised controlled trials are underway examining different therapeutics for the treatment of sarcopenia.¹

With the advent of the ICD-10-AM code, primary care clinicians, allied health staff, and members of the public will begin observing sarcopenia diagnoses on medical correspondence. Hospital funding models may adjust in line with the ICD-10-AM code and in recognition of the increased complexity and risk of complications that comes with caring for patients with sarcopenia. An understanding of this condition, its implications and treatment is key in providing evidence-based care to patients living with sarcopenia.

Acknowledgements: Jesse Zanker is supported by an Australian Government Research Training Program Scholarship. David Scott is supported by a National Health and Medical Research Council (NHMRC) RD Wright Biomedical Career Development Fellowship (GNT1123014). Sharon Brennan-Olsen is supported by an NHMRC Career Development Fellowship (1107510).

Competing interests: No relevant disclosures.

Author details

Jesse Zanker^{1,2}

David Scott^{1,3}

Sharon L Brennan-Olsen¹

Gustavo Duque^{1,2}

¹ Australian Institute of Musculoskeletal Science, University of Melbourne, Melbourne, VIC.

² Western Health, Melbourne, VIC.

³ Monash University, Melbourne, VIC.

jesse.zanker@unimelb.org.au

doi: 10.5694/mja19.00552

References

- 1 Cruz-Jentoft AJ, Sayer AA. Sarcopenia. *Lancet* 2019; 393: 2636-2646.
- 2 Rosenberg IH. Summary comments. *Am J Clin Nutr* 1989; 50: 1231-1233.
- 3 Cruz-Jentoft AJ, Baeyens JP, Bauer JM, et al. Sarcopenia: European consensus on definition and diagnosis: report of the European working group on sarcopenia in older people. *Age Ageing* 2010; 39: 412-23.
- 4 Cruz-Jentoft AJ, Bahat G, Bauer J, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing* 2019; 48: 16-31.
- 5 Zanker J, Scott D, Reijnierse EM, et al. Establishing an operational definition of sarcopenia in Australia and New Zealand: Delphi method based consensus statement. *J Nutr Health Aging* 2019; 23: 105-110.

[Insert box]

Diagnostic tools and measurements to diagnose sarcopenia* using the initial European Working Group on Sarcopenia in Older People (EWGSOP) definition^{†3}

Component	Thresholds and equipment
Low muscle strength	Hand grip strength using dynamometer: <ul style="list-style-type: none"> ▪ Men: < 30 kg ▪ Women: < 20 kg
Low physical performance	Men and women over 4 m course: <ul style="list-style-type: none"> ▪ Gait speed: ≤ 0.8 m/s
Low lean mass	ALM using whole-body DXA (adjusted for height, m ²): <ul style="list-style-type: none"> ▪ Men: < 7.26 kg/m² ▪ Women: < 5.50 kg/m²

ALM = appendicular lean mass; DXA = dual x-ray absorptiometry. * Diagnosis of sarcopenia is based on low lean mass **and** low physical performance **or** muscle strength. † The EWGSOP have developed a revised definition for sarcopenia (known as EWGSOP2);⁴ however, this has not yet been recommended for use in Australia.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Zanker, J;Scott, D;Brennan-Olsen, SL;Duque, G

Title:

Sarcopenia: a deserving recipient of an Australian ICD-10-AM code

Date:

2019-11-28

Citation:

Zanker, J., Scott, D., Brennan-Olsen, S. L. & Duque, G. (2019). Sarcopenia: a deserving recipient of an Australian ICD-10-AM code. MEDICAL JOURNAL OF AUSTRALIA, 212 (1), pp.45-+. <https://doi.org/10.5694/mja2.50432>.

Persistent Link:

<http://hdl.handle.net/11343/286672>