DR SIEW LIM (Orcid ID : 0000-0002-5333-6451)
DR SHARLEEN O'REILLY (Orcid ID : 0000-0003-3547-6634)
PROFESSOR TIMOTHY SKINNER (Orcid ID : 0000-0002-0018-6963)

Article type : Letter

Title: Diabetic Medicine
Created by: Maria Hale
Email proofs to: siew.lim1@monash.edu
Copyright: Diabetes UK
Article no.: DME-2019-00804
Article type: Letter
Figures: 0; Tables: 0; Equations:0; References: 10
Short title/Authors running head: Letter • Letter

Letter

Recommendations for promoting healthier lifestyles in postpartum women after gestational diabetes

We read with interest the study by Dennison *et al.* [1] which provided a qualitative synthesis on the facilitators and barriers to lifestyle changes from the perspective of women with recent gestational diabetes (GDM). We would like to provide some additional insights from our own work. Early intervention for diabetes prevention after GDM is important, but lifestyle intervention during the postpartum period can be challenging. Many of the barriers to a healthy lifestyle identified in this paper were also reported previously by young women and postpartum women in the general population [2,3]. These may be common experiences related to life stage and not unique to GDM, which could explain the high risk of weight gain in women between the ages of 18 and mid-30s [4]. Postpartum women or women with young children have additional barriers such as fatigue and the demands of an infant or young child.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as [doi: 10.1111/DME.14208](https://doi.org/10.1111/DME.14208)

This article is protected by copyright. All rights reserved

Considering the unique barriers in postpartum women, the development of interventions targeting this group requires careful consideration. Interventions need to be feasible, acceptable and fundable for this group and the health professionals, services and systems delivering them. Randomized controlled trial evidence shows that lifestyle intervention within the US diabetes prevention programme for women after GDM is efficacious [5]. We now need to progress the efficacy evidence into broader effectiveness by focusing on the 'how'. To address this, Dennison *et al.* [1] proposed 20 recommendations mapped against behaviour change techniques (BCT) based on the perceived needs of the women. Consideration of end-users' perspectives is critical in implementation research – we would like to commend the authors' effort in doing so. We offer perspectives gained from our systematic review and meta-analysis on effective elements of lifestyle intervention in postpartum women (33 studies, 4960 women), as a triangulation process to add value to the findings of Dennison and colleagues.

In our meta-analysis on intervention characteristics using the Template for Intervention Description and Replication (TIDieR) framework, we found that interventions delivered by health professionals were more effective than those delivered by non-health professionals [6], providing further support for Recommendation 19. Both suggest there is a need to establish models of care within the health system to facilitate goal setting, provide instructions on diet and exercise, and monitoring by health professionals for women after GDM.

Our meta-analysis did not find a significant effect on other intervention characteristics such as format of delivery, intensity or duration [6]. This aligns with Dennison *et al.* who described mixed responses on these characteristics [1]. The Consolidated Framework for Implementation Research (CFIR) [7] describes interventions as having core components and adaptable periphery. Core components are the 'active ingredients', which may be consistent findings across studies identified in systematic reviews and meta-analyses. The adaptable periphery, by contrast, depends on contextual needs that differ from one setting to another. Both Dennison *et al.* and our findings suggest that the format of delivery is part of the adaptable periphery, to be developed according to the needs of the provider and user within the local context and health system. A combination of efficacious core components and effective periphery is essential to maintain fidelity and efficacy, optimize engagement, feasibility and cost-effectiveness for scale-up. Co-production with stakeholders, including clinicians, service providers and consumers, could integrate the core components with adaptable periphery to develop an effective intervention.

This article is protected by copyright. All rights reserved

Dennison *et al.* had low confidence for Recommendation 13's evidence related to advising women on diabetes risk. We previously reported that women after GDM with higher baseline risk perception were more likely to be engaged in a diabetes prevention programme [8]. Higher engagement was also associated with greater intervention weight loss [8]. However, we are unclear if risk perception can be modified independent of absolute health risks, and if a modified risk perception will improve intervention outcomes.

Previous quantitative systematic reviews in other populations are consistent with Recommendations 16–20. They too identified that BCTs relating to self-regulation skills such as goal-setting and self-monitoring were significantly associated with behaviour change [9,10]. Recommendations 3–6 on practical social support were also previously found to be a significant BCT in diabetes care [10]. Our systematic review on the quantitative analysis of BCTs (currently under review) will be able to confirm if they were also significantly associated with diet and physical activity change in postpartum women.

In conclusion, diabetes prevention programmes for postpartum women with previous GDM should include practical support and instructions on how to exercise or have a healthy meal. These should be delivered by healthcare professionals within a model of care that allows for individualized goals and ongoing monitoring. To assist the convergence of these reductionist findings with systems thinking, these active ingredients need to be further developed into fit-for-purpose interventions based on a thorough understanding of the context and system, facilitated through co-production with the end-users.

Funding sources

Competing interests

Commented [MH1]: Author: Give any conflicts of interest or write 'None declared.'

S. S. Lim¹, S. O'Reilly⁴, V. Versace², E. Janus³, T. C. Skinner⁵, J. D. Best⁶, J. Dunbar²
and H. Teede¹

¹Monash Centre for Health Research and Implementation, Monash University, Clayton,

²Deakin Rural Health, School of Medicine, Deakin University, Geelong and ³General Internal Medicine Unit, Western Health and Department of Medicine, Melbourne Medical School, University of Melbourne, Melbourne, Australia, ⁴Institute of Food and Health, School of

Agriculture and Food Science, University College Dublin, Dublin, Ireland, ⁵Institut for Psykologi, Department of Psychology, University of Copenhagen, Copenhagen, Denmark and ⁶Lee Kong Chian School of Medicine, Nanyang Technology University, Singapore

References

- 1 Dennison RA, Ward RJ, Griffin SJ, Usher-Smith JA. Women's views on lifestyle changes to reduce the risk of developing Type 2 diabetes after gestational diabetes: a systematic review, qualitative synthesis and recommendations for practice. *Diabet Med* 2019; **36**: 702–717.
- 2 Pennington AVR, O'Reilly SL, Young D, Dunbar JA. Improving follow-up care for women with a history of gestational diabetes: perspectives of GPs and patients. *Aust J Prim Health* 2017; **23**: 66–74.
- 3 Carter-Edwards L, Ostbye T, Bastian LA, Yarnall KS, Krause KM, Simmons TJ. Barriers to adopting a healthy lifestyle: insight from postpartum women. *BMC Res Notes* 2009; **2**: 161.
- 4 Dutton GR, Kim Y, Jacobs DR Jr, Li X, Loria CM, Reis JP *et al*. 25-year weight gain in a racially balanced sample of U.S. adults: The CARDIA study. *Obesity (Silver Spring)* 2016; **24**: 1962–1968.
- 5 O'Reilly SL, Dunbar JA, Versace V, Janus E, Best JD, Carter R *et al*. Mothers after Gestational Diabetes in Australia (MAGDA): a randomised controlled trial of a postnatal diabetes prevention program. *PLoS Med* 2016; **13**: e1002092.
- 6 Lim S, Liang X, Hill B, Teede HJ, Moran L, O'Reilly S. A systematic review and meta-analysis of intervention characteristics in postpartum weight management using the TiDieR framework: A summary of evidence to inform implementation *Obes Rev* 2019; **20**: 1045–1056.
- 7 Damschroder LJ, Aron DC, Keith RE, Kirsh SR, Alexander JA, Lowery JC. Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implement Sci* 2009; **4**: 50.
- 8 Lim S, Dunbar JA, Versace VL, Janus E, Wildey C, Skinner T *et al*. Comparing a telephone- and a group-delivered diabetes prevention program: characteristics of engaged and non-engaged postpartum mothers with a history of gestational diabetes. *Diabetes Res Clin Pract* 2017; **126**: 254–262.

- 9 Samdal GB, Eide GE, Barth T, Williams G, Meland E. Effective behaviour change techniques for physical activity and healthy eating in overweight and obese adults; systematic review and meta-regression analyses. *Int J Behav Nutr Phys Act* 2017; **14**: 42.
- 10 Presseau J, Ivers NM, Newham JJ, Knittle K, Danko KJ, Grimshaw JM. Using a behaviour change techniques taxonomy to identify active ingredients within trials of implementation interventions for diabetes care. *Implement Sci* 2015; **10**: 55.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Lim, SS;O'Reilly, S;Versace, V;Janus, E;Skinner, TC;Best, JD;Dunbar, J;Teede, H

Title:

Recommendations for promoting healthier lifestyles in postpartum women after gestational diabetes

Date:

2019-12-22

Citation:

Lim, S. S., O'Reilly, S., Versace, V., Janus, E., Skinner, T. C., Best, J. D., Dunbar, J. & Teede, H. (2019). Recommendations for promoting healthier lifestyles in postpartum women after gestational diabetes. DIABETIC MEDICINE, 37 (4), pp.706-708. <https://doi.org/10.1111/dme.14208>.

Persistent Link:

<http://hdl.handle.net/11343/286773>