Development of a public audiology service in Southern Malawi: Profile of patients across two years.

Bhavisha Parmar^a, Mwanaisha Phiri^b, Courtney Caron^c, Tess Bright^d and Wakisa Mulwafu^e

^a UCL Ear Institute, University College London, 332-336 Grays Inn Road, London, WC1X 8EE, UK

^b Queen Elizabeth Central Hospital, Blantyre, Malawi

^c Veterans Health Administration, Veterans Affairs Southern Nevada Health Care System, Audiology Clinic, 6900 North Pecos Road, North Las Vegas, NV 89086, USA

^d International Centre for Evidence in Disability, London School of Hygiene & Tropical Medicine, Keppel St, London, WC1 E7HT, UK

^e Department of Surgery, College of Medicine, University of Malawi, Blantyre, Malawi

Corresponding author: bhavisha.parmar@ucl.ac.uk

Abstract

Objective: To describe the profile of patients attending the Queen Elizabeth Central Hospital (QECH) audiology clinic in Malawi, over a two-year period (2016-2017).

Design: A retrospective patient record review.

Study sample: There were 2299 patients assessed at the QECH audiology department between January 2016 to December 2017. Adult patients' ages ranged from 18 to 94 years ($M = 45.8$, $SD = 19.22$). The mean age of children included in this study was 7.7 years ($SD = 5.21$). Overall, 45.4% of patients were female.

Results: Of the 61.6% of adults and 41.7% of children found to have some degree of hearing loss, 28.3% and 15.4% were fitted with hearing aids, respectively. The number of patients seen in 2017 ($n=1385$) was 34% higher than that of 2016 ($n=914$).

Conclusion: This study found that demand for hearing services is increasing in this Malawian Audiology department but uptake of hearing aids for those in need is low. Future evaluation of service provision and treatment outcomes is needed. Results from this study can be used to inform the development of future audiology clinics in low resource settings.

Introduction

It is estimated that 466 million people in the world live with disabling hearing loss [1]. Over 80% of people with disabling hearing loss live in low- and middle- income countries (LMICs) [2]. The prevalence of hearing loss may be higher in LMICs due to the burden of infectious diseases, poorer access to health care and ear protection as well as increased use of unregulated ototoxic medications [3, 4]. It has been reported that public health measures could reduce the global prevalence of hearing loss by 50% in LMICs [5].

Unaddressed hearing loss can have a significant impact on an individual's communication, cognition, education, employment and overall wellbeing [6-9]. However, there are many barriers to addressing hearing loss in LMICs. Some of these include the access to ear and hearing services, including assistive technologies such as hearing aids or cochlear implants [5]. Of the 401 million people that need hearing aids globally, approximately 83% either do not use or have access to them. This increases to 90% for people residing in African countries [10]. Some significant barriers to the development of ear and hearing care services in these regions include the lack of local training opportunities and a dearth of contextual data regarding the prevalence and aetiology of hearing loss [11], alongside the lack of appropriate policy efforts needed to allocate the necessary resources [10].

Malawi is a landlocked country in Southern Africa with a population of approximately 17.5 million [12]. It is classified as a low income country by the World Bank [13]. The WHO estimates 4-5% of the population of sub-Saharan Africa is estimated to have a disabling hearing loss [14]. However, data on the prevalence and causes of hearing loss in the vast majority of countries in sub-Saharan Africa, including Malawi is very limited. A systematic review by Mulwafu et al (2016) investigated the prevalence of hearing loss in the region and found only eight population-based studies with others relying on data from school screening programmes [15]. Hunt et al (2017) carried out a community-based study in rural Malawi and reported a high prevalence of hearing loss in children aged 4-6 years (11.5%) [16]. A recent population- based longitudinal analysis of children with hearing loss in two districts of rural Malawi compared baseline assessment data obtained in 2013 with follow up data from 2016 [17]. The study found that over half of all children diagnosed with hearing loss (59.1%) were lost

to follow up and there was a low onward referral uptake, particularly for girls and younger children. Similarly, Bright et al (2017) found that 93% of children identified as needing ear and hearing care services in Southern Malawi could not attend their referral appointment [18].

A 2015 survey of Ear Nose and Throat (ENT) services revealed that Malawi, Kenya and South Africa have 0.1, 2, and 5.6 ENT surgeons per million, respectively [19]. When compared to the United Kingdom, where evidence suggests there are 24 ENT surgeons per million people, these numbers demonstrate the huge scarcity of human resources in the region [19]. There are currently three ENT surgeons and three audiologists [20, 21] in Malawi. The majority of ear and hearing care services are only available in urban areas and outreach support is provided to rural areas. This paper focusses on audiology services provided at Queen Elizabeth Central Hospital (QECH), Malawi's largest tertiary referral hospital located in Blantyre, Southern Malawi.

The Development of QECH Audiology Department, Blantyre, Malawi

The first resident ENT surgeon began work at QECH in 2007 [22]. In 2010 he developed a cadre of ENT clinical officers to work at central and district hospitals. This initiative encouraged the use of task shifting to overcome the significant shortage of specialised health professionals. A diploma in 'ENT and Audiology' was introduced to advance the training of clinical officers (secondary school graduates with a diploma in clinical medicine) [23].

Recognising the need for audiology services in the country, in 2014, Sound Seekers began work in Malawi to develop the audiology services and work alongside the pre-existing ENT services. Sound Seekers is a UK registered charity, established in 1959, which aims to "deliver practical solutions to support people with hearing loss to realise their rights by enabling access to healthcare and education" [24].

In November 2015, Sound Seekers signed a Memorandum of Understanding with the Malawian Ministry of Health (MoH) to work with QECH to advance audiology services. This ensured commitment and awareness of the need for ear and hearing care services at a government level and outlined the

objectives of both parties. Sound Seekers agreed to manage the construction of the clinic, establish the services and train personnel whilst the MoH committed to maintaining the services after the withdrawal of Sound Seekers' support and ensure the trained personnel were employed within the department with job roles corresponding to the relevant, specialist training they had received.

In 2015, with three-year funding from the UK Department for International Development (DFID) and Jersey Overseas Aid (JOA), Sound Seekers fully set up audiology services in QECH, including both clinic and outreach services. With the support of the MoH, the QECH Audiology department was opened in 2016. An experienced volunteer audiologist from the United States (CC) was appointed by Sound Seekers from 2015 – 2018 to manage the audiology clinic and train the new audiology staff at QECH.

Sound Seekers funded the training of the first two Malawian audiologists who successfully completed their Masters level (MSc) Audiology training at the University of Manchester (UK) in 2015. At the time of writing, there is one audiologist, five audiology officers, one ear mould technician and an office manager employed by the QECH audiology clinic. Audiology officers are nurses or clinical officers who have received diploma qualifications in audiology or hearing aid acoustics. In April 2018, the volunteer Audiologist handed the management and operations of the clinic and outreach services to the Malawian audiologists, Following completion of the project, all services were handed over to QECH in April 2019. Since the handover, as committed by the MoH, the clinic services continue. However, outreach services have been limited to periods when there is funding available either through the hospital or external organisations.

All clinical equipment to render comprehensive audiological services were procured with funding from donations to Sound Seekers. Audiologists and audiology officers at QECH provide diagnostic testing for patients of all-ages including: pure tone audiometry (PTA), visual reinforcement audiometry (VRA), auditory brainstem response (ABR), auditory steady state response (ASSR) and otoacoustic emissions (OAE). 9

For each clinical assessment, a detailed history is taken, followed by ear examination and formal audiological assessment. Occluding ear wax and/or foreign bodies are removed from the ear canal

before the hearing assessment. Intervention options at the service include hearing aid fittings, ear wax removal and onward referral to ENT and other specialist services.

Hearing aid provision at QECH, and in Malawi more broadly, is mostly donor dependent. In 2015, a 'Hearing Aid Refurbishment Programme (HARP)' was established by Sound Seekers. The HARP supports a small audiology lab based in Zambia to refurbish digital hearing aids donated from individuals and hospitals in the UK. Refurbished hearing aids are then distributed to all Sound Seekers project countries and custom ear moulds are made at each site. The QECH audiology department generates limited income through health insurance payments and patient fees for services and hearing aids. Those that cannot pay receive services without cost. There are no local providers of ear mould materials but QECH audiology clinic has been able to overcome this challenge by using locally available, low cost dental alginate for ear impressions and dental acrylic to create the custom ear moulds. The clinicians have also collaborated with the QECH's bioengineering department to make some of the consumables locally thus reducing the cost of the entire ear mould manufacturing process.

In association with other international charities, visiting ENT surgeons and a cochlear implant manufacturer (MED-EL), 17 children identified with severe to profound hearing loss have successfully received a cochlear implant since 2015 with follow up care provided at QECH audiology department [25]. In addition to clinic-based services, the clinic runs an outreach programme to districts in the Southern province of Malawi where patients are screened, simple treatments provided, hearing aids fitted and referrals made for more complex cases. A sound proofed hearing assessment and research clinic (HARK) vehicle is used for assessments during outreach activities. The audiology service has also developed a bespoke patient database, presenting an opportunity to analyse the profile of patients attending the clinic. Every patient's demographic and clinical data is recorded onto an Excel data collection sheet specifically designed to collate and facilitate analysis of data. Demographic data is collected including gender, place of residence and age. Clinical data includes information pertinent to potential causes of hearing loss, type of assessment carried out, clinical findings and interventions. Throughout the development of the clinic, audiology staff received data collection training and support from Sound Seekers to ensure the data was collected reliably and recorded accurately.

The World Health Assembly resolution on the prevention of deafness and hearing loss (WHA70.13), published in May 2017, provided nine action points for member states [26]. This research article is able to provide potential approaches for the following: “to integrate strategies for ear and hearing care within the framework of their primary health care system” and “to improve access to affordable, cost-effective, high-quality, assistive hearing technologies and products, including hearing aids”. Specifically, we focus on the development of the QECH audiology department and the profile of patients attending the clinic in the first two years of the service.

Methods

Study design

A retrospective patient record review of patient’s assessed at the QECH audiology department between January 2016 and December 2017 was conducted. A convenience sampling approach was taken to include all new patients, seen within QECH audiology department between January 2016 and December 2017 in the study sample.

Participants

A total of 2299 patients, 1521 adults and 778 children, were assessed at the QECH audiology department between January 2016 and December 2017. Adult patients’ ages ranged from 18 to 94 years (Mean = 45.8, SD = 19.22). The mean age of children included in this study was 7.7 years (SD= 5.21). Overall, 45.4% of patients identified as female.

Data collection procedures

Demographic and clinical data was extracted from the department's clinical records excel spreadsheet including patterns of diagnosis and interventions.

Definitions

The World Health Organization (WHO) definition of disabling hearing loss and hearing loss grading system was used to characterise each patient’s hearing loss [1].

As the database described the configuration and degree of hearing loss qualitatively rather than specifying the actual pure tone average threshold of each ear, the following assumptions were made around the degree of loss for the purposes of reporting in this study:

- For sloping losses that were minimal (slope from one degree to the next) the degree was coded according to the better category (e.g. normal to mild was coded as mild);
- For steeply sloping losses (jump two or more degree categories), the degree was coded according to the average of the two extremes (e.g. mild to severe was coded as moderate)
- For some losses the categorisation was less clear cut, for example “normal to moderately severe”, and these were therefore coded the same as for “normal to severe” (moderate).
- The type of hearing loss was coded according to the better ear

~~The type of hearing loss was coded according to the better ear and in line with the WHO definitions [27].~~

Ethical considerations

This study was approved by the College of Medicine Research and Ethics Committee (COMREC), University of Malawi (P.04/17/2153). No patient identifiable data was extracted or presented in this study.

Data Analysis

Data analysis was conducted in Stata (version 15.0) to arrange the data and give basic descriptive statistics (mean and standard deviation). *R* version 3.6.1 was used to perform a binary logistic regression analysis to predict year (2016 vs. 2017) in which a patient was examined from categorical variables described that patient: age (child, adult); sex (female, male); hearing loss diagnosis (yes, no); recommended intervention; degree of hearing loss. Patient demographics included age, gender and location. The main study outcomes included type of diagnostic assessment, degree and type of hearing loss and type of audiological intervention offered. Results for adults and children were disaggregated for each of the outcomes.

Results

The number of patients assessed in 2017 (n=1385) was 34% higher than that of 2016 (n=914). In terms of locality, 83% of patients originated from the Southern region of Malawi, with 8.8% from other Malawian regions and 1.3% were from other countries (the location data was only available for patients seen in 2016). Over the two-year period, 41.7% of children and 61.6% of adults attending the clinic were found to have some degree of hearing loss (Table 1).

Table 1. Patients with hearing loss according to age group

Age category (years)	Total number of patients with hearing loss Total % (n) (n= 1262)
0-5	6.6 (83)
6-10	7.5 (95)
11-17	11.7 (147)
18-29	17.9 (226)
30-39	13.6 (171)
40-49	8.9 (112)
50-59	8.8 (111)
60-69	11.9 (150)
70-79	7.5 (94)
80+	5.8 (73)

Of those with hearing loss, 80.9% of children and 77.0% of adults with hearing loss presented with a bilateral hearing loss. Sensorineural hearing loss was the most common type of loss affecting 66.8% of children and 67.8% of adults with hearing loss. Conductive hearing loss was present in 20% of children and 13.9% of adults with hearing loss. Table 2 describes the type, degree and laterality of hearing losses found in paediatric and adult patients assessed at QECH over the two-year period.

Table 2. Summary of the degree, type and laterality of hearing losses present in the adult and paediatric population		
	Children n= 325	Adults n= 937
	Total %(n)	Total %(n)
Degree of hearing loss (better ear)		
Normal	13.9 (45)	21.0 (206)
Mild	16.3 (53)	16.65 (156)
Moderate	23.1 (75)	33.1 (310)
Severe	11.7 (38)	15.7 (147)
Profound	26.8 (87)	7.4 (69)
Unknown	8.3 (27)	5.2 (49)
Type of hearing loss (better ear)		
Conductive	20.0 (65)	13.9 (130)
Sensorineural	66.8 (217)	67.8 (635)
Mixed	5.9 (19)	15.2 (142)
Unknown	1.2 (4)	0.5 (5)
Missing	6.2 (20)	2.7 (25)
Laterality of hearing loss		
Bilateral	80.9 (263)	76.9 (721)
Unilateral	15.4 (50)	22.5 (211)
Unknown	3.7 (12)	0.5 (5)

Table 3 highlights the causes of hearing loss in adults in children assessed at QECH. Overall, 18.5% of children with hearing loss presented with chronic suppurative otitis media (CSOM) and 8.6% had a history of ototoxic medication administration. Presbycusis was a major cause of hearing loss in adults (19.7%). CSOM was the cause for 14.4% of adults. The cause of hearing loss was unknown for 56.7% of adults with hearing loss, and 65.9% among children.

Table 3. Suspected causes of hearing loss for children and adult patient's diagnosed with hearing loss in January 2016- December 2017

Suspected cause of hearing loss	Children %(n) n=325	Adults %(n) n=937
ANSD	0.6 (2)	0.1 (1)
CSOM	18.8 (61)	14.4 (135)
Malaria	0.9 (3)	0
Meningitis	1.9 (6)	1.1 (11)
Mumps	0.9 (3)	0.1 (1)
OME	2.2 (7)	2.1 (20)
Ototoxicity	8.6 (29)	3.5 (33)
Presbycusis	0	19.7 (185)
Tuberculosis	0	1.3 (12)
Trauma	0.3 (1)	0.8 (7)
Noise	0	0.1 (1)
Unknown	65.9 (214)	56.7 (531)

ANSD: Auditory Neuropathy Spectrum Disorder

CSOM: Chronic Suppurative Otitis Media

OME: Otitis Media with Effusion

PTA was the most commonly used assessment method in 2016 and 2017 and was performed on 50% of all patients assessed in the two-year period (Table 4). The number of patients tested using electrophysiological methods increased from 20 patients in 2016 to 66 patients in 2017. The use of OAEs also increased from 187 patients in 2016 to 452 in 2017. VRA and play audiometry were reported to have only been used to assess a combined total of 32 patients across both years. Tympanometry was carried out on 84% of patients to assess middle ear function.

Table 4. Type of audiological assessment carried out for patients with and without hearing loss in 2016 and 2017.

Type of assessment	Patients with hearing loss n=1262		Patients without hearing loss n=1037		Total Patients 2016 n=914	Total Patients 2017 n= 1385
	2016 % (n)	2017 % (n)	2016 % (n)	2017 % (n)	2016 % (n)	2017 % (n)
Pure tone audiometry (PTA)	79.8 (423)	59.8 (438)	35.7 (137)	25.0 (163)	61.3(560)	43.4(601)
Play audiometry	0.2 (1)	0.6 (4)	0.3 (1)	0.2 (1)	0.2(2)	0.4(5)
VRA	0.2 (1)	0.3 (2)	3.1 (12)	1.5 (10)	1.4(13)	0.9(12)
OAE	15.7 (83)	31.3 (229)	27.1 (104)	34.2 (223)	20.5(187)	32.6(452)
ABR	0	0.1 (1)	0.3 (1)	0.8 (5)	0.1(1)	0.43(6)

ASSR	3.6 (19)	5.9 (43)	0	2.7 (17)	2(19)	4.3(60)
Data missing	0.6 (3)	2.0 (15)	33.6 (129)	35.8 (234)	14.4(132)	17.9(249)

VRA: Visual Reinforcement Audiometry, OAE: Otoacoustic Emissions, ABR: Auditory Brainstem Response, ASSR: Auditory Steady State Response

275
276

277 Table 5 summarises the recommended interventions for patients diagnosed with hearing loss. In
278 2016, 21.6% of children with hearing loss were fitted with hearing aids, compared to 12% in 2017.
279 Similarly, 39.4% of adults with hearing loss were fitted with hearing aids in 2016 and this decreased to
280 19.5% in 2017 (Table 5). Referrals to ENT increased over time. In 2016, 16.4% of children with
281 hearing loss and 9.9% of adults with hearing loss were referred to ENT compared to 21.5% and
282 17.2%, respectively in 2017. A further 11.4% of children with hearing loss were referred to a deaf
283 school or special education resource unit and 10.5% were recommended to have sign language
284 training. In order to statistically analyse the differences observed between the 2016 and 2017 patient
285 cohorts, a binary logistic regression was performed, predicting year (2016 vs. 2017) in which a patient
286 was examined from categorical variables described that patient: age (child, adult); sex (female, male);
287 hearing loss diagnosis (yes, no); recommended intervention; degree of hearing loss. . All two-way
288 interactions were included in the model to account for any differences in the distribution across
289 various levels of the categorical variables across the two years that should be attributed to other
290 non-significant differences in category membership. In this model, there were only two
291 coefficients that were significant predictors of year, other than the intercept, and both pertained to
292 recommended intervention (Table 5). The binomial variable indicating that sign language training
293 was recommended ($z = 2.781$, $p = 0.00542$) and the variable indicating that the patient was fitted
294 with a hearing aid ($z = -2.381$, $p = 0.0173$) were the two significant predictors of the year in which
295 a given patient was treated in the logistic regression model. The proportion of patients with
296 hearing loss that were referred to sign language training increased by 4.6% from 2016 to 2017.
297 There was a 12% decrease in the proportion of patients with hearing loss that were fitted with
298 hearing aids from 2016 to 2017.

299
300
301

Table 5. Recommended interventions for patients diagnosed with hearing loss

p-value column indicates the significance of the main effect of the individual treatment management plan variables, as regressors in a logistic regression, modeling the probability of a given observation being drawn from the 2016 vs. the 2017 sample.

* $p < .05$. ** $p < .01$.

— indicates data not included in the logistic regression

Individual Management Plan	Children (n=325)			Adults (n=937)			Combined (n=1262)			p-value
	2016 % (n)	2017 % (n)	Both % (n)	2016 % (n)	2017 % (n)	Both % (n)	2016 % (n)	2017 % (n)	Both % (n)	
Review in audiology	16.4 (19)	19.6 (41)	18.5 (60)	18.4 (76)	16.8 (88)	17.5 (164)	17.9 (95)	17.6 (129)	17.7 (224)	—
Hearing aid fitted	21.6 (25)	12 (25)	15.4 (50)	39.4 (163)	19.5 (102)	28.3 (265)	35.5 (188)	17.3 (127)	25 (315)	0.017 *
Hearing aid recommended	15.5 (18)	13.4 (28)	14.2 (46)	14.7 (61)	30 (157)	23.3 (218)	14.9 (79)	25.3 (185)	20.9 (264)	0.553
Medication	0 (0)	1.91 (4)	1.23 (4)	0 (0)	3.25 (17)	1.81 (17)	0 (0)	2.87 (21)	1.66 (21)	0.990
Refer to Ear, Nose, Throat	16.4 (19)	21.5 (45)	19.7 (64)	9.9 (41)	17.2 (90)	14 (131)	11.3 (60)	18.4 (135)	15.5 (195)	0.973
Refer to Deaf school	17.2 (20)	8.13 (17)	11.4 (37)	1.21 (5)	0.191 (1)	0.64 (6)	4.72 (25)	2.46 (18)	3.41 (43)	0.896
Refer elsewhere	1.72 (2)	3.35 (7)	2.77 (9)	0 (0)	0 (0)	0 (0)	0.377 (2)	0.956 (7)	0.713 (9)	0.573
Recommend sign language training	3.45 (4)	14.4 (30)	10.5 (34)	2.66 (11)	5.16 (27)	4.06 (38)	2.83 (15)	7.79 (57)	5.71 (72)	0.005 **
Recommend speech therapy	0 (0)	0.957 (2)	0.615 (2)	0 (0)	0 (0)	0 (0)	0 (0)	0.273 (2)	0.158 (2)	0.993
Tinnitus counselling	0 (0)	0 (0)	0 (0)	0.725 (3)	2.1 (11)	1.49 (14)	0.566 (3)	1.5 (11)	1.11 (14)	0.988
Other	0 (0)	0 (0)	0 (0)	0.725 (3)	0.765 (4)	0.747 (7)	0.566 (3)	0.546 (4)	0.555 (7)	0.781
No further action	7.76 (9)	4.78 (10)	5.85 (19)	11.8 (49)	4.97 (26)	8 (75)	10.9 (58)	4.92 (36)	7.45 (94)	0.106
Data missing	0 (0)	0 (0)	0 (0)	0.483 (2)	0 (0)	0.213 (2)	0.377 (2)	0 (0)	0.158 (2)	—

Discussion

This study provides a case study of a comprehensive, public sector audiology clinic in Malawi. It also details the profile and clinical findings of patients attending the clinic between 2016 and 2017. Due to the increasing awareness of the availability of specialist ear and hearing care services at QECH, the number of patients assessed in 2017 was double that of 2016 (Table 2).

Types and causes of hearing loss

The cause of hearing loss was unknown for the majority of patients with hearing loss (56.7% of adults and 65.9% of children). This could be due to several reasons. Firstly, genetic testing and other aetiological investigations of hearing loss were not available at QECH. Further, even if more resources were available, it is possible that the cause would remain unknown. For instance, if an individual developed hearing loss due to an associated viral infection, it may go unnoticed until later in life making it very difficult to ascertain the origin of hearing loss from the medical history at the time of assessment. As another example, for patients with multiple health conditions, there may be numerous potential causes of hearing loss present in the same individual.

In this study, the most common identifiable cause of hearing loss in children was CSOM (18.8%). Chronic suppurative otitis media (CSOM) is particularly prevalent, preventable cause of acquired hearing loss in school aged children [28, 29] and is characterised by persistent discharge from the middle ear through ear drum perforation often associated with varying levels of hearing loss [30]. Population-based studies have also found CSOM to be a very common cause of hearing loss in children in Swaziland and Botswana [31] and up to 90% of CSOM cases can occur in low income countries [29]. A systematic review reported that 24% of preschool children had a hearing loss due to ear wax impaction [15]. Due to the nature of the clinical data presented in this study, wax impaction is not cited as a reason for hearing loss because impacted wax is removed prior to hearing assessment at QECH audiology department. According to medical files and the individual patient's medical history, 8.6% of children with hearing loss had been administered with some form of ototoxic medication. This is a higher

than the data published by WHO that estimates 4% of childhood hearing losses to be attributed to ototoxic medication [32].

There are a limited number of studies detailing the clinical findings of an audiology clinic a low resource setting such as Malawi. Lasisi et al. (2006) reported results of a retrospective review of paediatric patients (n=713), seen at an outpatient otorhinolaryngologic clinic in Ibadan, Nigeria [33] and found 14% of children had some form of sensorineural hearing loss. This was considerably lower than the results reported in the present study. Lasisi et al (2006) recognised the unrepresentatively low number of people with sensorineural hearing losses attending the otorhinolaryngologic clinic and cited potential reasons including poor access to healthcare and widespread poverty [33]. The number of children presenting with CSOM was also lower (4.9% vs 18.8%) than reported in the present study but hearing losses associated with ototoxic medication was similar (12.6% vs 8.6%).

Banda et al (2018) described the clinical findings of children seen in a public sector audiology department in Botswana and reported 32% of children under 10 years of age presented with hearing loss [34]. This is similar to the proportion of similar aged children with hearing loss found in the present study (34%). However, the number of children that present with severe to profound levels of hearing loss at audiology clinics varies in LMICs. Banda et al (2018) reported 36.6% of hearing loss in children (under 10 years of age) was severe to profound in degree [34] which is much lower than in the present study (54.8%) and from clinic based data from rural Nicaragua (86%) [3], when comparing children of the same age range. It is likely that parents seek ear and hearing care services for their children once their difficulties appear more apparent which might explain the high levels of severe to profound hearing loss detected at the clinic level.

Audiological Assessment

The most commonly used assessment method was PTA. Tympanometry was used to assess the middle ear function of 84% of patients and screening OAEs were used to assess 27% of patients, most likely as they are known to be quick to administer and easy to interpret. ABR, ASSR and VRA methods were seldom used. Reasons for a low number of electrophysiological tests carried out include the presence of high electrical interference in the clinic, lack of the training for clinicians to carry out ABR threshold testing and lack of locally available consumables. This highlights the need for further training and a review of the types of audiological tests used to ensure they are suitable for the clinical skills, resources and environment available in low resource settings.

Hearing aid uptake

Overall, the uptake of hearing aids was low at QECH (28.3% of adults with hearing loss) and the number of patients fitted with hearing aids decreased from 2016 to 2017. This was surprising due to the adequate number of trained staff and suitable audiological equipment available throughout this time frame and the low-cost nature of providing refurbished hearing aids. However, there are a number of possible reasons to account for the poor uptake. Firstly, during this timeframe, the number of people with profound hearing loss increased, particularly in the paediatric population. People with very limited residual hearing are less likely to use or benefit from hearing aid technology [35]. Furthermore, and specific to this study's population, there are some practical barriers to uptake of healthcare as shown in a study by Bright et al. (2017) which found a 3% uptake of referrals to QECH ear and hearing care services for children residing in a district located 60km away [18]. Although healthcare is free at the point of access in Malawi, Bright et al (2017) reported that families faced challenges in acquiring the necessary funds to afford indirect costs of attending hospital appointments (food, travel and accommodation) as well as feeling fearful or uncertain about the hospital itself [18]. Limited knowledge of ear and hearing health in the general population has also been reported to be a significant barrier to referral uptake in Malawi [18]. These factors, as well as the prospect of needing funds for regular visits to the hospital for hearing aid aftercare and batteries, may have also affected hearing aid uptake in the current study.

Ramma and Sebothoma (2017) explored factors associated with seeking audiology services for people with hearing loss in Cape Town, South Africa. Despite, audiology services being available at minimal cost in South Africa, only 21% (35/166) of individuals with hearing loss reported seeking support for their hearing loss via audiology services. The researchers found that self-perceived hearing loss was associated with higher likelihood of seeking help but other factors including age, gender, severity of hearing loss, level of education and employment status were not found to be associated [36]. Other studies have suggested that certain cultural and traditional views of medical care [37] as well as the perception that hearing loss is not considered to be a life-threatening condition [38] should be taken into account when observing reluctance to seek help.

The provision of hearing aids is one of the most common management options for people with hearing loss but hearing aid uptake varies globally. It is estimated that the hearing aid adoption rate for developed countries like the United Kingdom and United States is between 20-25% of people with hearing loss whereas in developing countries adoption rates have been estimated to be 1-8% [39, 40]. Hearing aid uptake is a multi-faceted issue and can be affected by cultural opinions of hearing loss, social representation of hearing aids, stigma and a general lack of awareness regarding hearing difficulties and the benefit of using hearing aid technology [38, 41]. A retrospective study reviewing audiological data (n=3894) from an audiology clinic in Limpopo province, South Africa, found that 46% of patients (n=1778) were diagnosed with disabling hearing loss between 2012 and 2014. Of these, only 15% were fitted with hearing aids [42]. In contrast to the present study, the authors cited the distribution of funding within the healthcare system as a key factor affecting hearing aid uptake as the allocated hospital budget could only cover the purchase of 272 hearing aids.

Finally, in the present study, 14.2% of children and 23.3% of adults with hearing loss were recommended a hearing aid (Table 5) but had not been fitted. This may have occurred if a patient declined a hearing aid or a hearing aid fitting was booked for a later date. If hearing aid fitting was arranged for a later date these patients would not have been included in overall number of new patients fitted with hearing aids and this may have caused an underestimation in hearing aid uptake from our sample. The clinical database needs to be updated in a timely manner in order to capture a patient's complete journey through the audiology services.

429 *Recommendations for the strengthening of audiological services in Malawi*

430 The findings presented in this study highlight the need for further research to investigate long-term
431 hearing aid usage and reasons for poor uptake of hearing aids at QECH. Additionally, the development
432 of audiology clinics in rural communities could help strengthen support, raise awareness of the impact
433 of hearing loss and ensure patients are not lost to follow up.

434 The QECH audiology department is the product of collaboration between local, governmental and
435 international partners. An external organisation funded the infrastructure, equipment and training
436 opportunities and the MoH now employ all staff members and the department is embedded in the
437 hospital system. Despite this progress, there is still much work to do for these services to be sustainable.
438 Traditionally, audiology clinics have relied on very costly technology for hearing testing and hearing aid
439 fitting. The lack of local low-cost options makes the clinic reliant on external organisations for
440 refurbished hearing aids and equipment donations until other options become available. Recently, WHO
441 recommended a number of technological and workforce innovations that could lead to increased access
442 to hearing aids in low income countries [43]. These included the use of low-cost hearing aid technology
443 and models of community delivered hearing care [43-46]. It would be beneficial to pilot these innovations
444 in LMICs, particularly in clinics like QECH where there are trained professionals and appropriate testing
445 environments.

446 .

447 In this study, we found 6.7% of patients presented with profound hearing loss. These patients will not
448 gain significant benefit from hearing aids but due to lack of capacity, funds and human resources, many
449 will not have access to cochlear implantation. It is important for audiology services to maintain strong
450 links with the education sector to help teachers better support children with hearing difficulties and
451 allow for more timely referrals to ear , hearing care and sign language support..

452 There is still a significant shortage of ear and hearing care health professionals in Malawi and audiology
453 is not an established profession. Further work is needed to raise the profile of the audiology workforce
454 in Malawi to ensure health professionals obtain the appropriate employment positions, determined by
455 their specialist audiology training. Additionally, when choosing candidates for audiology training
456 programmes, candidates need the sufficient, relevant pre-training qualifications to allow them to be
457 employed by the MoH. Clinical staff at QECH audiology department still have regular access to remote

support and advice from Sound Seekers management team and experienced volunteer audiologists. The mentoring and continued clinical and counselling skills training of audiology health professionals and community health workers and the use of task shifting is vital for the continuity of healthcare in LMICs, especially in rural communities [47-49]. Bilateral mentorship between global partners and Malawian ear and hearing care professionals is particularly beneficial as the profession is emerging but still in its infancy [50]. The Vision 2020 Links programme has successfully established a network of global eye care training institutions and similar initiatives could be developed to facilitate ear care training [51].

Finally, in order for services to be sustainable, work should be carried out collaboratively between the MoH, existing hospital staff and external organisations to encourage expansion of the current healthcare system to include audiology staffing positions, resource budgeting and succession planning. Recent work has highlighted the benefits of training community health workers in hearing screening and delivery of basic treatments in LMICs [52, 53] as well as the use of validated hearing screening smartphone applications in these populations [54, 55]. These options should be explored further, with a thorough analysis of long-term cost effectiveness and reliability as well as follow up provision, to enhance access to sustainable, comprehensive ear and hearing care services.

Strengths, limitations and future work

The QECH audiology service has been successful in providing audiological care to patients in Southern Malawi since 2016. The department is supported by the Malawian MoH making it a pioneer in sub-Saharan Africa. A key strength of the current study is the amount of data presented including type of clinical assessment, clinical findings and intervention options for patients of all ages seen in the two-year period. Data of this kind, from an audiology clinic in a low resource setting, is relatively rare and can serve to strengthen the processes taking place within the clinic and help the development of other services in other LMICs.

There are also some limitations. Firstly, there is a potential underestimation of conductive hearing loss present in this study. Occluding ear wax is removed prior to hearing testing and patients with flat tympanograms, indicating presence of middle ear effusion, are immediately referred to ENT to be

administered with medical treatment. Therefore, those with possible conductive hearing loss associated with middle ear effusion and/or wax impaction are not recorded in the clinical database. Another limitation is the database is incomplete, fields including patient's residential location and type of assessment carried out were not completed for all patients included in this study. The data presented here provides a snapshot of the management options administered to patients after the initial diagnosis, including the fitting of hearing aids. However, it does not capture patients' hearing aid usage and progress. Future research should include the use of outcome measures of hearing aid benefit and analysis of the long-term impact of audiology services in LMICs, evidence in these areas is lacking in LMICs. Finally, formal impact and process evaluations of QECH Audiology services could help to understand what worked well and why in order to help others to implement similar programmes.

Conclusion

In this study, we provided a case study of a comprehensive audiology clinic in Blantyre, Malawi which could act as a service provision model for low resource settings. Results indicate a growing demand for ear and hearing care services and have identified specific training needs for clinical staff. Further research is needed to understand the reasons for the low hearing aid uptake found in this study in order to improve long term access to these services.

516

517 References

518

519 1. World Health Organization. Prevention of blindness and deafness: Grades of hearing

520 impairment 2019 [cited 06/02/2020]. Available from:

521 https://www.who.int/pbd/deafness/hearing_impairment_grades/en/.

522 2. Olusanya BO, Swanepoel DW, Chapchap MJ, Castillo S, Habib H, Mukari SZ, et al. Progress

523 towards early detection services for infants with hearing loss in developing countries. BMC Health

524 Serv Res. 2007;7:14-. doi: 10.1186/1472-6963-7-14. PubMed PMID: 17266763.

525 3. Saunders JE, Vaz S, Greinwald JH, Lai J, Morin L, Mojica K. Prevalence and etiology of

526 hearing loss in rural Nicaraguan children. Laryngoscope. 2007;117(3):387-98. Epub 2007/03/06. doi:

527 10.1097/MLG.0b013e31802e3726. PubMed PMID: 17334299.

528 4. Malema KM. The Working Conditions In The Mining Sector: The Case Of Mchenga And

529 Kaziwiziwi Coal Mines In Malawi. Open Science Journal. 2017;2.

530 5. Wilson BS, Tucci DL, Merson MH, O'Donoghue GM. Global hearing health care: new findings

531 and perspectives. The Lancet. 2017;390(10111):2503-15. doi: [https://doi.org/10.1016/S0140-](https://doi.org/10.1016/S0140-6736(17)31073-5)

532 [6736\(17\)31073-5](https://doi.org/10.1016/S0140-6736(17)31073-5).

533 6. Honeycutt A. Economic costs of mental retardation, cerebral palsy, hearing loss and vision

534 impairment. In: Grosse Scott D, Barbara MA, Sharon NB, Gerry EH, Sheryl AL, editors. Using Survey

535 Data to Study Disability: Results from the National Health Survey on Disability. Research in Social

536 Science and Disability. 3: Emerald Group Publishing Limited; 2003. p. 207-28.

537 7. Lin FR, Metter EJ, O'Brien RJ, Resnick SM, Zonderman AB, Ferrucci L. Hearing loss and

538 incident dementia. Arch Neurol. 2011;68(2):214-20. Epub 2011/02/16. doi:

539 10.1001/archneurol.2010.362. PubMed PMID: 21320988; PubMed Central PMCID:

540 PMCPMC3277836.

541 8. Bagli Z. Multicultural aspects of hearing loss. Communication Disorders in Multicultural and

542 International Populations. 2012:208-42. doi: 10.1016/B978-0-323-06699-0.00020-0.

543 9. Graydon K, Waterworth C, Miller H, Gunasekera H. Global burden of hearing impairment and

544 ear disease. The Journal of Laryngology & Otology. 2019;133(1):18-25. Epub 2018/07/26. doi:

545 10.1017/S0022215118001275.

546 10. Orji A, Kamenov K, Dirac M, Davis A, Chadha S, Vos T. Global and regional needs, unmet
547 needs and access to hearing aids. *Int J Audiol*. 2020;1-7. Epub 2020/02/06. doi:
548 10.1080/14992027.2020.1721577. PubMed PMID: 32011190.

549 11. Mulwafu W, Fagan JJ, Mukara KB, Ibekwe TS. ENT Outreach in Africa: Rules of
550 Engagement. *OTO Open*. 2018;2(2):2473974x18777220. Epub 2018/11/28. doi:
551 10.1177/2473974x18777220. PubMed PMID: 30480217; PubMed Central PMCID:
552 PMCPMC6239148.

553 12. National Statistics Office. Malawi Population and Housing Census: Main Report 2019
554 [updated May 2019]. 1]. Available from:
555 [http://www.nsomalawi.mw/images/stories/data_on_line/demography/census_2018/2018%20Malawi%](http://www.nsomalawi.mw/images/stories/data_on_line/demography/census_2018/2018%20Malawi%20Population%20and%20Housing%20Census%20Main%20Report.pdf)
556 [20Population%20and%20Housing%20Census%20Main%20Report.pdf](http://www.nsomalawi.mw/images/stories/data_on_line/demography/census_2018/2018%20Malawi%20Population%20and%20Housing%20Census%20Main%20Report.pdf).

557 13. The World Bank Group. Classifying countries by income 2019 [cited 05.05.2020]. Available
558 from: [https://datatopics.worldbank.org/world-development-indicators/stories/the-classification-of-](https://datatopics.worldbank.org/world-development-indicators/stories/the-classification-of-countries-by-income.html)
559 [countries-by-income.html](https://datatopics.worldbank.org/world-development-indicators/stories/the-classification-of-countries-by-income.html).

560 14. World Health Organization. Addressing the rising prevalence of hearing loss. Geneva: World
561 Health Organization, 2018 2018. Report No.: 9789241550260.

562 15. Mulwafu W, Kuper H, Ensink RJ. Prevalence and causes of hearing impairment in Africa.
563 *Trop Med Int Health*. 2016;21(2):158-65. Epub 2015/11/21. doi: 10.1111/tmi.12640. PubMed PMID:
564 26584722.

565 16. Hunt L, Mulwafu W, Knott V, Ndamala CB, Naunje AW, Dewhurst S, et al. Prevalence of
566 paediatric chronic suppurative otitis media and hearing impairment in rural Malawi: A cross-sectional
567 survey. *PLOS ONE*. 2017;12(12):e0188950. doi: 10.1371/journal.pone.0188950.

568 17. Mulwafu W, Tataryn M, Polack S, Viste A, Goplen FK, Kuper H. Children with hearing
569 impairment in Malawi, a cohort study. *Bull World Health Organ*. 2019;97(10):654-62. Epub
570 2019/10/28. doi: 10.2471/blt.18.226241. PubMed PMID: 31656330; PubMed Central PMCID:
571 PMCPMC6796677.

572 18. Bright T, Mulwafu W, Thindwa R, Zuurmond M, Polack S. Reasons for low uptake of referrals
573 to ear and hearing services for children in Malawi. *PLoS One*. 2017;12(12):e0188703. Epub
574 2017/12/21. doi: 10.1371/journal.pone.0188703. PubMed PMID: 29261683; PubMed Central PMCID:
575 PMCPMC5736203.

576 19. Mulwafu W, Ensink R, Kuper H, Fagan J. Survey of ENT services in sub-Saharan Africa: little
577 progress between 2009 and 2015. *Glob Health Action*. 2017;10(1):1289736. Epub 2017/05/10. doi:
578 10.1080/16549716.2017.1289736. PubMed PMID: 28485648; PubMed Central PMCID:
579 PMCPMC5496047.

580 20. Baum A, Mulwafu W, Phiri M, Polack S, Bright T. An Intervention to Improve Uptake of
581 Referrals for Children with Ear Disease or Hearing Loss in Thyolo District, Malawi: Acceptability and
582 Feasibility. *Int J Environ Res Public Health*. 2019;16(17):3144. doi: 10.3390/ijerph16173144. PubMed
583 PMID: 31466419.

584 21. Fagan JJ, Jacobs M. Survey of ENT services in Africa: need for a comprehensive
585 intervention. *Glob Health Action*. 2009;2. Epub 2009/12/23. doi: 10.3402/gha.v2i0.1932. PubMed
586 PMID: 20027268; PubMed Central PMCID: PMCPMC2779942.

587 22. Mulwafu W, Nyirenda TE, Fagan JJ, Bem C, Mlumbe K, Chitule J. Initiating and developing
588 clinical services, training and research in a low resource setting: the Malawi ENT experience. *Trop*
589 *Doct*. 2014;44(3):135-9. Epub 2014/02/27. doi: 10.1177/0049475514524393. PubMed PMID:
590 24569097.

591 23. Mulwafu W, Fagan J. Improving access to hearing care services and professionals in Africa
592 through task sharing: The Malawian experience. *East and Central African Journal of Surgery*.
593 2019;24(2):78-81.

594 24. Sound Seekers. Vision, Mission and Values 2019 [cited 05.05.2020]. Available from:
595 <https://www.sound-seekers.org.uk/about/>.

596 25. Mulwafu W, Strachan DR, Bartlett R, Caron C. Cochlear implantation in Malawi: report of the
597 first four cases. *J Laryngol Otol*. 2017;131(10):914-8. Epub 2017/08/16. doi:
598 10.1017/s0022215117001724. PubMed PMID: 28807077.

599 26. World Health Assembly. Prevention of deafness and hearing loss: WHA70.13. Geneva: 2017
600 Contract No.: 15.8.

601 27. World Health Organization. Deafness and hearing loss: World Health Organization; 2020
602 [cited 2020 7th April 2020]. Available from: [https://www.who.int/news-room/fact-](https://www.who.int/news-room/fact-sheets/detail/deafness-and-hearing-loss)
603 [sheets/detail/deafness-and-hearing-loss](https://www.who.int/news-room/fact-sheets/detail/deafness-and-hearing-loss).

28. Daly KA, Selviu RE, Lindgren B. Knowledge and attitudes about otitis media risk: implications for prevention. *Pediatrics*. 1997;100(6):931-6. Epub 1997/12/31. doi: 10.1542/peds.100.6.931. PubMed PMID: 9374559.

29. World Health Organization. Chronic suppurative otitis media: Burden of illness and Management Options Geneva, : Child and Adolescent Health and Development: Prevention of Blindness and Deafness, 2004.

30. Jacob A, Rupa V, Job A, Joseph A. Hearing impairment and otitis media in a rural primary school in South India. *International Journal of Pediatric Otorhinolaryngology*. 1997;39(2):133-8. doi: [https://doi.org/10.1016/S0165-5876\(96\)01479-6](https://doi.org/10.1016/S0165-5876(96)01479-6).

31. Smith A, Hatcher J. Preventing deafness in Africa's children. *Afr Health*. 1992;15(1):33-5. Epub 1992/11/01. PubMed PMID: 12285914.

32. World Health Organization. Childhood hearing loss: Strategies for Prevention and Care 2016 [updated 06/02/2020]. 28]. Available from: https://www.who.int/docs/default-source/imported2/childhood-hearing-loss--strategies-for-prevention-and-care.pdf?sfvrsn=cbbbb3cc_0.

33. Lasisi OA, Ayodele JK, Ijaluola GT. Challenges in management of childhood sensorineural hearing loss in sub-Saharan Africa, Nigeria. *Int J Pediatr Otorhinolaryngol*. 2006;70(4):625-9. Epub 2005/09/20. doi: 10.1016/j.ijporl.2005.08.009. PubMed PMID: 16168496.

34. Banda FM, Powis KM, Mokoka AB, Mmapetla M, Westmoreland KD, David T, et al. Hearing Impairment Among Children Referred to a Public Audiology Clinic in Gaborone, Botswana. *Glob Pediatr Health*. 2018;5:2333794x18770079. Epub 2018/05/16. doi: 10.1177/2333794x18770079. PubMed PMID: 29761140; PubMed Central PMCID: PMC5946350.

35. Hjaldaahl J, Widén S, Carlsson P-I. Severe to profound hearing impairment: factors associated with the use of hearing aids and cochlear implants and participation in extended audiological rehabilitation. *Hearing, Balance and Communication*. 2017;15(1):6-15. doi: 10.1080/21695717.2016.1242250.

36. Ramma L, Sebothoma B, editors. Exploring Factors Associated with a Higher Likelihood of Seeking Audiology services by Individuals with Hearing Impairment in Cape Town , South Africa 2017.

37. Zhao F, Manchaiah V, St Claire L, Danermark B, Jones L, Brandreth M, et al. Exploring the influence of culture on hearing help-seeking and hearing-aid uptake. *Int J Audiol*. 2015;54(7):435-43. Epub 2015/03/13. doi: 10.3109/14992027.2015.1005848. PubMed PMID: 25761498.

- 634 38. Davis A, Smith P, Ferguson M, Stephens D, Gianopoulos I. Acceptability, benefit and costs of
635 early screening for hearing disability: a study of potential screening tests and models. *Health Technol*
636 *Assess.* 2007;11(42):1-294. Epub 2007/10/12. doi: 10.3310/hta11420. PubMed PMID: 17927921.
- 637 39. National Sample Survey Organisation. Disabled Persons in India. India: Ministry of Statistics
638 and Programme Implementation Government of India, 2003.
- 639 40. Wong LLN, McPherson B. Universal Hearing Health Care: China. *ASHA Leader.*
640 2008;13(17):14-. doi: 10.1044/leader.wb2.13172008.14. PubMed PMID: pub.1101704615.
- 641 41. Manchaiah V, Danermark B, Vinay, Ahmadi T, Tomé D, Krishna R, et al. Social
642 representation of hearing aids: cross-cultural study in India, Iran, Portugal, and the United Kingdom.
643 *Clin Interv Aging.* 2015;10:1601-15. doi: 10.2147/CIA.S86108. PubMed PMID: 26504376.
- 644 42. Hlayisi VG, Ramma L. Rehabilitation for disabling hearing loss: evaluating the need relative to
645 provision of hearing aids in the public health care system. *Disabil Rehabil.* 2019;41(22):2704-7. Epub
646 2018/05/22. doi: 10.1080/09638288.2018.1473507. PubMed PMID: 29779397.
- 647 43. Yong MW, A; McMahon, C; McPherson, B; Nieman, C; Reed, N; Lin, F. Access to adults'
648 hearing aids: policies and technologies used in eight countries. *Bulletin of the World Health*
649 *Organization.* 2019;97:699-710.
- 650 44. Nieman CL, Marrone N, Mamo SK, Betz J, Choi JS, Contrera KJ, et al. The Baltimore HEARS
651 Pilot Study: An Affordable, Accessible, Community-Delivered Hearing Care Intervention.
652 *Gerontologist.* 2017;57(6):1173-86. Epub 2016/12/09. doi: 10.1093/geront/gnw153. PubMed PMID:
653 27927734; PubMed Central PMCID: PMC5881797.
- 654 45. Medtronix. Shruti Case Study: Aligning Value [cited 06/02/2020]. Available from:
655 [https://www.medtronic.com/za-en/transforming-healthcare/aligning-value/perspective/case-](https://www.medtronic.com/za-en/transforming-healthcare/aligning-value/perspective/case-studies/shruti.html)
656 [studies/shruti.html](https://www.medtronic.com/za-en/transforming-healthcare/aligning-value/perspective/case-studies/shruti.html).
- 657 46. World Wide Hearing. What we do: Our Solution 2017 [cited 06/02/2020]. Available from:
658 <https://www.wwhearing.org/en/our-solution/>.
- 659 47. Munoz K. Counseling Skill Development in Audiology: Clinical Instruction Considerations.
660 *Seminars in hearing.* 2018;39(1):9-12. Epub 2018/02/10. doi: 10.1055/s-0037-1613701. PubMed
661 PMID: 29422709; PubMed Central PMCID: PMC5802981.

48. Rohatinsky N, Cave J, Krauter C. Establishing a mentorship program in rural workplaces: connection, communication, and support required. *Rural and remote health*. 2020;20(1):5640. Epub 2020/01/14. doi: 10.22605/rrh5640. PubMed PMID: 31928037.
49. Ekberg K, Grenness C, Hickson L. Addressing patients' psychosocial concerns regarding hearing aids within audiology appointments for older adults. *Am J Audiol*. 2014;23(3):337-50. Epub 2014/07/19. doi: 10.1044/2014_aja-14-0011. PubMed PMID: 25036799.
50. Vandjelovic ND, Sugihara EM, Mulwafu W, Madgy DN. The Creation of a Sustainable Otolaryngology Department in Malawi. *Ear, Nose & Throat Journal*. 2019;0145561319855366. doi: 10.1177/0145561319855366.
51. London School of Hygiene and Tropical Medicine ICfEH. VISION 2020 LINKS programme [cited 07.04.2020]. Available from: <https://iceh.lshtm.ac.uk/vision-2020-links-programme/>.
52. Mulwafu W, Kuper H, Viste A, Goplen FK. Feasibility and acceptability of training community health workers in ear and hearing care in Malawi: a cluster randomised controlled trial. *BMJ Open*. 2017;7(10):e016457. doi: 10.1136/bmjopen-2017-016457.
53. O'Donovan J, Verkerk M, Winters N, Chadha S, Bhutta MF. The role of community health workers in addressing the global burden of ear disease and hearing loss: a systematic scoping review of the literature. *BMJ Global Health*. 2019;4(2):e001141. doi: 10.1136/bmjgh-2018-001141.
54. Yousuf Hussein S, Wet Swanepoel D, Biagio de Jager L, Myburgh HC, Eikelboom RH, Hugo J. Smartphone hearing screening in mHealth assisted community-based primary care. *J Telemed Telecare*. 2016;22(7):405-12. Epub 2015/10/16. doi: 10.1177/1357633x15610721. PubMed PMID: 26468215.
55. Sandstrom J, Swanepoel D, Laurent C, Umejford G, Lundberg T. Accuracy and Reliability of Smartphone Self-Test Audiometry in Community Clinics in Low Income Settings: A Comparative Study. *Ann Otol Rhinol Laryngol*. 2020;3489420902162. Epub 2020/01/23. doi: 10.1177/0003489420902162. PubMed PMID: 31965808.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Parmar, B;Phiri, M;Caron, C;Bright, T;Mulwafu, W

Title:

Development of a public audiology service in Southern Malawi: profile of patients across two years.

Date:

2021-10

Citation:

Parmar, B., Phiri, M., Caron, C., Bright, T. & Mulwafu, W. (2021). Development of a public audiology service in Southern Malawi: profile of patients across two years.. Int J Audiol, 60 (10), pp.789-796. <https://doi.org/10.1080/14992027.2020.1864486>.

Persistent Link:

<http://hdl.handle.net/11343/290232>