The nitrogen-potassium intersection: Membranes, metabolism, and mechanism

Running title: The N-K intersection

Devrim Coskun, Dev T. Britto, and Herbert J. Kronzucker*

Department of Biological Sciences and the Canadian Centre for World Hunger Research (CCWHR), University of Toronto, 1265 Military Trail, Toronto, Ontario, Canada, M1C 1A4

*Corresponding author: herbert.kronzucker@utoronto.ca

Keywords: ammonium, nitrate, influx, efflux, translocation, assimilation, plant productivity

Abstract

Nitrogen (N) and potassium (K) are the two most abundantly acquired mineral elements by plants and their acquisition pathways interact in complex ways. Here, we review pivotal interactions with respect to root acquisition, storage, translocation, and metabolism, between the K^+ ion and the two major N sources, ammonium (NH_4^+) and nitrate (NO_3^-). The intersections between N and K physiology are explored at a number of organizational levels, from molecular-genetic processes, to compartmentation, to whole plant physiology, and discussed in the context of both N-K cooperation and antagonism. Nutritional regulation and optimization of plant growth, yield, metabolism, and water-use efficiency are also discussed.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1111/pce.12671](https://doi.org/10.1111/pce.12671)

1. Introduction

Nitrogen (N) and potassium (K) are required for plants to complete their life cycles, and are the two nutrients acquired in greatest quantities by roots (Oosterhuis *et al.* 2014). The pathways by which these elements are transported and utilized by plants intersect in significant ways, including the influences exerted by potassium ions on nitrogen nutrition and physiology, and vice versa, in terms of primary uptake of N and K at the root plasma membrane, their transport, accumulation, and assimilation within the plant, and the regulation of these metabolic and transport pathways. In this paper we shall review what is known about these interactions, with special emphasis on mechanistic processes and plant productivity.

A few fundamental distinctions between these essential elements should be made at the outset. Firstly, while potassium is generally available to plants only as a simple monoatomic, monovalent cation, K^+ , nitrogen is available in the form of diverse compounds, *e.g.* as cationic ammonium (NH_4^+), anionic nitrate (NO_3^-), or as amino acids, which may be cationic, anionic, or zwitterionic, depending on the chemical species and soil pH. In addition, uncharged ammonia, NH_3 , is likely to rapidly enter plant root cells from the soil, particularly under conditions of high N supply and/or high pH (Coskun *et al.* 2013b). Because the dominant forms of N available to plants in most soils are the inorganic ions NH_4^+ and NO_3^- , however, they, with the K^+ ion, will make up the central focus of our review.

Another distinction is that K^+ ions, once taken up by plants, remain in this simple ionic state, while N-containing compounds undergo numerous chemical reactions and become covalently bonded within organic molecules throughout the plant. This fundamental difference is reflected in the major physiological roles the two elements play in plants. Nitrogen is an essential constituent of a vast array of metabolites and structural compounds, including proteins, nucleic acids, chlorophyll, co-enzymes, phytohormones, and secondary metabolites, while the main functions of K^+ are as a major osmolyte and source of positive charge for electrical homeostasis and enzyme activation (Evans & Wildes 1971; Leigh & Wyn Jones 1984; Britto & Kronzucker

2008; Marschner 2011). Thus, our discussion of the metabolic processes at the intersection of N and K physiology will focus on biochemical pathways involving the transformation of nitrogen.

Thirdly, it is interesting that N enters the terrestrial biosphere chiefly from the atmosphere as a result of enzymatic processes in N₂-fixing soil bacteria (and today, via the industrial Haber-Bosch process), although weathering of bedrock N (itself primarily atmospherically derived), which ties up about 20% of the global N pool, can sometimes produce ecologically significant rates of nitrogen release (Holloway & Dahlgren 2002; Xu *et al.* 2012). By contrast, K⁺ must be replenished by weathering of parent rock and release from exchangeable and non-exchangeable sources, if not supplied as fertilizer (Zhang *et al.* 2010).

Soil NH₄⁺ concentrations tend to range between 0.1 and 1 mM, while those of NO₃⁻ tend to be higher, often exceeding 1 mM and reaching 10 mM or even higher following fertilization or a burst of nitrification (Wolt 1994; Crawford & Glass 1998; Owen & Jones 2001; Miller *et al.* 2007). Thus, the molar ratios of soil NO₃⁻ to NH₄⁺ typically range between 10 and 100 (Wolt 1994; Miller *et al.* 2007). In some soils, however, such as rice paddies, bog lands, and boreal and montane forests, anaerobic, acidic, and reductive conditions reverse this scenario, with NH₄⁺ concentrations exceeding those of NO₃⁻ (Gillman & Bell 1978; Kronzucker *et al.* 1997, 2000). In the case of K⁺, soil concentrations tend to be similar to those of NH₄⁺, or about 0.1 - 1 mM (Wolt 1994; White *et al.* 2013). In some agricultural areas of the world, including China, India, and the Philippines, the relatively small K⁺ pools are replenished at rates far lower than those of N replenishment, leading to nutritional imbalances, reduced productivity, and eutrophication by N runoff (Dobermann *et al.* 1998; Hoa *et al.* 2006; Andrist-Rangel *et al.* 2007; Zhang *et al.* 2010). The concentrations of NO₃⁻, NH₄⁺, and K⁺ in soils can vary greatly, not only over regional scales, but even within relatively small patches (Wolt 1994; Miller *et al.* 2007; Cramer *et al.* 2009). Nitrate levels also tend to show greater heterogeneity in soils, and can range over two orders of magnitude across relatively short distances (*e.g.* 4 m or less; Lark *et al.* 2004), whereas NH₄⁺ and K⁺ levels tend to vary more narrowly, about one order of magnitude over a similar scale. This is

partially due to the negative charge on NO_3^- , which results in a high degree of mobility within soils (Miller & Cramer 2004). Plant roots themselves directly increase soil heterogeneity within the rhizosphere via N- and K-uptake processes, producing zones of depletion (Scherer & Ahrens 1996; Hinsinger *et al.* 2005; Kayser & Isselstein 2005; Moody & Bell 2006; Andrist-Rangel *et al.* 2007).

Soil concentrations of N and K are important not only because plant productivity can be limited by their scarcity, but also because their excess can bring about toxicity and repress growth. Thus, optimum growth and yield curves, which can vary greatly with plant species and environmental factors, are typically seen with respect to these nutrients (Asher & Ozanne 1967; MacLeod 1969; Loué 1980; Britto & Kronzucker 2013). The shapes of such curves can be greatly influenced by the presence of other potentially limiting nutrients. In Fig. 1, this type of influence is shown for K supply upon N optima (1A), and vice-versa (1B). It is notable that the lowest provision of K (1A) or N (1B) results in a relatively moderate N or K concentration (respectively) required to reach maximum yield (*i.e.* about 6 mM N, or 70 kg ha⁻¹ K), but this yield is still much lower than can be reached with higher provision of the companion nutrient.

2. Transport

It is rather remarkable that the transport functions involved in the uptake of NO_3^- , NH_4^+ , and K^+ from soil solution, moving these ions across the plasma membrane and into the cytosol of the root cell, share a common feature: all have been characterized using two-mechanism models, which consist of saturable, high-affinity and linear, low-affinity transport systems (“HATS” and “LATS”) generally operating at low and high external substrate concentrations, respectively (Miller *et al.* 2007; Szczerba *et al.* 2009; Zhang *et al.* 2010). For the cations K^+ and NH_4^+ , these two types of transport systems mechanistically contend with the two main thermodynamic scenarios encountered by roots engaged in cation uptake: those that require active uptake (HATS conditions), and those that allow passive uptake (LATS conditions). In general, the active uptake

of a cationic substrate is driven by an energetic coupling between the passive flow of H^+ down its transmembrane electrochemical potential gradient, and the substrate's thermodynamically "uphill" influx into the cell, while passive uptake proceeds "downhill" via ion channels (Hedrich & Schroeder 1989; Szczerba *et al.* 2009; Coskun *et al.* 2013a). It is worth noting that the distinction between HATS and LATS transporters is somewhat blurred, and ion channels can at times function at low substrate concentrations, while active-transporting carriers can operate at high concentrations (see below). In the case of NO_3^- , it is likely that, under most conditions, uptake is thermodynamically active, due to the inside-negative electrical potential difference across the plasma membrane (Wang *et al.* 2012). The plethora of transport proteins catalyzing the fluxes of NO_3^- , NH_4^+ , and K^+ under both HATS and LATS scenarios, many of which have been identified at the genetic level, is consistent with the spatially and temporally heterogeneous distributions of these substrates in the soil (see Introduction). However, it has been pointed out that the transport of NO_3^- under many field conditions might be catalyzed primarily by low-affinity transporters, given the relatively high NO_3^- concentrations in soils, and because the expression and activities of high-affinity NO_3^- transporters become downregulated as soil $[NO_3^-]$ rises (Miller *et al.* 2007).

One of the best known interactions between N and K physiology in plants is the marked inhibitory effect that NH_4^+ exerts on the high-affinity K^+ uptake system (Smith & Epstein 1964; Deane-Drummond & Glass 1983; Pettersson *et al.* 1984; Scherer *et al.* 1984; Vale *et al.* 1987, 1988; Wang *et al.* 1996; Spalding *et al.* 1999; Nielsen and Schjoerring 1998; Santa-Maria *et al.* 2000; Szczerba *et al.* 2008a; ten Hoopen *et al.* 2010; Coskun *et al.* 2010, 2013a; Zhang *et al.* 2010). In barley seedlings, the inhibition of K^+ translocation to the shoot by NH_4^+ appears to be even more pronounced than the inhibition of primary K^+ uptake at the plasma membrane, with suppressions as high as 90% (Kronzucker *et al.* 2003). This inhibition appears to be at least partially reciprocal, resulting in an "antagonism" between the two ions, with both competitive and non-competitive components. The competitive effects likely arise from the similarities between the two ions, in terms of their hydrated diameters, charge, and influence on membrane

potentials (Wang *et al.* 1996; ten Hoopen *et al.* 2010). Non-competitive effects include a suppression of NH_4^+ uptake in tobacco seedlings by K^+ that persists even after K^+ removal (Scherer *et al.* 1984), the lack of suppression of NH_4^+ uptake by a wide range of external $[\text{K}^+]$ in sweet pepper (Xu *et al.* 2002), a stimulation of NH_4^+ transport by K^+ in *Arabidopsis* (ten Hoopen *et al.* 2010; also see Szczerba *et al.* 2008b, for a LATS-range stimulation), and the inhibition of K^+ uptake by the acidification of the rhizosphere that is associated with NH_4^+ uptake in many species (Findenegg 1987). It has also been shown that the induction, by K^+ starvation, of *AtHAK5* (the gene encoding the dominant component of high-affinity K^+ transport in *Arabidopsis* roots; Gierth & Mäser 2007; Rubio *et al.* 2008; see Fig. 2) expression was almost completely suppressed by the presence of only 100 μM NH_4^+ (Qi *et al.* 2008; *cf.* Rubio *et al.* 2008; see Section 3). In another study, it was shown that the removal of NH_4^+ from solutions bathing the roots of barley and *Arabidopsis* plants resulted in dramatic increases (4.5- and 6-fold in the two species, respectively) in K^+ uptake (Coskun *et al.* 2013a). Interestingly, this effect occurred under both low- and high- K^+ conditions, and was partially attributed to the hyperpolarization of the plasma-membrane electrical potential that was observed upon NH_4^+ withdrawal.

A lower accumulation of K^+ in NH_4^+ -grown (relative to NO_3^- -grown) plants is thus partially attributable to the suppression, by NH_4^+ , of K^+ influx in the HATS range. However, it may also be due to an enhancement of K^+ efflux by NH_4^+ , which has been demonstrated in several studies (Munn & Jackson 1978; Rufty *et al.* 1982; Topa & Jackson 1988; Coskun *et al.* 2010; *cf.* Deane-Drummond & Glass 1983). In the study by Coskun *et al.* (2010), it was shown that NH_4^+ -stimulated K^+ efflux in roots of barley could be completely counteracted by the channel-blocking agents tetraethylammonium (TEA^+) and cesium (Cs^+), suggesting that it is catalyzed by an as yet unidentified ion channel, possibly a member of the Shaker family of K^+ -specific channels. This study also showed that K^+ efflux could be stimulated by the co-presence of NH_4^+ and NO_3^- (as NH_4NO_3), but NO_3^- as the sole N source reduced K^+ efflux, an effect also observed by Pettersson (1984).

A few studies, however, have provided evidence that NH_4^+ can actually stimulate high-affinity K^+ transport under some conditions (Nieves-Cordones *et al.* 2008; Alvarez-Pizarro *et al.* 2011). These studies have suggested that the stimulation was due to a more electrically hyperpolarized plasma membrane in the presence of NH_4^+ , a condition, however, that is rarely seen in other studies (*e.g.* Coskun *et al.* 2013a; see Section 3). Ammonium-induced hyperpolarization was also inferred in sorghum roots based on an increase in expression of the plasma-membrane H^+ -ATPase (Alvarez-Pizarro *et al.* 2011), and such a hyperpolarization was measured in roots of tomato (Nieves-Cordones *et al.* 2008). The latter study suggested that membrane hyperpolarization is a regulatory component of HAK5 expression in tomato (see Section 3).

By contrast, few effects of NH_4^+ on K^+ transport in the low-affinity range have been reported (*cf.* Coskun *et al.* 2013a). However, a study using T-DNA insertion lines showed that transport of K^+ not only via HAK5 but also via AKT1, the dominant low-affinity K^+ -specific influx channel in plant roots (Ivashikina *et al.* 2001; Gierth & Mäser 2007; Fig. 2), is sensitive to NH_4^+ (Rubio *et al.* 2008). Nevertheless, the powerful suppression of HAK-mediated K^+ transport by NH_4^+ , coupled to the lack of such an effect on AKT1, has been used as a parsing tool to distinguish between the two systems. By this means, it was shown that AKT1, normally associated with LATS, is capable of transporting K^+ at external concentrations as low as $10\ \mu\text{M}$ in *Arabidopsis* (Hirsch *et al.* 1998; Spalding *et al.* 1999), and $100\ \mu\text{M}$ in barley (Coskun *et al.* 2013a), given a sufficiently negative membrane potential. The “dual-affinity” nature of AKT1 is not unique; it can also be seen in the *Arabidopsis* high-affinity K^+ transporter AtKUP1 (Fu & Luan 1998) and the NO_3^- transporter AtNRT1.1, in which a molecular switch between high- and low-affinity functions via changes in phosphorylation and conformational states has been demonstrated (Ho *et al.*, 2009; Sun *et al.* 2014; Parker & Newstead 2014; see also Section 3).

Nevertheless, some studies have shown effects of NH_4^+ supply on low-affinity K^+ transport. Vale *et al.* (1987) found that both a saturating and a linear phase of $[\text{K}^+]$ -dependent K^+ transport in corn were suppressed by NH_4^+ , but the latter phase, while kinetically resembling

“classic” low-affinity transport due to its linearity (Szczerba *et al.* 2009), was only investigated between 50 and 200 μM K^+ , at which the flux may have still been dominated by HAK systems. An interesting aspect of this study was that the maximal suppression of K^+ influx was seen at 100 μM NH_4^+ , with no further suppression seen at 200–500 μM . At higher K^+ and NH_4^+ concentrations (5 and 10 mM, respectively), however, it was clearly demonstrated in a study on barley roots that steady-state NH_4^+ supply can indeed inhibit LATS-range K^+ transport (Coskun *et al.* 2013a). In this study, removal of NH_4^+ during measurement of $^{42}\text{K}^+$ uptake resulted in a flux more than twice as high as in control plants (*i.e.* those in the presence of 10 mM NH_4^+). Moreover, when NH_4^+ was removed and replaced by NO_3^- (also at 10 mM), K^+ influx was more than three times that of controls. Interestingly, the NH_4^+ -withdrawal effect was generally short-term, and essentially disappeared over a 10-hr period, while similar experiments at low K^+ (20 and 100 μM) showed a longer-lasting effect of NH_4^+ withdrawal, resulting in fluxes 4–5 times higher than control, as well as substantially higher K^+ accumulation, even after 24 hr. This suggests that under some conditions, such as that of low K^+ supply, the influence of other ions on K^+ transport may override regulation by internal K^+ status or growth demands. Under other conditions, the reverse may be true, and K^+ uptake and content may be relatively unaffected by the supply of other nutrients (including N), as has been frequently observed (Pitman 1972; Pettersson *et al.* 1984; Vale *et al.* 1987; Zhang *et al.* 2010).

At least one study has reported the stimulation of low-affinity K^+ uptake under steady-state provision of high (10 mM) NH_4^+ , relative to equimolar NO_3^- (Szczerba *et al.* 2008b). At or below 100 μM K^+ , the characteristic suppression of K^+ influx by NH_4^+ was observed in this study, but, at 1.5 mM K^+ and above, growth on NH_4^+ pronouncedly stimulated both K^+ influx and accumulation in the shoots of rice plants. Because of the upregulation of NH_4^+ transport known to occur under high NH_4^+ conditions (*e.g.* Wang *et al.* 1993; Rawat *et al.* 1999; Cerezo *et al.* 2001), it was suggested that the stimulation of K^+ transport under these conditions was mediated by the enhanced activity of a low-affinity NH_4^+ transporter. However, it should be noted that this effect was found in tropical lowland rice, considered to be an “ammonium

specialist”, and might not be common among species sensitive to NH_4^+ toxicity. Moreover, the increased K^+ accumulation found under NH_4^+ nutrition in rice did not always translate into increased growth; maximal growth was found at 1.5 mM K^+ , but growth declined by 40% when K^+ was raised to 40 mM (the condition at which the largest stimulation of NH_4^+ transport and accumulation was observed). Under NO_3^- nutrition, the optimal growth concentration was even lower, with a fresh-weight decline of about 20% observed at 1.5 mM K^+ (relative to 100 μM) despite the higher shoot K^+ content in these plants.

It is well known that NH_4^+ nutrition is often associated with substantial declines in tissue levels of essential cations including K^+ as well as Ca^{2+} and Mg^{2+} (Kirkby & Knight 1977; Kurvits & Kirkby 1980; van Beusichem *et al.* 1988); this is a central aspect of the NH_4^+ toxicity syndrome, and is largely attributable to the inhibition of fluxes of other cations by NH_4^+ (Britto & Kronzucker 2002). The converse of this is that NH_4^+ toxicity is frequently found to be relieved by an increase in K^+ provision (Lips *et al.* 1990; Cao *et al.* 1993; Britto & Kronzucker 2002; ten Hoopen *et al.* 2010). In addition, elevated K^+ provision can reduce the inhibition of NO_3^- transport by NH_4^+ , when N is provided as NH_4NO_3 , and improve growth on this mixed-N source (Rufty *et al.* 1982; Hagin *et al.* 1990). The alleviation of NH_4^+ toxicity by K^+ is in part a result of the upregulation, by increased K^+ , of enzymes involved in NH_4^+ assimilation (see Section 4), but also because of the pronounced, dose-dependent reductions in low-affinity NH_4^+ transport and accumulation brought about by elevated K^+ (Szczerba *et al.* 2008a, b; Balkos *et al.* 2010). This contrasts with high-affinity NH_4^+ uptake via the AMT1 transporter (possibly involving a H^+ - NH_4^+ symport mechanism; Ortiz-Ramirez *et al.* 2011; Xu *et al.* 2012), which appears to be quite selective for NH_4^+ and against K^+ (Ninnemann *et al.* 1994). At high (10 mM) NH_4^+ , however, a switch in external $[\text{K}^+]$ ($[\text{K}^+]_{\text{ext}}$) from 0.1 to 1.5 mM reduced NH_4^+ influx into barley roots by nearly 60% (Szczerba *et al.* 2008a), with similar results seen in rice (Szczerba *et al.* 2008b; Balkos *et al.* 2010). In addition, elevated $[\text{K}^+]_{\text{ext}}$ reduced the efflux of NH_4^+ from roots of barley and rice, to an even greater extent than influx, resulting in a lower extent of futile N cycling across the plasma membrane, which has been linked to NH_4^+ toxicity (Britto *et al.* 2001; Chen *et*

al. 2013). Interestingly, these effects were seen both instantaneously and over the long term, and, although K^+ elevation did not generally reduce the net flux of NH_4^+ into the plant (indeed, it was substantially increased in rice), the increased metabolism of NH_4^+ resulted in greatly reduced accumulation of this toxic compound (of up to 75-80%) in both barley and rice (Szczerba *et al.* 2008a; Balkos *et al.* 2010). Subsequent work has shown that, under these low-affinity NH_4^+ -transport conditions, the major transported species may in fact be the deprotonated, uncharged ammonia molecule (NH_3), which crosses the plasma membrane via aquaporins (Jahn *et al.* 2004; Coskun *et al.* 2013b). The ability of elevated $[K^+]_{ext}$ to greatly diminish transport via aquaporins may reflect the significance of this ion in the area of plant water relations (Quintero *et al.* 2007; Britto *et al.* 2014; see below).

In contrast to the antagonistic relationship between K^+ and NH_4^+ nutrition, the acquisition rates of K^+ and NO_3^- are often found to be positively correlated, and enhance one another, likely because of improved charge balance (Minotti *et al.* 1968; Blevins *et al.* 1974; Kirkby & Knight 1977; Kurvits & Kirkby 1980; Pettersson *et al.* 1984; Kochian *et al.* 1985; Le Bot & Kirkby 1991; Macduff *et al.* 1997; Marschner *et al.* 1996; Siebrecht & Tischner 1999; Coskun *et al.* 2013a; Delaire *et al.* 2014) and the activation, by K^+ , of enzymes involved in NO_3^- assimilation (Hagin *et al.* 1990; Zhang *et al.* 2010; Roosta & Schjoerring 2009; Balkos *et al.* 2010; see Section 4). Thus, it is commonly observed that plants take up and accumulate substantially more K^+ under NO_3^- nutrition than with NH_4^+ , or in the presence of other cations such as Mg^{2+} , Na^+ , and Ca^{2+} (Ivashikina & Feyziev 1998; Coskun *et al.* 2013a). However, the direct influences exerted by one ion on the transport of the other are poorly understood. Nitrate and potassium can both be stored in the vacuole at high concentrations (Martinoia *et al.* 1981; 2007; Walker *et al.* 1996; Oosterhuis *et al.* 2014), where they may electrically balance one another, and can be dynamically retrieved according to plant demand (Blumwald & Poole 1985; Walker *et al.* 1996). The situation for NH_4^+ in the vacuole is less clear. It is not often accumulated in high amounts in plant tissues, except under toxic conditions, under which it may enter the vacuole as NH_3 , via aquaporins (Jahn *et al.* 2004; Loqué *et al.* 2005; Martinoia *et al.* 2007; Coskun *et al.* 2013).

An intriguing example of the co-operative use of K^+ and NO_3^- in plants is what is referred to as the “Dijkshoorn-Ben Zioni model” of internal ion cycling via the vasculature, in which NO_3^- is transported from root to shoot in the xylem, using K^+ as a counterion (Dijkshoorn *et al.* 1968; Ben Zioni *et al.* 1971; Kirkby & Knight 1977; Pettersson 1984; Touraine *et al.* 1988; Fig. 3A). NO_3^- is reduced and assimilated into amino acids in the shoot, with the concomitant transfer of negative charge to organic (carboxylic) acids. These acids, particularly in the form of malate, are then transported to the roots via the phloem, accompanied by K^+ as counterion, where they are decarboxylated and release HCO_3^- to the external medium. Indeed, it has been estimated that as much as half the K^+ found in the xylem stream is not directly introduced via uptake processes in the root, but comes from recirculated K^+ in the phloem (Engels & Kirkby 2001; Chérel *et al.* 2014).

The Dijkshoorn-Ben Zioni model has proven useful in a large number of studies, and provides one of the few mechanisms known to link shoot demand for nutrients with their uptake by the roots (Casadesús *et al.* 1995; Marschner *et al.* 1996). However, its applicability is limited by several conditions, and has been contradicted in some studies. First, the model pertains to plants that reduce NO_3^- mainly in the shoot, which is a common but by no means universal occurrence (Pate, 1973; Andrews, 1986; Touraine *et al.* 1990). Interestingly, a low supply of K^+ can increase the degree of NO_3^- reduction in roots relative to shoots, possibly due to the lack of a xylem-mobile cation to accompany NO_3^- moving to the shoot (Rufty *et al.* 1981; Forster & Jeschke 1993; see Section 4). Second, even when shoot reduction is pronounced, and the upward movement of K^+ as a counterion with NO_3^- is observed, a significant downward recirculation of K-malate in the phloem is not always seen (Kirkby and Knight 1977). This appears to be related to conditions where there is a balanced uptake of cations and anions by roots, as in many herbaceous dicotyledons growing under nutrient-replete conditions (van Egmond 1978; *cf.* Touraine *et al.* 1990). Third, the regulation of diurnal variations in NO_3^- uptake may depend less on the synthesis, transport, or addition of malate than on light-dark transitions affecting the shoot (Delhon *et al.* 1996; also see Casadesús *et al.* 1995; Fig. 3B).

Nevertheless, in many cases, the model appears to be fairly robust in terms of the co-ordinated translocation of the two ions from root to shoot. Significant progress in this area includes demonstrations of K^+ and NO_3^- interdependence at the molecular level, in terms of alterations of transporters that load NO_3^- into the xylem, and resulting effects on K^+ translocation (Lin *et al.* 2008; Xia *et al.* 2015; see Section 3). However, it may be possible for other cations, in particular Mg^{2+} , to at least partially substitute for K^+ in the charge balancing of NO_3^- in the xylem (Forster & Jeschke 1993). In addition, there appears to be a regulatory decoupling between the uptake of K^+ and NO_3^- across the plasma membrane, on the one hand, and their translocation to the shoot (Kirkby & Armstrong 1980; Touraine & Grignon 1982; Casadesús *et al.* 1995; Ivashikina & Feyziev 1998).

3. Sensing, Signaling, and Co-regulation

Plants have evolved complex sensing, signaling, and regulatory mechanisms to manage the acquisition of ions in fluctuating environments, and key similarities can be found in mechanisms involving K^+ and N transport and assimilation. These similarities include the co-regulation at the transcriptional level of K^+ on N transporters and vice versa, the post-translational modifications of K^+ and N transporters by the same regulatory proteins, as well as more systemic changes, such as in membrane potential, the accumulation of reactive oxygen species (ROS), and phytohormones.

One of the clearest examples of K-N interaction at the transport level, *i.e.* the inhibition of high-affinity K^+ transport via HAK transporters by NH_4^+ (see Section 2; Fig. 2), manifests itself at the transcriptional level as well. NH_4^+ has been shown to downregulate *AtHAK5* and *CaHAK1* transcription in K^+ -deprived *Arabidopsis* and pepper plants, respectively (Martínez-Cordero *et al.* 2005; Qi *et al.* 2008). It is well documented that K^+ deprivation leads to hyperpolarization of the root plasma membrane (Etherton & Higinbotham 1960; Maathuis & Sanders 1993; Amtmann *et al.* 2006; Schachtman & Shin 2007; Britto & Kronzucker 2008). It

has recently been suggested, based on a correlation analysis, that membrane hyperpolarization could trigger an as yet unknown signaling cascade inducing the expression of high-affinity HAK transporters (*LeHAK5* and *AtHAK5* in tomato and *Arabidopsis*, respectively; Nieves-Cordones *et al.* 2008; Rubio *et al.* 2014). Interestingly, this induction appears to occur even under K^+ -replete conditions, contrary to reports indicating that HAK induction is dependent on K^+ starvation (Armengaud *et al.* 2004; Gierth & Mäser 2007; Qi *et al.* 2008). One such condition that has been tested is that of NO_3^- deprivation, in which membrane hyperpolarization was seen to coincide with an increased expression of *AtHAK5* and *LeHAK5* (Rubio *et al.* 2014; see also Wang *et al.* 2001; Shin *et al.* 2005). Perhaps critically, however, it does not coincide with increased root (net) K^+ uptake, in contrast to K^+ -deprivation conditions, suggesting a specific regulatory role of low K^+ in HAK5 activity (Rubio *et al.* 2014). Also intriguingly, Nieves-Cordones *et al.* (2008) found that tomato plants grown in the presence of NH_4^+ displayed hyperpolarized membrane potentials relative to NH_4^+ -free conditions and increased *LeHAK5* expression. This is in apparent contradiction to many reports of NH_4^+ -induced membrane depolarization, at least in the short term (Ullrich *et al.* 1984; Ayling 1993; Wang *et al.* 1994). In tomato, transcript levels of *LeHAK5* were also found to quickly (2–6 h) decline with membrane-depolarization events (*e.g.* K^+ re-supply, or initial exposure to NH_4^+ , Rb^+ , vanadate, or CCCP; Nieves-Cordones *et al.* 2008). It would be interesting to see whether other means of short-term membrane hyperpolarization (*e.g.* with bicarbonate (Poole 1969), fusaric acid (Ullrich & Novacky 1990), or NH_4^+ withdrawal from NH_4^+ -grown plants (Coskun *et al.* 2013a; see above)) would also result in increased HAK5 expression. Overall, this is an interesting, but as yet speculative, regulatory mechanism that requires further investigation.

Another important link between K^+ and N nutrition is the involvement of calcineurin B-like (CBL)-interacting protein kinase 23 (CIPK23) in high-affinity K^+ and NO_3^- uptake (Xu *et al.* 2006; Ho *et al.* 2009; Castaings *et al.* 2011; Tsay *et al.* 2011; see also Fig. 2). CIPK23 is involved in activating high-affinity uptake of both K^+ and NO_3^- via AKT1 and NPF6.3 (NRT1.1), respectively (Xu *et al.* 2006; Cheong *et al.* 2007; Lee *et al.* 2007; Li *et al.* 2006; Ho *et*

al. 2009). Under K^+ deprivation, a Ca^{2+} signaling cascade (triggered perhaps by ethylene and ROS; see below) results in its binding to CBL1 and CBL9, which in turn activate CIPK23, which phosphorylates and thus activates AtAKT1 (Xu *et al.* 2006; Cheong *et al.* 2007; Li *et al.* 2006; Lee *et al.* 2007; Luan *et al.* 2009; Wang and Wu 2013). Like AtAKT1, AtNPF6.3 is a dual-affinity transporter, conducting both high- and low-affinity NO_3^- uptake (Wang *et al.* 1998; Liu *et al.* 1998; Liu & Tsay 2003; Ho *et al.* 2009; Tsay *et al.* 2011; Lran *et al.* 2014). At low external nitrate concentrations (*i.e.* HATS conditions), the same signaling cascade triggers AtCIPK23 to phosphorylate AtNPF6.3 (at T101), and thus switches the transporter into ‘high-affinity mode’ (Ho *et al.* 2009; Sun *et al.* 2014; Parker & Newstead 2014). By contrast, unphosphorylated AtNPF6.3 functions as a low-affinity transporter (Hu *et al.* 2009). The crystal structure of AtNPF6.3 (AtNRT1.1) and the allosteric effects of phosphorylation have been determined (Sun *et al.* 2014; Parker & Newstead 2014); to our knowledge, such a study has yet to be conducted for AKT1. Recently, a developmental role of the CBL1-CBL9-CIPK23-AKT1/NRT1.1 signaling module was proposed, identifying specific parameters related to root system architecture as targets (Kellermeier *et al.* 2014). For example, it was found that under K^+ and NO_3^- deficiency, phosphorylated AKT1 and NRT1.1 stimulate second-order lateral root emergence. This is an important new development since conditions arise that exclude AKT1 and NRT1.1 in high-affinity K^+ and NO_3^- uptake, respectively (*e.g.* under conditions where high-affinity HAK transporters are functional (Rubio *et al.* 2008; Coskun *et al.* 2013a, 2014) or where high-affinity NRT2 transporters predominate (Okamoto *et al.* 2003; Wang *et al.* 2012)).

The apparent coupling of K^+ and NO_3^- transport in the xylem (Section 2) appears to play out at the molecular level as well. In *nrt1.5* knock-out mutants for a transporter responsible for xylem NO_3^- loading in *Arabidopsis*, K^+ translocation was reduced along with that of NO_3^- (Lin *et al.* 2008). However, a K^+ -coupled mechanism for this transporter was ruled out in a heterologous expression system in favour of one of H^+ coupling. Furthermore, K^+ deprivation was shown to downregulate the expression of *AtNRT1.5* (*AtNPF7.3*), suggesting NO_3^- translocation was controlled by plant K^+ levels (Lin *et al.* 2008). Similar results were found in rice with respect to

the nitrate transporter OsNPF2.4 (Xia *et al.* 2015), and other nitrate transporters have been implicated in NO₃⁻ xylem loading (e.g. AtNPF6.3 (AtNRT1.1), AtNPF2.3; OsNPF2.2, and LeNRT2.3; Lran *et al.* 2013; Taochy *et al.* 2015; Li *et al.* 2015; Fu *et al.* 2015), but their relationships to K⁺ nutrition have yet to be elucidated. The expression of the gene for the K⁺ xylem-loading transporter SKOR was shown to be upregulated by nitrate supply (Wang *et al.* 2004), demonstrating a similar response as *AtNRT1.5* and *OsNPF2.4* to K⁺ supply. These data strongly support a co-regulation at the level of xylem transport that maintains the balance between NO₃⁻ and K⁺ (Wang *et al.* 2012). It will be interesting to see if such co-regulation at the molecular level exists at the level of primary uptake, where, for example, the stimulation of low-affinity K⁺ uptake in the presence of NO₃⁻ is observed (Kochian *et al.* 1985; Coskun *et al.* 2013a; see Section 2; Fig. 2).

In addition to nutritional regulation, K⁺ and NO₃⁻ xylem-loading transporters and root-to-shoot translocation also show diurnal regulation (Fig. 3B). *AtNPF7.3* (*AtNRT1.5*) expression peaked at the night-to-day transition and fell to a minimum at the day-to-night transition (Lin *et al.* 2008). In the legume *Samanea saman*, *SPORK1* (a *SKOR* homolog) expression showed similar diel patterns (Moshelion *et al.* 2002). Such patterns are consistent with xylem loading of NO₃⁻ and K⁺ being highest during the day and lowest during the night (Mattson *et al.* 1988; Schurr & Schulze 1995; Macduff & Bakken 2003; Siebrecht *et al.* 2003), as they are closely tied to transpiration rates (Le Bot & Kirkby 1992; Siebrecht *et al.* 2003; Fig. 3C). Moreover, leaf nitrate reductase (NR) activity (the first enzyme involved in nitrate reduction; Campbell 1999) also rises to a maximum during the first half of the light period (Scheible *et al.* 1997; Lillo *et al.* 2001). This highly suggests a coordinated and concerted mechanism at play related to NO₃⁻ and K⁺ translocation. Insight into the underlying signaling and regulatory network is eagerly anticipated.

K⁺ deprivation has been demonstrated to result in strong upregulation of the nitrate transporters *LeNRT1.2* and *LeNRT2.1* in tomato roots (Wang *et al.* 2001). This parallels the well-

documented effects of the induction of high-affinity HAK transporters in response to K^+ deprivation (Martínez-Cordero *et al.* 2005; Gierth & Mäser, 2007; Qi *et al.* 2008). By contrast, K^+ deprivation resulted in the *downregulation* of *AtNRT2.1*, as well as *AtNRT2.3*, and *AtNRT2.6*, in *Arabidopsis* (Armengaud *et al.* 2004). Expression of *AtNRT1.1*, however, was upregulated with K^+ deprivation in this study. To our knowledge, these discrepancies have yet to be thoroughly investigated or resolved. However, the differences may be the result of varying timeframes of K^+ deficiency in each study, as the responses observed for tomato were made over a relatively short time (within 1–3 h), whereas the responses observed for *Arabidopsis* took place after 2 weeks of K^+ starvation (although re-supply of K^+ resulted in *AtNRT2* genes being upregulated within 6 h; Armengaud *et al.* 2004). Nevertheless, these reports clearly indicate a “crosstalk” between the nutritional status of one nutrient and the expression of membrane transporters for the other.

Another common observance to K^+ and NO_3^- deficiency is the accumulation of ROS as a signaling component (Shin & Schachtman 2004; Shin *et al.* 2005; Schachtman & Shin 2007). While the consequences of ROS accumulation are not well understood (Schachtman & Shin 2007), they include the upregulation of *AtHAK5* and *AtNRT2.1* expression under K^+ and NO_3^- deprivation, respectively (Shin *et al.* 2005; Kim *et al.* 2010, 2012). In contrast to the study by Armengaud *et al.* (2004), *AtNRT2.1* expression was not affected by K^+ deprivation in the study by Shin *et al.* (2005). This may also be due to timing, as K^+ deprivation was maintained for up to 30 h in the latter study, as opposed to 2 weeks in the former. A more comprehensive study on the role of ROS in this crosstalk would be most interesting. Ca^{2+} may be part of this signaling cascade both upstream (Torres & Dangl 2005; Li *et al.* 2006; Lebaudy *et al.* 2007) and downstream (Mori & Schroeder 2004) of ROS production. At least in the case of K^+ deprivation, ethylene acts upstream of ROS (Shin & Schachtman 2004; Jung *et al.* 2009). Mutants defective in ethylene or ROS production, such as ethylene insensitive2-1 (*ein2-1*), an NADPH oxidase (*rhd2*), or a type III peroxidase (*rci3*), all showed reduced *AtHAK5* expression (Shin &

Schachtman 2004; Jung *et al.* 2009; Kim *et al.* 2010; Fig. 2). The expression and function of nitrate transporters in these mutants, however, remain unknown.

4. Metabolism

Although K^+ itself is not metabolized, it plays a vital role in many aspects of plant metabolism. K^+ is involved in the activity of some 46 enzymes (Evans & Sorger 1966; Leigh & Wyn Jones 1984; Hagin *et al.* 1990; Britto & Kronzucker 2008; Armengaud *et al.* 2009), including as cofactor to such critical enzymes as pyruvate kinase, starch synthase, Rubisco, and nitrate reductase (NR; Beevers & Hageman 1969; Sorger *et al.* 1965; Evans & Sorger 1966; Nitsos & Evans 1966; 1969; Peoples & Koch 1979; Wyn Jones & Pollard 1983). Moreover, K^+ is crucial to protein synthesis, not only with respect to enzyme activation, but also to ribosome synthesis and mRNA turnover (Blevins 1985; Evans & Wildes 1971; Pettigrew 2008). Its importance in cellular metabolism is reflected in its high, homeostatically set concentration in the cytosol of ~100 mM (Leigh & Wyn Jones 1984; Walker *et al.* 1996; Kronzucker *et al.* 2003).

The close relationships between K^+ supply and N metabolism is evident from many studies. One such connection involves the partitioning of NR activity between the root and shoot (Blevins *et al.* 1978; Barneix & Breteler 1985; Förster & Jeschke 1993; Casadesús *et al.* 1995; Armengaud *et al.* 2009), which also depends on plant species, external nitrate supply, temperature, and light intensity (Pate 1973; Smirnoff & Stewart 1985; Andrews 1986). Generally, with high external K^+ supply, the co-translocation of K^+ and NO_3^- to the shoot increases (Ben Zioni *et al.* 1971; Blevins *et al.* 1978; see Section 2; Fig. 3), and both storage of NO_3^- and NR activity increase in leaves, while less N assimilation is found in roots (Blevins *et al.* 1978; Rufty *et al.* 1981). In contrast, with K^+ deprivation, less translocation of NO_3^- is observed and hence higher N assimilation occurs in roots (Förster and Jeschke 1993; Rufty *et al.* 1981; Wang *et al.* 2003). Interestingly, this has not been borne out in the model species *Arabidopsis thaliana*; Armengaud *et al.* (2009) observed significant decreases in root NR

activity, relative to K^+ sufficiency, with prolonged (2-week) K^+ deprivation. By contrast, the authors observed significant upregulation in the activity of key enzymes involved in ammonium assimilation, *i.e.* glutamine synthetase (GS), ferredoxin-glutamine-2-oxoglutarate aminotransferase (Fd-GOGAT), and glutamate dehydrogenase (GDH). It was concluded that the downregulation of NR (as well as nitrate transporters; Armengaud *et al.* 2004), could be as a result of decreased carbon-skeleton production via the TCA cycle (decreased glycolysis and the direct inhibition of pyruvate kinase due to low cytoplasmic K^+ , was observed). On the other hand, the upregulation of GS, GOGAT, and GDH was interpreted as a compensatory response to maintain C flux through the TCA cycle and into amino acids and proteins (increased NADP-malic enzyme activity due to low cytoplasmic pH was also observed).

In rice, root GS activity was found to increase with increasing $[K^+]_{ext}$ (again, in apparent contradiction to the *Arabidopsis* model; see above), reaching its maximum at 5 mM in an NH_4^+ (10 mM) background (Balkos *et al.* 2010). The same pattern was also observed for phosphoenolpyruvate carboxylase (PEPC) activity, the key anapleurotic C-fixing enzyme, as well as with total protein content (Fig. 4). This corresponded with tremendous gains in plant biomass, with growth 160% higher than NO_3^- -grown plants and 220% higher than plants grown at 0.1 mM K^+ (Balkos *et al.* 2010). Similar effects were observed in cucumber, although they were not as dramatic, which may reflect the lower tolerance to NH_4^+ in this species, relative to rice (Roosta & Schjoerring 2008). Other reports of the stimulation of expression and activity of N-assimilatory enzymes by K^+ can be found in Ali *et al.* (1991) and Mohammad & Naseem (2006). Taken together, these findings, including the controversial findings with *Arabidopsis* (above), point to a distinct reprogramming, by K^+ , of C and N metabolism (Armengaud *et al.* 2009; Balkos *et al.* 2010; Pettigrew 2008). A similar reconfiguration of primary (and secondary) metabolic machinery was observed by transcriptomic analysis in *Arabidopsis*, in response to N. After two days of N deprivation, the majority of genes associated with photosynthesis, chlorophyll synthesis, and plastid protein synthesis were repressed, and those associated with secondary metabolism were induced (Scheible *et al.* 2004).

Lastly, phytohormone responses involving K⁺ and N nutrition include auxin and cytokinin (CK) biosynthesis and signaling. Cao *et al.* (1993) showed that growth suppression of *Arabidopsis* seedling roots in the presence of 6 mM NH₄⁺ and 20 μM K⁺ could be relieved by elevation of K⁺ to 200 μM, but not in the presence of supplementary auxin (indole acetic acid) or cytokinin (6-benzylaminopurine), or a combination of the two. Moreover, growth suppression by NH₄⁺ was also much less pronounced in auxin-resistant mutants. Cytokinins have been implicated in K⁺ and N signaling and the regulation of genes encoding K⁺ and N transporters (Brenner *et al.* 2005; Kiba *et al.* 2005; Schachtman & Shin, 2007; Shin 2011; Nam *et al.* 2012). Under low-K⁺ stress, root CK levels are reduced, which result in ROS accumulation, root hair growth, and *AtHAK5* expression (Nam *et al.* 2012). Similarly, low N results in lowered CK levels (Takei *et al.* 2002, 2004; Scheible *et al.* 2004; Miyawaki *et al.* 2006; Nam *et al.* 2012) which have been linked to increased expression of genes encoding nitrate and ammonium transporters (*NRT2* and *AMT1*, respectively; Brenner *et al.* 2005; Kiba *et al.* 2005).

5. Conclusion

On a planet where the human population continues to greatly expand (forecasts estimates growth of up to 12.3 billion by 2100; Gerland *et al.* 2014), ever-increasing demands on the world's agricultural systems are expected to produce immense strains on land, water, and nutrient resources in the coming decades. Rapidly changing climates and environmental degradation in many parts of the world will put additional stresses on crop production, while a dangerous feed-forward cycle may play itself out, given that food systems are themselves responsible for 19-29% of anthropogenic greenhouse gas emissions globally (Vermeulen *et al.* 2012). Plant physiologists have applied their knowledge and skills to produce remarkable increases in crop yields over the past century, preventing the starvation of millions, and must be called upon once again to address new challenges.

Clearly, to reduce the amount of new land required to meet growing demands, and thereby reduce strains upon “marginal” or ecologically fragile environments, the efficiency of crop production will need to be increased. One approach to this would be to precisely determine species- and site-specific growth optima with respect to nutrients such as NO_3^- , NH_4^+ , and K^+ , not only singly, but in relation to one another. In this review, we have discussed some of the key physiological issues pertaining to these relationships, from the often observed antagonism between NH_4^+ and K^+ , to the alleviation of NH_4^+ toxicity by K^+ , to the co-operative uptake, internal transport, and utilization of NO_3^- and K^+ . Other interactions, such as those involving synergies and antagonisms between NO_3^- and NH_4^+ , and their relationships to K^+ supply, as well as interactions involving other essential macro- and micronutrients in optimizing growth and yield, will greatly improve the science of plant nutrition. In addition, the examination of nutrient interactions with water use in plants (and its efficiency), such as the influences of NO_3^- and K^+ on aquaporin and stomatal function (Guo *et al.* 2003; Cramer *et al.* 2009; Fig. 2), and as major osmotica, will become increasingly important as water scarcity and soil salinity become more widespread. Moreover, the world’s most important crop species, rice, is also the world’s most water consumptive, further highlighting the necessity to understand and optimize plant water use (Hoekstra & Chapagain 2007; Chapagain & Hoekstra 2011; Britto *et al.* 2014).

While the complexities of nutrient sensing, signaling mechanisms, and transport regulation cannot be fully unraveled without the benefits of molecular biology, there is still much room for physiological methods in the pursuit of more efficient and productive crop systems (Pettigrew 2008; Cramer *et al.* 2009; Britto *et al.* 2014). This is important in part because of the current public reaction against recombinant DNA technology, particularly in Europe. However, even while “genetically modified” crop plants are out of favour, the vast amount of data provided by modern molecular biology (*e.g.* Armengaud *et al.* 2009) has the potential to inform and direct physiologists and agronomists seeking to improve crop production by nutritional means. These approaches, combined with gradual improvements in remote and local sensing of soil fertility

and crop mineral content, and in spatially and temporally precise application of water and nutrients, can help bring about yield increases even in GMO-free zones.

Acknowledgements

Funding was provided by the Natural Sciences and Engineering Research Council of Canada (NSERC).

Author Manuscript

References

- Ali A.A., Ikeda M., Yamada Y. (1991) Effects of the supply of K, Ca, and Mg on the absorption and assimilation of ammonium-nitrogen and nitrate-nitrogen in tomato plants. *Soil Science & Plant Nutrition* **37**, 283-289.
- Alvarez-Pizarro J.C., Gomes-Filho E., Prisco J.T., Grossi-de-Sá M.F., de Oliveira-Neto O.B. (2011) NH_4^+ -stimulated low- K^+ uptake is associated with the induction of H^+ extrusion by the plasma membrane H^+ -ATPase in sorghum roots under K^+ deficiency. *Journal of Plant Physiology* **168**, 1617-1626.
- Amtmann A., Hammond J.P., Armengaud P., White P.J. (2006) Nutrient sensing and signalling in plants: potassium and phosphorus. *Advances in Botanical Research* **43**, 209-257.
- Andrews M. (1986) The partitioning of nitrate assimilation between root and shoot of higher plants. *Plant, Cell & Environment* **9**, 511-519.
- Andrist-Rangel Y., Edwards A.C., Hillier S., Öborn I. (2007) Long-term K dynamics in organic and conventional mixed cropping systems as related to management and soil properties. *Agriculture, Ecosystems & Environment* **122**, 413-426.
- Armengaud P., Breitling R., Amtmann A. (2004) The potassium-dependent transcriptome of *Arabidopsis* reveals a prominent role of jasmonic acid in nutrient signaling. *Plant Physiology* **136**, 2556-2576.
- Armengaud P., Sulpice R., Miller A.J., Stitt M., Amtmann A., Gibon Y. (2009) Multilevel analysis of primary metabolism provides new insights into the role of potassium nutrition for glycolysis and nitrogen assimilation in *Arabidopsis* roots. *Plant Physiology* **150**, 772-785.
- Asher C.J., Ozanne P.G. (1967) Growth and potassium content of plants in solution cultures maintained at constant potassium concentrations. *Soil Science* **103**, 155-161.

- Ayling S.M. (1993). The effect of ammonium ions on membrane potential and anion flux in roots of barley and tomato. *Plant, Cell & Environment* **16**, 297-303.
- Balkos K.D., Britto D.T., Kronzucker H.J. (2010) Optimization of ammonium acquisition and metabolism by potassium in rice (*Oryza sativa* L. cv. IR-72). *Plant, Cell & Environment* **33**, 23-34.
- Barneix A.J., Breteler H. (1985) Effect of cations on uptake, translocation and reduction of nitrate in wheat seedlings. *New Phytologist* **99**, 367-379.
- Beevers L., Hageman R.H. (1969) Nitrate reduction in higher plants. *Annual Review of Plant Physiology* **20**, 495-522.
- Ben Zioni A., Vaadia Y., Herman Lips S. (1971) Nitrate uptake by roots as regulated by nitrate reduction products of the shoot. *Physiologia Plantarum* **24**, 288-290.
- Blevins D.G. (1985) Role of potassium in protein metabolism in plants. *Potassium in Agriculture* 413-424.
- Blevins D.G., Barnett N.M., Frost W.B. (1978) Role of potassium and malate in nitrate uptake and translocation by wheat seedlings. *Plant Physiology* **62**, 784-788.
- Blevins D.G., Hiatt A.J., Lowe R.H. (1974) The influence of nitrate and chloride uptake on expressed sap pH, organic acid synthesis, and potassium accumulation in higher plants. *Plant Physiology* **54**, 82-87.
- Blumwald E., Poole R.J. (1985) Nitrate storage and retrieval in *Beta vulgaris*: Effects of nitrate and chloride on proton gradients in tonoplast vesicles. *Proceedings of the National Academy of Sciences, USA* **82**, 3683-3687.
- Brenner W.G., Romanov G.A., Köllmer I., Bürkle L., Schmölling T. (2005) Immediate-early and delayed cytokinin response genes of *Arabidopsis thaliana* identified by genome-wide

- expression profiling reveal novel cytokinin-sensitive processes and suggest cytokinin action through transcriptional cascades. *Plant Journal* **44**, 314-333.
- Britto D.T., Balkos K.D., Becker A., Coskun D., Huynh W.Q., Kronzucker H.J. (2014) Potassium and nitrogen poisoning: Physiological changes and biomass gains in rice and barley. *Canadian Journal of Plant Science* **94**, 1085-1089.
- Britto D.T., Kronzucker H.J. (2002) NH_4^+ toxicity in higher plants: a critical review. *Journal of Plant Physiology* **159**, 567-584.
- Britto D.T., Kronzucker H.J. (2008) Cellular mechanisms of potassium transport in plants. *Physiologia Plantarum* **133**, 637-650.
- Britto D.T., Kronzucker H.J. (2013) Ecological significance and complexity of N-source preference in plants. *Annals of Botany* **112**, 957-963.
- Britto D.T., Siddiqi M.Y., Glass A.D.M., Kronzucker H.J. (2001) Futile transmembrane NH_4^+ cycling: a cellular hypothesis to explain ammonium toxicity in plants. *Proceedings of the National Academy of Sciences, USA* **98**, 4255-4258.
- Campbell W.H. (1999) Nitrate reductase structure, function and regulation: bridging the gap between biochemistry and physiology. *Annual Review of Plant Biology* **50**, 277-303.
- Cao Y., Glass A.D., Crawford N.M. (1993) Ammonium inhibition of *Arabidopsis* root growth can be reversed by potassium and by auxin resistance mutations *aux1*, *axr1*, and *axr2*. *Plant Physiology* **102**, 983-989.
- Casadesús J., Tapia L., Lambers H. (1995) Regulation of K^+ and NO_3^- fluxes in roots of sunflower (*Helianthus annuus*) after changes in light intensity. *Physiologia Plantarum* **93**, 279-285.

- Castaings L., Marchive C., Meyer C., Krapp A. (2011) Nitrogen signalling in *Arabidopsis*: how to obtain insights into a complex signalling network. *Journal of Experimental Botany* **62**, 1391-1397.
- Cerezo M., Tillard P., Gojon A., Primo-Millo E., Garcia-Agustin P. (2001) Characterization and regulation of ammonium transport systems in *Citrus* plants. *Planta* **214**, 97-105.
- Chapagain A.K., Hoekstra A.Y. (2011) The blue, green and grey water footprint of rice from production and consumption perspectives. *Ecological Economics* **70**, 749-758.
- Chen G., Guo S., Kronzucker H.J., Shi W.M. (2013) Nitrogen use efficiency (NUE) in rice links to NH_4^+ toxicity and futile NH_4^+ cycling in roots. *Plant and Soil* **369**, 351-363.
- Chérel I., Lefoulon C., Boeglin M., Sentenac H. (2014) Molecular mechanisms involved in plant adaptation to low K^+ availability. *Journal of Experimental Botany* **65**, 833-848.
- Cheong Y. H., Pandey G.K., Grant J.J., Batistic O., Li L., Kim B.G., ..., Luan S. (2007) Two calcineurin B-like calcium sensors, interacting with protein kinase CIPK23, regulate leaf transpiration and root potassium uptake in *Arabidopsis*. *Plant Journal* **52**, 223-239.
- Coskun D., Britto D.T., Li M., Oh S., Kronzucker H.J. (2013a) Capacity and plasticity of potassium channels and high-affinity transporters in roots of barley and *Arabidopsis*. *Plant Physiology* **162**, 496-511.
- Coskun D., Britto D.T., Li M., Becker A., Kronzucker H.J. (2013b) Rapid ammonia gas transport accounts for futile transmembrane cycling under $\text{NH}_3/\text{NH}_4^+$ toxicity in plant roots. *Plant Physiology* **163**, 1859-1867.
- Coskun D., Britto D.T., Kronzucker H.J. (2010) Regulation and mechanism of potassium release from barley roots: an in planta $^{42}\text{K}^+$ analysis. *New Phytologist* **188**, 1028-1038.
- Cramer M.D., Hawkins H.-J., Verboom G.A. (2009) The importance of nutritional regulation of plant water flux. *Oecologia* **161**, 15-24.

- Crawford N.M., Glass A.D. (1998) Molecular and physiological aspects of nitrate uptake in plants. *Trends in Plant Science* **3**, 389-395.
- Deane-Drummond C.E., Glass A.D. (1983) Short term studies of nitrate uptake into barley plants using ion-specific electrodes and $^{36}\text{ClO}_3^-$ II. Regulation of NO_3^- efflux by NH_4^+ . *Plant Physiology* **73**, 105-110.
- Delaire M., Maugé J-C., Beaujard F. (2014) Evidence for a strong correlation between season-dependent nitrate and potassium uptake in two deciduous trees. *Trees* **28**, 769-776.
- Delhon P., Gojon A., Tillard P., Passama L. (1996) Diurnal regulation of NO_3^- uptake in soybean plants III. Implication of the Dijkshoorn-Ben Zioni model in relation with the diurnal changes in NO_3^- assimilation. *Journal of Experimental Botany* **47**, 885-892.
- Dijkshoorn W., Lathwell D.J., De Wit C.T. (1968) Temporal changes in carboxylate content of ryegrass with stepwise change in nutrition. *Plant and Soil* **29**, 369-390.
- Dobermann A., Cassman K.G., Mamaril C.P., Sheehy J.E. (1998) Management of phosphorus, potassium, and sulfur in intensive, irrigated lowland rice. *Field Crops Research* **56**, 113-138.
- Engels C., Kirkby E.A. (2001) Cycling of nitrogen and potassium between shoot and roots in maize as affected by shoot and root growth. *Journal of Plant Nutrition and Soil Science* **164**, 183-191.
- Etherton B., Higinbotham N. (1960) Transmembrane potential measurements of cells of higher plants as related to salt uptake. *Science* **131**, 409-410.
- Evans H.J., Sorger G.J. (1966) Role of mineral elements with emphasis on the univalent cations. *Annual Review of Plant Physiology* **17**, 47-76.
- Evans H.J., Wildes R.A. (1971) Potassium and its role in enzyme activation. In *Proc. 8th Colloq. Int. Potash Inst. Bern* (pp. 13-39).

- Findenegg G.R. (1987) A comparative study of ammonium toxicity at different constant pH of the nutrient solution. *Plant and Soil* **103**, 239-243.
- Förster J.C., Jeschke W. D. (1993) Effects of potassium withdrawal on nitrate transport and on the contribution of the root to nitrate reduction in the whole plant. *Journal of Plant Physiology* **141**, 322-328.
- Fu H.H., Luan S. (1998) AtKUP1: a dual-affinity K⁺ transporter from *Arabidopsis*. *Plant Cell* **10**, 63-73.
- Fu Y., Yi H., Bao J., Gong J. (2015) LeNRT2.3 functions in nitrate acquisition and long-distance transport in tomato. *FEBS Letters* **589**, 1072-1079.
- Gerland P., Raftery A.E., Ševčíková H., Li N., Gu D., Spoorenberg T., ..., Wilmoth J. (2014) World population stabilization unlikely this century. *Science* **346**, 234-237.
- Gierth M., Mäser P. (2007) Potassium transporters in plants—involvement in K⁺ acquisition, redistribution and homeostasis. *FEBS Letters* **581**, 2348-2356.
- Gillman G.P., Bell L.C. (1978) Soil solution studies on weathered soils from tropical North Queensland. *Australian Journal of Soil Research* **16**, 67-77.
- Guo F.Q., Young J., Crawford N.M. (2003) The nitrate transporter AtNRT1.1 (CHL1) functions in stomatal opening and contributes to drought susceptibility in *Arabidopsis*. *Plant Cell* **15**, 107-117.
- Hagin J., Olsen S.R., Shaviv A. (1990) Review of interaction of ammonium-nitrate and potassium nutrition of crops. *Journal of Plant Nutrition* **13**, 1211-1226.
- Hedrich R., Schroeder J.I. (1989) The physiology of ion channels and electrogenic pumps in higher plants. *Annual Review of Plant Physiology* **40**, 539-569.

- Hinsinger P., Gobran G.R., Gregory P.J., Wenzel W.W. (2005) Rhizosphere geometry and heterogeneity arising from root-mediated physical and chemical processes. *New Phytologist* **168**, 293-303.
- Hirsch R.E., Lewis B.D., Spalding E.P., Sussman M.R. (1998) A role for the AKT1 potassium channel in plant nutrition. *Science* **280**, 918-921.
- Ho C.H., Lin S.H., Hu H.C., Tsay Y.F. (2009) CHL1 functions as a nitrate sensor in plants. *Cell* **138**, 1184-1194.
- Ho C.H., Tsay Y.F. (2010) Nitrate, ammonium, and potassium sensing and signaling. *Current Opinion in Plant Biology* **13**, 604-610.
- Hoa N.M., Janssen B.H., Oenema O., Dobermann A. (2006) Comparison of partial and complete soil K budgets under intensive rice cropping in the Mekong Delta, Vietnam. *Agriculture, Ecosystems & Environment* **116**, 121-131.
- Hoekstra A.Y., Chapagain A.K. (2007) Water footprints of nations: water use by people as a function of their consumption pattern. *Water Resources Management* **21**, 35-48.
- Holloway J.M., Dahlgren R.A. (2002) Nitrogen in rock: occurrences and biogeochemical implications. *Global Biogeochemical Cycles* **16**, 65-1.
- Ivashikina N.V., Feyziev Y.M. (1998) Regulation of nitrate uptake in maize seedlings by accompanying cations. *Plant Science* **131**, 25-34.
- Ivashikina N., Becker D., Ache P., Meyerhoff O., Felle H.H., Hedrich R. (2001) K⁺ channel profile and electrical properties of *Arabidopsis* root hairs. *FEBS Letters* **508**, 463-469.
- Jahn T.P., Møller A.L., Zeuthen T., Holm L.M., Klærke D.A., Mohsin B., ..., Schjoerring J.K. (2004) Aquaporin homologues in plants and mammals transport ammonia. *FEBS Letters* **574**, 31-36.

- Jung J.Y., Shin R., Schachtman D.P. (2009) Ethylene mediates response and tolerance to potassium deprivation in Arabidopsis. *Plant Cell* **21**, 607-621.
- Jungk A., Claassen N. (1986) Availability of phosphate and potassium as the result of interactions between root and soil in the rhizosphere. *Zeitschrift für Pflanzenernährung und Bodenkunde* **149**, 411-427.
- Kayser M., Isselstein J. (2005) Potassium cycling and losses in grassland systems: a review. *Grass and Forage Science* **60**, 213-224.
- Kellermeier F., Armengaud P., Seditas T.J., Danku J., Salt D.E., Amtmann A. (2014) Analysis of the root system architecture of Arabidopsis provides a quantitative readout of crosstalk between nutritional signals. *Plant Cell* **26**, 1480-1496.
- Kiba T., Naitou T., Koizumi N., Yamashino T., Sakakibara H., Mizuno T. (2005) Combinatorial microarray analysis revealing Arabidopsis genes implicated in cytokinin responses through the His→Asp phosphorelay circuitry. *Plant and Cell Physiology* **46**, 339-355.
- Kim M.J., Ciani S., Schachtman D.P. (2010) A peroxidase contributes to ROS production during Arabidopsis root response to potassium deficiency. *Molecular Plant* **3**, 420-427.
- Kirkby E.A., Armstrong M.J. (1980) Nitrate uptake by roots as regulated by nitrate assimilation in the shoot of castor oil plants. *Plant Physiology* **65**, 286-290.
- Kirkby E.A., Knight A.H. (1977) Influence of the level of nitrate nutrition on ion uptake and assimilation, organic acid accumulation and cation balance in whole tomato plants. *Plant Physiology* **60**, 249-253.
- Kirkby E.A., Mengel K. (1967) Ionic balance in different tissues of the tomato plant in relation to nitrate, urea, or ammonium nutrition. *Plant Physiology* **42**, 6-14.
- Kochian L.V., Xin-Zhi J., Lucas W.J. (1985) Potassium transport in corn roots IV. Characterization of the linear component. *Plant Physiology* **79**, 771-776.

- Kronzucker H.J., Britto D.T. (2011) Sodium transport in plants: a critical review. *New Phytologist* **189**, 54-81.
- Kronzucker H.J., Glass A.D.M., Siddiqi M.Y., Kirk G.J.D. (2000) Comparative kinetic analysis of ammonium and nitrate acquisition by tropical lowland rice: implications for rice cultivation and yield potential. *New Phytologist* **145**, 471-476.
- Kronzucker H.J., Schjoerring J.K., Erner Y., Kirk G.J., Siddiqi M.Y., Glass A.D.M. (1998) Dynamic interactions between root NH_4^+ influx and long-distance N translocation in rice: insights into feedback processes. *Plant and Cell Physiology* **39**, 1287-1293.
- Kronzucker H.J., Siddiqi M.Y., Glass A.D. (1997) Conifer root discrimination against soil nitrate and the ecology of forest succession. *Nature* **385**, 59-61.
- Kronzucker H.J., Szczerba M.W., Britto D.T. (2003) Cytosolic potassium homeostasis revisited: ^{42}K -tracer analysis in *Hordeum vulgare* L. reveals set-point variations in $[\text{K}^+]$. *Planta* **217**, 540-546.
- Kurvits A., Kirkby E.A. (1980) The uptake of nutrients by sunflower plants (*Helianthus annuus*) growing in a continuous flowing culture system, supplied with nitrate or ammonium as nitrogen source. *Zeitschrift für Pflanzenernährung und Bodenkunde* **143**, 140-149.
- Lark R.M., Milne A.E., Addiscott T.M., Goulding K.W.T., Webster C.P., O'Flaherty S. (2004) Scale-and location-dependent correlation of nitrous oxide emissions with soil properties: an analysis using wavelets. *European Journal of Soil Science* **55**, 611-627.
- Lebaudy A., Véry A.A., Sentenac H. (2007) K^+ channel activity in plants: genes, regulations and functions. *FEBS Letters* **581**, 2357-2366.
- Le Bot J., Kirkby E.A. (1992) Diurnal uptake of nitrate and potassium during the vegetative growth of tomato plants. *Journal of Plant Nutrition* **15**, 247-264.

- Lee S.C., Lan W.Z., Kim B.G., Li L., Cheong Y.H., Pandey G.K., ..., Luan S. (2007) A protein phosphorylation/dephosphorylation network regulates a plant potassium channel. *Proceedings of the National Academy of Sciences, USA* **104**, 15959-15964.
- Leigh R.A., Wyn Jones R.G. (1984) A hypothesis relating critical potassium concentrations for growth to the distribution and functions of this ion in the plant cell. *New Phytologist* **97**, 1-13.
- Léran S., Edel K.H., Pervent M., Hashimoto K., Corratgé-Faillie C., Offenborn J.N., ..., Lacombe B. (2015) Nitrate sensing and uptake in *Arabidopsis* are enhanced by ABI2, a phosphatase inactivated by the stress hormone abscisic acid. *Science Signaling* **8**, ra43.
- Léran S., Muños S., Brachet C., Tillard P., Gojon A., Lacombe B. (2013) *Arabidopsis* NRT1.1 is a bidirectional transporter involved in root-to-shoot nitrate translocation. *Molecular Plant* **6**, 1984-1987.
- Léran S., Varala K., Boyer J.C., Chiurazzi M., Crawford N., Daniel-Vedele F., ..., Lacombe B. (2014) A unified nomenclature of NITRATE TRANSPORTER 1/PEPTIDE TRANSPORTER family members in plants. *Trends in Plant Science* **19**, 5-9.
- Li L., Kim B.G., Cheong Y.H., Pandey G.K., Luan S. (2006) A Ca^{2+} signaling pathway regulates a K^{+} channel for low-K response in *Arabidopsis*. *Proceedings of the National Academy of Sciences, USA* **103**, 12625-12630.
- Li Y., Ouyang J., Wang Y.Y., Hu R., Xia K., Duan J., ..., Zhang M. (2015) Disruption of the rice nitrate transporter OsNPF2.2 hinders root-to-shoot nitrate transport and vascular development. *Scientific Reports* **5**, 9635
- Lillo C., Meyer C., Ruoff P. (2001) The nitrate reductase circadian system. The central clock dogma contra multiple oscillatory feedback loops. *Plant Physiology* **125**, 1554-1557.

- Lin S.H., Kuo H.F., Canivenc G., Lin C.S., Lepetit M., Hsu P.K., ..., Tsay Y.F. (2008) Mutation of the Arabidopsis NRT1. 5 nitrate transporter causes defective root-to-shoot nitrate transport. *Plant Cell* **20**, 2514-2528.
- Lips S.H., Leidi E.O., Silberbush M., Soares M.I.M., Lewis O.E.M. (1990) Physiological aspects of ammonium and nitrate fertilization. *Journal of Plant Nutrition* **13**, 1271-1289.
- Liu K.H., Huang C.Y., Tsay Y.F. (1999) CHL1 is a dual-affinity nitrate transporter of *Arabidopsis* involved in multiple phases of nitrate uptake. *Plant Cell* **11**, 865-874.
- Liu K.H., Tsay Y.F. (2003) Switching between the two action modes of the dual-affinity nitrate transporter CHL1 by phosphorylation. *EMBO Journal* **22**, 1005-1013.
- Loqué D., Ludewig U., Yuan L., von Wirén N. (2005) Tonoplast Intrinsic Proteins AtTIP2;1 and AtTIP2;3 facilitate NH_3 transport into the vacuole. *Plant Physiology* **137**, 671-680.
- Loué A. (1980) The interaction of potassium with other growth factors, particularly with other nutrients. *Potassium Fertilization in Agricultural Practice. IPI Research Topics* **8**, 67-93.
- Luan S. (2009) The CBL–CIPK network in plant calcium signaling. *Trends in Plant Science* **14**, 37-42.
- Maathuis F.J., Sanders D. (1993) Energization of potassium uptake in *Arabidopsis thaliana*. *Planta* **191**, 302-307.
- Macduff J.H., Bakken A.K., Dhanoa M.S. (1997) An analysis of the physiological basis of commonality between diurnal patterns of NH_4^+ , NO_3^- and K^+ uptake by *Phleum pratense* and *Festuca pratensis*. *Journal of Experimental Botany* **48**, 1691-1701.
- MacLeod L.B. (1969) Effects of N, P, and K and their interactions on the yield and kernel weight of barley in hydroponic culture. *Agronomy Journal* **61**, 26-29.

- Marschner H., Kirkby E.A., Cakmak I. (1996) Effect of mineral nutritional status on shoot-root partitioning of photoassimilates and cycling of mineral nutrients. *Journal of Experimental Botany* **47**, 1255-1263.
- Marschner H. (2011) Marschner's mineral nutrition of higher plants. Academic Press.
- Martínez-Cordero M.A., Martínez V., Rubio F. (2005) High-affinity K⁺ uptake in pepper plants. *Journal of Experimental Botany* **56**, 1553-1562.
- Martinoia E., Heck U., Wiemken A. (1981) Vacuoles as storage compartments for nitrate in barley leaves. *Nature* **289**, 292-294.
- Martinoia E., Maeshima M., Ekkehard H.E. (2007) Vacuolar transporters and their essential role in plant metabolism. *Journal of Experimental Botany* **58**, 83-102.
- Mattson W.J., Palmer S.R. (1988). Changes in levels of foliar minerals and phenolics in trembling aspen, *Populus tremuloides*, in response to artificial defoliation. In *Mechanisms of woody plant defenses against insects* (pp. 157-169). Springer New York.
- Mengel K., Viro M., Hehl G. (1976) Effect of potassium on uptake and incorporation of ammonium-nitrogen of rice plants. *Plant and Soil* **44**, 547-558.
- Miller A.J., Cramer M.D. (2004) Root nitrogen acquisition and assimilation. *Plant and Soil* **274**, 1-36.
- Miller A.J., Fan X., Orsel M., Smith S.J., Wells D.M. (2007) Nitrate transport and signalling. *Journal of Experimental Botany* **58**, 2297-2306.
- Minotti P.L., Williams D.C., Jackson W.A. (1968) Nitrate uptake and reduction as affected by calcium and potassium. *Soil Science Society of America Journal* **32**, 692-698.

- Miyawaki K., Tarkowski P., Matsumoto-Kitano M., Kato T., Sato S., Tarkowska D., ..., Kakimoto T. (2006) Roles of Arabidopsis ATP/ADP isopentenyltransferases and tRNA isopentenyltransferases in cytokinin biosynthesis. *Proceedings of the National Academy of Sciences, USA* **103**, 16598-16603.
- Mohammad F., Naseem U. (2006) Effect of K application on leaf carbonic anhydrase and nitrate reductase activities, photosynthetic characteristics, NPK and NO_3^- contents, growth and yield of mustard. *Photosynthetica* **44**, 471-473.
- Moody P.W., Bell M.J. (2006) Availability of soil potassium and diagnostic soil tests. *Soil Research* **44**, 265-275.
- Mori I.C., Schroeder J.I. (2004) Reactive oxygen species activation of plant Ca^{2+} channels. A signaling mechanism in polar growth, hormone transduction, stress signaling, and hypothetically mechanotransduction. *Plant Physiology* **135**, 702-708.
- Moshelion M., Becker D., Czempinski K., Mueller-Roeber B., Attali B., Hedrich R., Moran N. (2002) Diurnal and circadian regulation of putative potassium channels in a leaf moving organ. *Plant Physiology* **128**, 634-642.
- Munn D.A., Jackson W.A. (1978) Nitrate and ammonium uptake by rooted cuttings of sweet potato. *Agronomy Journal* **70**, 312-316.
- Nam Y.J., Tran L.S.P., Kojima M., Sakakibara H., Nishiyama R., Shin R. (2012) Regulatory roles of cytokinins and cytokinin signaling in response to potassium deficiency in *Arabidopsis*. *PLoS ONE* **7**, e47797
- Nielsen K.H., Schjoerring J.K. (1998) Regulation of apoplastic NH_4^+ concentration in leaves of oilseed rape. *Plant Physiology* **118**, 1361-1368.

- Nieves-Cordones M., Miller A.J., Alemán F., Martínez V., Rubio F. (2008) A putative role for the plasma membrane potential in the control of the expression of the gene encoding the tomato high-affinity potassium transporter HAK5. *Plant Molecular Biology* **68**, 521-532.
- Ninnemann O., Jauniaux J.C., Frommer W.B. (1994) Identification of a high affinity NH_4^+ transporter from plants. *EMBO Journal* **13**, 3464-3471.
- Nitsos R.E., Evans H.J. (1966) Effects of univalent cations on the inductive formation of nitrate reductase. *Plant Physiology* **41**, 1499-1504.
- Okamoto M., Vidmar J.J., Glass A.D.M. (2003) Regulation of *NRT1* and *NRT2* gene families of *Arabidopsis thaliana*: responses to nitrate provision. *Plant and Cell Physiology* **44**, 304-317.
- Oosterhuis D.M., Loka D.A., Kawakami E.M., Pettigrew W.T. (2014) The physiology of potassium in crop production. *Advances in Agronomy* **126**, 203-233.
- Ortiz-Ramirez C., Mora S.I., Trejo J., Pantoja O. (2011) *PvAMT1;1*, a highly selective ammonium transporter that functions as H^+/NH_4^+ symporter. *The Journal of Biological Chemistry* **286**, 31113-31122.
- Ourry A., Macduff J.H., Prudhomme M.P., Boucaud J. (1996) Diurnal variation in the simultaneous uptake and 'sink' allocation of NH_4^+ and NO_3^- by *Lolium perenne* in flowing solution culture. *Journal of Experimental Botany* **47**, 1853-1863.
- Owen A.G., Jones D.L. (2001) Competition for amino acids between wheat roots and rhizosphere microorganisms and the role of amino acids in plant N acquisition. *Soil Biology and Biochemistry* **33**, 651-657.
- Parker J.L., Newstead S. (2014) Molecular basis of nitrate uptake by the plant nitrate transporter NRT1. 1. *Nature* **507**, 68-72.

- Pate J.S. (1973) Uptake, assimilation and transport of nitrogen compounds by plants. *Soil Biology and Biochemistry* **5**, 109-119.
- Peoples T.R., Koch D.W. (1979) Role of potassium in carbon dioxide assimilation in *Medicago sativa* L. *Plant Physiology* **63**, 878-881.
- Pettersson S. (1984) Effects of nitrate on influx, efflux and translocation of potassium in young sunflower plants. *Physiologia Plantarum* **61**, 663-669.
- Pettigrew W.T. (2008) Potassium influences on yield and quality production for maize, wheat, soybean and cotton. *Physiologia Plantarum* **133**, 670-681.
- Pitman M.G. (1972) Uptake and transport of ions in barley seedlings III. Correlation between transport to the shoot and relative growth rate. *Australian Journal of Biological Sciences* **25**, 905-920.
- Poole R.J. (1969). Carrier-mediated potassium efflux across the cell membrane of red beet. *Plant Physiology* **44**, 485-490.
- Qi Z., Hampton C.R., Shin R., Barkla B.J., White P.J., Schachtman D.P. (2008) The high affinity K^+ transporter AtHAK5 plays a physiological role *in planta* at very low K^+ concentrations and provides a caesium uptake pathway in *Arabidopsis*. *Journal of Experimental Botany* **59**, 595-607.
- Quintero J.M., Fournier J.M., Benlloch M. (2007) Na^+ accumulation in shoot is related to water transport in K^+ -starved sunflower plants but not in plants with a normal K^+ status. *Journal of Plant Physiology* **164**, 60-67.
- Rawat S.R., Silim S.N., Kronzucker H.J., Siddiqi M.Y., Glass A.D.M. (1999) *AtAMT1* gene expression and NH_4^+ uptake in roots of *Arabidopsis thaliana*: Evidence for regulation by root glutamine levels. *Plant Journal* **19**, 143-152.

- Roosta H.R., Schjoerring J.K. (2008) Effects of nitrate and potassium on ammonium toxicity in cucumber plants. *Journal of Plant Nutrition* **31**, 1270-1283.
- Rubio F., Fon M., Ródenas R., Nieves-Cordones M., Alemán F., Rivero R.M., Martínez V. (2014) A low K^+ signal is required for functional high-affinity K^+ uptake through HAK5 transporters. *Physiologia Plantarum* **152**, 558-570.
- Rubio F., Nieves-Cordones M., Alemán F., Martínez V. (2008) Relative contribution of AtHAK5 and AtAKT1 to K^+ uptake in the high-affinity range of concentrations. *Physiologia Plantarum* **134**, 598-608.
- Rufty T.W., Jackson W.A., Raper C.D. (1981) Nitrate reduction in roots as affected by the presence of potassium and by flux of nitrate through the roots. *Plant Physiology* **68**, 605-609.
- Rufty T.W., Jackson W.A., Raper C.D. (1982) Inhibition of nitrate assimilation in roots in the presence of ammonium: the moderating influence of potassium. *Journal of Experimental Botany* **33**, 1122-1137.
- Santa-María G.E., Danna C.H., Czibener C. (2000) High-affinity potassium transport in barley roots. Ammonium-sensitive and -insensitive pathways. *Plant Physiology* **123**, 297-306.
- Schachtman D.P., Shin R. (2007) Nutrient sensing and signaling: NPKS. *Annual Review of Plant Biology* **58**, 47-69.
- Scherer H.W., Ahrens G. (1996) Depletion of non-exchangeable NH_4^+ -N in the soil—root interface in relation to clay mineral composition and plant species. *European Journal of Agronomy* **5**, 1-7.
- Scherer H.W., Mackown C.T., Leggett J.E. (1984) Potassium-ammonium uptake interactions in tobacco seedlings. *Journal of Experimental Botany* **35**, 1060-1070.

- Scheible W.R., Gonzalez-Fontes A., Lauerer M., Muller-Rober B., Caboche M., Stitt M. (1997) Nitrate acts as a signal to induce organic acid metabolism and repress starch metabolism in tobacco. *Plant Cell* **9**, 783-798.
- Scheible W.R., Morcuende R., Czechowski T., Fritz C., Osuna D., Palacios-Rojas N., ..., Stitt M. (2004) Genome-wide reprogramming of primary and secondary metabolism, protein synthesis, cellular growth processes, and the regulatory infrastructure of *Arabidopsis* in response to nitrogen. *Plant Physiology* **136**, 2483-2499.
- Schurr U., Schulze E.D. (1995) The concentration of xylem sap constituents in root exudate, and in sap from intact, transpiring castor bean plants (*Ricinus communis* L.). *Plant, Cell & Environment* **18**, 409-420.
- Shin R. (2011) Transcriptional regulatory components responding to macronutrient limitation. *Journal of Plant Biology* **54**, 286-293.
- Shin R., Berg R.H., Schachtman D.P. (2005) Reactive oxygen species and root hairs in *Arabidopsis* root response to nitrogen, phosphorus and potassium deficiency. *Plant and Cell Physiology* **46**, 1350-1357.
- Shin R., Schachtman D.P. (2004) Hydrogen peroxide mediates plant root cell response to nutrient deprivation. *Proceedings of the National Academy of Sciences, USA* **101**, 8827-8832.
- Siebrecht S., Herdel K., Schurr U., Tischner R. (2003) Nutrient translocation in the xylem of poplar—diurnal variations and spatial distribution along the shoot axis. *Planta* **217**, 783-793.
- Siebrecht S., Tischner R. (1999) Changes in the xylem exudate composition of poplar (*Populus tremula* x *P. alba*)—dependent on the nitrogen and potassium supply. *Journal of Experimental Botany* **50**, 1797-1806.

- Smith R.C., Epstein E. (1964) Ion absorption by shoot tissue: kinetics of potassium and rubidium absorption by corn leaf tissue. *Plant Physiology* **39**, 992.
- Smirnoff N., Stewart G.R. (1985) Nitrate assimilation and translocation by higher plants: comparative physiology and ecological consequences. *Physiologia Plantarum* **64**, 133-140.
- Sorger G.J., Ford R.E., Evans H.J. (1965) Effects of univalent cations on the immunoelectrophoretic behavior of pyruvic kinase. *Proceedings of the National Academy of Sciences, USA* **54**, 1614-1621.
- Spalding E.P., Hirsch R.E., Lewis D.R., Qi Z., Sussman M.R., Lewis B.D. (1999) Potassium uptake supporting plant growth in the absence of AKT1 channel activity. Inhibition by ammonium and stimulation by sodium. *Journal of General Physiology* **113**, 909-918.
- Sun J., Bankston J.R., Payandeh J., Hinds T.R., Zagotta W.N., Zheng N. (2014) Crystal structure of the plant dual-affinity nitrate transporter NRT1. 1. *Nature* **507**, 73-77.
- Szczerba M.W., Britto D.T., Balkos K.D., Kronzucker H.J. (2008a) Alleviation of rapid, futile ammonium cycling at the plasma membrane by potassium reveals K⁺-sensitive and-insensitive components of NH₄⁺ transport. *Journal of Experimental Botany* **59**, 303-313.
- Szczerba M.W., Britto D.T., Ali S.A., Balkos K.D., Kronzucker H.J. (2008b) NH₄⁺-stimulated and-inhibited components of K⁺ transport in rice (*Oryza sativa* L.). *Journal of Experimental Botany* **59**, 3415-3423.
- Szczerba M.W., Britto D.T., Kronzucker H.J. (2009) K⁺ transport in plants: physiology and molecular biology. *Journal of Plant Physiology* **166**, 447-466.
- Takei K., Takahashi T., Sugiyama T., Yamaya T., Sakakibara H. (2002) Multiple routes communicating nitrogen availability from roots to shoots: a signal transduction pathway mediated by cytokinin. *Journal of Experimental Botany* **53**, 971-977.

- Takei K., Ueda N., Aoki K., Kuromori T., Hirayama T., Shinozaki K., ..., Sakakibara H. (2004) AtIPT3 is a key determinant of nitrate-dependent cytokinin biosynthesis in *Arabidopsis*. *Plant and Cell Physiology* **45**, 1053-1062.
- ten Hoopen F., Cuin T.A., Pédas P., Hegelund J.N., Shabala S., Schjoerring J.K., Jahn T.P. (2010) Competition between uptake of ammonium and potassium in barley and *Arabidopsis* roots: molecular mechanisms and physiological consequences. *Journal of Experimental Botany* **61**, 2303-2315.
- Taochy C., Gaillard I., Ipotesi E., Oomen R., Leonhardt N., Zimmermann S., ..., Boyer J.C. (2015) The *Arabidopsis* root stele transporter NPF2.3 contributes to nitrate translocation to shoots under salt stress. *Plant Journal* **83**, 466-479.
- Topa M.A., Jackson W.A. (1988) Influence of ambient ammonium on net potassium uptake by decapitated maize seedlings. *New Phytologist* **110**, 135-141.
- Torres M.A., Dangel J.L. (2005) Functions of the respiratory burst oxidase in biotic interactions, abiotic stress and development. *Current Opinion in Plant Biology* **8**, 397-403.
- Touraine B., Grignon C. (1982) Energetic coupling of nitrate secretion into the xylem of corn roots. *Physiologie Vegetale* **20**, 33-39.
- Touraine B., Grignon N., Grignon C. (1988) Charge balance in NO_3^- fed soybean estimation of K^+ and carboxylate recirculation. *Plant Physiology* **88**, 605-612.
- Touraine B., Grignon N., Grignon C. (1990) Interaction between nitrate assimilation in shoots and nitrate uptake by roots of soybean (*Glycine max*). *Plant and Soil* **124**, 169-174.
- Tsay Y-F., Ho C-H., Chen H-Y., Lin S-H. (2011) Integration of nitrogen and potassium signaling. *Annual Review of Plant Biology* **62**, 207-226.

- Ullrich W.R., Larsson M., Larsson C.M., Lesch S., Novacky A. (1984) Ammonium uptake in *Lemna gibba* G 1, related membrane potential changes, and inhibition of anion uptake. *Physiologia Plantarum* **61**, 369-376.
- Ullrich C.I., Novacky A.J. (1990) Extra-and intracellular pH and membrane potential changes induced by K^+ , Cl^- , $H_2PO_4^-$, and NO_3^- uptake and fusicoccin in root hairs of *Limnobium stoloniferum*. *Plant Physiology* **94**, 1561-1567.
- Vale F.R., Jackson W.A., Volk R.J. (1987) Potassium influx into maize root systems influence of root potassium concentration and ambient ammonium. *Plant Physiology* **84**, 1416-1420.
- Vale F.R., Volk R.J., Jackson W.A. (1988) Simultaneous influx of ammonium and potassium into maize roots: kinetics and interactions. *Planta* **173**, 424-431.
- Van Beusichem M.L., Kirkby E.A., Baas R. (1988) Influence of nitrate and ammonium nutrition on the uptake, assimilation, and distribution of nutrients in *Ricinus communis*. *Plant Physiology* **86**, 914-921.
- Van Egmond F. (1978) Nitrogen nutritional aspects of the ionic balance of plants. In *Nitrogen in the Environment, Vol. 2: Soil-Plant-Nitrogen Relationships* (eds D. R. E. Nielsen, J. G. MacDonald), pp. 171-189. Academic Press, New York.
- Vermeulen S.J., Campbell B.M., Ingram J.S.I. (2012) Climate change and food systems. *Annual Review of Environment and Resources* **37**, 195-222.
- Walker D.J., Leigh R.A, Miller A.J. (1996) Potassium homeostasis in vacuolate plant cells. *Proceedings of the National Academy of Sciences, USA* **93**, 10510-10514.
- Wang M.Y., Siddiqi M.Y., Glass A.D.M. (1996) Interactions between K^+ and NH_4^+ : effects on ion uptake by rice roots. *Plant, Cell & Environment* **19**, 1037-1046.
- Wang M.Y., Siddiqi M.Y., Ruth T.J., Glass A.D.M. (1993) Ammonium uptake by rice roots. II. Kinetics of $^{13}NH_4^+$ influx across the plasmalemma. *Plant Physiology* **103**, 1259-1267.

- Wang M.Y., Glass A.D.M., Shaff J.E., Kochian L.V. (1994) Ammonium uptake by rice roots (III. Electrophysiology). *Plant Physiology* **104**, 899-906.
- Wang R., Liu D., Crawford N.M. (1998) The Arabidopsis CHL1 protein plays a major role in high-affinity nitrate uptake. *Proceedings of the National Academy of Sciences, USA* **95**, 15134-15139.
- Wang Y.H., Garvin D.F., Kochian L.V. (2001). Nitrate-induced genes in tomato roots. Array analysis reveals novel genes that may play a role in nitrogen nutrition. *Plant Physiology* **127**, 345-359.
- Wang Y.Y., Hsu P.K., Tsay Y.F. (2012) Uptake, allocation and signaling of nitrate. *Trends in Plant Science* **17**, 458-467.
- Wang R., Tischner R., Gutiérrez R. A., Hoffman M., Xing X., Chen M., ..., Crawford N.M. (2004) Genomic analysis of the nitrate response using a nitrate reductase-null mutant of *Arabidopsis*. *Plant Physiology* **136**, 2512-2522.
- Wang Y., Wu W-H. (2013) Potassium transport and signaling in higher plants. *Annual Review of Plant Biology* **64**, 451-476.
- White P.J. (2013) Improving potassium acquisition and utilisation by crop plants. *Journal of Plant Nutrition and Soil Science* **176**, 305-316.
- Wolt J.D. (1994) *Soil solution chemistry: applications to environmental science and agriculture*. John Wiley and Sons.
- Wyn Jones R.G., Pollard A. (1983) Proteins, enzymes and inorganic ions. *Encyclopedia of Plant Physiology* **15B**, 528-562.
- Xia X., Fan X., Wei J., Feng H., Qu H., Xie D., ..., Xu G. (2015) Rice nitrate transporter OsNPF2.4 functions in low-affinity acquisition and long-distance transport. *Journal of Experimental Botany* **66**, 317-331.

- Xu G., Fan X., Miller A.J. (2012) Plant nitrogen assimilation and use efficiency. *Annual Review of Plant Biology* **63**, 153-182.
- Xu G., Wolf S., Kafkafi U. (2002) Ammonium on potassium interaction in sweet pepper. *Journal of Plant Nutrition* **25**, 719-734.
- Xu J., Li H.D., Chen L.Q., Wang Y., Liu L.L., He L., Wu W.H. (2006) A protein kinase, interacting with two calcineurin B-like proteins, regulates K⁺ transporter AKT1 in Arabidopsis. *Cell* **125**, 1347-1360.
- Zhang F., Niu J., Zhang W., Chen X., Li C., Yuan L., Xie J. (2010) Potassium nutrition of crops under varied regimes of nitrogen supply. *Plant and Soil* **335**, 21-34.
- Zsoldos F., Haunold E., Herger P., Vasfaegy A. (2006) Effects of sulfate and nitrate on K⁺ uptake and growth of wheat and cucumber. *Physiologia Plantarum* **80**, 425-430.

Author Manuscript

Fig 1

Fig 2

Fig 3

Author

Fig 4

Author Mar

This review summarizes fundamental intersections between the pathways of inorganic nitrogen (NH_4^+ and NO_3^-) and potassium (K^+) acquisition in plants. Uptake, storage, translocation, and metabolism are discussed at levels of organization ranging from molecular-genetic processes to whole-plant physiology. The regulation and optimization of plant growth, yield, metabolism, and water-use efficiency are discussed in this nutritional context.

Author Manuscript

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Coskun, D;Britto, DT;Kronzucker, HJ

Title:

The nitrogen-potassium intersection: membranes, metabolism, and mechanism

Date:

2017-10-01

Citation:

Coskun, D., Britto, D. T. & Kronzucker, H. J. (2017). The nitrogen-potassium intersection: membranes, metabolism, and mechanism. PLANT CELL AND ENVIRONMENT, 40 (10), pp.2029-2041. <https://doi.org/10.1111/pce.12671>.

Persistent Link:

<http://hdl.handle.net/11343/290842>