TITLE

Brain Attacks and Stroke in Children

ARTICLE TYPE

Invited review

AUTHORS

Doctor Mark T. Mackay, MBBS FRACP^{1, 3, 4, 5}

Professor Paul Monagle, MD FRACP^{3,5}

Associate Professor Franz Babl, MD FRACP^{2, 3, 5}

AFFILIATIONS

1. Department of Neurology, Royal Children's Hospital, Parkville Australia
2. Emergency Department, Royal Children's Hospital Melbourne, Parkville Australia
3. Murdoch Childrens Research Institute, Parkville Australia
4. Florey Institute of Neurosciences and Mental Health, Parkville, Australia
5. Department of Paediatrics, University of Melbourne, Parkville Australia

CORRESPONDENCE TO

Dr. Mark Mackay, Department of Neurology, Royal Children's Hospital Melbourne, 50
Flemington Road, Parkville, Victoria, Australia. 3052

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/jpc.13086](https://doi.org/10.1111/jpc.13086)

JPCH EM Mackay_ Brain Attacks in Children_R1_22072015

Email: mark.mackay@rch.org.au

Phone: +61-3-9345 5661 Fax: +61-3-9345 5977

Version 4

Title character count: 12 words.

Abstract count: 150 words.

Manuscript count: 1990 words.

References: 56.

Tables: 1.

Abstract

Emergency physicians are often the first point of contact in children presenting with acute neurological disorders. Differentiating serious disorders such as stroke, from benign disorders such as migraine can be challenging. Clinical assessment influences decision making, in particular the need for emergent neuroimaging to confirm diagnosis. This review describes the spectrum of disorders causing “brain attack” symptoms, or acute onset focal neurological dysfunction, with particular emphasis on childhood stroke, because early recognition is essential to improve access to thrombolytic treatments which have improved outcomes in adults. Clues to diagnosis of specific conditions are discussed. Symptoms and signs which discriminate stroke from mimics are described, highlighting differences to adults. Haemorrhagic and ischaemic stroke have different presenting features which influence choice of the most appropriate imaging modality to maximise diagnostic accuracy. Improvements in the care of children with brain attacks requires coordinated approaches and system improvements similar to those developed in adults.

Key words

Neurological, emergency, cerebrovascular disease, childhood stroke, clinical neurology.

Additional key words: brain attack.

Key points

1. Triage nurses and emergency physicians play key roles in differentiating uncommon but serious conditions such as stroke, from more common and benign conditions such as migraine, in children presenting with acute focal neurological deficits or brain attacks.
2. Clinical diagnosis influences decisions about timing of investigations including emergent neuroimaging and the most appropriate treatments. Bedside tools have been developed to assist paramedics and emergency physicians to discriminate stroke from mimics but their diagnostic utility has not been adequately tested in children.
3. Development of paediatric brain attack protocols are urgently needed to improve care of children with neurological emergencies and increase access to time critical interventions such as thrombolytic therapy in ischaemic stroke.

Introduction: Brain attacks in children

Paediatric neurological emergencies include status epilepticus, stroke, acute raised intracranial pressure, traumatic brain injury, central nervous system infections, acute demyelinating disorders and acute flaccid paralysis. Some of these conditions present with altered conscious state whereas others present with sudden onset focal neurological or “brain attack” symptoms which may include weakness, speech, sensory or visual disturbance, headache or ataxia. In adult public education campaigns the term brain attack is used synonymously with stroke because of mechanistic similarities to heart attacks. This review will focus on differentiating uncommon but serious causes of brain attack symptoms such as stroke, from more benign conditions such as migraine, and discuss approaches to investigation and management.

The spectrum of paediatric conditions causing brain attack symptoms differ from adults.

Whereas adults with brain attacks are diagnosed with stroke in three quarter of emergency department (ED) presentations⁽¹⁻³⁾ children presenting with brain attack symptoms generally do not have a stroke.⁽⁴⁾ Yet, it is critical to identify these children who require urgent neuroimaging to improve access to hyper-acute therapies such as tissue plasminogen activator and endovascular treatments.⁽⁵⁻⁹⁾ A recent study of children presenting to an Australian tertiary paediatric ED found that migraine was the most common cause of sudden onset focal neurological symptoms and signs, accounting for 28% of cases. First febrile or afebrile seizures

were the second most common diagnosis in 15% of children, followed by Bells Palsy in 10%, ischaemic or haemorrhagic stroke in 7%, and conversion disorders in 6% of children.⁽⁴⁾

In addition the paediatric conditions which mimic stroke differ from adults, and disorders such as post infectious cerebellitis are only encountered in children.⁽⁴⁾ Another study describing the spectrum of disorders mimicking paediatric stroke in an inpatient setting found that serious neurological conditions accounted for almost two-thirds of cases, with diagnoses including posterior reversible leucoencephalopathy, vascular abnormalities, inflammatory disease, intracranial infection, metabolic stroke, tumours, drug toxicity and idiopathic intracranial hypertension.⁽¹⁰⁾

Clues to specific neurological diagnoses in children presenting with brain attack symptoms (Table)

Migraine is a common neurological disorder of children and teenagers^(11, 12) and is typically associated with unilateral throbbing headache. Associated symptoms may include nausea, photophobia and phonophobia.⁽⁵⁻⁹⁾ Migraine headaches can be of shorter duration than adults and are frequently bilateral in children. Auras are less common than adults and typically consist of a mix of positive and negative visual symptoms, and less commonly sensory symptoms.⁽¹³⁾ Simple analgesics are the mainstay of acute treatment in younger children but triptans can be considered in teenagers.⁽¹⁴⁾ Clues to childhood migraine diagnosis include a positive family history, *gradual* onset of neurological symptoms, which usually resolve within 15-30 minutes,

followed by headache which builds in intensity. Focal weakness, speech disturbance, confusion and altered consciousness are unusual. Red flags which warrant urgent neuroimaging include sudden onset severe headache associated with stroke and focal neurological deficits which are not resolving.

Focal seizures can be followed by a Todd's paresis and may be difficult to differentiate from stroke. Todd's paresis typically occurs in the context of known focal epilepsy and resolves within a few hours. Therefore children with no prior history of seizures whose neurological deficits are not resolving should undergo urgent neuroimaging because seizures are a common presenting symptom of stroke,⁽¹⁵⁻¹⁸⁾ encephalitis, and acute disseminated encephalomyelitis.

There are well developed protocols for emergency management of seizures

(<https://www.apls.org.au/page/about-us>) based on good evidence from randomised controlled trials.⁽¹⁹⁻²²⁾

Diagnosis of Bells palsy is fairly straightforward but requires careful examination by the emergency physician to ensure weakness involves the upper and lower facial muscles and *absence* of other neurological signs referable to the brainstem.⁽²³⁾ There is good evidence to support use of steroids in adults^(24, 25) but data are lacking in children.⁽²⁶⁾

Conversion disorders are probably the most challenging group of patients for non-neurologists because the child's neurological symptoms are by definition unexplainable from a neuroanatomical perspective. Conversion disorders typically occur in children over the age of 8

years⁽²⁷⁾ and present with multiple neurological complaints including weakness, speech, sensory or visual disturbance, seizures, collapse, inability to walk or abnormal gait.^(28, 29) Clues to diagnosis include examination findings which are inconsistent with the complaint, signs which vary from one examination to the next or which do not conform to neuroanatomical pathways. Input from colleagues with specific expertise in neurological examination and knowledge of neuroanatomy is often required and non-urgent neuroimaging is sometimes indicated to exclude more serious organic conditions.⁽²⁷⁾

Childhood brain tumors can present with focal neurological deficits, often referable to posterior fossa structures but they are usually associated with non-abrupt onset of symptoms unless there is a bleed into the lesion. Similarly acute disseminated encephalomyelitis (ADEM) typically has non-abrupt onset, often following an identifiable prodromal infection. Clues to the clinical diagnosis of ADEM include encephalopathy, seizures and *multifocal* neurological deficits referable to more than one location within the central nervous system.⁽³⁰⁾ Magnetic resonance imaging is the neuroimaging investigation of choice and typically shows multifocal lesions involving white matter, deep grey matter and sometimes the cerebral cortex. High dose corticosteroids are the mainstay of treatment.⁽³⁰⁾

Childhood stroke is the most time critical of all neurological emergencies

Childhood stroke may be less common than other causes of brain attack symptoms but it is associated with high personal cost to affected children and their families,⁽³¹⁾ because more than

half of survivors have long-term impairments that interfere with normal development and lifestyle.⁽³²⁻³⁴⁾ Licencing of tissue plasminogen activator in 1996 was a watershed moment for treatment of stroke in adults. The ischaemic penumbra is the biological target of thrombolytic treatments in acute stroke. Brain tissue which is destined to go onto infarction can be salvaged if blood flow is restored by vascular recanalisation.⁽³⁵⁾ As a consequence there is a major focus on reducing time delay to stroke diagnosis. Thrombolysis is established as an effective treatment if given within 4 and a half hours of from symptom onset, based on evidence from multiple randomised controlled trials.⁽³⁶⁾ Very recently several adult studies have confirmed the effectiveness of endovascular treatments (such as intra-arterial lysis and mechanical thrombectomy), when given within 6 to 12 hours from symptoms onset.⁽⁵⁻⁹⁾ Therefore rapid radiological confirmation of diagnosis within 6 to 12 hours of symptom onset is the key to accessing thrombolytic or endovascular treatments to reduce the long term impact of stroke. Adult emergency stroke management guidelines recommend radiological confirmation of diagnosis within 45 minutes, decision to thrombolyse within 60 minutes and admission to a stroke unit within 3 hours.⁽³⁷⁾

Delays to stroke diagnosis in children are much longer than adults with three studies reporting delays of more than 20 hours.⁽³⁸⁻⁴⁰⁾ Systems of care do not currently exist for childhood stroke. In contrast to well-developed protocols for other neurological emergencies such as status epilepticus the Advanced Paediatric Life Support (APLS) guidelines do not contain algorithms for

identification, investigation or management of children with brain attack symptoms relevant for stroke. (<https://www.apls.org.au/page/about-us>) Reasons for delayed presentation of children with stroke are poorly elucidated but include failure to call an ambulance, short duration⁽⁴¹⁾ or mild symptom severity⁽³⁸⁾ and poor awareness of stroke amongst paediatric physicians.⁽³⁹⁾ Strategies are urgently required to reduce long term morbidity by improving identification and acute treatment of childhood stroke.

Clinical features which distinguish childhood stroke from mimics

Clinical differentiation of stroke from mimics is the first step in reducing treatment delays.

More common presenting features of stroke include hemiparesis (22-100%), headache (16-45%), altered mental state (12-24%), speech disturbance (28-55%), altered consciousness (24-52%) and seizures (11-58%).^(16, 41-45) Age at presentation influences the clinical presentation with seizures, altered mental state and non-focal signs being more likely in infants.^(16, 43)

The frequent occurrence of seizures in childhood stroke⁽¹⁵⁻¹⁸⁾ is a notable difference to adults.^(1, 46) This difference may reflect age related cortical hyperexcitability or decreased inhibitory influences in the developing brain. In adults seizures are more likely to occur with mimics and are used as a predictor variable for mimic diagnosis in prehospital and emergency department bedside stroke recognition tools.⁽⁴⁶⁻⁴⁸⁾

Sudden onset symptoms of focal weakness, speech disturbance and headache and signs including face, arm, hand and leg weakness, dysphasia, dysarthria, inability to walk and altered

conscious state are associated with increased odds of stroke. In contrast absence of focal signs on examination is with decreased odds of stroke.⁽⁴⁹⁾ Presenting clinical features differ by stroke subtype. Arterial ischaemic stroke is more likely to present with focal limb or face weakness, speech disturbance, limb incoordination or ataxia whereas haemorrhagic stroke is more likely to present with headache, vomiting and altered mental state.⁽⁵⁰⁾

Bedside recognition tools improve accuracy of stroke diagnosis in adults

Whereas stroke recognition tools have been developed to improve paramedic^(47, 48, 51) and emergency physician accuracy in differentiating stroke from mimics in adults⁽⁴⁶⁾ no such tools exist in children and a preliminary analysis of adult bedside tools in childhood arterial ischaemic stroke found that they had lower sensitivity than reported in adult studies.⁽⁵²⁾

Emergent imaging is the key to confirming diagnosis and selecting appropriate treatments

Standard adult stroke protocols recommend CT imaging to identify intracranial haemorrhage because it is a contraindication to thrombolysis.⁽³⁷⁾ The higher a priori probability of stroke in adults means there is a low likelihood of inappropriately administering thrombolytic treatment to someone with a stroke mimic. In contrast mimics are more common in children and therefore specificity of brain imaging is very important. Computed tomography has the advantage of greater accessibility to emergency physicians. It is adequate for identification of intracranial haemorrhage but has poor sensitivity for detection of acute brain ischaemia,

ranging from 16% to 56% in paediatric studies.^(38-41, 50) Therefore MRI is the diagnostic modality of choice for suspected arterial ischaemic stroke.⁽⁵³⁾

Rapid MRI in adult stroke in the ED may allow more accurate positive identification of stroke and mimics.⁽⁵⁴⁾ The latter finding is particularly relevant to children given the lower a priori probability of stroke. However the need for sedation in young children presents an additional challenge in the paediatric emergency setting.

Computed tomography is also relatively insensitive to detection of demyelinating lesions, acute encephalitis and detection of posterior fossa tumors due to bony artefact from the base of skull. Therefore MRI is also the neuroimaging investigation of choice for children with suspected ADEM, viral encephalitis and posterior fossa tumors.

Management of neurological emergencies in the ED

Urgent interventions are required for several serious neurological disorders. Treatments may include antibiotics for meningitis, antiviral agents for encephalitis, anticonvulsants for status epilepticus, immunotherapies for acute disseminated encephalomyelitis and transverse myelitis and mannitol, steroids or neurosurgical interventions for raised intracranial pressure. Acute treatment of childhood stroke currently focuses on normalisation of physiological parameters and early initiation of secondary preventative anticoagulant or antithrombotic treatment. Paediatric stroke management guidelines do not currently recommend tPA usage in children because there is a paucity of efficacy and safety data for tPA in children.^(55, 56)

Conclusions and future directions

The spectrum of disorders causing paediatric brain attack symptoms differ to adults with migraine, seizures and Bell's palsy being the most common conditions. The presenting symptoms and signs which discriminate stroke from mimics are similar to adults but seizures are a common presenting feature of stroke in children and therefore they are not discriminatory for mimics. This represents an important difference from adults and means that bedside stroke recognition tools, which are widely used by paramedics and in the ED, may not be as accurate in children to differentiate stroke from mimics. Haemorrhagic and ischaemic stroke have different presenting features which influence the choice of the most appropriate imaging modality to maximise accuracy of radiological diagnosis. Children presenting with sudden onset headache, vomiting and coma have a higher probability of haemorrhagic stroke and CT is the most appropriate acute neuroimaging modality. Children presenting with sudden onset focal symptoms and signs have a higher probability of ischaemic stroke and MRI is the preferred first imaging modality.

Better and earlier diagnosis of brain attacks, in particular the differentiation between stroke and non-strokes, are crucial for developing the capacity to offer thrombolysis to select children with ischaemic stroke to improve long term outcomes.

References

1. Hand PJ, Kwan J, Lindley RI, Dennis MS, Wardlaw JM. Distinguishing between stroke and mimic at the bedside: the brain attack study. *Stroke*. 2006 Mar;37(3):769-75.
2. Libman RB, Wirkowski E, Alvir J, Rao TH. Conditions that mimic stroke in the emergency department. Implications for acute stroke trials. *Archives of Neurology*. 1995 Nov;52(11):1119-22.
3. Whiteley WN, Wardlaw JM, Dennis MS, Sandercock PA. Clinical scores for the identification of stroke and transient ischaemic attack in the emergency department: a cross-sectional study. *Journal of neurology, neurosurgery, and psychiatry*. 2011 Sep;82(9):1006-10.
4. Mackay MT, Chua ZK, Lee M, Yock-Corrales A, Churilov L, Monagle P, et al. Stroke and nonstroke brain attacks in children. *Neurology*. 2014 Apr 22;82(16):1434-40.
5. Berkhemer OA, Fransen PS, Beumer D, van den Berg LA, Lingsma HF, Yoo AJ, et al. A randomized trial of intraarterial treatment for acute ischemic stroke. *The New England journal of medicine*. 2015 Jan 1;372(1):11-20.
6. Goyal M, Demchuk AM, Menon BK, Eesa M, Rempel JL, Thornton J, et al. Randomized Assessment of Rapid Endovascular Treatment of Ischemic Stroke. *The New England journal of medicine*. 2015 Feb 11. PubMed PMID: 25671798. Epub 2015/02/12. Eng.
7. Campbell BC, Mitchell PJ, Kleinig TJ, Dewey HM, Churilov L, Yassi N, et al. Endovascular Therapy for Ischemic Stroke with Perfusion-Imaging Selection. *The New England journal of medicine*. 2015 Feb 11. PubMed PMID: 25671797. Epub 2015/02/12. Eng.
8. Jovin TG, Chamorro A, Cobo E, de Miquel MA, Molina CA, Rovira A, et al. Thrombectomy within 8 hours after symptom onset in ischemic stroke. *The New England journal of medicine*. 2015 Jun 11;372(24):2296-306.
9. Saver JL, Goyal M, Bonafe A, Diener HC, Levy EI, Pereira VM, et al. Stent-retriever thrombectomy after intravenous t-PA vs. t-PA alone in stroke. *The New England journal of medicine*. 2015 Jun 11;372(24):2285-95.
10. Shellhaas RA, Smith SE, O'Tool E, Licht DJ, Ichord RN. Mimics of childhood stroke: characteristics of a prospective cohort. *Pediatrics*. 2006 Aug;118(2):704-9.
11. Winner P. Pediatric headache. *Curr Opin Neurol*. 2008 Jun;21(3):316-22.
12. Goadsby PJ, Lipton RB, Ferrari MD. Migraine--current understanding and treatment. *The New England journal of medicine*. 2002 Jan 24;346(4):257-70.

13. Hershey AD, Winner P, Kabbouche MA, Gladstein J, Yonker M, Lewis D, et al. Use of the ICHD-II criteria in the diagnosis of pediatric migraine. *Headache*. 2005 Nov-Dec;45(10):1288-97.
14. Lewis D, Ashwal S, Hershey A, Hirtz D, Yonker M, Silberstein S, et al. Practice parameter: pharmacological treatment of migraine headache in children and adolescents: report of the American Academy of Neurology Quality Standards Subcommittee and the Practice Committee of the Child Neurology Society. *Neurology*. 2004 Dec 28;63(12):2215-24.
15. Chadehumbe MA, Khatrri P, Khoury JC, Alwell K, Szaflarski JP, Broderick JP, et al. Seizures are common in the acute setting of childhood stroke: a population-based study. *J Child Neurol*. 2009 Jan;24(1):9-12.
16. Mallick AA, Ganesan V, Kirkham FJ, Fallon P, Hedderly T, McShane T, et al. Childhood arterial ischaemic stroke incidence, presenting features, and risk factors: a prospective population-based study. *The Lancet Neurology*. 2014 Jan;13(1):35-43.
17. Abend NS, Beslow LA, Smith SE, Kessler SK, Vossough A, Mason S, et al. Seizures as a presenting symptom of acute arterial ischemic stroke in childhood. *Journal of Pediatrics*. 2011;159(3):479-83.
18. Singh RK, Zecavati N, Singh J, Kaulas H, Nelson KB, Dean NP, et al. Seizures in acute childhood stroke. *Journal of Pediatrics*. 2012 Feb;160(2):291-6.
19. Scott RC, Besag FM, Neville BG. Buccal midazolam and rectal diazepam for treatment of prolonged seizures in childhood and adolescence: a randomised trial. *Lancet*. 1999 Feb 20;353(9153):623-6.
20. McIntyre J, Robertson S, Norris E, Appleton R, Whitehouse WP, Phillips B, et al. Safety and efficacy of buccal midazolam versus rectal diazepam for emergency treatment of seizures in children: a randomised controlled trial. *Lancet*. 2005 Jul 16-22;366(9481):205-10.
21. Fisgin T, Gurer Y, Tezic T, Senbil N, Zorlu P, Okuyaz C, et al. Effects of intranasal midazolam and rectal diazepam on acute convulsions in children: prospective randomized study. *J Child Neurol*. 2002 Feb;17(2):123-6.
22. Chin RF, Neville BG, Peckham C, Wade A, Bedford H, Scott RC. Treatment of community-onset, childhood convulsive status epilepticus: a prospective, population-based study. *The Lancet Neurology*. 2008 Aug;7(8):696-703.
23. Manning JJ, Adour KK. Facial paralysis in children. *Pediatrics*. 1972 Jan;49(1):102-9.
24. Engstrom M, Berg T, Stjernquist-Desatnik A, Axelsson S, Pitkaranta A, Hultcrantz M, et al. Prednisolone and valaciclovir in Bell's palsy: a randomised, double-blind, placebo-controlled, multicentre trial. *The Lancet Neurology*. 2008 Nov;7(11):993-1000.

25. Gilden DH. Clinical practice. Bell's Palsy. The New England journal of medicine. 2004 Sep 23;351(13):1323-31.
26. Pitaro J, Waissbluth S, Daniel SJ. Do children with Bell's palsy benefit from steroid treatment? A systematic review. Int J Pediatr Otorhinolaryngol. 2012 Jul;76(7):921-6.
27. de Gusmao CM, Guerriero RM, Bernson-Leung ME, Pier D, Ibeziako PI, Bujoreanu S, et al. Functional neurological symptom disorders in a pediatric emergency room: diagnostic accuracy, features, and outcome. Pediatr Neurol. 2014 Aug;51(2):233-8.
28. Dula DJ, DeNaples L. Emergency department presentation of patients with conversion disorder. Academic emergency medicine : official journal of the Society for Academic Emergency Medicine. 1995 Feb;2(2):120-3.
29. Kozłowska K, Nunn KP, Rose D, Morris A, Ouvrier RA, Varghese J. Conversion disorder in Australian pediatric practice. Journal of the American Academy of Child and Adolescent Psychiatry. 2007 Jan;46(1):68-75.
30. Tenenbaum S, Chitnis T, Ness J, Hahn JS, International Pediatric MSSG. Acute disseminated encephalomyelitis. Neurology. 2007 Apr 17;68(16 Suppl 2):S23-36.
31. Gardner MA, Hills NK, Sidney S, Johnston SC, Fullerton HJ. The 5-year direct medical cost of neonatal and childhood stroke in a population-based cohort. Neurology. 2010;74(5):372-8.
32. deVeber GA, MacGregor D, Curtis R, Mayank S. Neurologic outcome in survivors of childhood arterial ischemic stroke and sinovenous thrombosis. Journal of child neurology. 2000;15(5):316-24.
33. Ganesan V, Hogan A, Shack N, Gordon A, Isaacs E, Kirkham FJ. Outcome after ischaemic stroke in childhood. Developmental Medicine & Child Neurology. 2000;42(7):455-61.
34. Gordon AL, Ganesan V, Towell A, Kirkham FJ. Functional outcome following stroke in children. Journal of child neurology. 2002;17(6):429-34.
35. Schlaug G, Benfield A, Baird AE, Siewert B, Lovblad KO, Parker RA, et al. The ischemic penumbra: operationally defined by diffusion and perfusion MRI. Neurology. 1999 Oct 22;53(7):1528-37.
36. Wardlaw JM, Murray V, Berge E, del Zoppo G, Sandercock P, Lindley RL, et al. Recombinant tissue plasminogen activator for acute ischaemic stroke: an updated systematic review and meta-analysis. Lancet. 2012 Jun 23;379(9834):2364-72.
37. Jauch EC, Cucchiara B, Adeoye O, Meurer W, Brice J, Chan YY, et al. Part 11: adult stroke: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. Circulation. 2010 Nov 2;122(18 Suppl 3):S818-28.
38. Rafay MF, Pontigon AM, Chiang J, Adams M, Jarvis DA, Silver F, et al. Delay to diagnosis in acute pediatric arterial ischemic stroke. Stroke. 2009 Jan;40(1):58-64.

39. Srinivasan J, Miller SP, Phan TG, Mackay MT. Delayed recognition of initial stroke in children: need for increased awareness. *Pediatrics*. 2009;124(2):e227-34.
40. Mallick AA, Ganesan V, Kirkham FJ, Fallon P, Hedderly T, McShane T, et al. Diagnostic delays in paediatric stroke. *Journal of neurology, neurosurgery, and psychiatry*. 2014 Oct 23.
41. McGlennan C, Ganesan V. Delays in investigation and management of acute arterial ischaemic stroke in children. *Developmental Medicine & Child Neurology*. 2008;50(7):537-40.
42. Mackay MT, Wiznitzer M, Benedict SL, Lee KJ, Deveber GA, Ganesan V, et al. Arterial Ischemic Stroke Risk Factors: The International Pediatric Stroke Study. *Annals of Neurology*. 2011;69(1):130-40.
43. Zimmer JA, Garg BP, Williams LS, Golomb MR. Age-related variation in presenting signs of childhood arterial ischemic stroke. *Pediatric neurology*. 2007;37(3):171-5.
44. Steinlin M, Pfister I, Pavlovic J, Everts R, Boltshauser E, Capone Mori A, et al. The first three years of the Swiss Neuropaediatric Stroke Registry (SNPSR): a population-based study of incidence, symptoms and risk factors. *Neuropediatrics*. 2005;36(2):90-7.
45. Gabis LV, Yangala R, Lenn NJ. Time lag to diagnosis of stroke in children. *Pediatrics*. 2002 Nov;110(5):924-8.
46. Nor AM, Davis J, Sen B, Shipsey D, Louw SJ, Dyker AG, et al. The Recognition of Stroke in the Emergency Room (ROSIER) scale: development and validation of a stroke recognition instrument. *Lancet Neurology*. 2005 Nov;4(11):727-34.
47. McQuillen PS, Miller SP. Congenital heart disease and brain development. *Annals of the New York Academy of Sciences*. 2010 Jan;1184:68-86.
48. Bray JE, Coughlan K, Barger B, Bladin C. Paramedic diagnosis of stroke: examining long-term use of the Melbourne Ambulance Stroke Screen (MASS) in the field. *Stroke*. 2010;41(7):1363-6.
49. Mackay MT, Lee M, Churilov L, Yock-Corrales A, Donnan G, Monagle PB, F. Pediatric Brain Attacks: Differentiating Between Stroke and Mimics in the Emergency Room. *Stroke*. 2014;45(Suppl 1):A37.
50. Yock-Corrales A, Mackay MT, Mosley I, Maixner W, Babl FE. Acute childhood arterial ischemic and hemorrhagic stroke in the emergency department. *Annals of Emergency Medicine*. 2011 Aug;58(2):156-63.
51. Kidwell CS, Starkman S, Eckstein M, Weems K, Saver JL. Identifying stroke in the field. Prospective validation of the Los Angeles prehospital stroke screen (LAPSS). *Stroke*. 2000;31(1):71-6.
52. Yock-Corrales A, Babl FE, Mosley IT, Mackay MT. Can the FAST and ROSIER adult stroke recognition tools be applied to confirmed childhood arterial ischemic stroke? *BMC Pediatrics*. 2011;11(Journal Article):93.

53. Chalela JA, Kidwell CS, Nentwich LM, Luby M, Butman JA, Demchuk AM, et al. Magnetic resonance imaging and computed tomography in emergency assessment of patients with suspected acute stroke: a prospective comparison. *Lancet*. 2007 Jan 27;369(9558):293-8.
54. Solling C, Ashkanian M, Hjort N, Gyldensted C, Andersen G, Ostergaard L. Feasibility and logistics of MRI before thrombolytic treatment. *Acta Neurologica Scandinavica*. 2009;120(3):143-9.
55. Roach ES, Golomb MR, Adams R, Biller J, Daniels S, Deveber G, et al. Management of stroke in infants and children: a scientific statement from a Special Writing Group of the American Heart Association Stroke Council and the Council on Cardiovascular Disease in the Young. *Stroke*. 2008;39(9):2644-91.
56. Monagle P, Chan AK, Goldenberg NA, Ichord RN, Journeycake JM, Nowak-Gottl U, et al. Antithrombotic therapy in neonates and children: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest*. 2012 Feb;141(2 Suppl):e737S-801S.

Formatted: Font: Not Bold, Font color: Auto

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Mackay, MT;Monagle, P;Babl, FE

Title:

Brain attacks and stroke in children

Date:

2016-02-01

Citation:

Mackay, M. T., Monagle, P. & Babl, F. E. (2016). Brain attacks and stroke in children. JOURNAL OF PAEDIATRICS AND CHILD HEALTH, 52 (2), pp.158-163. <https://doi.org/10.1111/jpc.13086>.

Persistent Link:

<http://hdl.handle.net/11343/290873>