The chronology and tectonic style of landscape evolution along the elevated Atlantic continental margin of South Africa resolved by joint apatite fission track and (U-Th-Sm)/He thermochronology

Mark Wildman¹, Roderick Brown¹, Romain Beucher^{1*}, Cristina Persano¹, Fin Stuart², Kerry Gallagher³, James Schwanethal⁴ and Andrew Carter⁵

1. School of Geographical and Earth Sciences, College of Science and Engineering, University of Glasgow, Gregory Building, Glasgow, G12 8QQ, Scotland.
2. Scottish Universities Environmental Research Centre, Rankine Avenue, Scottish Enterprise Technology Park, East Kilbride, G75 0QF, UK
3. Géosciences Rennes, Université de Rennes 1, Campus de Beaulieu, 35042 Rennes, France
4. Department of Earth Sciences, University College London, Gower Street, London, WC1E 6BT
5. Department of Earth and Planetary Sciences, Birkbeck, University of London, Malet Street, WC1E 7HX, United Kingdom

* Current address: Department of Earth Science, Faculty of Mathematics and Natural Sciences, University of Bergen, 5020 Bergen, Norway.

Corresponding author e-mail address: markwildman88@hotmail.com

Contact telephone number: +44 141 330 5469

Key Points:

- (1) First joint apatite fission track and (U-Th-Sm)/He data for the western continental margin of South Africa;
- (2) Present topography is a result of km-scale erosion during the Cretaceous with <1km denudation during the Cenozoic
- (3) Localised faulting onshore as well as regional uplift of the margin inferred during the mid to Late Cretaceous.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1002/2015TC004042](https://doi.org/10.1002/2015TC004042)

Abstract

Atlantic-type continental margins have long been considered “passive” tectonic settings throughout the entire post-rift phase. Recent studies question the long-term stability of these margins and have shown that post-rift uplift and reactivation of pre-existing structures may be a common feature of a continental margin’s evolution. The Namaqualand sector of the western continental margin of South Africa is characterised by a ubiquitously faulted basement but lacks preservation of younger geological strata to constrain post-rift tectonic fault activity. Here we present the first systematic study using joint apatite fission track and apatite (U-Th-Sm)/He thermochronology to achieve a better understanding on the chronology and tectonic style of landscape evolution across this region. Apatite fission-track ages range from 58.3 ± 2.6 to 132.2 ± 3.6 Ma, with mean track lengths between 10.9 ± 0.19 and 14.35 ± 0.22 μm , and mean (U-Th-Sm)/He sample ages range from 55.8 ± 31.3 to 120.6 ± 31.4 Ma. Joint inverse modelling of these data reveals two distinct episodes of cooling at c. 150-130 Ma and 110-90 Ma with limited cooling during the Cenozoic. Estimates of denudation based on these thermal histories predict c. 1-3 km of denudation coinciding with two major tectonic events. The first event, during the Early Cretaceous, was driven by continental rifting and the development and removal of syn-rift topography. The second event, during the Late Cretaceous, includes localised reactivation of basement structures as well as regional mantle driven uplift. Relative tectonic stability prevailed during the Cenozoic and regional denudation over this time is constrained to be less than 1 km.

1. INTRODUCTION

Passive continental margins, as the term explicitly implies, have been characterised by an apparent tectonic stability following their formation by intracontinental rifting and break-up (Braun and Beaumont, 1989; Lister et al., 1991; Ziegler and Cloetingh, 2004). However, a growing body of work is questioning this view and providing better insight into the thermal and structural processes that operate during rifting and influence the long-term geomorphic development of extensional continental margins and their interior hinterlands (Péron-Pinvidic and Manatschal, 2009; Bronner et al., 2011; Paton, 2012; Dauteuil et al., 2013; Masini et al., 2013; Karl et al., 2013; Péron-Pinvidic et al., 2013; Brune et al., 2014; Huismans and Beaumont, 2014; Koopmann et al., 2014; Salomon et al., 2014).

Plate reconstructions have primarily focussed on in-plane restorations of large plate blocks and have used overlaps and gaps in plate reconstructions as evidence for extensional and compressional deformation of the lithosphere, respectively (Eagles and König, 2008; Aslanian et al., 2009; Torsvik et al., 2009; Moulin et al., 2010; Aslanian and Moulin, 2013; Gaina et al., 2013; Heine et al., 2013; Pérez-Díaz and Eagles, 2014). The nature of this lithospheric deformation has been explored by numerical modelling approaches that have revealed complex styles of rifted margin development, involving rift migration (Brune et al., 2014; Naliboff and Buiter, 2015), hyper-extension (Lundin and Doré, 2011; Péron-Pinvidic et al., 2013; Redfield and Osmundsen, 2013), and/or multi-phase rifting (Reston, 2005; Blaich et al., 2011; Huismans and Beaumont, 2014). These models have predicted different styles and mechanisms for intracontinental deformation (Pérez-Gussinyé et al., 2007; Cloetingh and Burov, 2011; Kennett and Iaffaldano, 2013; Calignano et al., 2015).

Alongside geodynamical modelling of rifted margin evolution, but treated independently, a growing body of work has investigated regional and local vertical motions of the Earth's crust as a result of “dynamic” uplift or subsidence stemming from vertical stresses imposed on the base of the lithosphere by convection of the underlying mantle (Moucha et al., 2008; Braun, 2010; Forte et al., 2010; Flament et al., 2014). Recently, new thermo-mechanical

modelling approaches that have coupled the vertical forces acting on the base of the lithosphere, in-plane stresses from plate movements and the deformational response of a rheologically and structurally heterogeneous lithosphere over different length scales (Guillou-Frottier et al., 2012; Cloetingh et al., 2013; Buiter and Torsvik, 2014; Burov and Gerya, 2014; Colli et al., 2014; Koptev et al., 2015). A fundamental outcome of these modelling approaches is that plate motions, mantle flow and vertical motions of the surface are all intrinsically linked (François et al., 2013; Colli et al., 2014).

Tectonic uplift, as a consequence of deformation of the lithosphere, will trigger a response in surface processes (Willet, 1999; Burbank and Anderson, 2012). Therefore, new models of how the crust responds to intracontinental rifting and deep Earth processes will have implications for the origin and longevity of topography at high elevation passive continental margins. This issue is particularly pertinent for the South African case, where the antiquity of the present day first-order topography is still ardently debated. One end-member scenario argues that the South African landscape is “young”, having been formed by major river incision and erosion of a flat initial topography in response to regional epeirogenic uplift during the late Cenozoic (e.g. Partridge and Maud, 1987; Burke, 1996; Burke and Gunnell, 2008; Roberts and White, 2010; Paul et al., 2014; Rudge et al., 2015). Contrary to this, another end-member scenario proposes that the present day landscape is an “old” remnant of Cretaceous topography that was produced by deep erosion of uplifted rift flanks following continental break-up, with minimal erosion during the Cenozoic (Gallagher and Brown, 1999; Bierman and Caffee, 2001; Brown et al., 2002; van der Beek et al., 2002; Decker et al., 2013). However, several thermochronometry studies have argued for a more eventful post-rift history involving tectonic reactivation in the mid-Late Cretaceous (Raab et al., 2002; Tinker et al., 2008a; Kounov et al., 2009; Brown et al., 2014; Wildman et al., 2015) or multiple phases of regional uplift/exhumation and subsidence/burial (e.g. Green et al., 2013; Green et al., 2015).

Due to a lack of post-rift geological markers and poor age constraints on the timing of deformation on major structures, the tectonic history of the southwest African continental

margin is incomplete. Recent investigations have alluded to the potential importance of post-rift structural reactivation and neo-tectonic activity to the development of the South African margin (Viola et al., 2005, 2012; Brandt et al., 2003, 2005; Andreoli et al., 1996, 2009; de Beer, 2012; Kounov et al., 2009; Wildman et al., 2015), and post-rift reactivation has been advocated by investigations of other so-called “passive” continental margins (Redfield et al., 2005; Cogné et al., 2011; Holford et al., 2014; Franco-Magalhaes et al., 2014; Ksienzyk et al., 2014; Leprêtre et al., 2015). Constraining the surface response to post-rift deformation over different length scales will have major implications for conceptual geomorphic models of high-elevation continental margins that envisage regional patterns of denudation triggered by regional base level fall and augmented by long wavelength epeirogenic/flexural uplift.

In this paper we report new quantitative constraints on the timing and magnitude of major episodes of denudation across the continental margin obtained from a dense array of joint apatite fission track (AFT) and apatite (U-Th-Sm)/He (AHe) low-temperature thermochronometry samples. The sampling strategy was to collect samples from the coastline to the elevated continental interior with high-density sampling through the high-relief escarpment zone referred to as the Namaqualand Highlands (Fig. 1). Samples from the Namaqualand Highlands were collected with a specific focus on constraining brittle offsets on basement structures. This approach was employed to detect short and long wavelength patterns of crustal cooling. By using combined AFT and AHe thermochronometry, we achieve tighter thermal-history constraints than has previously been obtained by studies using either system in isolation. In particular, including AHe data helps constrain cooling in the very shallow crust (c. 1-2 km), which has hitherto been left poorly constrained by AFT analysis. Robust thermal histories were obtained by jointly inverting the analytical data using a Bayesian transdimensional Markov Chain Monte Carlo (MCMC) approach described by Gallagher (2012). We demonstrate that these cooling episodes relate to crustal denudation and a multi-stage development of the continental margin is proposed that requires significant (km scale) differential displacement on onshore brittle structures well inland (100s of kms) of the present coastline.

2. GEOLOGY

2.1 Tectonic setting of the Namaqualand basement rocks

Rocks of the Meso-Proterozoic Namaqua Metamorphic Province dominate the geology of the study area (Fig. 2). These basement rocks are comprised mainly of granitic gneisses and granite bodies emplaced during multiple periods of orogeny and terrane accumulation (Tankard et al., 1982; Groenewald et al., 1991; Jacobs et al., 1993; Cornell et al., 2006). The Namaqua Metamorphic Province therefore records a complicated history of magma emplacement and metamorphism that has produced a variety of supracrustal and intrusive rock types (Thomas et al., 1994; Eglington, 2006; Voordouw and Rajesh, 2012). A detailed description of these rock types can be found in Eglington (2006).

What is more pertinent in the context of this study is the tectonic fabric that was established at this time, as this basement structure may have been important in controlling the style of pre-, syn- and post-rift deformation (e.g. Daly et al., 1989; Clemson et al., 1997; Ziegler and Cloetingh, 2004; Corti et al., 2007; Frimmel et al., 2013). At least four compressional deformation events have been documented in gneissic foliations and folds and are thought to have been driven by processes coeval with the regional Kibaran Orogeny (c. 1200 – 1000 Ma) and global Grenville Orogeny (c. 1300 – 950 Ma) (Jacobs et al., 1993; Clifford et al., 2004; Eglington, 2006; Viola et al., 2012; Colliston et al., 2014). These deformation events were followed by a phase of extension in western Namaqualand that has been attributed to the collapse of the Namaqualand orogenic belt (Dewey et al., 2006). Dykes, folds and lineations ascribed to these deformation events typically strike NW-SE or NE-SW and occasionally E-W (Cornell et al., 2006). The main phase of metamorphism of the Namaqua Metamorphic Province was largely granulite-facies metamorphism (T: 800–860 °C and P: 5–6 kbar/15–18 km) and was completed by 1000 – 800 Ma (Waters, 1989; Eglington, 2006).

The Vanrhynsdorp Group unconformably overlies the metamorphic basement and was deposited during the latter stages of the Pan-African orogeny (c. 700 – 500 Ma) in a foreland

basin setting and subsequently experienced very low-grade metamorphism (Gresse and Germs, 1993). These rocks crop out in the south of the study area, close to the town of Bitterfontein, in half-graben structures trending N-S as gently folded, fault parallel synclines that overlie an easterly dipping unconformity across the basement rocks (Gresse et al., 2006; Macey et al., 2011) (Fig. 2). The influence of the pan-African Orogeny is observed in the Namaqua Metamorphic Province as brittle overprint structures (Viola et al., 2012). Cambrian to Early Carboniferous rocks of the Cape Supergroup only crop out in the very southeastern part of our study area and overlie the Vanrhynsdorp group. The Cape Supergroup, which dominates the geology south of our study area, was deformed during Permian to Triassic compression to form the Cape Fold Belt (Newton et al., 2009).

2.2 The Karoo Supergroup

The sedimentary rocks of the Karoo Supergroup were deposited in a sub-continental foreland basin, north of the developing Cape Fold Belt, during the Carboniferous to Triassic period (Tankard et al., 2009). The Karoo basin covers an area of approximately $7 \times 10^5 \text{ km}^2$ across central South Africa (Fig. 2). The Karoo Supergroup rocks that crop out within the study area include the lowermost Permian Dwyka and Ecca Group rocks (Johnson et al., 2006), which are intruded by Triassic-Jurassic Karoo dolerite sills and dykes. Dwyka glacial diamictites were unconformably deposited on the metamorphic basement and Vanrhynsdorp units during Late Carboniferous to Early Permian glaciations (Visser, 1990). The thickness and lithofacies of the Dwyka Group are highly variable and are attributed to erosion of an irregular pre-Karoo relief (Visser, 1983, 1987, 1989; Johnson et al., 2006). Conformably overlying the Dwyka Group glacial deposits is the Permian Eccca Group, consisting of shale units that are thinly laminated and fossiliferous (Johnson et al., 2006). These rocks were deposited in a shallow-marine environment and are the dominant sedimentary unit in the northwest Karoo Basin. The presence of Karoo Supergroup sedimentary rocks are important as they represent the last time the South African interior was unequivocally below sea level (Summerfield, 1985). The intrusion of Karoo dolerite sills and dykes was coeval with the eruption of thick continental flood basalts ($> c. 1.5 - 2 \text{ km}$) and marks the termination of Karoo deposition and

the onset of continental rifting in eastern Gondwana (c. 180 ± 5 Ma; Duncan et al., 1997; Jourdan et al., 2005, 2007; Moulin et al., 2011; Svensen et al., 2012). Karoo dolerite intrusions are widely preserved in the eastern part of the study area as prominent isolated summits above the flat lying plateau.

2.3 Syn and post-rift intrusive activity

Post-karoo igneous rocks, coeval with Atlantic rifting (c. 160 – 130 Ma), can be found in the most southern region of the study area (Fig. 2). The Koegel Fontein complex near the towns of Kotzesrus and Biesiesfontein is comprised of a variety of intrusive bodies including tholeiitic basalt plugs, syenite and granite plutons and dykes of varying composition (de Beer et al., 2002; Curtis et al., 2011). A suite of NNW-SSE trending dolerite dykes cross cut the Koegel Fontein complex and is believed to have the same origin as the False Bay dolerite suite in the Southern Cape based on their age and composition (Day, 1987; Reid, 1990; Reid et al., 1991; Trumbull et al., 2007; de Beer, 2010).

Early Cenozoic intrusions are commonly olivine melilitite plugs and alkali and carbonatite dykes. The Biesiesfontein plugs intrude into the Reitpoort Granite and have been dated at c. 55 Ma (Moore and Verwoerd, 1985). The Sandkopsdrif Complex north of Rietpoort is made up of numerous alkaline and carbonitite dykes (de Beer, 2010; Curtis et al., 2013) and is thought to post-date olivine melilitites in Biesiesfontein (Moore and Verwoerd, 1985). An important observation that can be made from these intrusions is their coherent trend along a N to NNW striking fracture zone, suggesting a link between reactivation of major structures during the Early Cretaceous and the Early Cenozoic creating pathways for intrusive rock (e.g. Moore et al., 2008; Jelsma et al., 2009).

The Gamoep intrusive suite (Fig. 2), consisting of olivine melilitite plugs and possible kimberlite intrusions, is found inland on the interior plateau north of Banke farm close to the Vaalputs nuclear repository site. The intrusions have been radiometrically dated at 54 – 77 Ma (Moore and Verwoerd, 1985) while terrestrial fossil and palynological evidence

discovered in the sediment infill confirm a Late Cretaceous to Eocene age for these intrusions (Haughton, 1931; Estes, 1977; Scholtz, 1985; de Wit et al., 1992).

2.4 Late Cretaceous and Tertiary deposits

The preservation of crater-lake sedimentary successions over intrusive bodies is highly significant as they provide insights into the margin's post-rift geological history, a record of which is otherwise absent. Up to 260 m of carbonaceous shales and tuff bands, commonly topped with conglomerates and arenites, are preserved within the diatremes of Late Cretaceous – Early Cenozoic intrusions (Moore and Verwoerd, 1985). These sediments are interpreted as lacustrine deposits and imply limited erosion has taken place since the emplacement of the igneous intrusive rocks (Cornelissen and Verwoerd, 1975; Moore and Verwoerd, 1985; Hanson et al; 2009; Stanley et al., 2013). Additional post-rift sedimentary units include silicified kaolinitised conglomerates and coarse grained, cross-bedded sandstones of the Dasdap and Vaalputs formation, which form prominent mesas on the plateau near Banke Farm, south of Vaalputs (Brandt et al., 2003, 2005) (Fig. 2). Clast compositions within the conglomerate horizons and the occurrence of widespread cross laminations in the Dasdap sediments suggest that these sediments were deposited in a fluvial environment and derived from the surrounding basement and Dwyka Group rocks (Brandt et al., 2003; 2005). The volcanic pipes in the region discussed above are overlain by these units in places and therefore provide an upper age limit of c. 67 Ma for the Dasdap sequence (Brandt et al., 2003, 2005; Viola et al., 2012).

The Vaalputs formation and adjacent Santab-se-Vloer Basin overlie the Dasdap sequence and are intersected and bounded by prominent NNW – SSW trending faults. The orientation of these structures is in keeping with the regional NNW – SSE structural trend (Viola et al., 2012). Although these sediments are severely lacking in age constraints, their similarity to nearby crater-lake deposits and their relationship with Late Cretaceous igneous intrusions suggests a Late Cretaceous-Early Cenozoic age is possible.

Cenozoic deposits along the coastal plain are typically preserved in the form of elevated marine terraces, semi- and unconsolidated aeolian sand, and as channel fill sediments preserved in valleys within the high-relief escarpment zone. The coastal marine terraces are best represented by the Alexander Bay Formation, which records Miocene and younger cycles of erosion and weathering attributed to sea-level fluctuations driven by both tectonic and eustatic processes (Roberts et al., 2006). Three distinct packages have been defined within the Alexander Bay Formation at 90 m, 50 m and 30 m above sea level. The de Toren Formation is present on the Koegel Fontein intrusive complex and on the periphery of the Namaqualand Highlands at Quaggafontein farm. The sediments are proposed to have been deposited in a similar environment to the Dasdap Formation and have a similarly uncertain age, which is tentatively placed in the Oligocene by de Beer (2010, 2012).

3. APATITE FISSION TRACK AND (U-Th-Sm)/He ANALYSIS

3.1 Apatite Fission Track Results

Apatite fission track analysis was performed on all 42 samples using the external detector method (Hurford and Green, 1982) (Table 1; SI-1). Central AFT ages range from 58.3 ± 2.6 to 132.2 ± 3.6 Ma and do not show a correlation with elevation. Indeed, some of the youngest measured AFT ages can be found at the highest elevations (Fig. 3a). Mean horizontal confined track lengths (MTL) range from 10.9 ± 0.19 to 14.35 ± 0.22 μm with the majority being greater than 13 μm ; the standard deviation of length measurements range from 0.97 to 2.50. In cases where MTLs are relatively short or length distributions are relatively broad, the shape of the distribution is either normal around the mean value or is negatively skewed with a large proportion of longer tracks and tail of shorter tracks (see SI-2 for all Track Length Distribution plots).

Radial plots graphically illustrating single-grain dispersion can be found in SI-3. A chi-squared (χ^2) age homogeneity test was performed on all samples to assess whether the measured single-grain ages represent a single population (Galbraith, 2010). Nine samples

yielded $P(\chi^2)$ equal to or less than 0.05. Despite failing the chi-squared test, a mixture model (Galbraith and Green, 1990; SI-3) does not define a clear distinction of two or more populations. For the majority of samples, single-grain FT age dispersion is moderate to low (i.e. $\leq 20\%$). Samples with higher levels of dispersion are typically associated with samples with low track densities. It is likely that every sample is best approximated by a single population, and any apparent dispersion in single-grain ages is predominantly induced by inter-grain variations in compositionally controlled track annealing rates.

The classic ‘boomerang’ relationship (e.g. Green, 1986; Gallagher and Brown, 1997) is not seen for this dataset, as almost all MTLs are reasonably long and similar in value (i.e. $>13\ \mu\text{m}$) and because there is a fairly narrow age range for most samples (c. 80 – 120 Ma) (Fig. 3b). As there are no clear peaks in the Boomerang plot, it is not possible to define discrete thermal events, and it is likely that most samples resided at elevated temperatures and were entirely, or almost entirely, reset prior to the Cretaceous. However, several samples have short ($< 13\ \mu\text{m}$) MTL and young AFT ages (c. $< 80\ \text{Ma}$), implying that these samples may have resided at temperatures within the PAZ following an initial cooling event and then either cooled during a distinct cooling event or by slow, protracted cooling.

Dpar values range from 1.45 to 3.42 μm with most values (76% of samples) falling between 1.7 and 2.3 μm . While many samples have comparable Dpar measurements with Durango apatite ($2.05 \pm 0.16\ \mu\text{m}$; Sobel and Seward, 2010), many fall out with the Durango range. This variation means that some samples have compositionally controlled annealing properties different from Durango standard. Any significant compositional control over annealing is treated appropriately during thermal-history inversion.

3.2 Apatite (U-Th-Sm)/He Results

Multiple single-grain (U-Th-Sm)/He analyses were performed on a subset of 16 samples that were chosen based on the high quality of the apatite separates and optimal geographical location for the study (Table 2). Between 5 and 21 single-grain ages were determined for

each sample in an attempt to quantify and utilise the inter sample-grain date dispersion (see below). Details on analytical techniques can be found in SI-4. Mean AHe ages, uncorrected for alpha-ejection, range from 55.8 ± 31.34 to 120.6 ± 31.4 Ma and alpha-ejection corrected ages (after Farley et al., 1996) range from 74.0 ± 43.9 to 156.9 ± 40.9 Ma. Although “mean” AHe sample ages provide a useful frame of reference for comparison with AFT ages and the wider geological context, they do not necessarily correspond to a specific geological event at that time. In a similar manner to an AFT age, an AHe age is a product of a thermal history that is initially unknown. Depending on the form of this thermal history and the chemical and physical properties of individual apatite grains, mean AHe ages can be associated with large degrees of single-grain age dispersion. Dispersion (standard deviation of age/mean age) is large for samples in our data set; ranging from 12 – 56%. As thermal diffusion of ^4He dominates at elevated temperatures (Brown et al., 2013), the style of thermal history experienced by a sample will almost always lead to an overcorrection of the ^4He age using a single alpha-correction factor (Farley et al., 1996; Meesters and Dunai, 2002 a, b; Herman et al., 2007; Gautheron et al., 2012). This effect may be a contributing cause of some of the corrected AHe ages being older than their corresponding AFT age (Fig. 3c) and the magnitude of single-grain age dispersion (Fig. 4; SI-5). However, as the interaction between alpha-ejection, radiation damage effects and He diffusion remain poorly understood, we have chosen to quote raw, uncorrected, AHe ages and we deal with the effect of alpha-ejection by accounting for ^4He lost through alpha-ejection during each time step of the modelling approach following the approach proposed by Meesters and Dunai (2002 a, b).

Variation in effective uranium (eU) content, grain size and whether or not crystals analysed are fragments of larger crystals have been shown to be key factors that control the measured AHe date for a particular grain (Table 3; Flowers et al., 2009; Gautheron et al., 2009; Reiners and Farley, 2001; Brown et al., 2013; Beucher et al., 2013). If a sample has experienced relatively slow cooling through the partial He retention zone, then positive correlations of spherical equivalent grain radius (R^*) and/or eU with AHe age may be evident (Reiners and Farley, 2001; Flowers et al., 2009). However, as demonstrated by Brown et al. (2013), the combined effects of these influences may decouple the expected correlation. Moreover, a

large number of grains analysed in this study were fragments of larger grains and had only 1 (1T) or no (0T) terminations and therefore simple 2-D correlations between eU and R^* and grain date will be disrupted (Fig. 4). Despite this, many samples do show an apparent positive age- R^* correlation (see SI-5 for all R^* and eU – AHe date plots). Sample NQ12-15 is the only sample for which we can observe a negative date- R^* correlation; although, this sample has the fewest number of grains and therefore the correlation is poorly constrained.

Plots of eU versus single-grain age are more complex, and different samples show correlations that are positive, negative or entirely absent. However, a simple linear correlation between eU and grain age should be not expected even for ‘well’ behaved samples, because the effect of radiation damage accumulation and annealing is not linear, and it is dependent on the thermal history. For example, Flowers et al. (2009) report that for relatively slow cooling rates (c. 0.1 – 1 °C/Myr), the age-eU relationship will first be positively correlated but will then plateau above eU concentrations of c. 50 ppm. Our understanding of how the accumulation and annealing of radiation defects occurs, and the impact this has on alpha-trapping and helium diffusion, is still developing (Gautheron et al., 2013; Mbongo-Djimbi et al., 2015). Expecting to observe simple correlations in single-grain ages and eU may also be ambitious considering that grain radius and the presence of fragmented grains influence the effective closure temperature and preservation of the He diffusion profile within grains, respectively. However, we emphasise that the observed inter sample grain-age dispersion is likely a real and natural effect that contains useful information about the sample’s thermal history irrespective of whether the simple 2D plots of grain-age versus eU and R^* show clear correlations or not (Flowers and Kelley, 2011; Brown et al., 2013).

Other factors that can cause single-grain age dispersion (Table 3) must also be considered though, such as the possible presence of U and Th zonation (Flowers and Kelley, 2011; Ault and Flowers, 2012; Farley et al., 2011), unidentified U and Th bearing inclusions (House et al., 1997; Stockli et al., 2000; Fitzgerald et al., 2006; Vermeesch et al., 2007), implantation (Farley et al., 1996; Spiegel et al., 2009; Gautheron et al., 2012), the contribution of Sm

(Hansen and Reiners, 2006; Vermeesch, 2008) and the dominant influence of the thermal history of the sample in controlling He diffusion.

3.3 Summary of joint AFT and AHe data

The AFT data presented here include AFT ages that range from the Early Cretaceous to Early Cenozoic with moderate to long MTLs (c. 13 μm) and track-length distributions that are generally narrow to moderately broad (Std. Dev. 0.97 to 1.95 μm). The qualitative relationship of the track-length and AFT age data indicates that the dominant phase of crustal cooling occurred throughout the Cretaceous. However, only by using inverse modelling can we resolve the temporal and spatial variability of crustal cooling across the study area.

In general, the AHe ages are in agreement with AFT ages in that they are predominantly Early Cretaceous to Early Cenozoic. It is not uncommon to observe AHe ages equal to or exceeding their corresponding AFT ages (Hendriks and Redfield, 2005; Fitzgerald et al., 2006; Green and Duddy, 2006; Flowers and Kelley, 2011; Danišík et al., 2012; Ksienzyk et al., 2014) for reasons discussed above. Only three samples have mean AHe ages that are older and do not overlap within the uncertainty of the mean AFT age (Fig. 3c).

When comparing mean sample AHe ages with mean (or central or pooled) sample AFT ages, it is not clear what meaning to ascribe to any differences in these values because of the wide range of factors that affect single-grain ages in both systems. For many cases, where a sample has experienced a protracted low-temperature thermal history, it would be expected that some single grains yield AFT ages older than AHe and also the reverse, depending on the relative eU, grain size and Cl composition of the different grains. Large, U-, Th- and F-rich grains may yield 'old' AHe ages and 'young' AFT ages whereas small, U-poor, Th-poor but Cl-rich grains may yield 'young' AHe ages and 'old' AFT ages. For this reason, we present and examine the single-grain date distributions of both methods to try and avoid problems that may arise from interpreting sample mean ages directly (Fig. 4b; SI-6).

If a sample yields concordant mean AHe and mean AFT ages, this is often interpreted as indicative of a sample cooling rapidly through both the partial annealing zone (PAZ) and partial retention zone (PRZ) at a time indicated by the measured ages. However, because AFT and AHe analyses are not performed on the same grains, and the thermal sensitivity of each grain is different due to different grain sizes and compositions, it does not necessarily follow that concordant mean sample ages indicates rapid cooling at that time. For the same reasons, samples having AFT and AHe data that appear at first complex, or even incompatible, may well contain coherent, useful thermal-history information once the real uncertainties on each measurement are formally acknowledged and accounted for when modelling the data.

4. THERMAL-HISTORY INVERSION

4.1 Approach

In this study, QTQt, a Bayesian transdimensional approach to data inversion, as described in detail by Gallagher (2012), is used for the inference of thermal-history models. Fission-track ages and track-length data are modelled using the multi-kinetic fission track annealing model of Ketcham et al. (2007). For samples that also have AHe data, all single-grain ages are jointly inverted alongside the corresponding fission-track data using the model and parameters of Gautheron et al., (2009), which accounts for the effect of radiation damage accumulation and its annealing on He diffusion.

The Bayesian method requires us to define the model space as a prior probability distribution (i.e. range of temperature and time over which to search). The model space is then sampled using a MCMC approach (e.g. Gilks, 2005; Sambridge et al., 2006; Gallagher et al., 2009), whereby the current model is perturbed to produce a proposed model. This model is then accepted or rejected based on a combination of the data fit (likelihood) of the proposed model and the prior probability, relative to the same measures for the current model. The transdimensional Bayesian aspect of the modelling technique penalises complex models proposed during sampling in favour of models with fewer T-t points that adequately fit the

observed data. An additional novel aspect of QTQt is the ability to model samples from vertical profiles together. This function is also employed below for a subset of samples that form an “elevation profile” over a relatively short distance of c. 10 km. Such a multi-sample profile inversion also incorporates an additional model parameter representing the temperature difference or offset between the top and bottom sample, with a prior determined by an assumed range for the geothermal gradient.

The MCMC algorithm was run for a minimum of c. 200,000 iterations after discarding an initial c. 50,000 “burn in” runs (see Gallagher et al., 2009). In most cases, however, the number of runs far exceeds this as short c. 10,000 iteration runs are performed in order to choose appropriate values for the MCMC search parameters before performing longer runs. This choice is based on the acceptance rates of proposed model parameters (being between 20 and 60%), and the parameter sampling being stationary (i.e. no obvious trends as a function of iteration). The output of the Bayesian approach is an ensemble of accepted thermal history models, each with an associated posterior probability. From this collection of thermal histories, a mean thermal-history model (weighted by the posterior probability of each individual thermal history), termed the expected model, is produced with 95% credible intervals that provide the uncertainty on the model. The nature of the expected model is that it will retain well constrained features (i.e. features common to many individual models) while more complex deviations observed in only a small number of viable models are averaged out. The expected model, and its associated uncertainty measures, provides the most robust albeit conservative insight to the overall thermal history of the sample.

4.2 Additional model constraints

For each sample, the oldest sample age (i.e. the AFT age or oldest AHe single-grain age for the sample) is used to define the centre of a uniform prior on time, with a range of $\pm$ oldest sample age. The prior information on temperature range was centred on 70 °C with a range of $\pm$ 70 °C. The number of time-temperature points sampled from this prior is allowed to vary from 2 (including the present data temperature) to 50. The thermal history is constructed by

interpolating linearly between these sampled time-temperature points. This sampling strategy imposes no constraints about the form of the thermal history (e.g. monotonic cooling) but the form is determined by the data themselves. Additional prior information, if available from geological constraints, can be entered in a similar manner as a range of time and temperature. All samples from this region are outcrop samples taken from basement lithologies apart from SA12-27, which was collected from a boulder clast of basement origin from within the glacial tillite of the Permian Dwyka Group. During modelling, SA12-27 was assigned a specific initial constraint of 300 ± 10 Ma, 5 ± 5 °C to represent that the sample was at the surface at this time. For basement samples, an initial constraint of 550 ± 50 Ma and 100 ± 100 °C was used to reflect the uncertainty surrounding the palaeo-temperature of the sample during the Pan-African Orogeny. For each model, only the post-300 Ma thermal history is presented. Before 300 Ma, thermal history models are unconstrained because the data requires that all samples temperatures close to or above the base of the apatite PAZ (i.e. c. 110 ± 10 °C) prior to cooling in the Mesozoic. The present-day temperature value is assumed to be 20 ± 10 °C for all samples.

A lack of geological information limits the amount of post-rift constraints that can be added to the models. However, the implications of sporadic fossil evidence, sedimentary deposits and igneous intrusions across the region on nearby samples are also investigated. The thermal history for the cluster of samples collected at Hondeklip Bay can be constrained using fossil evidence in thin sedimentary formations that overly the Namaqua Metamorphic Province rocks across the coastal plain. Fossil hominoid teeth from the marine sand Alexander Bay formation assign an early Miocene age to the deposit (Senut et al., 1997). The Alexander Bay formation overlies the calcified and silicified gravels and sandstones of the Koignass Formation, which is presumed to have been deposited in a fluvial environment during a Late Oligocene to Early Miocene marine regression (de Beer, 2010). Coastal samples are therefore assigned an additional constraint of 20 ± 10 Ma, 20 ± 10 °C to reflect near surface temperatures of the basement rock prior to deposition of these Cenozoic deposits.

Sediments in and around the Vaalputs area and fossiliferous crater-lake sediments preserved in the Gamoep intrusive suite, and in particular the Arnot pipe at Banke farm (Fig. 1 and 2), provide Late Mesozoic and Cenozoic constraints for samples on the plateau. The Dasdap and Vaalputs sediments are proposed to be Late Cretaceous to Early Palaeocene in age. This age is supported by palynological evidence dated at 71 to 64 Ma (Scholtz, 1985) and preservation of fossilised frogs and wood in clay deposits (Adamson, 1931; Haughton, 1931). Estes (1977) reported younger pollen and fossil leaf evidence from the same lacustrine sediments aged from the late Eocene to early Oligocene. A constraint box of 50 ± 20 Ma, 20 ± 10 °C was included for samples SA12-51, SA12-52 and SA12-27.

4.3 Results

4.3.1 Early Cretaceous cooling histories

A signal of Early Cretaceous cooling is evident in many samples across the study area. Figure 5 shows representative thermal histories of this form (see SI-7 for all thermal history models). The path of the expected thermal history indicates cooling through the PAZ beginning at c. 150 – 130 Ma and reaches near surface temperatures by c. 50 – 30 Ma (Fig 5a, b). Due to the unconstrained initial conditions of the model, it is unclear whether the sample was already undergoing cooling or still being heated prior to the cooling episode predicted by the model. Cooling rates for the main phase of cooling predicted by these thermal histories is c. 0.5 – 2.5 °C/Myr, over a period of c. 50 Myr (150 – 100 Myr).

More rapid cooling during the early Cretaceous (c. 150 – 130 Ma) time interval is rarely observed. However, in occasional samples (e.g. JN2, Fig. 5c), cooling rates of c. 4.5 – 5.5 °C/Myr over a 10 Myr period are predicted. Following this period of rapid cooling, the cooling rate rapidly decreases to c. 0.2 – 0.3 °C/Myr during the Late Cretaceous and is maintained through the Cenozoic. These models imply that crustal cooling persisted throughout the early Cretaceous across the entire margin, predominantly at a moderate and

steady rate; although, local variations exist. A consistent observation is that near surface temperatures ($\leq 40\text{ }^{\circ}\text{C}$) were reached and maintained by the Early Cenozoic.

4.3.2 *Late Cretaceous cooling histories*

Cooling initiating during the mid-Late Cretaceous (c. 110 – 90 Ma) time interval, characterised by rapid cooling rates, is the dominant style of thermal history across the margin (e.g. Fig. 6a, b, see SI-7). Cooling rates generally range from c. 1.5 – 5 $^{\circ}\text{C}/\text{Myr}$ but in certain samples can be higher at 7.4 – 8.1 $^{\circ}\text{C}/\text{Myr}$ (e.g. NQ12-04). Following rapid cooling during the Late Cretaceous, samples are predicted to reside at temperatures $\leq 40\text{ }^{\circ}\text{C}$ and cool steadily through the Cenozoic towards present-day surface temperatures. As these samples cooled from temperatures of c. $110\pm 10\text{ }^{\circ}\text{C}$ during the Late Cretaceous, any record of earlier cooling (i.e. the early Cretaceous cooling history described above) is not preserved. The absence of a Late Cretaceous pulse of cooling in ‘early Cretaceous cooling histories’ could be due to a lack of information in the data; meaning that the approach adopted by QTQt does not need, or cannot resolve a two-stage cooling history from the available data. However, we interpret the monotonic Early Cretaceous thermal histories as a real feature with geological significance, as it is the most probable cooling path given the data available, and any alternative would involve forcing the model to satisfy a preferred thermal history.

A second Late Cretaceous cooling episode is poorly preserved but is present in the study area. The thermal history for one sample clearly shows cooling initiating at between 70 and 65 Ma (Fig. 6c). The style of cooling for NQ12-23 is very rapid, occurring at a rate of 10 $^{\circ}\text{C}/\text{Myr}$. This sample resides in a complex structural setting and is c. 50 km SW of the Koegel Fontein igneous complex, which has intrusive bodies dated to c. 70 – 50 Ma (de Beer, 2002). Fission-track data for this sample are robust with a large number of track lengths and single-grain ages. The regional significance of this cooling history cannot be inferred from this sample alone and is therefore discussed in more detail below alongside other regional AFT data.

4.3.3 *Multi-sample inversion*

A subset of samples, comprised of 8 outcrop samples from the western flank of the “Kamiesberg Mountain”, including two samples from the dataset of Wildman et al. (2015), cover an elevation range of 1220 m and are separated by a distance of less than 10 km (N to S) and 5 km (E to W) (Fig. 7). Data from these samples are treated as an “elevation profile” and are modelled together as described in section 4.1. Contrary to the expected relationship of AFT age and elevation, the oldest AFT ages in this profile are at the base. As expected, these samples have the poorest data fit when all samples in the profile are modelled together (Fig. 7a). To resolve this issue, the profile is better split into (i) a lower profile consisting of three samples with AFT ages of c. 115 Ma (Fig. 7b) and (ii) an upper profile consisting of five AFT ages that are c. 90 Ma (Fig. 7c). A rapidly cooled, tightly constrained cooling event at c. 110 – 90 Ma is defined for the upper profile while a slower cooling history initiating at c. 150 – 130 Ma is defined for the lower profile. As the age-elevation relationship of the entire profile is inconsistent with that of a profile sampled from a single coherent block and the data are better reproduced by two distinct thermal histories, it is possible that displacement on an unmapped fault has resulted in the lower profile (i.e. down-thrown block) retaining thermal-history information from the syn-rift period while the upper profile (i.e. up-thrown block) has cooled during later erosion.

It is difficult to reproduce all of the single-grain AHe ages for samples in the elevation profile regardless of whether the data are modelled as a single profile or two profiles. In fact, for most models with AHe data, it is predominantly the younger AHe single-grain ages that are predicted best. The poor fit of the AHe data relative to that of AFT data is likely to be due to a combination of our limited understanding of the He system and the treatment of the data in the modelling approach to compensate for this. Although we have attempted to account for both grain size and radiation damage effects, the latter is still not ideally parameterised, particularly for high eU samples and potentially for additional compositional heterogeneity (Gautheron et al., 2013; Mbongo-Djimbi et al., 2015). Moreover, the effect of other factors that could produce observed AHe ages that were ‘too old’ (e.g. grain fragmentation, zonation, implantation, etc.) are unaccounted for and may, in the future, improve data fits. For this

reason, less weight was placed on AHe ages, by assigning an uncertainty to AHe single-grain ages greater than the analytical uncertainty and allowing the AHe single-grain age to be resampled from this uncertainty range.

5. TIMING AND MECHANISMS OF CRUSTAL COOLING

5.1 AFT and AHe data from the SW African continental margin and interior plateau

Aside from new AFT data presented in this study and previously published by the same authors (e.g. Brown et al., 1990; Wildman et al., 2015), AFT analysis in the Namaqualand region has been limited (Fig. 8). Data from Kounov et al. (2009) consists of two transects across the Namaqualand continental margin and extend onto the inland plateau (Fig. 8). Combining all available datasets provides an extremely detailed low-temperature thermochronology database across this sector of the margin by improving the regional and local sample coverage as well as improving the temporal resolution by including AHe analysis.

Data from earlier studies is largely consistent with the new data presented here. AFT ages are generally Early – Late Cretaceous with moderate to long MTLs. The exceptions to this are one sample of Late Jurassic age reported by Wildman et al. (2015) (8732-46) in the heart of the Namaqualand Highlands and an Early Jurassic age recorded by Kounov et al. (2009) from their southern transect near to the town Calvinia. Kounov et al. (2009) propose that inverse modelling of their data supports a two stage thermal history across the margin. The major phase of cooling is ascribed to the mid-Cretaceous (115 – 90 Ma) and is attributed to a tectonically induced period of enhanced denudation. Moreover, they suggest that discrete fault block reactivation during this time results in differential denudation over major structures. The earlier event was driven by either rift-related tectonic denudation or thermal relaxation of the surface following widespread Karoo magmatism at c. 180 Ma (Jourdan et al., 2005). Kounov et al. (2009) give preference to the latter mechanism, as this cooling

episode is only recorded in samples from the plateau region of the southern transect that were collected in the Karoo basin. However, data from Brown et al. (1990), Wildman et al. (2015) and data presented here clearly preserved a record of Early Cretaceous cooling across the coastal plain, escarpment zone and plateau. Moreover, offshore sediment-volume analysis has predicted syn-rift sediment accumulation in offshore basins at this time (Guillocheau et al., 2012; see section 7.1). Although a thermal effect of the Karoo may have contributed to the recorded thermal history, an erosional response to rifting is likely to have been a major factor.

The latest Jurassic – Early Cretaceous cooling at c. 160 – 130 Ma is recorded regionally but preserved only locally. The discrete, structurally controlled nature of the second cooling event from 115 – 90 Ma across the margin is somewhat poorly constrained by Kounov et al. (2009), due to the uncertainty in their fission-track data and uncertainties in models. The distinction of these two cooling episode is much more prominent in our study through the combination of AFT and AHe analysis and high resolution sampling.

Evidence for a 110 – 90 Ma cooling event spanning the entire sub-continent of South Africa can be found in (i) AFT data from the southwestern cape (Wildman et al., 2015), southern margin (Tinker et al., 2008a; Green et al., 2015), southeastern margin (Brown et al., 2002) and along the southern Orange River valley (de Wit, 1988); (ii) AHe data from the southeastern Kaapvaal Craton (Flowers and Schoene, 2010) and from samples collected from kimberlites on the south African plateau (Stanley et al., 2013); and (iii) from a joint AFT and AHe study from the Augrabies Falls and Fish River Canyon along the Orange River (Kounov et al., 2013) (Fig. 8).

AFT or AHe data advocating a Late Cretaceous – Early Cenozoic (or even younger) cooling event is still limited despite the growing number of studies occurring in single outcrop samples, if at all (e.g. de Wit, 1988; Kounov et al., 2013). However, in northern Namibia, many AFT ages of c. 60 – 80 Ma are widely observed on the coastal plain as well as penetrating well inland (Raab et al., 2002; Brown et al., 2014). These AFT ages agree with the single Early Cenozoic AFT age presented here, implying that structurally controlled

denudation in Namibia may have also been manifest in parts of Southern Africa. AFT data is used by Green et al. (2015) to infer that parts of the southern margin, particularly the southwestern cape, have cooled from c. 60 – 70 °C since 30 Myr. This cooling equates to c. 1.5 to 2.5 km of denudation for a geothermal gradient of 20 to 25 °C/km, which is greater than the estimates derived from other data sets from the region (c. 1 – 1.5 km; Tinker et al., 2008a; Stanley et al., 2013; Wildman et al., 2015). Although local variations in the amount of Cenozoic denudation may exist, Green et al. (2015) utilise a different modelling approach and geological framework for interpreting their data (e.g. Green et al., 2013). While our Bayesian approach yields simple cooling histories that adequately fit the data without invoking unconstrained geological events, Green et al. (2015) use likelihood theory (Gallagher, 1995) and a modelling procedure that explicitly seeks discrete temperature maxima and thus invariably produces thermal histories with multiple discrete episodes of inferred burial and erosion.

5.2 Timing and spatial patterns of denudation

The last major thermal event expressed regionally across Southern Africa is the emplacement of the Karoo igneous intrusions and continental flood basalts at c. 182 – 183 Ma (Svensen et al., 2012). Evidence of younger igneous bodies in Southern Africa is found in the form of localised alkaline intrusive pipes emplaced during the Late Cretaceous and Early Cenozoic (Moore and Verwoerd, 1985; de Beer, 2010; Curtis et al., 2013). In Namibia, early Cretaceous flood basalts are preserved as part of the Parana-Etendeka igneous province dated at c. 130 Ma (Reid et al., 1990; Reid and Rex, 1994). However, due to the lack of any preservation in South Africa their southern extent is unknown. The lack of significant thermal activity following the Jurassic combined with the coherence of the timing of cooling with peaks in offshore sediment accumulation in the offshore basin (Rouby et al., 2009; Guillocheau et al., 2012; Wildman et al., 2015) suggests that cooling was primarily driven by onshore denudation.

We performed simple calculations of the amount of denudation required to drive cooling over distinct time intervals: $D_{t_0-t_1} = (\Delta T_{t_0-t_1})/G$, where D is the estimate of denudation over the desired time interval (t_0-t_1) and G is the estimated geothermal gradient. The amount of denudation required to cool the sample from elevated temperatures to surface temperatures depends on the geothermal gradient of the region. Present-day geothermal gradients across much of Southern Africa are c. 20 – 25 °C/km (Ballard and Pollack, 1987; Jones, 1987). Maintaining the conservative approach adopted during modelling, the upper limit of this temperature range (i.e. 25 °C/km) was assumed to best represent the palaeogeothermal gradient. The influence of geothermal gradient on denudation estimates is non-linear and most significant for low geothermal gradients. A low geothermal gradient is unlikely to be the case in the study area as the basement rocks are particularly enriched in heat-producing radiogenic elements (U, Th, K) (Andreoli et al., 2006), which will have helped to maintain moderate-high heat flow values across the Namaqualand Highlands (Jones, 1987; Andreoli et al., 2006).

Estimates of denudation are made using the expected thermal-history model and assuming a geothermal gradient of 25 °C/km. Figure 9 shows three coast perpendicular transects highlighting the spatial patterns of denudation over major time intervals. The timing for the onset of cooling that is predicted by sample thermal histories are only recorded when the sample cools below c. 110 °C. Therefore, in cases where much of the early portion of the thermal history is unconstrained, cooling and the mechanism driving cooling may have been ongoing prior to the predicted onset of cooling. Estimates of denudation for a particular time interval should be consulted alongside the thermal history and assessed in terms of how well the thermal history is constrained at this time.

During the early Cretaceous (c. 150 – 110 Ma) total denudation is generally between 1 to 2 km and appears to be fairly consistent across the length of the transect. However, as many samples do not record cooling until the late Cretaceous, fewer samples can be used to estimate earlier denudation. Two samples along transect YY' predict estimates of c. 4 km of denudation in the early Cretaceous with c. 2 km occurring during the 150 – 130 Ma interval.

Localised zones of higher denudation at this time may have been a result of a variable pre-rift topography or localised structural offsets. The furthest sample inland, SA12-27, was collected from a basement clast within the Dwyka Group stratigraphic unit and is therefore constrained at the surface at c. 300 Ma. The thermal history predicts consistent heating to temperatures of c. 120 °C at c. 100 Ma, which would imply progressive burial until the late Cretaceous. This burial event could reflect internal drainage of an eroding rift flank resulting in inland deposition. However, what the maximum post-Dwyka palaeotemperature was for this sample and when that palaeotemperature was attained is uncertain, and therefore there is the possibility that burial was completed at almost any time between the stratigraphic age and c. 100 Ma (SI-7).

The most significant period of denudation occurred during the Late Cretaceous (c. 110 – 70 Ma) with a denudation thickness of c. 2 – 4 km, and in some cases close to 5 km, being predicted across all three transects. This thickness of denudation is not seen uniformly across the transects, with some sample thermal histories observing no inflection in their cooling history in the Late Cretaceous but preserving the early Cretaceous record of cooling. Although ‘early Cretaceous’ thermal histories do not record the onset of rapid Late Cretaceous cooling, these thermal history paths still imply that slow cooling, and denudation on the order of 1 km, continued throughout the Late Cretaceous across the entire margin. This pattern of denudation may have been driven by a combination of localised fault block uplift across major and minor faults combined with a regional uplift of the entire margin.

Denudation during the Cenozoic is predicted to be low in comparison to the Cretaceous; less than c. 1 km in most samples over this time. Exceptions to this appear to be predominantly found far onto the plateau (see thermal histories for NQ Plateau and SA12-27), where denudation in the early Cenozoic exceeds 1 – 2 km from c. 70 – 30 Ma. Over the last 30 Ma, denudation is predicted to be less than 0.5 km. Low amounts of Cenozoic denudation are supported by cosmogenic nuclide studies in southwestern Africa (Kounov et al., 2007), Namibia (Cockburn et al., 2000; van der Wateren and Dunai, 2001; Bierman and Caffee, 2001; Codilean et al., 2008, 2014) and the southern cape (Decker et al., 2011; Erlanger et al.,

2012; Scharf et al., 2013; Bierman et al., 2014; Kounov et al., 2015), which all report a decrease in denudation rate by an order of magnitude during the Cenozoic relative to the Cretaceous.

7. DISCUSSION

7.1 Sediment accumulation in the Orange Basin

The catchment of the present-day Orange River covers an area of c. 1×10^6 km² and transports material eroded from the continent to the offshore Orange Basin. It is suggested that, since break-up, the main drainage outlet to the Orange Basin was located in a position similar to that of the present-day Orange River, Buffels River or Olifants River (e.g. Partridge and Maud, 1987; Dollar, 1998; de Wit, 1999; de Wit et al., 2000; Goudie, 2005, Stevenson and McMillan, 2004; Fig. 8). Therefore, most of the sediment derived from erosion of the southwestern continental margin and interior hinterland following break-up has been transported to the Orange Basin via the proto-Orange River and smaller rivers along the coast (e.g. Gilchrist et al., 1994; Kooi and Beaumont, 1994; van der Beek et al., 2002; de Wit et al., 2000; Goudie, 2005). Due to the long-term stability of the Orange River catchment, the record of sediment accumulation preserved in the Orange Basin provides independent constraints on the timing of onshore erosion (e.g. Rouby et al., 2009; Guillocheau et al., 2012).

Previous studies have provided detailed descriptions of the Orange Basin stratigraphy (Muntingh and Brown, 1993; Brown et al., 1995; McMillan, 2003; Paton et al., 2008; Kuhlmann et al., 2010; Wildman et al., 2015) and chronology of sediment accumulation over the last c. 150 Myr (Rust and Summerfield, 1990; Rouby et al., 2009; Guillocheau et al., 2012). Here we simply highlight the correlation between offshore accumulation in the Orange Basin as constrained primarily by the comprehensive investigation by Guillocheau et al., (2012) and the timing of denudation as inferred from our thermal history models. Two major peaks in accumulated sediment volume characterise the record of post-rift sediment influx to

the Orange Basin. The first peak records accumulated sediment volumes of c. $10 \times 10^{14} \text{ m}^3$ from 150 to 130 Ma (i.e. syn-rift) and the second peak, the maximum recorded sediment volume, of c. $11 \times 10^{14} \text{ m}^3$ is observed at c. 80 – 70 Ma. In addition to a clear peak in accumulated volume being defined during the Campanian, there is a well-defined period of increasing sediment volume preceding this peak that begins at c. 110 – 100 Ma. Another noticeable feature of the offshore sediment accumulation history is the minimal accumulation predicted throughout the Cenozoic ($5 \times 10^{14} \text{ m}^3$).

We reiterate and lend further support to the conclusions drawn from our previous AFT study on the southern Atlantic margin (Wildman et al., 2015) that denudation episodes inferred from the thermal histories derived from joint inversion of AFT and AHe data are consistent with the offshore accumulation history in the Orange Basin. Coherent features include an initial episode of denudation/sedimentation during the syn-rift (c. 165 – 130 Ma; Jackson et al., 2000; Eagles, 2007; Heine et al., 2013) period; an episode of post-rift denudation at c. 110 – 90 Ma, which coincides with increasing sediment volumes during the middle to Late Cretaceous; and minimal denudation throughout the Cenozoic, which is reflected in low-levels of accumulation from the end Cretaceous to the present day. The lag time between the onset of denudation in the mid-Cretaceous and the peak volume of sediment in the basin may be related to a disruption of internal drainage networks during the creation of fault-related relief, transient storage within the large proto-Orange catchment, or due to regional tilting of the African continent (e.g. Braun et al., 2014).

7.2 Evidence for post-rift faulting along the southwest African margin

The structural evolution of the Namaqualand sector of the South Atlantic continental margin is complex and poorly understood (Andreoli et al., 2009). Deformation associated with major pre-rift tectonic events defined the pre-existing structural grain in both Namaqualand and in

the Cape Fold Belt. These ancient fractures likely provided sites of pre-existing weakness in the lithosphere that controlled the style of fault propagation during rifting (e.g. Ziegler and Cloetingh, 2004; Autin et al., 2013; Corti et al., 2013; Will and Frimmel, 2013) and may have provided the location for post-rift deformation during later tectonic events (e.g. Sykes, 1978; Daly et al., 1989; Saintot et al., 2011; Viola et al., 2012).

By combining remote sensing techniques to identify major structural lineaments with a detailed field based analysis of brittle fault structures, Viola et al. (2012) suggest that a record of post-rift reactivation is preserved in the Namaqualand basement rocks. The earliest syn- or post-rift palaeo-stress tensor identified by Viola et al., (2012) is suggested to have been produced by sub-horizontal extension orientated NE-SW, causing reactivation of older faults during the Early Cretaceous. This phase of extension is assigned to the initiation of rifting and opening of the Atlantic due to the coherence of the orientation of extension and the perpendicular orientation of major Early Cretaceous dyke swarms (Reid and Rex, 1994; Will and Frimmel, 2013).

Post-rift palaeo-tensors identified by Viola et al., (2012) are suggested to represent alternating episodes of regional extension and compression throughout the middle to Late Cretaceous. However, the timing of these events is poorly constrained. Moreover, palaeo-stress analysis in NW Namibia suggests ENE-WSW and NE-SW extension prevailed during rifting and Late Cretaceous, respectively, and provides no clear evidence of compression (Salomon et al., 2014, 2015). In the Vaalputs region, inland of the Namaqualand Highlands, normal fault-slip structures extend into the silicified and kaolinitised weathering profiles of the Dasdap alluvial fan sediments (Brandt et al., 2003; 2005) and into the volcanic breccias preserved at the Gamoep melilite pipe (77 – 54 Ma; Phillips et al., 2000). Despite the limited robust information on the timing of faulting from these studies, they provide important field evidence that post-rift faulting has occurred. Thermochronology data reported here and in Kounov et al., (2009) suggest that this faulting may have begun during the mid-Cretaceous. The thermal history predictions made from the Kamiesberg profile (see section 4.3.3 above)

suggests that differential denudation may have occurred across prominent NNW-SSE lineaments dissecting the profile, as a result of mid-Cretaceous fault movement.

Within the offshore post-rift sedimentary successions, gravitational tectonics is described involving extension-driven slumping in the proximal margin and toe-thrusting in the outer-shelf (Brown et al., 1995; Paton et al., 2008; de Vera et al., 2010; Hirsch et al., 2010). The major zone of extension and compression has a basal detachment in the marine shales of the Turonian-Cenomanian boundary (Muntingh and Brown, 1993; Séranne and Anka, 2005; de Vera et al., 2010). The development of growth strata and stratigraphic relationships led de Vera et al. (2010) to conclude that gravitational tectonics were periodically active in short-lived episodes. The regional mechanisms that trigger movement on these faults are still uncertain, but have been related to the influx of high volumes of sediment to the basin (Jungslager, 1999; Paton et al., 2007; Kuhlmann et al., 2010), basin inversion and the development of enhanced structural relief; or margin uplift (Séranne and Anka, 2005; Paton et al., 2008; de Vera et al., 2010). The timing of offshore gravitationally driven tectonic episodes (c. 100 – 80 Ma) overlaps with the timing of major denudation of the continental margin inferred from low-temperature thermochronology. It is now possible to explicitly link this period of structurally controlled denudation onshore, involving the removal of several kilometres of material from the continental margin and interior, to the onset of offshore slumping. Viola et al. (2005) have documented younger neotectonic inversion structures in the Orange Basin that are linked to the onshore structural network in southern Namibia and Namaqualand. In the southern offshore basins, detailed seismic reflection have provided additional evidence of discrete episodes of basin inversion and/or reactivation (Bate and Malan, 1992; Boyd et al., 2011; Hartwig et al., 2012; Paton et al., 2008; Tinker et al., 2008b; van der Merwe and Fouche, 1992).

7.3 Drivers of post-rift tectonic activity: Regional and local mechanisms

Apatite fission track and AHe data from continental margins have commonly been interpreted in terms of simple conceptual models of escarpment evolution following break-up (e.g. Ollier

and Pain, 1997; Gallagher and Brown, 1999; Gunnel et al., 2003; Campanile et al., 2008; Persano et al., 2002, 2005; Braun and van der Beek, 2004). However, the temporal and spatial pattern of the data presented here is incompatible with that predicted by any of these conceptual models (Gallagher and Brown, 1999). The discrepancy primarily lies in the common occurrence of “young” post-rift AFT and AHe ages far inland of the escarpment. Abrupt age variations across basement structures also suggest a more complex evolution of the margin. We advocate that while a phase of syn-rift erosion of elevated rift flanks occurred (Fig. 10a), an episode of post-rift denudation during the middle to Late Cretaceous is more significant in terms of the geomorphic evolution of the margin (Fig. 10b). Based on our study, we suggest that post-rift denudation is triggered by deformation of the upper crust driven by a combination of regional horizontal and vertical stresses that are focused by local thermal and mechanical properties of the crust.

The present-day stress field across the southwestern African continental margin is characterised by a NW-SE/NNW-SSE orientation of horizontal principal compressive stress, which has been attributed to lithospheric resistance to plate and microplate movements (Bird et al., 2006). The stress field is proposed to be induced by overall net NE-SW extension and has been linked to neotectonics and present day seismic activity in the area (Andreoli et al., 1996; Ben-Avraham et al., 2002; Brandt et al., 2005; Viola et al., 2005; White et al., 2009; de Beer, 2012). This stress field will have evolved over time with changes in plate direction and spreading velocity and, as is the case today, would have promoted faulting across the margin. Loading of the offshore Orange Basin during the syn and post-rift phase, causing flexure of the lithosphere, would have enhanced coast perpendicular extensional stresses at this time (Kusznir et al., 1981; Wernicke and Axen, 1988; Redfield et al., 2005; Salomon et al., 2014; Redfield and Osmundsen, 2013). However, the style and location of fault reactivation will be dependent on the orientation of faults to the regional stress field and may have involved local compression within an overall extension regime (e.g. Holford et al., 2014).

The timing of post-rift denudation proposed here is coeval with a maximum Atlantic seafloor spreading rate (Fig. 11) of 7 – 8 cm/yr during the middle Cretaceous (c. 120 – 80 Ma) (Coffin

et al., 1998; Müller et al., 2008; Labails et al., 2009; Torsvik et al., 2009; Moulin et al., 2010). The start of this maximum spreading rate coincides with the onset of seafloor spreading in the southern Atlantic at c. 120 Ma (chron - CM0). At c. 100 Ma, a change in plate motion linked to the final break-up of South America and Africa after seafloor spreading propagated northward to the Equatorial Atlantic (Eagles, 2007; Heine et al., 2013; Jones et al., 1995; Moulin et al., 2010; Nürnberg and Müller, 1991; Torsvik et al., 2009; Granot and Dymment, 2015); although a change in spreading rate velocity is only recorded in one model (Africa-NGU; Colli et al., 2014) of Atlantic spreading rates (Fig. 11). A sharp decrease in plate velocity occurs from 80 – 70 Ma, dropping to rates of c. 3 – 4 cm/yr (Cande et al., 1988; Müller et al., 2008; Torsvik et al., 2009; Moulin et al., 2010), which is coincident with another shift in plate spreading direction (C31; Fig. 11) and the cessation of major denudation predicted by our thermal history models.

Changes to the orientation of plate movement can be observed in the geometry of Late Cretaceous to Early Cenozoic magnetic anomalies in the Mid-Atlantic and SW Indian Ocean (Fig. 11). The slow-down in plate velocity at this time has been linked to the development of new transform faults along the ridge crest and the arrival of the Reunion Plume beneath the India-African plate boundary (Cande and Stegman, 2011; Pérez-Díaz and Eagles, 2014). The magnitude of stress required to reactivate faults is dependent on fault geometry, fault orientation relative to the regional stress field, and the frictional resistance along the fault plane (Sibson, 1985; Turner and Williams, 2004). However, it is possible that even relatively small in-plane stresses may promote deformation (Cloetingh and Burov, 2011). Recent plate reconstruction models have predicted deformation along intracontinental rift systems in South America and West and Central Africa during the Cretaceous normal polarity superchron (chron CM0-C34) (Torsvik et al., 2009; Heine et al., 2013; Pérez-Díaz and Eagles, 2014), which was largely completed by c. 90 Ma (Granot and Dymment, 2015). However, kinematic modelling has predicted that between 33 – 58% of intracontinental deformation is accommodated away from major intracontinental rift basins (e.g. Guiraud and Maurin, 1992; Loule and Pospisil, 2013; Pérez-Díaz and Eagles, 2014). It is suggested in our study that mid-

to Late Cretaceous post-rift fault reactivation at the southwest African continental margin contributes to this additional deformation.

Regional vertical movements of the African plate are driven by isostatic adjustments to unloading and loading of the crust (Gilchrist and Summerfield, 1990; Hartley et al., 1996; Molnar et al., 2015) and the presence of vertical stresses at the base of the lithosphere exerted by a buoyant mantle anomaly beneath the continent (Lithgow-Bertelloni and Silver, 1998; Gurnis et al., 2000; Nyblade and Sleep, 2003; Braun et al., 2014). The latter of these has received particular attention in an attempt to constrain the spatial extent and magnitude of “dynamic” uplift and the creation of “dynamic” topography over time (Forte et al., 2010; Moucha and Forte, 2011; Braun, 2010; Flament et al., 2014). Recent numerical surface process modelling has shown that modest amounts of “dynamic” uplift can trigger kilometres of erosion (Braun et al., 2013). The relative contribution of truly “dynamic” topography (i.e. topography supported only by active flow within the mantle) in creating southern Africa’s high elevation is arguably small, while the isostatic compensation of the South African crust and mantle lithosphere arising from the relative buoyancy of material comprising the flow plays a far more important role (Molnar et al., 2015). Changes in the thermal structure and chemical and mineralogical composition of the lithosphere during the middle Cretaceous (c. 100 – 90 Ma) have been documented (Bell et al., 2003; Griffin et al., 2003; Janney et al., 2010; Stanley et al., 2013, 2015) and could have triggered isostatically driven uplift of the South African landscape at this time (Molnar et al., 2015).

The coupled dynamics of horizontal and vertical motions has been explored by mantle flow modelling by Colli et al., (2014). The intrinsic linkage between horizontal and vertical motions illustrated by these models suggests that the coeval changes at c. 90 Ma and 30 Ma in the South Atlantic spreading rates (Fig. 11) and uplift of the southwest African continental margin are geodynamically linked to one another. The interaction between in-plane stresses in the lithosphere and thermal convection in the mantle has been investigated by recent three-dimensional thermo-mechanical models using realistic lithospheric rheologies and predict complex patterns of short wavelength deformation of a brittle crust overlying a ductile

mantle-lithosphere (Burov, 2011; Guillou-Frottier et al., 2012; Burov and Gerya, 2014; Koptev et al., 2015). This work offers the interesting possibility of a regional, long-wave length mechanism of uplift, like mantle flow, causing quite local deformation at the surface, and would explain why we observe both regional and local signals of tectonic activity within southern Africa during the mid-Cretaceous. The emplacement of mid-Cretaceous alkaline intrusions along major tectonic boundaries may also indicate a link between mantle convection and tectonic readjustments (Jelsma et al., 2009).

Regional denudation during the mid- to Late Cretaceous, predicted by thermochrometry data, (Brown et al., 2002; Tinker et al., 2008a; Kounov et al., 2009; Flowers and Schoene, 2010; Stanley et al., 2013, 2015; Kounov et al., 2013; Brown et al., 2014; Wildman et al., 2015) has underpinned the timing of uplift events proposed by numerical models (e.g. van der beek et al., 2002; Braun et al., 2014; Molnar et al., 2015). By jointly inverting AFT and AHe data, our modelling results limit the thickness of overburden removed during the Cenozoic to less than 1 km or, in rare cases, less than 1.5 km. This observation is therefore consistent with models invoking a Cretaceous age for the present day landscape and is difficult to reconcile with models that predict the development of the present-day topography from an initially flat surface over the last 30 Myr (e.g. Partridge and Maud, 1987; Burke, 1996; Burke and Gunnell, 2008; Roberts and White, 2010; Paul et al., 2014; Rudge et al., 2015). Our interpretation does not preclude modest amounts surface uplift and channel incision occurring during the Cenozoic, but requires that subsequent erosion was minimal, likely due to the prevailing arid to semi-arid climate at the time (de Wit, 1999; Pickford and Senut, 1999; Bamford, 2000; Gutzmer and Beukes, 2000). In addition to limiting the contribution of Cenozoic erosion, our new data reveal short-wavelength variations in the spatial pattern of mid-Cretaceous denudation, which are attributed to post-rift brittle deformation of the margin and continental interior. Our data are incapable of resolving the specific kinematics of post-rift deformation that is caused by a complex interaction of horizontal and vertical induced stresses and crustal heterogeneity. However, these data provides important new quantitative constraints that can help to constrain future numerical models that link surface process, mantle geodynamics and global plate tectonics.

6. CONCLUSION

This study provides the first suite of joint AFT and AHe data across the Namaqualand sector of the southwest African continental margin. The data provide evidence for major crustal cooling of the margin during the Cretaceous with only minor cooling (less than 30 – 40 °C) occurring through the Cenozoic. Joint thermal-history inversion of the data provides evidence for two distinct major cooling episodes, which caused kilometre-scale denudation (c. 2 – 5 km). The early denudation event is associated with syn-rift erosion of pre-existing topography, possibly augmented by tectonic uplift during rifting, while the later episode was triggered by post-rift tectonic reactivation of the margin and regional uplift of the continent in the mid-Late Cretaceous. Any contribution of Cenozoic tectonic or surface processes did not cause enough erosion to be reflected in the thermochronology data, and the total Cenozoic erosion is restricted to less than c. 1 – 1.5 km.

The overall spatial and temporal pattern of erosion as documented here requires an explanation that can accommodate both a regional, long-wavelength geomorphic response extending well inland of the major escarpment zone, as well as more localised, short-wavelength fault-related erosion. We believe the observed pattern of topographic evolution, as inferred through the erosional response documented here, implies that several mechanisms operating at or certainly dominating over different wavelengths are involved.

In addition to erosion of the regional, long-wavelength topography linked to chemical and thermal mantle-lithosphere changes, we propose that intracontinental deformation and related relief was an important feature of the post-rift development of the margin. The transfer of plate-boundary stresses arising from significant changes in the plate kinematics within the South Atlantic and West Indian oceans, combined with an enhanced tensional stress field across the margin induced by flexure of the lithosphere, triggered fault reactivation and led to localised erosion patterns during the mid – Late Cretaceous. It is possible that some of the short-wavelength deformation is itself linked to the deeper mantle mechanism through a more

complex response of the lithosphere to vertical stresses, although this possibility remains speculative at this stage.

While the exact geometry of structures and the precise location of faults that accommodated major offsets are still unknown, the data presented here indicate a major surface response to significant mid-Late Cretaceous tectonic activity, which occurred along the so-called “passive” continental margin of the southwest Africa.

7. ACKNOWLEDGMENTS

Full analytical details, data tables, QTQt models and QTQt data files can be accessed through the online supporting information. We thank Dr. Marco Andreoli, Dr. Olivier Datueil, Prof. Francois Guillocheau, Dr. Tim Redfield, Dr. Delphine Rouby, and Dr. Giulio Viola, for insightful discussions at various stages of the project. Dr. Marco Andreoli and staff at NECSA are thanked for logistic support during fieldwork in Namaqualand. Prof. Rebecca Flowers, Dr. Taylor Schildgen and Prof. Peter van der Beek are thanked for their constructive comments on this work. This work was funded by the Natural Environment Research Council, UK, grant number NE/H008276/1.

8. REFERENCES

- Adamson, R. S. (1931). Note on some petrified wood from Banke, Namaqualand. Transactions of the Royal Society of South Africa, 19, (3), 255-258, doi: 10.1080/00359193109518838
- Andreoli, M. A. G., Doucouré, M., Van Bever Donker, J., Brandt, D., & Andersen, N. J. B. (1996). Neotectonics of southern Africa. Africa Geoscience Review, 3, 1-16.
- Andreoli, M. A., Hart, R. J., Ashwal, L. D., & Coetzee, H. (2006). Correlations between U, Th content and metamorphic grade in the western Namaqualand Belt, South Africa, with implications for radioactive heating of the crust. Journal of Petrology, 47(6), 1095-1118, doi: 10.1093/petrology/egl004.

- Andreoli, M. A., Viola, G., Kounov, A., Scheepers, J., Heidbach, O., & Stengel, I. (2009, September). History of stress at Vaalputs, Namaqualand, South Africa: evidence for a Mid-Cretaceous “Wegener-type Orogeny” in western southern Africa. 11th SAGA Biennial Technical Meeting and Exhibition, Swaziland, 16 - 18 September 2009, p. 515 - 520
- Aslanian, D., Moulin, M., Olivet, J. L., Unternehr, P., Matias, L., Bache, F., Rabineau, M., Nouzé, H., Klingelhoefer, F., Contrucci, I. & Labails, C. (2009). Brazilian and African passive margins of the Central Segment of the South Atlantic Ocean: Kinematic constraints. *Tectonophysics*, 468(1), 98-112, doi:10.1016/j.tecto.2008.12.016.
- Aslanian, D., & Moulin, M. (2013). Palaeogeographic consequences of conservative models in the South Atlantic Ocean. Geological Society, London, Special Publications, 369(1), 75-90, doi: 10.1144/SP369.5.
- Ault, A. K., & Flowers, R. M. (2012). Is apatite U–Th zonation information necessary for accurate interpretation of apatite (U–Th)/He thermochronometry data?. *Geochimica et Cosmochimica Acta*, 79, 60-78, doi:10.1016/j.gca.2011.11.037.
- Autin, J., Bellahsen, N., Leroy, S., Husson, L., Beslier, M. O., & d'Acremont, E. (2013). The role of structural inheritance in oblique rifting: insights from analogue models and application to the Gulf of Aden. *Tectonophysics*, 607, 51-64, doi:10.1016/j.tecto.2013.05.041.
- Ballard, S., & Pollack, H. N. (1987). Diversion of heat by Archean cratons: a model for southern Africa. *Earth and Planetary Science Letters*, 85(1), 253-264, doi:10.1016/0012-821X(87)90036-7.
- Bamford, M. K. (2000). Fossil woods of Karoo age deposits in South Africa and Namibia as an aid to biostratigraphical correlation. *Journal of African Earth Sciences*, 31(1), 119-132, doi:10.1016/S0899-5362(00)00077-4.
- Bate, K. J., & Malan, J. A. (1992). Tectonostratigraphic evolution of the Algoa, Gamtoos and Pletmos basins, offshore South Africa. In *Inversion tectonics of the Cape Fold Belt, Karoo and cretaceous basins of Southern Africa* (pp. 61-73). Balkema Rotterdam, ISBN: 9054100478, 9789054100478.

- Bell, D. R., Schmitz, M. D., & Janney, P. E. (2003). Mesozoic thermal evolution of the southern African mantle lithosphere. *Lithos*, 71(2), 273-287.
- Ben-Avraham, Z., Smith, G., Reshef, M., & Jungslager, E. (2002). Gas hydrate and mud volcanoes on the southwest African continental margin off South Africa. *Geology*, 30(10), 927-930, doi:10.1016/S0024-4937(03)00117-8.
- Beucher, R., Brown, R. W., Roper, S., Stuart, F., & Persano, C. (2013). Natural age dispersion arising from the analysis of broken crystals: Part II. Practical application to apatite (U–Th)/He thermochronometry. *Geochimica et Cosmochimica Acta*, 120, 395-416, doi:10.1016/j.gca.2013.05.042.
- Bierman, P. R., & Caffee, M. (2001). Slow rates of rock surface erosion and sediment production across the Namib Desert and escarpment, southern Africa. *American Journal of Science*, 301(4-5), 326-358, doi: 10.2475/ajs.301.4-5.326.
- Bierman, P. R., Coppersmith, R., Hanson, K., Neveling, J., Portenga, E. W., & Rood, D. H. (2014). A cosmogenic view of erosion, relief generation, and the age of faulting in southern Africa. *GSA Today*, 24(9), 4-11, doi: 10.1130/GSATG206A.1.
- Bird, P., Ben-Avraham, Z., Schubert, G., Andreoli, M., & Viola, G. (2006). Patterns of stress and strain rate in southern Africa. *Journal of Geophysical Research: Solid Earth* (1978–2012), 111, B08402, doi: 10.1029/2005JB003882.
- Blaich, O. A., Faleide, J. I., & Tsikalas, F. (2011). Crustal breakup and continent-ocean transition at South Atlantic conjugate margins. *Journal of Geophysical Research: Solid Earth* (1978–2012), 116(B1), doi: 10.1029/2010JB007686..
- Boyd, D., Anka, Z., Di Primio, R., Kuhlmann, G., & De Wit, M. J. (2011). Passive margin evolution and controls on natural gas leakage in the Orange Basin, South Africa. *South African Journal of Geology*, 114(3-4), 415-432.
- Brandt, D., Andreoli, M. A. G., & McCarthy, T. S. (2003). Mesozoic fluvial deposits on a rifted continental margin near Vaalputs, Namaqualand, South Africa. *South African Journal of Geology*, 106(1), 11-16, doi: 10.2113/1060011.
- Brandt, D., Andreoli, M. A. G., & McCarthy, T. S. (2005). The late Mesozoic palaeosoils and Cenozoic fluvial deposits at Vaalputs, Namaqualand, South Africa: Possible

- depositional mechanisms and their bearing on the evolution of the continental margin. *South African Journal of Geology*, 108(2), 271-284, doi: 10.2113/108.2.271.
- Braun, J. (2010). The many surface expressions of mantle dynamics. *Nature Geoscience*, 3(12), 825-833, doi:10.1038/ngeo1020.
- Braun, J., & Beaumont, C. (1989). A physical explanation of the relation between flank uplifts and the breakup unconformity at rifted continental margins. *Geology*, 17(8), 760-764, doi: 10.1130/0091-7613.
- Braun, J., & Van Der Beek, P. (2004). Evolution of passive margin escarpments: What can we learn from low-temperature thermochronology?. *Journal of Geophysical Research: Earth Surface* (2003–2012), 109, F04009, doi:10.1029/2004JF000147.
- Braun, J., Robert, X., & Simon-Labric, T., (2013), Eroding dynamic topography. *Geophysical Research Letters*. 40, 8, 1494-1499, doi: 10.1002/grl.50310.
- Braun, J., Guillocheau, F., Robin, C., Baby, G., & Jelsma, H. (2014). Rapid erosion of the Southern African Plateau as it climbs over a mantle superswell. *Journal of Geophysical Research: Solid Earth*, 119(7), 6093-6112, doi: 10.1002/2014JB010998.
- Bronner, A., Sauter, D., Manatschal, G., Péron-Pinvidic, G., & Munsch, M. (2011). Magmatic breakup as an explanation for magnetic anomalies at magma-poor rifted margins. *Nature geoscience*, 4(8), 549-553, doi:10.1038/ngeo1201.
- Brown, L. F. (Ed.). (1995). *Sequence Stratigraphy in Offshore South African Divergent Basins: An Atlas on Exploration for Cretaceous Lowstand Traps* by Soekor (Pty) Ltd, AAPG Studies in Geology 41 (No. 41). AAPG, ISBN: 0891810498.
- Brown, R. W., Beucher, R., Roper, S., Persano, C., Stuart, F., & Fitzgerald, P. (2013). Natural age dispersion arising from the analysis of broken crystals. Part I: Theoretical basis and implications for the apatite (U–Th)/He thermochronometer. *Geochimica et Cosmochimica Acta*, 122, 478-497, doi:10.1016/j.gca.2013.05.041.
- Brown, R. W., Rust, D. J., Summerfield, M. A., Gleadow, A. J., & De Wit, M. C. (1990). An Early Cretaceous phase of accelerated erosion on the south-western margin of Africa: Evidence from apatite fission track analysis and the offshore sedimentary record. *International Journal of Radiation Applications and Instrumentation. Part D. Nuclear*

- Tracks and Radiation Measurements, 17(3), 339-350, doi:10.1016/1359-0189(90)90056-4.
- Brown, R. W., Summerfield, M. A., & Gleadow, A. J. (2002). Denudational history along a transect across the Drakensberg Escarpment of southern Africa derived from apatite fission track thermochronology. *Journal of Geophysical Research: Solid Earth* (1978–2012), 107(B12), ETG-10, doi: 10.1029/2001JB000745.
- Brown, R., Summerfield, M., Gleadow, A., Gallagher, K., Carter, A., Beucher, R., & Wildman, M. (2014). Intracontinental deformation in southern Africa during the Late Cretaceous. *Journal of African Earth Sciences*, 100, 20-41, doi:10.1016/j.jafrearsci.2014.05.014.
- Brune, S., Heine, C., Pérez-Gussinyé, M., & Sobolev, S. V. (2014). Rift migration explains continental margin asymmetry and crustal hyper-extension. *Nature communications*, 5, 1 – 9.
- Buiter, S. J., & Torsvik, T. H. (2014). A review of Wilson Cycle plate margins: A role for mantle plumes in continental break-up along sutures?. *Gondwana Research*, 26(2), 627-653, doi:10.1038/ncomms5014.
- Burbank, D.W., and Anderson, R.S., 2012, *Tectonic Geomorphology*; 2nd edition: Wiley, 454 pp., ISBN : 978-1-4443-3886-7.
- Burke, K. (1996). The African plate. *South African journal of geology*, 99(4), 341-409.
- Burke, K., & Gunnell, Y. (2008). The African erosion surface: a continental-scale synthesis of geomorphology, tectonics, and environmental change over the past 180 million years. *Geological Society of America Memoirs*, 201, 1-66, doi: 10.1130/2008.1201.
- Burov, E. B. (2011). Rheology and strength of the lithosphere. *Marine and Petroleum Geology*, 28(8), 1402-1443, doi:10.1016/j.marpetgeo.2011.05.008.
- Burov, E., & Gerya, T. (2014). Asymmetric three-dimensional topography over mantle plumes. *Nature*, 513(7516), 85-89, doi:10.1038/nature13703.
- Calignano, E., Sokoutis, D., Willingshofer, E., Gueydan, F., & Cloetingh, S. (2015). Strain localization at the margins of strong lithospheric domains: Insights from analog models. *Tectonics*, 34(3), 396-412, doi: 10.1002/2014tc003756.

- Campanile, D., Nambiar, C. G., Bishop, P., Widdowson, M., & Brown, R. (2008). Sedimentation record in the Konkan–Kerala Basin: implications for the evolution of the Western Ghats and the Western Indian passive margin. *Basin Research*, 20(1), 3-22, doi: 10.1111/j.1365-2117.2007.00341.x.
- Cande, S. C., & Stegman, D. R. (2011). Indian and African plate motions driven by the push force of the Reunion plume head. *Nature*, 475(7354), 47-52, doi: 10.1038/nature10174.
- Cande, S. C., J. L. LaBrecque, and W. F. Haxby (1988), Plate kinematics of the South Atlantic: Chron C34 to present, *J. Geophys. Res.*, 93(B11), 12,479–13,492, doi: 10.1029/jb093ib11p13479.
- Clemson, J., Cartwright, J., & Booth, J. (1997). Structural segmentation and the influence of basement structure on the Namibian passive margin. *Journal of the Geological Society*, 154(3), 477-482, doi: 10.1144/gsjgs.154.3.0477.
- Clifford, T. N., Barton, E. S., Stern, R. A., & Duchesne, J. C. (2004). U–Pb zircon calendar for Namaquan (Grenville) crustal events in the granulite-facies terrane of the O'okiep copper district of South Africa. *Journal of Petrology*, 45(4), 669-691, doi: 10.1093/petrology/egg097.
- Cloetingh, S., & Burov, E. (2011). Lithospheric folding and sedimentary basin evolution: a review and analysis of formation mechanisms. *Basin Research*, 23(3), 257-290.
- Cloetingh, S., Burov, E., & Francois, T. (2013). Thermo-mechanical controls on intra-plate deformation and the role of plume-folding interactions in continental topography. *Gondwana Research*, 24(3), 815-837, doi: 10.1111/j.1365-2117.2010.00490.x.
- Cockburn, H. A. P., Brown, R. W., Summerfield, M. A., & Seidl, M. A. (2000). Quantifying passive margin denudation and landscape development using a combined fission-track thermochronology and cosmogenic isotope analysis approach. *Earth and Planetary Science Letters*, 179(3), 429-435, doi: 10.1016/s0012-821x(00)00144-8.
- Codilean, A. T., Bishop, P., Stuart, F. M., Hoey, T. B., Fabel, D., & Freeman, S. P. (2008). Single-grain cosmogenic ^{21}Ne concentrations in fluvial sediments reveal spatially variable erosion rates. *Geology*, 36(2), 159-162, doi: 10.1130/g24360a.1.
- Codilean, A. T., Fenton, C. R., Fabel, D., Bishop, P., & Xu, S. (2014). Discordance between cosmogenic nuclide concentrations in amalgamated sands and individual fluvial

- pebbles in an arid zone catchment. *Quaternary Geochronology*, 19, 173-180, doi: 10.1016/j.quageo.2012.04.007.
- Coffin, M. F., Gahagan, L. M., & Lawver, L. A. (1998). Present-day plate boundary digital data compilation. University of Texas Institute for Geophysics Technical Report, 174(5).
- Cogné, N., Gallagher, K., & Cobbold, P. R. (2011). Post-rift reactivation of the onshore margin of southeast Brazil: Evidence from apatite (U–Th)/He and fission-track data. *Earth and Planetary Science Letters*, 309(1), 118-130, doi: 10.1016/j.epsl.2011.06.025.
- Colli, L., Stotz, I., Bunge, H. P., Smethurst, M., Clark, S., Iaffaldano, G., Tassara, A., Guillocheau, F. & Bianchi, M. C. (2014). Rapid South Atlantic spreading changes and coeval vertical motion in surrounding continents: Evidence for temporal changes of pressure-driven upper mantle flow. *Tectonics*, 33(7), 1304-1321, doi: 10.1002/2014tc003612.
- Colliston, W. P., Schoch, A. E., & Cole, J. (2014). The Grenvillian Namaqua-Natal fold belt adjacent to the Kaapvaal Craton: 1. Distribution of Mesoproterozoic collisional terranes deduced from results of regional surveys and selected profiles in the western and southern parts of the fold belt. *Journal of African Earth Sciences*, 100, 7-19, doi: 10.1016/j.jafrearsci.2014.05.017.
- Cornelissen, A. K., & Verwoerd, W. J. (1975). The Bushmanland kimberlites and related rocks. *Physics and Chemistry of the Earth*, 9, 71-80, doi: 10.1016/b978-0-08-018017-5.50011-0.
- Cornell, D.H., Thomas, R.J., Gibson, R., Moen, H.F.G., Reid, D.L., Moore, J.M., Gibson, R.L., 2006. The Namaqua-Natal Province. Geological Society of South Africa, pp. 325–379, ISBN: 9781919908779.
- Corti, G., Ranalli, G., Agostini, A., & Sokoutis, D. (2013). Inward migration of faulting during continental rifting: Effects of pre-existing lithospheric structure and extension rate. *Tectonophysics*, 594, 137-148, doi: 10.1016/j.tecto.2013.03.028.
- Corti, G., van Wijk, J., Cloetingh, S., & Morley, C. K. (2007). Tectonic inheritance and continental rift architecture: Numerical and analogue models of the East African Rift system. *Tectonics*, 26(6), doi: 10.1029/2006tc002086.

- Curtis, C. G., Harris, C., Trumbull, R. B., de Beer, C., & Mudzanani, L. (2013). Oxygen Isotope Diversity in the Anorogenic Koegel Fontein Complex of South Africa: a Case for Basement Control and Selective Melting for the Production of Low- $\delta^{18}\text{O}$ Magmas. *Journal of Petrology*, 54(7), 1259-1283, doi: 10.1093/petrology/egt011.
- Curtis, C. G., Trumbull, R. B., de Beer, C. H., Harris, C., Reid, D. L., & Romer, R. L. (2011). Geochemistry of the early Cretaceous Koegel Fontein anorogenic igneous complex, South Africa. *South African Journal of Geology*, 114(3-4), 353-378, doi: 10.2113/gssajg.114.3-4.353.
- Daly, M. C., Chorowicz, J., & Fairhead, J. D. (1989). Rift basin evolution in Africa: the influence of reactivated steep basement shear zones. Geological Society, London, Special Publications, 44(1), 309-334, doi: 10.1144/gsl.sp.1989.044.01.17.
- Danišik, M., Štěpančíková, P., & Evans, N. J. (2012). Constraining long-term denudation and faulting history in intraplate regions by multisystem thermochronology: An example of the Sudetic Marginal Fault (Bohemian Massif, central Europe). *Tectonics*, 31(2), TC2003, doi: 10.1029/2011tc003012.
- Dauteuil, O., Deschamps, F., Bourgeois, O., Mocquet, A., & Guillocheau, F. (2013). Post-breakup evolution and palaeotopography of the North Namibian Margin during the Meso-Cenozoic. *Tectonophysics*, 589, 103-115, doi: 10.1016/j.tecto.2012.12.022.
- Day, R. W. (1987). False Bay dolerites. *Annals Geological Survey of South Africa*, 21, 1-7.
- de Beer, C. H. (2010). The geology of the Garies area: Explanation of 1:250 000-scale Sheet 3017 Garies. South Africa: Council for Geoscience.
- de Beer, C. H. (2012). Evidence of Neogene to Quaternary faulting and seismogenic deformation along the Namaqualand coast, South Africa. *South African Journal of Geology*, 115(2), 117-136, doi: 10.2113/gssajg.115.2.117.
- de Beer, C. H., & Council for geoscience. (2002). The Geology of the Calvinia Area: Explanation of 1: 250 000-scale Sheet 3118 Calvinia. South Africa: Council for Geoscience.
- de Vera, J., Granado, P., & McClay, K. (2010). Structural evolution of the Orange Basin gravity-driven system, offshore Namibia. *Marine and Petroleum Geology*, 27(1), 223-237, doi: 10.1016/j.marpetgeo.2009.02.003.

- de Wit, M. C. J. (1988). Aspects of the geomorphology of the north-western Cape, South Africa. In Dardis, G. F. and Moon, B. P. (eds.), *Geomorphological studies in southern Africa*, p. 57-69, Rotterdam, Netherlands: CRC Press, ISBN: 1-86812-072-4.
- de Wit, M. C. J., Ward, J. D. and Spaggiari, R., 1992, A reappraisal of the Kangnas dinosaur site, Bushmanland, South Africa., *South African Journal of Science*. v. 88 (9-10), p. 504-507
- de Wit, M. C. J. (1999). Post-Gondwana drainage and the development of diamond placers in western South Africa. *Economic Geology*, 94(5), 721-740, doi: 10.2113/gsecongeo.94.5.721.
- de Wit, M. J. C., Marshall, T. R., and Partridge, T. C., (2000), Fluvial deposits and drainage evolution. In: Partridge, T. C. and Maud, R. R. (Editors), *The Cenozoic of southern Africa*, Oxford University Press, United Kingdom, p. 55 – 72, ISBN: 978-0195125306.
- Decker, J. E., Niedermann, S., & de Wit, M. J. (2011). Soil erosion rates in South Africa compared with cosmogenic ³He-based rates of soil production. *South African Journal of Geology*, 114(3-4), 475-488, doi: 10.2113/gssajg.114.3-4.475.
- Decker, J. E., Niedermann, S., & De Wit, M. J. (2013). Climatically influenced denudation rates of the southern African plateau: Clues to solving a geomorphic paradox. *Geomorphology*, 190, 48-60, doi: 10.1016/j.geomorph.2013.02.007.
- Dewey, J. F., Robb, L., & Van Schalkwyk, L. (2006). Did Bushmanland extensionally unroof Namaqualand?. *Precambrian Research*, 150(3), 173-182, doi: 10.1016/j.precamres.2006.07.007.
- Dollar, E. S. (1998). Palaeofluvial geomorphology in southern Africa: a review. *Progress in Physical Geography*, 22(3), 325-349, doi: 10.1177/030913339802200302.
- Donelick, R. A., O'Sullivan, P. B., & Ketcham, R. A. (2005). Apatite fission-track analysis. *Reviews in Mineralogy and Geochemistry*, 58(1), 49-94, doi: 10.2138/rmg.2005.58.3.
- Duncan, R. A., Hooper, P. R., Rehacek, J., Marsh, J. S., & Duncan, A. R. (1997). The timing and duration of the Karoo igneous event, southern Gondwana. *Journal of Geophysical Research: Solid Earth* (1978–2012), 102(B8), 18127-18138, doi: 10.1029/97jb00972.

- Dunkl, I. (2002). TRACKKEY: a Windows program for calculation and graphical presentation of fission track data. *Computers & Geosciences*, 28(1), 3-12, doi: 10.1016/s0098-3004(01)00024-3.
- Eagles, G. (2007). New angles on South Atlantic opening. *Geophysical Journal International*, 168(1), 353-361, doi: 10.1111/j.1365-246x.2006.03206.x.
- Eagles, G., & König, M. (2008). A model of plate kinematics in Gondwana breakup. *Geophysical Journal International*, 173(2), 703-717, doi: 10.1111/j.1365-246x.2008.03753.x.
- Eglington, B. M. (2006). Evolution of the Namaqua-Natal Belt, southern Africa—A geochronological and isotope geochemical review. *Journal of African Earth Sciences*, 46(1), 93-111, doi: 10.1016/j.jafrearsci.2006.01.014.
- Erlanger, E. D., Granger, D. E., & Gibbon, R. J. (2012). Rock uplift rates in South Africa from isochron burial dating of fluvial and marine terraces. *Geology*, 40(11), 1019-1022, doi: 10.1130/g33172.1.
- Estes, R. (1977). Relationships of the South African fossil frog *Eoxenopoides reuningi* (Anura, Pipidae). *South African Museum, Annals*. 73(2), 1-36.
- Farley, K. A., Shuster, D. L., & Ketcham, R. A. (2011). U and Th zonation in apatite observed by laser ablation ICPMS, and implications for the (U–Th)/He system. *Geochimica et Cosmochimica Acta*, 75(16), 4515-4530, doi: 10.1016/j.gca.2011.05.020.
- Farley, K. A., Wolf, R. A., & Silver, L. T. (1996). The effects of long alpha-stopping distances on (U–Th)/He ages. *Geochimica et Cosmochimica Acta*, 60(21), 4223-4229, doi: 10.1016/s0016-7037(96)00193-7.
- Fitzgerald, P. G., Baldwin, S. L., Webb, L. E., & O'sullivan, P. B. (2006). Interpretation of (U–Th)/He single grain ages from slowly cooled crustal terranes: a case study from the Transantarctic Mountains of southern Victoria Land. *Chemical Geology*, 225(1), 91-120, doi: 10.1016/j.chemgeo.2005.09.001.
- Flament, N., Gurnis, M., Williams, S., Seton, M., Skogseid, J., Heine, C., & Müller, R. D. (2014). Topographic asymmetry of the South Atlantic from global models of mantle

- flow and lithospheric stretching. *Earth and Planetary Science Letters*, 387, 107-119, doi: 10.1016/j.epsl.2013.11.017.
- Flowers, R. M., & Kelley, S. A. (2011). Interpreting data dispersion and —inverted dates in apatite (U–Th)/He and fission-track datasets: an example from the US midcontinent. *Geochimica et Cosmochimica Acta*, 75(18), 5169-5186, doi: 10.1016/j.gca.2011.06.016.
- Flowers, R. M., & Schoene, B. (2010). (U–Th)/He thermochronometry constraints on unroofing of the eastern Kaapvaal craton and significance for uplift of the southern African Plateau. *Geology*, 38(9), 827-830, doi: 10.1130/g30980.1.
- Flowers, R. M., Ketcham, R. A., Shuster, D. L., & Farley, K. A. (2009). Apatite (U–Th)/He thermochronometry using a radiation damage accumulation and annealing model. *Geochimica et Cosmochimica Acta*, 73(8), 2347-2365.
- Forte, A. M., Moucha, R., Simmons, N. A., Grand, S. P., & Mitrovica, J. X. (2010). Deep-mantle contributions to the surface dynamics of the North American continent. *Tectonophysics*, 481(1), 3-15, doi: 10.1016/j.gca.2009.01.015.
- Franco-Magalhaes, A. O. B., Cuglieri, M. A. A., Hackspacher, P. C., & Saad, A. R. (2014). Long-term landscape evolution and post-rift reactivation in the southeastern Brazilian passive continental margin: Taubaté basin. *International Journal of Earth Sciences*, 103(2), 441-453, doi: 10.1007/s00531-013-0967-4.
- François, T., Burov, E., Meyer, B., & Agard, P. (2013). Surface topography as key constraint on thermo-rheological structure of stable cratons. *Tectonophysics*, 602, 106-123, doi: 10.1016/j.tecto.2012.10.009.
- Frimmel, H. E., Basei, M. A., Correa, V. X., & Mbangula, N. (2013). A new lithostratigraphic subdivision and geodynamic model for the Pan-African western Saldania Belt, South Africa. *Precambrian Research*, 231, 218-235, doi: 10.1016/j.precamres.2013.03.014.
- Gaina, C., Torsvik, T. H., van Hinsbergen, D. J., Medvedev, S., Werner, S. C., & Labails, C. (2013). The African Plate: A history of oceanic crust accretion and subduction since the Jurassic. *Tectonophysics*, 604, 4-25, doi: 10.1016/j.tecto.2013.05.037.

- Galbraith, R. F. (2010). Statistics for fission track analysis, Chapman and Hall/CRC, p. 240, ISBN: 9781584885337.
- Galbraith, R. F., & Green, P. F. (1990). Estimating the component ages in a finite mixture. *International Journal of Radiation Applications and Instrumentation. Part D. Nuclear Tracks and Radiation Measurements*, 17(3), 197-206, doi: 10.1016/1359-0189(90)90035-v.
- Gallagher, K. (2012). Transdimensional inverse thermal history modeling for quantitative thermochronology. *Journal of Geophysical Research: Solid Earth* (1978–2012), 117(B2), doi: 10.1029/2011jb008825.
- Gallagher, K. (1995). Evolving temperature histories from apatite fission-track data. *Earth and Planetary Science Letters*, 136(3), 421-435, doi: 10.1016/0012-821x(95)00197-k.
- Gallagher, K. & Brown, R. (1999). Denudation and uplift at passive margins: the record on the Atlantic Margin of southern Africa. *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 357(1753), 835-859, doi: 10.1098/rsta.1999.0354.
- Gallagher, K. & Brown, R., (1997), The onshore record of passive margin evolution. *Journal of the Geological Society*, 154(3), 451-457, doi: 10.1144/gsjgs.154.3.0451.
- Gallagher, K., Charvin, K., Nielsen, S., Sambridge, M., & Stephenson, J. (2009). Markov chain Monte Carlo (MCMC) sampling methods to determine optimal models, model resolution and model choice for Earth Science problems. *Marine and Petroleum Geology*, 26(4), 525-535, doi: 10.1016/j.marpetgeo.2009.01.003.
- Gautheron, C., Barbarand, J., Ketcham, R. A., Tassan-Got, L., van der Beek, P., Pagel, M., Pinna-Jamme, R., Couffignal, F. & Fialin, M. (2013). Chemical influence on α -recoil damage annealing in apatite: Implications for (U–Th)/He dating. *Chemical Geology*, 351, 257-267, doi: 10.1016/j.chemgeo.2013.05.027.
- Gautheron, C., Tassan-Got, L., Barbarand, J., & Pagel, M. (2009). Effect of alpha-damage annealing on apatite (U–Th)/He thermochronology. *Chemical Geology*, 266(3), 157-170, doi: 10.1016/j.chemgeo.2009.06.001.
- Gautheron, C., Tassan-Got, L., Ketcham, R. A., & Dobson, K. J. (2012). Accounting for long alpha-particle stopping distances in (U–Th–Sm)/He geochronology: 3D modeling of

- diffusion, zoning, implantation, and abrasion. *Geochimica et Cosmochimica Acta*, 96, 44-56, doi: 10.1016/j.gca.2012.08.016.
- Gilks, W. R. (2005). *Markov chain monte carlo*. John Wiley & Sons, Ltd, doi: 10.1002/0470011815.b2a14021.
- Gilchrist, A. R., Kooi, H., & Beaumont, C. (1994). Post-Gondwana geomorphic evolution of southwestern Africa: Implications for the controls on landscape development from observations and numerical experiments. *Journal of Geophysical Research: Solid Earth* (1978–2012), 99(B6), 12211-12228, doi: 10.1029/94jb00046.
- Gilchrist, A. R., & Summerfield, M. A. (1990). Differential denudation and flexural isostasy in formation of rifted-margin upwarps, 346, 739 – 742, doi:10.1038/346739a0.
- Goudie, A. S. (2005). The drainage of Africa since the Cretaceous. *Geomorphology*, 67(3), 437-456, doi: 10.1016/j.geomorph.2004.11.008.
- Granot, R., & Dymant, J. (2015). The Cretaceous opening of the South Atlantic Ocean. *Earth and Planetary Science Letters*, 414, 156-163, doi: 10.1016/j.epsl.2015.01.015.
- Green, P. F. (1986). On the thermo-tectonic evolution of Northern England: evidence from fission track analysis. *Geological Magazine*, 123(5), 493-506, doi: 10.1017/s0016756800035081.
- Green, P. F., & Duddy, I. R. (2006). Interpretation of apatite (U–Th)/He ages and fission track ages from cratons. *Earth and Planetary Science Letters*, 244(3), 541-547, doi: 10.1016/j.epsl.2006.02.024.
- Green, P. F., Duddy, I. R., Japsen, P., Bonow, J. M., & Malan, J. A. (2015). Post-breakup burial and exhumation of the southern margin of Africa. *Basin Research*, doi: 10.1111/bre.12167.
- Green, P. F., Lidmar-Bergström, K., Japsen, P., Bonow, J. M., & Chalmers, J. A. (2013). Stratigraphic landscape analysis, thermochronology and the episodic development of elevated, passive continental margins. Geological Survey of Denmark and Greenland. Danish Ministry of climate, energy and building, ISBN: 978-87-7871-372-8.
- Gresse, P. G., & Germs, G. J. B. (1993). The Nama foreland basin: sedimentation, major unconformity bounded sequences and multisided active margin advance. *Precambrian Research*, 63(3), 247-272, doi: 10.1016/0301-9268(93)90036-2.

- Gresse, P.G., Von Veh, M.W., Frimmel, H.E., 2006. Namibian (Neoproterozoic) to early Cambrian successions. *The Geology of South Africa*. Geological Society of South Africa, Johannesburg, pp. 395–420, ISBN: 9781919908779.
- Griffin, W. L., O'Reilly, S. Y., Natapov, L. M., & Ryan, C. G. (2003). The evolution of lithospheric mantle beneath the Kalahari Craton and its margins. *Lithos*, 71(2), 215–241, doi: 10.1016/j.lithos.2003.07.006.
- Groenewald, P. B., Grantham, G. H., & Watkeys, M. K. (1991). Geological evidence for a Proterozoic to Mesozoic link between southeastern Africa and Dronning Maud Land, Antarctica. *Journal of the Geological Society*, 148(6), 1115–1123, doi: 10.1144/gsjgs.148.6.1115.
- Guillocheau, F., Rouby, D., Robin, C., Helm, C., Rolland, N., Le Carlier de Veslud, C., & Braun, J. (2012). Quantification and causes of the terrigenous sediment budget at the scale of a continental margin: a new method applied to the Namibia–South Africa margin. *Basin Research*, 24(1), 3–30, doi: 10.1111/j.1365-2117.2011.00511.x.
- Guillou-Frottier, L., Burov, E., Cloetingh, S., Le Goff, E., Deschamps, Y., Huet, B., & Bouchot, V. (2012). Plume-induced dynamic instabilities near cratonic blocks: Implications for P–T–t paths and metallogeny. *Global and Planetary Change*, 90, 37–50, doi: 10.1016/j.gloplacha.2011.10.007.
- Guiraud, R., & Maurin, J. C. (1992). Early Cretaceous rifts of Western and Central Africa: an overview. *Tectonophysics*, 213(1), 153–168, doi: 10.1016/0040-1951(92)90256-6.
- Gunnell, Y., Gallagher, K., Carter, A., Widdowson, M., & Hurford, A. J. (2003). Denudation history of the continental margin of western peninsular India since the early Mesozoic—reconciling apatite fission-track data with geomorphology. *Earth and Planetary Science Letters*, 215(1), 187–201, doi: 10.1016/s0012-821x(03)00380-7.
- Gurnis, M., Mitrovica, J. X., Ritsema, J., & van Heijst, H. J. (2000). Constraining mantle density structure using geological evidence of surface uplift rates: The case of the African superplume. *Geochemistry, Geophysics, Geosystems*, 1(7), doi: 10.1029/1999gc000035.

- Gutzmer, J., & Beukes, N. J. (2000). Asbestiform manjiroite and todorokite from the Kalahari manganese field, South Africa. *South African Journal of Geology*, 103(3-4), 163-174, doi: 10.2113/1030163.
- Hansen, K., & Reiners, P. W. (2006). Low temperature thermochronology of the southern East Greenland continental margin: evidence from apatite (U–Th)/He and fission track analysis and implications for intermethod calibration. *Lithos*, 92(1), 117-136, doi: 10.1016/j.lithos.2006.03.039.
- Hanson, E.K., Moore, J.M., Bordy, E.M., Marsh, J.S., Howarth, G., and Robey, J.V.A., 2009, Cretaceous erosion in central South Africa: Evidence from upper-crustal xenoliths in kimberlite diatremes: *South African Journal of Geology*, v. 112, p. 125–140, doi:10.2113/gssajg.112.2.125.
- Hartley, R., Watts, A. B., & Fairhead, J. D. (1996). Isostasy of Africa. *Earth and Planetary Science Letters*, 137(1), 1-18, doi: 10.1016/0012-821x(95)00185-f.
- Hartwig, A., Anka, Z., & di Primio, R. (2012). Evidence of a widespread paleo-pockmarked field in the Orange Basin: An indication of an early Eocene massive fluid escape event offshore South Africa. *Marine Geology*, 332, 222-234, doi: 10.1016/j.margeo.2012.07.012.
- Haughton, S. H. (1931). On a collection of fossil frogs from the clays at Banke. *Transactions of the Royal Society of South Africa*, 19(3), 233-249, doi: 10.1080/00359193109518836.
- Heine, C., Zoethout, J., and Müller, R. D., (2013), Kinematics of the South Atlantic rift, *Solid Earth*, 4, 215-253, doi: 10.5194/se-4-215-2013.
- Hendriks, B. W. H., & Redfield, T. F. (2005). Apatite fission track and (U-Th)/He data from Fennoscandia: An example of underestimation of fission track annealing in apatite. *Earth and Planetary Science Letters*, 236(1), 443-458, doi: 10.1016/j.epsl.2005.05.027.
- Hirsch, K. K., Scheck-Wenderoth, M., van Wees, J. D., Kuhlmann, G., & Paton, D. A. (2010). Tectonic subsidence history and thermal evolution of the Orange Basin. *Marine and Petroleum Geology*, 27(3), 565-584, doi: 10.1016/j.marpetgeo.2009.06.009.
- Holford, S. P., Tuitt, A. K., Hillis, R. R., Green, P. F., Stoker, M. S., Duddy, I. R., Sandiford, M. & Tassone, D. R. (2014). Cenozoic deformation in the Otway Basin, southern

- Australian margin: implications for the origin and nature of post-breakup compression at rifted margins. *Basin Research*, 26(1), 10-37, doi: 10.1111/bre.12035.
- House, M. A., Wernicke, B. P., Farley, K. A., & Dumitru, T. A. (1997). Cenozoic thermal evolution of the central Sierra Nevada, California, from (UTh)/He thermochronometry. *Earth and Planetary Science Letters*, 151(3), 167-179, doi: 10.1016/s0012-821x(97)81846-8.
- Huisman, R. S., & Beaumont, C. (2014). Rifted continental margins: The case for depth-dependent extension. *Earth and Planetary Science Letters*, 407, 148-162, doi: 10.1016/j.epsl.2014.09.032.
- Hurford, A. J., & Green, P. F. (1982). A users' guide to fission track dating calibration. *Earth and Planetary Science Letters*, 59(2), 343-354, doi: 10.1016/0012-821x(82)90136-4.
- Jackson, M., Cramez, C., & Fonck, J. M. (2000). Role of subaerial volcanic rocks and mantle plumes in creation of South Atlantic margins: implications for salt tectonics and source rocks. *Marine and Petroleum Geology*, 17(4), 477-498, doi: 10.1016/s0264-8172(00)00006-4.
- Jacobs, J., Thomas, R. J., & Weber, K. (1993). Accretion and indentation tectonics at the southern edge of the Kaapvaal craton during the Kibaran (Grenville) orogeny. *Geology*, 21(3), 203-206, doi: 10.1130/0091-7613(1993)021<0203:aaitat>2.3.co;2.
- Janney, P. E., Shirey, S. B., Carlson, R. W., Pearson, D. G., Bell, D. R., Le Roex, A. P., Ishikawa, A., Nixon, P. H. & Boyd, F. R. (2010). Age, composition and thermal characteristics of South African off-craton mantle lithosphere: Evidence for a multi-stage history. *Journal of Petrology*, doi: 10.1093/petrology/egq041.
- Jelsma, H., Barnett, W., Richards, S., & Lister, G. (2009). Tectonic setting of kimberlites. *Lithos*, 112, 155-165, doi: 10.1016/j.lithos.2009.06.030.
- Johnson, M. R., Anhaeuser, C. R., & Thomas, R. J. (2006). *The Geology of South Africa*. Geological Society of South Africa, ISBN: 9781919908779.
- Jones, E. J. W., Cande, S. C., & Spathopoulos, F. (1995). Evolution of a major oceanographic pathway: the equatorial Atlantic. Geological Society, London, Special Publications, 90(1), 199-213, doi: 10.1144/gsl.sp.1995.090.01.12.

- Jones, M. Q. W. (1987). Heat flow and heat production in the Namaqua mobile belt, South Africa. *Journal of Geophysical Research: Solid Earth* (1978–2012), 92(B7), 6273–6289, doi: 10.1029/jb092ib07p06273.
- Jourdan, F., Féraud, G., Bertrand, H., & Watkeys, M. K. (2007). From flood basalts to the inception of oceanization: Example from the $^{40}\text{Ar}/^{39}\text{Ar}$ high-resolution picture of the Karoo large igneous province. *Geochemistry, Geophysics, Geosystems*, 8(2), doi: 10.1029/2006gc001392.
- Jourdan, F., Féraud, G., Bertrand, H., Kampunzu, A. B., Tshoso, G., Watkeys, M. K., & Le Gall, B. (2005). Karoo large igneous province: Brevity, origin, and relation to mass extinction questioned by new $^{40}\text{Ar}/^{39}\text{Ar}$ age data. *Geology*, 33(9), 745–748, doi: 10.1130/g21632.1.
- Jungslager, E. H. (1999). Petroleum habitats of the Atlantic margin of South Africa. *Geological Society, London, Special Publications*, 153(1), 153–168, doi: 10.1144/gsl.sp.1999.153.01.10.
- Karl, M., Glasmacher, U. A., Kollenz, S., Franco-Magalhaes, A. O., Stockli, D. F., & Hackspacher, P. C. (2013). Evolution of the South Atlantic passive continental margin in southern Brazil derived from zircon and apatite (U–Th–Sm)/He and fission-track data. *Tectonophysics*, 604, 224–244, doi: 10.1016/j.tecto.2013.06.017.
- Kennett, B. L. N., & Iaffaldano, G. (2013). Role of lithosphere in intra-continental deformation: Central Australia. *Gondwana Research*, 24(3), 958–968, doi: 10.1016/j.gr.2012.10.010.
- Ketcham, R. A. (2005). The role of crystallographic angle in characterizing and modeling apatite fission-track length data. *Radiation measurements*, 39(6), 595–601, doi: 10.1016/j.radmeas.2004.07.008.
- Ketcham, R. A., Carter, A., Donelick, R. A., Barbarand, J., & Hurford, A. J. (2007). Improved modeling of fission-track annealing in apatite. *American Mineralogist*, 92(5–6), 799–810, doi: 10.2138/am.2007.2281.
- Kooi, H., & Beaumont, C. (1994). Escarpment evolution on high-elevation rifted margins: Insights derived from a surface processes model that combines diffusion, advection,

- and reaction. *Journal of Geophysical Research: Solid Earth* (1978–2012), 99(B6), 12191-12209, doi: 10.1029/94jb00047.
- Koopmann, H., Franke, D., Schreckenberger, B., Schulz, H., Hartwig, A., Stollhofen, H., & di Primio, R. (2014). Segmentation and volcano-tectonic characteristics along the SW African continental margin, South Atlantic, as derived from multichannel seismic and potential field data. *Marine and Petroleum Geology*, 50, 22-39, doi: 10.1016/j.marpetgeo.2013.10.016.
- Koptev, A., Calais, E., Burov, E., Leroy, S., & Gerya, T. (2015). Dual continental rift systems generated by plume-lithosphere interaction. *Nature Geoscience*, 8 (5), 388 – 392, doi: 10.1038/ngeo2401.
- Kounov, A., Niedermann, S., de Wit, M. J., Codilean, A. T., Viola, G., Andreoli, M., & Christl, M. (2015). Cosmogenic ^{21}Ne and ^{10}Be reveal a more than 2 ma alluvial fan flanking the Cape Mountains, South Africa. *South African Journal of Geology*, 118(2), 129-144, doi: 10.2113/gssajg.118.2.129.
- Kounov, A., Niedermann, S., de Wit, M. J., Viola, G., Andreoli, M., & Erzinger, J. (2007). Present denudation rates at selected sections of the South African escarpment and the elevated continental interior based on cosmogenic ^3He and ^{21}Ne . *South African Journal of Geology*, 110(2-3), 235-248, doi: 10.2113/gssajg.110.2-3.235.
- Kounov, A., Viola, G., De Wit, M., & Andreoli, M. A. G. (2009). Denudation along the Atlantic passive margin: new insights from apatite fission-track analysis on the western coast of South Africa. *Geological Society, London, Special Publications*, 324(1), 287-306, doi: 10.1144/sp324.19.
- Kounov, A., Viola, G., Dunkl, I., & Frimmel, H. E. (2013). Southern African perspectives on the long-term morpho-tectonic evolution of cratonic interiors. *Tectonophysics*, 601, 177-191, doi: 10.1016/j.tecto.2013.05.009.
- Ksienzyk, A. K., Dunkl, I., Jacobs, J., Fossen, H., & Kohlmann, F. (2014). From orogen to passive margin: constraints from fission track and (U–Th)/He analyses on Mesozoic uplift and fault reactivation in SW Norway. *Geological Society, London, Special Publications*, 390, SP390-27, doi: 10.1144/sp390.27.

- Kuhlmann, G., Adams, S., Campher, C., van der Spuy, D., di Primio, R., & Horsfield, B. (2010). Passive margin evolution and its controls on natural gas leakage in the southern Orange Basin, blocks 3/4, offshore South Africa. *Marine and Petroleum Geology*, 27(4), 973-992, doi: 10.1016/j.marpetgeo.2010.01.010.
- Kusznir, N.J., Marsden, G., Egan, S.S., 1991. A flexural-cantilever simple-shear/pure shear model of continental lithosphere extension: applications to the Jeanne d'Arc basin, Grand Banks and Viking Graben, North sea. In: Roberts, A.M., Yielding, G., Freeman, B. (Eds.), *The Geometry of Normal Faults*, Spec. Publ. - Geol. Soc. Lond., vol. 56, pp. 41-60, doi: 10.1144/gsl.sp.1991.056.01.04.
- Labails, C., T. H. Torsvik, C. Gaina, and R. M. Cocks (2009), Global plate polygons 2009. SPlates Model (version 2.0), Tech. Rep. 2009.047, NGU, Postboks 6315 Sluppen, 7491 Trondheim. [Available at <http://www.ngu.no/en-gb/>.]
- Leprêtre, R., Missenard, Y., Barbarand, J., Gautheron, C., Saddiqi, O., & Pinna-Jamme, R. (2015). Post-rift history of the eastern Central Atlantic passive margin: insights from the Saharan region of South Morocco. *Journal of Geophysical Research: Solid Earth*, 120 (6), 4645 – 4666, doi: 10.1002/2014jb011549.
- Lister, G. S., Etheridge, M. A., & Symonds, P. A. (1991). Detachment models for the formation of passive continental margins. *Tectonics*, 10(5), 1038-1064, doi: 10.1029/90tc01007.
- Lithgow-Bertelloni, C., & Silver, P. G. (1998). Dynamic topography, plate driving forces and the African superswell. *Nature*, 395(6699), 269-272, doi: 10.1038/26212.
- Loule, J. P., & Pospisil, L. (2013). Geophysical evidence of Cretaceous volcanics in Logone Birni Basin (Northern Cameroon), Central Africa, and consequences for the West and Central African Rift System. *Tectonophysics*, 583, 88-100, doi: 10.1016/j.tecto.2012.10.021.
- Lundin, E. R., & Doré, A. G. (2011). Hyperextension, serpentinization, and weakening: A new paradigm for rifted margin compressional deformation. *Geology*, 39(4), 347-350, doi: 10.1130/g31499.1.

- Macey, P. H., Siegfried, H. P., Minnaar, H., Almond, J., & Botha, P. M. W. (2011). The geology of the Loeriesfontein area. Explanation of 1:250,000-scale Loeriesfontein. South Africa: Council for Geoscience.
- Marais, J. A. H., Agenbacht, A. L. D., Prinsloo, M., & Basson, W. A. (2001). The geology of the Springbok area. Explanation of 1:250 000-scale 2916 Loeriesfontein. South Africa: Council for Geoscience
- Masini, E., Manatschal, G., & Mohn, G. (2013). The Alpine Tethys rifted margins: reconciling old and new ideas to understand the stratigraphic architecture of magma-poor rifted margins. *Sedimentology*, 60(1), 174-196, doi: 10.1111/sed.12017.
- Mbongo-Djimbi, D., Gautheron, C., Roques, J., Tassan-Got, L., Gerin, C., & Simoni, E. (2015). Impact of apatite chemical composition on (U-Th)/He thermochronometry: An atomistic point of view. *Geochimica et Cosmochimica Acta*, 167, 162-176, doi: 10.1016/j.gca.2015.06.017.
- McMillan, I. K. (2003). Foraminiferally defined biostratigraphic episodes and sedimentation pattern of the Cretaceous drift succession (Early Barremian to Late Maastrichtian) in seven basins on the South African and southern Namibian continental margin. *South African Journal of Science*, 99(11 & 12), p-537, ISSN: 00382353.
- Meesters, A. G. C. A., & Dunai, T. J. (2002a). Solving the production–diffusion equation for finite diffusion domains of various shapes: Part I. Implications for low-temperature (U-Th)/He thermochronology. *Chemical Geology*, 186(3), 333-344, doi: 10.1016/s0009-2541(01)00422-3.
- Meesters, A. G. C. A., & Dunai, T. J. (2002b). Solving the production–diffusion equation for finite diffusion domains of various shapes: Part II. Application to cases with α -ejection and nonhomogeneous distribution of the source. *Chemical Geology*, 186(3), 347-363, doi: 10.1016/s0009-2541(02)00073-6.
- Molnar, P., England, P. C., & Jones, C. H. (2015). Mantle dynamics, isostasy, and the support of high terrain. *Journal of Geophysical Research: Solid Earth*, 120(3), 1932-1957, doi: 10.1002/2014jb011724.

- Moore, A. E., & Verwoerd, W. J. (1985). The olivine melilitite-" kimberlite"-carbonatite suite of Namaqualand and Bushmanland, South Africa. *South African Journal of Geology*, 88(2), 281-294.
- Moore, A., Blenkinsop, T., & Cotterill, F. W. (2008). Controls on post-Gondwana alkaline volcanism in Southern Africa. *Earth and Planetary Science Letters*, 268(1), 151-164, doi: 10.1016/j.epsl.2008.01.007.
- Moucha, R., Forte, A. M., Mitrovica, J. X., Rowley, D. B., Quéré, S., Simmons, N. A., & Grand, S. P. (2008). Dynamic topography and long-term sea-level variations: There is no such thing as a stable continental platform. *Earth and Planetary Science Letters*, 271(1), 101-108, doi: 10.1016/j.epsl.2008.03.056.
- Moucha, R., & Forte, A. M. (2011). Changes in African topography driven by mantle convection. *Nature Geoscience*, 4(10), 707-712, doi: 10.1038/ngeo1235.
- Moulin, M., Aslanian, D., & Unternehr, P. (2010). A new starting point for the South and Equatorial Atlantic Ocean. *Earth-Science Reviews*, 98(1), 1-37, doi: 10.1016/j.earscirev.2009.08.001.
- Moulin, M., Fluteau, F., Courtillot, V., Marsh, J., Delpech, G., Quidelleur, X., Gerard, M. & Jay, A. E. (2011). An attempt to constrain the age, duration, and eruptive history of the Karoo flood basalt: Naude's Nek section (South Africa). *Journal of Geophysical Research: Solid Earth* (1978–2012), 116(B7), doi: 10.1029/2011jb008210.
- Müller, R. D., M. Sdrolias, C. Gaina, and W. R. Roest (2008), Age, spreading rates, and spreading asymmetry of the world's ocean crust, *Geochem. Geophys. Geosyst.*, 9(4), Q04006, doi:10.1029/2007GC001743.
- Muntingh, A., & Brown Jr, L. F. (1993). Sequence Stratigraphy of Petroleum Plays, Post-Rift Cretaceous Rocks (Lower Aptian to Upper Maastrichtian), Orange Basin, Western Offshore, South Africa: Chapter 4: Recent Applications of Siliciclastic Sequence Stratigraphy, In *Siliciclastic Sequence Stratigraphy: Recent Developments and Applications*, Weimer, P. and Posamentier, H. (Eds.), ISBN: 978-1-58861-449-0.
- Naliboff, J., & Buiter, S. J. (2015). Rift reactivation and migration during multiphase extension. *Earth and Planetary Science Letters*, 421, 58-67, doi: 10.1016/j.epsl.2015.03.050.

- Nürnberg, D., & Müller, R. D. (1991). The tectonic evolution of the South Atlantic from Late Jurassic to present. *Tectonophysics*, 191(1), 27-53, doi: 10.1016/0040-1951(91)90231-g.
- Nyblade, A. A., & Sleep, N. H. (2003). Long lasting epeirogenic uplift from mantle plumes and the origin of the Southern African Plateau. *Geochemistry, Geophysics, Geosystems*, 4(12), 10.1029/2003gc000573.
- Ollier, C. D., & Pain, C. F. (1997). Equating the basal unconformity with the palaeoplain: a model for passive margins. *Geomorphology*, 19(1), 1-15, doi: 10.1016/s0169-555x(96)00048-7.
- Partridge, T. C., & Maud, R. R. (1987). Geomorphic evolution of southern Africa since the Mesozoic. *South African Journal of Geology*, 90(2), 179-208.
- Paton, D. (2012). Post-Rift Deformation of the North East and South Atlantic Margins: Are “Passive Margins” Really Passive? *Tectonics of sedimentary basins: recent advances*, 249-269, doi: 10.1002/9781444347166.ch12.
- Paton, D. A., Di Primio, R., Kuhlmann, G., Van Der Spuy, D., & Horsfield, B. (2007). Insights into the petroleum system evolution of the southern Orange Basin, South Africa. *South African Journal of Geology*, 110(2-3), 261-274, doi: 10.2113/gssajg.110.2-3.261.
- Paton, D. A., van der Spuy, D., di Primio, R., & Horsfield, B. (2008). Tectonically induced adjustment of passive-margin accommodation space; influence on the hydrocarbon potential of the Orange Basin, South Africa. *AAPG bulletin*, 92(5), 589-609, doi: 10.1306/12280707023.
- Paul, J. D., Roberts, G. G., & White, N. (2014). The African landscape through space and time. *Tectonics*, 33(6), 898-935, doi: 10.1002/2013tc003479.
- Pérez-Díaz, L., & Eagles, G. (2014). Constraining South Atlantic growth with seafloor spreading data. *Tectonics*, 33(9), 1848-1873, doi: 10.1002/2014tc003644.
- Péron-Pinvidic, G., & Manatschal, G. (2009). The final rifting evolution at deep magma-poor passive margins from Iberia-Newfoundland: a new point of view. *International Journal of Earth Sciences*, 98(7), 1581-1597, doi: 10.1007/s00531-008-0337-9.

- Péron-Pinvidic, G., Manatschal, G., & Osmundsen, P. T. (2013). Structural comparison of archetypal Atlantic rifted margins: a review of observations and concepts. *Marine and Petroleum Geology*, 43, 21-47, doi: doi:10.1016/j.marpetgeo.2013.02.002.
- Pérez-Gussinyé, M., Lowry, A. R., & Watts, A. B. (2007). Effective elastic thickness of South America and its implications for intracontinental deformation. *Geochemistry, Geophysics, Geosystems*, 8(5), doi: 10.1029/2006gc001511.
- Persano, C., Stuart, F. M., Bishop, P., & Barfod, D. N. (2002). Apatite (U–Th)/He age constraints on the development of the Great Escarpment on the southeastern Australian passive margin. *Earth and Planetary Science Letters*, 200(1), 79-90, doi: 10.1016/s0012-821x(02)00614-3.
- Persano, C., Stuart, F. M., Bishop, P., & Dempster, T. J. (2005). Deciphering continental breakup in eastern Australia using low-temperature thermochronometers. *Journal of Geophysical Research: Solid Earth* (1978–2012), 110(B12), doi: 10.1029/2004jb003325.
- Phillips, D., Kiviets, G.B., Biddulph, M.G., Madav, M.K., 2000. Cenozoic volcanism. In: Partridge, T.C., Maud, R.R. (Eds.), *The Cenozoic of Southern Africa*, p. 182–197. Oxford University Press. ISBN: 978-0195125306.
- Pickford, M., and B. Senut (1999), *Geology and Palaeobiology of the Namib Desert Southwest Africa*, Memoir, vol. 18, 155 pp., Geological Survey of Namibia, Windhoek, Namibia.
- Raab, M. J., Brown, R. W., Gallagher, K., Carter, A., & Weber, K. (2002). Late Cretaceous reactivation of major crustal shear zones in northern Namibia: constraints from apatite fission track analysis. *Tectonophysics*, 349(1), 75-92, doi: 10.1016/s0040-1951(02)00047-1.
- Redfield, T. F., Osmundsen, P. T., & Hendriks, B. W. H. (2005). The role of fault reactivation and growth in the uplift of western Fennoscandia. *Journal of the Geological Society*, 162(6), 1013-1030, doi: 10.1144/0016-764904-149.
- Redfield, T. F., & Osmundsen, P. T. (2013). The long-term topographic response of a continent adjacent to a hyperextended margin: A case study from Scandinavia. *Geological Society of America Bulletin*, 125(1-2), 184-200, doi: 10.1130/b30691.1.

- Reid, D. L. (1990). The Cape Peninsula dolerite dyke swarm, South Africa, In *Mafic Dikes and Emplacement Mechanisms*, Parker, A. J., Rickwood, P. C., and Tucker, D. H. (Eds.), p. 325-334. Rotterdam, The Netherlands: Balkema, ISBN: 978-9061911586.
- Reid, D. L., & Rex, D. C. (1994). Cretaceous dykes associated with the opening of the South Atlantic; the Mehlberg Dyke, northern Richtersveld. *South African Journal of Geology*, 97(2), 135-145.
- Reid, D. L., Cooper, A. F., Rex, D. C., & Harmer, R. E. (1990). Timing of post-Karoo alkaline volcanism in southern Namibia. *Geological Magazine*, 127(5), 427-433, doi: 10.1017/s001675680001517x.
- Reid, D. L., Erlank, A. J., & Rex, D. C. (1991). Age and correlation of the False Bay dolerite dyke swarm, south-western Cape, Cape Province. *South African Journal of Geology*, 94(2-3), 155-158, doi: 10.2113/gssajg.114.3-4.335.
- Reiners, P. W., & Farley, K. A. (2001). Influence of crystal size on apatite (U-Th)/He thermochronology: an example from the Bighorn Mountains, Wyoming. *Earth and Planetary Science Letters*, 188(3), 413-420, doi: 10.1016/s0012-821x(01)00341-7.
- Reston, T. J. (2005). Polyphase faulting during the development of the west Galicia rifted margin. *Earth and Planetary Science Letters*, 237(3), 561-576, doi: 10.1016/j.epsl.2005.06.019.
- Roberts, D. L., Botha, G. A., Maud, R. R., & Pether, J. (2006). Coastal cenozoic deposits. In Johnson, M. R., Anhaeusser, C. R., and Thomas, R. J. (eds.), *The Geology of South Africa*. Geological Society of South Africa, p. 605-628, ISBN: 9781919908779.
- Roberts, G. G., & White, N. (2010). Estimating uplift rate histories from river profiles using African examples. *Journal of Geophysical Research: Solid Earth* (1978–2012), 115(B2), doi: 10.1029/2009jb006692.
- Rouby, D., Bonnet, S., Guillocheau, F., Gallagher, K., Robin, C., Biancotto, F., Dauteuil, O. & Braun, J. (2009). Sediment supply to the Orange sedimentary system over the last 150My: An evaluation from sedimentation/denudation balance. *Marine and Petroleum Geology*, 26(6), 782-794, doi: 10.1016/j.marpetgeo.2008.08.004.

- Rudge, J. F., Roberts, G. G., White, N., & Richardson, C. N. (2015). Uplift histories of Africa and Australia from linear inverse modeling of drainage inventories. *Journal of Geophysical Research: Earth Surface*, 120 (5), 894 – 914, doi: 10.1002/2014jf003297.
- Rust, D. J., & Summerfield, M. A. (1990). Isopach and borehole data as indicators of rifted margin evolution in southwestern Africa. *Marine and Petroleum Geology*, 7(3), 277-287, doi: 10.1016/0264-8172(90)90005-2.
- Saintot, A., Stephens, M. B., Viola, G., & Nordgulen, Ø. (2011). Brittle tectonic evolution and paleostress field reconstruction in the southwestern part of the Fennoscandian Shield, Forsmark, Sweden. *Tectonics*, 30(4), doi: 10.1029/2010tc002781.
- Salomon, E., Koehn, D., & Passchier, C. (2015). Brittle reactivation of ductile shear zones in NW Namibia in relation to South Atlantic rifting. *Tectonics*, 34(1), doi: 10.1002/2014tc003728.
- Salomon, E., Koehn, D., Passchier, C., Hackspacher, P. C., & Glasmacher, U. A. (2014). Contrasting stress fields on correlating margins of the South Atlantic. *Gondwana Research*, 28 (3), 1152 - 1167, doi: 10.1016/j.gr.2014.09.006.
- Sambridge, M., Gallagher, K., Jackson, A., & Rickwood, P. (2006). Trans-dimensional inverse problems, model comparison and the evidence. *Geophysical Journal International*, 167(2), 528-542, doi: 10.1111/j.1365-246x.2006.03155.x.
- Scharf, T. E., Codilean, A. T., de Wit, M., Jansen, J. D., & Kubik, P. W. (2013). Strong rocks sustain ancient postorogenic topography in southern Africa. *Geology*, 41(3), 331-334, doi: 10.1130/g33806.1.
- Scholtz, A. (1985). The palynology of the upper lacustrine sediments of the Arnot Pipe, Banke, Namaqualand, The South African Museum, 95. p. 109, ISBN: 0868130664, 9780868130668.
- Senut, B., Pickford, M., & Wessels, D. (1997). Panafrican distribution of lower Miocene Hominoidea. *Comptes Rendus de l'Académie des Sciences-Series IIA-Earth and Planetary Science*, 325(9), 741-746, doi: 10.1016/s1251-8050(97)89119-7.
- Séranne, M., & Anka, Z. (2005). South Atlantic continental margins of Africa: a comparison of the tectonic vs climate interplay on the evolution of equatorial west Africa and SW

- Africa margins. *Journal of African Earth Sciences*, 43(1), 283-300, doi: 10.1016/j.jafrearsci.2005.07.010.
- Sibson, R. H. (1985). A note on fault reactivation. *Journal of Structural Geology*, 7(6), 751-754, doi: 10.1016/0191-8141(85)90150-6.
- Sobel, E. R., & Seward, D. (2010). Influence of etching conditions on apatite fission-track etch pit diameter. *Chemical Geology*, 271(1), 59-69, doi: 10.1016/j.chemgeo.2009.12.012.
- Spiegel, C., Kohn, B., Belton, D., Berner, Z., & Gleadow, A. (2009). Apatite (U–Th–Sm)/He thermochronology of rapidly cooled samples: the effect of He implantation. *Earth and Planetary Science Letters*, 285(1), 105-114, doi: 10.1016/j.epsl.2009.05.045.
- Stanley, J. R., Flowers, R. M., & Bell, D. R. (2013). Kimberlite (U–Th)/He dating links surface erosion with lithospheric heating, thinning, and metasomatism in the southern African Plateau. *Geology*, 41(12), 1243-1246, doi: 10.1130/g34797.1.
- Stanley, J. R., Flowers, R. M., & Bell, D. R. (2015). Erosion patterns and mantle sources of topographic change across the southern African Plateau derived from the shallow and deep records of kimberlites. *Geochemistry, Geophysics, Geosystems*, 16(9), 3235-3256, doi: 10.1002/2015gc005969.
- Stevenson, I. R., & McMillan, I. K. (2004). Incised valley fill stratigraphy of the Upper Cretaceous succession, proximal Orange Basin, Atlantic margin of southern Africa. *Journal of the Geological Society*, 161(2), 185-208, doi: 10.1144/0016-764902-003.
- Stockli, D. F., Farley, K. A., & Dumitru, T. A. (2000). Calibration of the apatite (U–Th)/He thermochronometer on an exhumed fault block, White Mountains, California. *Geology*, 28(11), 983-986, doi: 10.1130/0091-7613(2000)28<983:cotaht>2.0.co;2.
- Summerfield, M. A. (1985). Plate tectonics and landscape development on the African continent. In Morisawa, M., and Hack, J. T., *Tectonic Geomorphology*, 27-51. Boston: Allan and Unwin, ISBN: 9780045510986.
- Svensen, H., Corfu, F., Polteau, S., Hammer, Ø., & Planke, S. (2012). Rapid magma emplacement in the Karoo large igneous province. *Earth and Planetary Science Letters*, 325, 1-9, doi:10.1016/j.epsl.2012.01.015.

- Sykes, L. R. (1978). Intraplate seismicity, reactivation of preexisting zones of weakness, alkaline magmatism, and other tectonism postdating continental fragmentation. *Reviews of Geophysics*, 16(4), 621-688, doi: 10.1029/rg016i004p00621.
- Tankard, A., Welsink, H., Aukes, P., Newton, R., Stettler, E., 2009. Tectonic evolution of the Cape and Karoo basins of South Africa. *Mar. Pet. Geol.* 26 (8), 1379–1412, doi: 10.1016/j.marpetgeo.2009.01.022.
- Tankard, A.J., Eriksson, S.C., Hunter, D.R., Jackson, M.P.A., Hobday, D.K., Minter, W.E.L., 1982. *Crustal Evolution of Southern Africa: 3.8 Billion Years of Earth History*. Springer-Verlag, New York, p. 523, ISBN: 978-1-4613-8149-5.
- Thomas, R. J., Agenbacht, A. L. D., Cornell, D. H., & Moore, J. M. (1994). The Kibaran of southern Africa: tectonic evolution and metallogeny. *Ore Geology Reviews*, 9(2), 131-160, doi: 10.1016/0169-1368(94)90025-6.
- Tinker, J., de Wit, M., & Brown, R. (2008a). Mesozoic exhumation of the southern Cape, South Africa, quantified using apatite fission track thermochronology. *Tectonophysics*, 455(1), 77-93, doi: 10.1016/j.tecto.2007.10.009.
- Tinker, J., de Wit, M., & Brown, R. (2008b). Linking source and sink: evaluating the balance between onshore erosion and offshore sediment accumulation since Gondwana break-up, South Africa. *Tectonophysics*, 455(1), 94-103, doi: 10.1016/j.tecto.2007.11.040.
- Torsvik, T. H., Rouse, S., Labails, C., & Smethurst, M. A. (2009). A new scheme for the opening of the South Atlantic Ocean and the dissection of an Aptian salt basin. *Geophysical Journal International*, 177(3), 1315-1333, doi: 10.1111/j.1365-246x.2009.04137.x.
- Trumbull, R. B., Reid, D. L., de Beer, C., Van Acken, D., & Romer, R. L. (2007). Magmatism and continental breakup at the west margin of southern Africa: A geochemical comparison of dolerite dikes from northwestern Namibia and the Western Cape. *South African Journal of Geology*, 110(2-3), 477-502, doi: 10.2113/gssajg.110.2-3.477.
- Turner, J. P., & Williams, G. A. (2004). Sedimentary basin inversion and intra-plate shortening. *Earth-Science Reviews*, 65(3), 277-304, doi: 10.1016/j.earscirev.2003.10.002.

- van der Beek, P., Summerfield, M. A., Braun, J., Brown, R. W., & Fleming, A. (2002). Modeling postbreakup landscape development and denudational history across the southeast African (Drakensberg Escarpment) margin. *Journal of Geophysical Research: Solid Earth* (1978–2012), 107(B12), ETG-11, doi: 10.1029/2001jb000744.
- van der Merwe, R., & Fouche, J. (1992). Inversion tectonics in the Bredasdorp basin, offshore South Africa. *Inversion Tectonics of the Cape Fold Belt, Karoo and Cretaceous Basins of Southern Africa*, de Wit, M. J. and Ransome, I. G. D. (Eds.), 49-59, ISBN: 9054100478, 9789054100478.
- van der Wateren, F. M., & Dunai, T. J. (2001). Late Neogene passive margin denudation history—cosmogenic isotope measurements from the central Namib desert. *Global and Planetary Change*, 30(3), 271-307, doi: 10.1016/s0921-8181(01)00104-7.
- Vermeesch, P. (2008). Three new ways to calculate average (U–Th)/He ages. *Chemical Geology*, 249(3), 339-347, doi: 10.1016/j.chemgeo.2008.01.027.
- Vermeesch, P., Seward, D., Latkoczy, C., Wipf, M., Günther, D., & Baur, H. (2007). α -Emitting mineral inclusions in apatite, their effect on (U–Th)/He ages, and how to reduce it. *Geochimica et Cosmochimica Acta*, 71(7), 1737-1746, doi: 10.1016/j.gca.2006.09.020.
- Viola, G., Andreoli, M., Ben-Avraham, Z., Stengel, I., & Reshef, M. (2005). Offshore mud volcanoes and onland faulting in southwestern Africa: neotectonic implications and constraints on the regional stress field. *Earth and Planetary Science Letters*, 231(1), 147-160, doi: 10.1016/j.epsl.2004.12.001.
- Viola, G., Kounov, A., Andreoli, M. A. G., & Mattila, J. (2012). Brittle tectonic evolution along the western margin of South Africa: more than 500Myr of continued reactivation. *Tectonophysics*, 514, 93-114, doi: 10.1016/j.tecto.2011.10.009.
- Visser, J. N. J. (1983). Submarine debris flow deposits from the Upper Carboniferous Dwyka tillite formation in the Kalahari Basin, South Africa. *Sedimentology*, 30(4), 511-523, doi: 10.1111/j.1365-3091.1983.tb00689.x.
- Visser, J. N. J. (1990). Glacial bedforms at the base of the Permo-Carboniferous Dwyka Formation along the western margin of the Karoo Basin, South Africa. *Sedimentology*, 37(2), 231-245, doi: 10.1111/j.1365-3091.1990.tb00957.x.

- Visser, J. N. J. (1987). The palaeogeography of part of southwestern Gondwana during the Permo-Carboniferous glaciation. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 61, 205-219, doi: 10.1016/0031-0182(87)90050-2.
- Visser, J. N. J. (1989). The Permo-Carboniferous Dwyka Formation of southern Africa: deposition by a predominantly subpolar marine ice sheet. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 70(4), 377-391, doi: 10.1016/0031-0182(89)90115-6.
- Voordouw, R. J., & Rajesh, H. M. (2012). Granitoids from the Margate Terrane and their implications for tectono-magmatic models of the Natal Metamorphic Province (South Africa). *South African Journal of Geology*, 115(1), 47-64, doi: 10.2113/gssajg.115.1.47.
- Waters, D. J. (1989). Metamorphic evidence for the heating and cooling path of Namaqualand granulites. Geological Society, London, Special Publications, 43(1), 357-363, doi: 10.1144/gsl.sp.1989.043.01.31.
- Wernicke, B., & Axen, G. J. (1988). On the role of isostasy in the evolution of normal fault systems. *Geology*, 16(9), 848-851, doi: 10.1130/0091-7613(1988)016<0848:otroii>2.3.co;2.
- White, S., Stollhofen, H., Stanistreet, I. G., & Lorenz, V. (2009). Pleistocene to recent rejuvenation of the Hebron Fault, SW Namibia. Geological Society, London, Special Publications, 316(1), 293-317, doi: 10.1144/sp316.18.
- Wildman, M., Brown, R., Watkins, R., Carter, A., Gleadow, A., & Summerfield, M. (2015). Post break-up tectonic inversion across the southwestern cape of South Africa: new insights from apatite and zircon fission track thermochronometry, *Tectonophysics*, 654, 30 – 55, doi: 10.1016/j.tecto.2015.04.012.
- Will, T. M., & Frimmel, H. E. (2013). The Influence of Inherited Structures on Dike Emplacement during Gondwana Breakup in Southwestern Africa. *The Journal of Geology*, 121(5), 455-474, doi: 10.1086/671398.
- Willett, S. D. (1999). Orogeny and orography: The effects of erosion on the structure of mountain belts. *Journal of Geophysical Research: Solid Earth* (1978–2012), 104(B12), 28957-28981, doi: 10.1029/1999jb900248.

Ziegler, P. A., & Cloetingh, S. (2004). Dynamic processes controlling evolution of rifted basins. *Earth-Science Reviews*, 64(1), 1-50, doi: 10.1016/s0012-8252(03)00041-2.

FIGURE AND TABLE CAPTIONS

Figure 1: Inset figure shows the first-order topography of southern Africa and location of the Namaqualand Highlands study area. Main location map DEM created using SRTM90m data. Elevation map is draped over Landsat ETM+ RGB:321 satellite images to enhance local relief and geomorphic features. Elevation profiles for three coast-perpendicular transects are shown and are used in Figure 9 with projected data.

Figure 2: Geological map of the Namaqualand Highlands study area. Geological map was redrawn to highlight the major geological units, structures and features relevant to this study, using 1:250,000 maps produced by the Council for Geoscience, South Africa. (Garies, sheet 3017: de Beer, 2010; Loriesfontein, sheet 3018: Macey et al., 2011; Springbok, sheet 2916: Marais et al., 2001). The dashed line around the Dasdap and Vaalputs formations represents the proposed extent of these units (after Brandt et al., 2005 and de Beer, 2010).

Figure 3: (a) Plot of AFT and AHe age against elevation. White circles are Central AFT ages with uncertainty bars representing 1σ standard error on the central age. Red squares are uncorrected mean AHe ages with uncertainty bars representing 1σ standard deviation (standard deviation is used in this instance to highlight the dispersion in single-grain AHe ages), red crosses and x's represent the minimum and maximum AHe single-grain age. (b) AFT age against MTLs uncorrected for their c-axis orientation; In AFT ages are Central AFT ages with 1σ standard error; MTLs are uncorrected for their c-axis orientation with 1σ standard error; (c) Plot of Central AFT ages against mean AHe age. Vertical error bars are 1σ standard error on the central age and horizontal error bars are 1σ standard deviation on the mean AHe age. Dashed line represents the one-to-one relationship.

Figure 4: (a) Relationships of AHe age (uncorrected) against spherical equivalent radius (R^*) (upper row) and against effective uranium ($[eU] = [U] + [0.235*Th]$). All uncertainties

include 1σ analytical uncertainty plus an additional 10% uncertainty observed in the dispersion of Durango standards analysed. The two plots in the final column are cartoons illustrating the relative influence on AHe ages by the different factors causing natural AHe single-grain age dispersion (R^* -spherical equivalent radius; eU – effective uranium; F – crystal fragment length; after Brown et al., 2013). The competing influence of all dispersion contributors perturbs simple 2D relationships. Age increases with increasing eU and R^* . Larger fragment lengths of broken crystals are typically older than small fragment lengths. (b) Probability density functions of a normal distribution centred on the mean AFT and mean AHe age with 1σ the standard deviation on the mean. (c) Location map of samples used for dispersion plots.

Figure 5: Thermal history modelling results for three samples from the Namaqualand Highlands highlighting Early Cretaceous cooling (upper plots), their data predictions (lower plots) and their location on the topographic map of the study area. Upper Plots: Blue line shows the expected model (i.e. average of all models weighted for their posterior probability); magenta lines indicate 95% credible intervals for the expected model. Red box indicates the prior information on temperature and time. Green box is used to indicate the timing of the syn-rift cooling episode predicted across the entire margin (i.e. c. 150 – 130 Ma). Lower Plots: Histogram shows the observed track-length distributions; red curve indicates the predicted track-length distributions, grey curves indicate 95% credible intervals (i.e. uncertainty) for track-length distribution prediction. Inset plot shows the relationship of observed data against model predicted data; green circles represent single-grain AHe ages and blue circles represent AFT ages.

Figure 6: Thermal history modelling results for three samples from the Namaqualand Highlands highlighting mid-Late Cretaceous cooling (upper plots) and their data predictions (lower plots) and their location on the topographic map of the study area. Upper Plots: Blue line shows the expected model (i.e. average of all models weighted for their posterior probability); magenta lines indicate 95% credible intervals for the expected model. Red box indicates the prior information on temperature and time. Yellow and red boxes are used to

indicate the timing of the post-rift cooling episodes: c. 110 – 90 Ma (structurally controlled denudation across the entire margin and regional denudation of the interior plateau) and c. 80 – 60 Ma (structurally controlled denudation, potential caused by compression driven inversion in parts of South Africa and in Namibia), respectively. Lower Plots: Histogram shows the observed track-length distributions; red curve indicates the predicted track-length distributions, grey curves indicate 95% credible intervals (i.e. uncertainty) for track-length distributions prediction. Inset plot shows the relationship of observed data against model predicted data; green circles represent single-grain AHe ages and blue circles represent AFT age.

Figure 7: Thermal history modelling results for an elevation profile of samples in the Namaqualand Highlands (upper plots) and their data predictions (lower plots). Upper Plots: Blue line shows the expected model (i.e. average of all models weighted for their posterior probability) for the upper sample; red lines shows the expected model for the bottom sample in the profile; grey lines show expected model for intervening samples. Cyan and magenta lines indicate 95% credible intervals for the top and bottom sample expected model, respectively. Red box indicates the prior information on temperature and time. Middle and lower plots show model predictions. Legend for data prediction plots can be found in the figure.

Figure 8: Location map of published low temperature thermochronology data for southwest Africa: AFT data only – Wildman et al., 2015, Green et al., 2015, Brown et al., 2014, Kounov et al., 2009, Tinker et al., 2008a, de Wit, 1998; AHe data only – Stanley et al., 2013; AFT and AHe data - Wildman et al. (this study), Kounov et al., 2013. Drainage network taken from Dollar, (1998).

Figure 9: Coast perpendicular sections (see Fig. 1) with predictions on magnitudes of denudation over time intervals since 150 Ma and sample/profile models used to derive these estimates. Data within 7.5km either side of the section trace was projected at 90° onto the line of section. Denudation estimates are made directly from thermal history models generated by

inverting data from this study and from Wildman et al., (2015). Denudation is estimated using an assumed geothermal gradient of 25 °C/km. Samples comprising multi-sample inversions are as follows, Kamiesberg Upper: 8832-78, 8832-79, SA12-38, SA12-37 and SA12-36; Kamiesberg Lower: SA12-35 and SA12-33; NQ Plateau: 8832-75 and SA12-52; NQ Coast: NQ12-15, NQ12-16, NQ12-17 and NQ12-18.

Figure 10: Cartoon representation of the post-Jurassic tectonic evolution of the southwest African continental margin. (a) Rifting during the early Cretaceous creates deep fault-bounded grabens in the proximal part of the margin and a fault bounded uplifted rift flank. Syn-rift erosion of uplifted fault blocks and elevated rift flanks occurs during extension due to the drop of regional base-levels to the nascent Atlantic Ocean with deposition resulting in developing basins. The uplifted rift flank is eroded by escarpment downwearing most likely with an interior drainage divide causing denudation to extend far inland. (b) Prolonged extension across the South Atlantic and regional intraplate stresses are augmented with extensional stresses induced from vertical motions related to the loading and unloading of the margin and the possible presence of a buoyant mantle upwelling beneath the continental interior also causing regional uplift of South Africa. The pre-existing structure and preferential orientation of these structures with the regional stress field at this time primes the faults for reactivation. Brittle tectonics may have extended into the offshore domain with gravitational slumping occurring further ocean-ward due to a regional uplift of the continent. Palaeostress tensors taken after Viola et al (2012).

Figure 11: Top left and top right: key Late Cretaceous – Early Cenozoic magnetic anomalies on WGM2012 global model gravity maps of ocean flow lines for the Mid-Atlantic Ridge and Southwest Indian Ridge. Bottom left: Spreading rates of South Atlantic opening for different plate reconstruction models from 150 Myr to the present for a point currently located at 57.2°W, 36°S in a reference frame that keeps Africa fixed (after Colli et al., 2014). Africa (NGU) and Africa (UTIG) show the absolute velocities of Africa in a mantle reference frame for an approximately conjugate point in the Orange Basin, at 15°E, 27.5°S. See Colli et al., (2014), and references there in for a full discussion of each individual plate reconstruction

model. Bottom right: location of the African plate finite rotation pole over time (after Pérez-Díaz and Eagles, 2014). Ellipses indicate 95% confidence intervals on location. Coloured lines correspond to time intervals beneath histogram of spreading rate velocity.

Table 1: Results of apatite fission track analysis. a: $p_{i,s,d}$ are track density of induced, spontaneous, dosimeter tracks. b: $N_{i,s,d}$ are the number of induced, spontaneous and dosimeter tracks counted. c: p-value of the chi-sq age homogeneity test (Galbraith, 2010). d: Dpar measurements are etch pit diameters used as a proxy for the influence of chemical composition on track annealing (Donelick et al., 2005). Between three to five Dpar measurements were measured for each dated single grain. e: Uranium content estimated using EDM. f: Central AFT ages calculated with TrackKey (Dunkl, 2002) with 1σ standard error. Ages were calculated using a $\zeta = 316.7 \pm 10.5$ (317.3 ± 11.1 for samples JN2 and JN3) for a standard IRMM540 glass g: Dispersion is the standard deviation of the true single-grain ages as a percentage of their central age (Galbraith, 2010); h: SD is the standard deviation of measured horizontal confined track lengths. i: mean track length after individual track length measurements are corrected for their orientation to the c-axis after Ketcham., (2005). j: HCT = Horizontal Confined Track. Analytical details can be found in SI-1 and SI-2. For details on sample lithology see SI-8.

Table 2: Results of apatite (U-Th-Sm)/He analysis. a: eU (effective uranium) is calculated as $eU_{ppm} = [U_{ppm}] + (0.235 * [Th_{ppm}])$. b: T = Number of terminations identified on crystal. c: L & W = Length and Width of crystal or crystal fragment. d: R^* = spherical equivalent radius calculated as $R^* = (3 * (RL)) / (2 * (R + L))$ where $R = W/2$. e: correction factor after Farley et al. (1996), assuming homogeneous distribution U and Th. f: Estimate Uncertainty is equal to 1σ analytical uncertainty, which include error propagated from U, Th, Sm and He measurement uncertainties, plus an additional 10% which is the standard deviation (reproducibility) of repeat analysis of Durango apatite standards. g: Corrected AHe age = Raw AHe age/Ft). Analytical details can be found in SI-4. For details on sample lithology see SI-8.

Table 3: Summary of the different factors that contribute to single-grain AHe age dispersion and the estimated magnitude of dispersion introduced by each of these factors.

Author Manuscript

(a)

(b)

(c)

(a) Early Cretaceous (syn-rift)

(b) mid-Late Cretaceous (post-rift)

This article is protected by copyright. All rights reserved.

Sample	Long. (°)	Lat. (°)	Elev. (m)	ρ_s^a (10^6cm^{-2})	N_s^b	ρ_i^a (10^6cm^{-2})	N_i^b	ρ_d^a (10^6cm^{-2})	N_d^b	$P(\chi^2)^c$	D_{par}^d (μm)	$[U]^e$ (ppm)	C. AFT Age ^f (Ma)	$\pm 1\sigma$	Disp. ^g (%)	#Xtls	Measured			c-axis correction ⁱ			#HCT ^j
																	MTL (μm)	$\pm 1\sigma$	SD ^h	MTL (μm)	$\pm 1\sigma$	SD ^h	
JN2	18.05	-30.68	376	42.7	3605	100.2	8451	19.8	14137	0.01	2.13	67.6	132.2	3.6	13%	20	13.73	0.11	1.08	14.74	0.12	0.79	103
JN3	17.38	-29.90	418	24.2	1790	64.7	4796	19.9	14137	0.81	1.91	44.1	116.8	3.2	10%	20	13.62	0.11	1.10	14.72	0.12	0.78	103
NQ12-01	17.99	-30.55	238	15.0	1365	40.4	3675	14.6	16086	0.04	1.49	37.6	85.4	3.4	19%	20	13.39	0.13	1.32	14.65	0.14	0.80	100
NQ12-03	17.95	-30.46	582	11.1	1163	29.8	3127	14.7	16086	0.57	1.83	27.0	86.0	3.0	17%	20	14.01	0.11	1.10	15.03	0.12	0.76	101
NQ12-04	17.93	-30.40	720	7.2	1324	21.2	3898	16.6	16348	0.00	1.65	27.8	89.6	4.1	20%	19	13.85	0.10	0.98	14.95	0.11	0.63	100
NQ12-06	17.86	-30.34	690	26.7	2080	76.9	5989	16.5	16348	0.32	1.62	65.3	90.2	2.4	12%	19	12.50	0.13	1.39	13.73	0.14	1.04	115
NQ12-07	17.82	-30.32	688	29.6	2511	72.0	6095	14.8	16086	0.12	1.92	65.0	96.0	2.6	14%	20	13.63	0.10	1.23	14.74	0.11	0.82	151
NQ12-08	17.80	-30.34	598	23.1	2283	60.9	6012	16.5	16348	0.06	2.98	49.5	98.7	3.0	14%	19	13.96	0.13	1.23	14.95	0.14	0.90	73
NQ12-09	17.77	-30.34	355	12.6	1153	31.8	2906	17.1	16086	0.49	2.49	25.8	106.6	3.8	15%	20	13.60	0.19	1.92	14.74	0.21	1.20	101
NQ12-10	17.75	-30.36	239	5.9	208	19.2	678	16.5	16348	0.50	3.42	16.3	79.7	6.3	25%	9	14.35	0.22	1.05	15.18	0.23	0.83	22
NQ12-11	17.72	-30.37	416	4.6	477	11.2	1167	17.2	16086	0.82	2.07	10.1	110.4	6.0	23%	20	13.51	0.16	1.43	14.64	0.17	0.90	38
NQ12-12	17.68	-30.36	388	31.2	1649	96.3	5096	16.5	16348	0.10	2.89	83.4	83.7	2.7	16%	18	13.45	0.15	1.50	14.67	0.16	0.97	106
NQ12-13	17.62	-30.36	354	7.8	747	21.8	2082	14.9	16086	0.28	1.45	19.3	84.0	4.1	20%	20	13.59	0.13	1.27	14.77	0.14	0.91	100
NQ12-15	17.30	-30.32	38	4.5	373	11.7	966	15.0	16086	0.17	2.06	10.8	91.5	6.3	35%	20	14.05	0.15	1.45	15.10	0.16	0.91	94
NQ12-16	17.27	-30.29	11	8.5	643	22.0	1654	15.2	16086	0.01	1.87	19.5	93.5	6.1	34%	20	13.70	0.15	1.22	14.80	0.16	0.83	65
NQ12-17	17.29	-30.36	5	8.6	914	21.7	2316	16.8	16086	0.14	1.82	17.5	104.9	4.6	23%	20	13.02	0.19	1.91	14.07	0.21	1.33	100
NQ12-18	17.28	-30.34	5	8.9	1161	21.8	2846	16.9	16086	0.02	2.11	16.6	108.4	4.9	23%	20	13.88	0.19	1.30	14.92	0.20	0.88	100
NQ12-19	17.64	-30.28	250	5.2	754	11.8	1723	15.4	16086	0.64	1.90	10.8	105.8	4.6	22%	20	13.40	0.19	1.48	14.55	0.20	1.06	82
NQ12-20	17.72	-30.20	473	22.3	1556	53.5	3723	15.6	16086	0.18	2.28	45.3	102.5	3.6	15%	19	13.77	0.12	1.28	14.82	0.13	0.87	108
NQ12-21	17.77	-30.21	665	24.6	1468	80.9	4831	17.3	16086	0.00	1.82	54.6	83.2	4.0	23%	17	13.67	0.19	1.41	14.57	0.20	0.98	102
NQ12-23	18.52	-31.23	300	9.0	779	40.3	3474	16.5	16348	0.18	1.56	32.0	58.3	2.6	29%	20	13.83	0.13	1.31	14.56	0.14	0.90	103
NQ12-24	18.35	-31.13	400	20.8	1164	58.3	3259	16.5	16348	0.93	2.27	46.8	92.7	3.2	12%	20	14.20	0.09	0.97	15.15	0.10	0.66	108
NQ12-25	17.93	-30.21	740	7.1	772	15.6	1700	15.8	16086	0.64	2.46	14.3	112.6	4.9	19%	20	14.01	0.14	1.45	14.98	0.15	0.96	112
NQ12-26	17.93	-30.13	850	37.4	2822	90.4	6823	16.5	16348	0.17	2.43	79.7	107.2	2.7	12%	20	13.94	0.12	1.21	14.93	0.13	0.80	103
NQ12-27	17.88	-30.03	600	18.5	2167	44.5	5213	16.2	16086	0.43	1.91	38.0	105.8	2.7	15%	23	13.37	0.13	1.35	14.61	0.14	0.85	104
NQ12-28	17.90	-30.17	650	35.6	2282	95.2	6107	17.4	16086	0.35	2.15	76.0	102.1	2.5	12%	20	12.89	0.16	1.83	14.28	0.18	1.11	128
NQ12-29	17.78	-30.09	720	28.7	586	74.8	1527	16.5	16348	0.17	2.20	61.6	99.1	5.6	17%	10	13.81	0.11	1.15	14.85	0.12	0.80	111
NQ12-30	17.83	-30.15	550	31.6	2426	74.9	5758	16.4	16348	0.40	1.75	58.2	108.5	2.6	12%	20	13.86	0.11	1.13	14.99	0.12	0.75	110
NQ12-33	18.08	-30.17	1050	13.9	1319	33.4	3174	16.3	16086	0.03	1.64	29.1	105.8	4.5	21%	20	13.56	0.14	1.40	14.76	0.15	0.97	100
NQ12-34	18.16	-30.18	1000	15.0	949	34.8	2203	16.3	16438	0.05	1.93	30.6	110.0	5.3	22%	20	13.21	0.15	1.49	14.49	0.16	0.98	106
SA12-22	19.08	-29.93	922	35.9	2565	81.3	5812	14.0	16086	0.01	2.20	76.6	97.2	3.1	15%	20	14.04	0.14	1.48	15.07	0.15	0.97	108
SA12-27	18.70	-30.23	987	5.6	334	24.1	1434	16.6	16348	0.71	1.84	22.4	60.9	3.7	27%	18	12.66	0.37	1.95	14.16	0.41	1.38	27
SA12-30	18.00	-30.53	258	9.4	1012	33.4	3600	16.6	16348	0.06	2.94	27.0	73.2	3.3	19%	20	14.17	0.22	1.20	15.10	0.23	0.80	100
SA12-32	18.06	-30.53	351	6.0	106	20.4	360	16.2	16348	0.85	1.70	18.6	75.1	8.3	8%	3	10.90	0.19	1.54	13.28	0.23	0.95	2
SA12-33	18.06	-30.49	400	5.2	353	12.3	836	16.6	16348	0.75	1.71	9.9	110.0	7.0	27%	23	13.48	0.28	1.33	14.63	0.30	0.95	23
SA12-35	18.06	-30.45	605	3.4	232	6.5	442	14.1	16086	0.62	1.87	6.1	119.0	10.0	43%	17	13.71	0.22	1.31	14.85	0.24	0.86	35
SA12-36	18.06	-30.45	707	3.6	288	10.4	822	16.6	16348	0.57	2.03	9.2	91.4	6.3	33%	21	13.90	0.17	1.58	14.98	0.18	0.93	82
SA12-37	18.06	-30.43	807	12.4	409	38.0	1256	16.6	16348	0.61	1.74	32.5	85.0	4.8	19%	12	13.43	0.34	1.68	14.64	0.37	1.13	24
SA12-38	18.07	-30.40	959	15.3	808	53.9	2839	16.6	16348	0.61	1.68	45.3	74.4	3.0	17%	20	13.83	0.13	1.31	14.88	0.14	0.89	100
SA12-47	18.23	-30.39	1064	5.0	484	11.2	1086	14.3	16086	0.22	2.06	12.5	100.8	5.9	34%	21	12.40	0.22	1.80	13.89	0.25	1.22	74
SA12-51	18.42	-30.32	1066	34.9	3341	75.0	7183	14.4	16086	0.00	2.30	67.7	107.7	3.8	16%	20	12.81	0.15	1.19	14.24	0.17	1.19	139
SA12-52	18.46	-30.32	1065	3.2	208	7.1	461	14.5	16086	0.99	2.20	7.3	102.8	8.6	24%	16	13.67	0.19	1.42	14.68	0.20	1.01	55

This article is protected by copyright. All rights reserved.

Dispersion factor	Contribution to dispersion	Comments	References
Grain size (R^*)	$\pm 50 - 100$ %	R^* of c. 50–100 μm , depending on the thermal history.	Reiners and Farley, 2001; Brown et al., 2013.
Radiation Damage (eU)	$\pm 50 - 100$ %	Depending on thermal history and difference in the eU content of individual grains can exceed 200%.	Gautheron et al., 2009; Flowers et al., 2009; Brown et al., 2013.
Fragment length	$\pm 7 - 60\%$	Depending on length of fragment and original position of fragment relative to the whole grain.	Brown et al., 2013.
Zonation	$\pm 10 - 15\%$,	30–40% dispersion for some thermal histories and extreme heterogeneity, true age may be older or younger depending on whether the crystal has an enriched core or rim, respectively.	Meesters and Dunai, 2002a ; Fitzgerald et al., 2006;
Implantation	older by 60%	Effect will vary depending on the size of 'bad neighbour', [U] and [Th] content of 'bad neighbour' and whether or not the apatite is in contact with one or many 'bad neighbours'.	Farley et al., 2011; Ault and Flowers, 2012.
alpha-ejection factor (Ft)	older age by 3–8%	Depending on thermal history and zonation patterns.	Spiegel et al., 2009; Gautheron et al., 2012.
Sm	0.1–10%	In certain situations (low [U] and [Th], anomalously high [Sm]) can exceed 25%.	Meesters and Dunai, 2002a,b; Gautheron et al., 2012;
Mineral Inclusions	older by < 10%	More than a few % only when inclusions are quite large (<0.1 of the grain size) and/or have an unusually high U and Th content (i.e. >1000 times that of the apatite).	Brown et al., 2013.
			Fitzgerald et al., 2006; Vermeesch et al., 2008.
			Vermeesch et al., 2007.