1

2 Received Date : 05-May-2015

3 Revised Date : 30-Jan-2016

4 Accepted Date : 03-Feb-2016

5 Article type : Original Article

6

7

8 **Trends and management implications of human-influenced life-history changes in**
9 **marine ectotherms**

10 **Human-induced life-history changes in marine ectotherms: trends and management**
11 **implications**

12

13 *Asta Audzijonyte¹, Elizabeth Fulton², Malcolm Haddon², Fay Helidoniotis², Alistair J. Hobday², Anna*
14 *Kuparinen¹, John Morrongiello³, Anthony D.M. Smith², Judy Upston² and Robin S. Waples⁴*

15

16 ¹ Department of Environmental Sciences, POB 65, University of Helsinki, FIN-00014, Finland

17 ² CSIRO Oceans and Atmosphere, Castray Esplanade, Hobart, Tasmania, 7001, Australia

18 ³ School of BioSciences, University of Melbourne, Victoria, 3010, Australia

19 ⁴ NOAA Fisheries, Northwest Fisheries Science Center, 2725 Montlake Blvd. East Seattle, WA 98112,
20 USA

21

22 **Corresponding author:**

23 Asta Audzijonyte

24 Department of Environmental Sciences

25 University of Helsinki

26 PO Box 65

27 FIN-00014 Helsinki Finland

28 Telephone: +61450782631

29 Fax: +358 5 2250570

30 Email: asta.audzijonyte@helsinki.fi

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/faf.12156](https://doi.org/10.1111/faf.12156)

This article is protected by copyright. All rights reserved

Running title: marine ectotherm life-history changes

Abstract

Evidence is accumulating that many marine ectotherms are undergoing rapid changes in their life-history characteristics. These changes have been variously attributed to fisheries-induced evolution, inhibited adult growth rate due to oxygen limitation at higher temperatures, and plastic responses to density dependence or changes in ocean productivity. Here we review the diverse underlying mechanisms by which plastic and evolutionary responses to climate change and fisheries are likely to produce similar life-history trends in harvested marine ectotherms, leading to faster life-histories with earlier maturation and smaller adult size-at-age. While mechanistically understanding these growth and maturation changes may be difficult, it is becoming clear that changing life-histories will lead to modified population dynamics, productivity and natural mortality of the affected species. We discuss how the observed and expected life-history changes could affect the assumptions and uncertainty within single and multi-species models currently used in marine ecosystem management, highlighting that models which allow for dynamic life-history traits often report significantly different estimates of stock biomass. Given that both climate and harvest induced life-history changes are likely to intensify and possibly amplify each other, there is an urgent need to adequately assess the implications of faster life-histories for marine ecosystem management. This is especially true for data poor stocks, where growth and maturation are not regularly assessed. Targeted monitoring can be used to inform responsive management, but for improved sustainability outcomes, a precautionary approach to management that is robust to life-history trends is advised.

Keywords: climate change, evolution, fisheries, phenotypic response, plasticity, physiology

Table of contents

Introduction

Plastic and genetic determinants of life-history strategies and important trade-offs

Drivers and empirical evidence on life-history changes in marine ectotherms

Harvesting: short-term density-dependent growth changes

Global warming: response to productivity change

Global warming: response to temperature change

Harvesting: evolutionary response

Summary: climate change and fishing are likely to lead to faster life-histories

67 Implications for population persistence: life-history changes will lead to increased stochasticity

68 Management implications of changes in life-history characteristics and diversity

69 *Effects of life-history changes on fish stock dynamics*

70 *Impact of life-history trends in single species management*

71 *Life-history trends in multi-species models*

72 How to address changes in life-histories of marine organisms - a plan of action

73 Acknowledgements

74 References

75

76 **Introduction**

77

78 Most of the world's oceans are now believed to be affected by human activities at a medium to very high
79 level (Chevin *et al.* 2010; Halpern *et al.* 2012). These anthropogenic pressures, such as fishing, climate
80 change, pollution, and habitat degradation, are increasing in distribution and intensity (Merrie *et al.* 2014;
81 IPCC-AR5 2014). Organisms exposed to such rapid changes in their environment and to new or
82 intensified selective pressures are forced to adapt to new conditions through shifts in distributions,
83 phenotypic plasticity and contemporary evolution (Hoffmann and Sgro 2011). Among other aspects, such
84 adaptive responses will be strongly manifested through changes in growth and reproductive schedules, or
85 life-histories.

86

87 Life-histories determine species energy allocation to growth and reproduction, and describe traits such as
88 age and size of maturation, reproductive effort and maximum adult size and age. Some of the most rapid
89 changes in life-history traits have been observed in commercially harvested fishes (Jørgensen *et al.* 2007;
90 Devine *et al.* 2012), where the rate of change can be as high as 1-4% per year for size and age at maturity
91 (Audzijonyte *et al.* 2013a) and 2.5% per year for growth (Morrongiello and Thresher 2015). The relative
92 importance of genetic and plastic components in this change and the drivers have been intensely debated
93 (e.g. Olsen *et al.* 2004; Kraak 2007; Therkildsen *et al.* 2013; Rogers *et al.* 2011; Baudron *et al.* 2014), yet
94 regardless of the underlying mechanisms, the observed trends will have important implications for marine
95 ecosystem management, including management of individual species. Given that life-history parameters
96 for harvested stocks will be affected by both fishing and climate change simultaneously, understanding
97 the combined effects is an important but neglected priority (Planque *et al.* 2010).

98

99 This review aims to highlight how global environmental change and fishing are affecting life-histories of
100 marine ectotherms. Naturally, responses to climate change or fishing involve many other features, such as
101 changes in distributions, phenology or behaviour, but these have been discussed elsewhere (e.g. Brander
102 2013; Poloczanska *et al.* 2013). In contrast, while there have been important studies on how either

harvesting or climate change are affecting life-histories separately (e.g. Heino *et al.* 2013; 2015; Holt and Jørgensen 2015), we believe that insufficient attention has been paid to the fact that both forces are likely to lead to convergent trends and possibly amplify each other. Hence the main message of this review is that the diverse mechanisms caused by harvesting and climate change will, in many species, lead to faster ectotherm life-histories, with earlier maturation and smaller adult body sizes. To demonstrate this point we first summarize the dominant trade-offs determining life-history strategy and explore how these trade-offs might be useful to predict likely changes in life-history traits. Second, we explore how the different contemporary environmental drivers, in particular climate change and fishing, are likely to produce similar life-history trends. Third, we describe how these trends might increase fluctuations in population abundance and hence reduce resilience to environmental change and exploitation. Finally, we discuss how contemporary life-history changes might affect the management advice and outcomes of single and multiple species models, and suggest areas for further scientific investigation.

Plastic and genetic determinants of life-history strategies and important trade-offs

A study of directional life-history change in natural populations cannot avoid questions about the underlying nature (plastic, genetic or both) and hence reversibility of the observed trends. From an evolutionary perspective, life-histories are constantly being tuned to optimise fitness under a given juvenile and adult mortality regime; for example, different life-histories are expected in stable versus highly stochastic environments (Roff 1992). Studies of heritability of life-history traits in aquaculture, laboratory experiments and wild populations typically estimate that about 20 to 40% of the phenotypic variability is genetically determined (Conover and Munch 2002; Carlson and Seamons 2008). Rapid evolutionary changes have also been demonstrated in wild populations and linked to ecological changes in aquatic systems (Reznick *et al.* 1997; Palkovacs *et al.* 2012). At the same time, growth rates and maturation schedules are highly plastic, showing rapid responses to changes in resource availability, temperature, or density (Enberg *et al.* 2012; Baudron *et al.* 2014; Morrongiello and Thresher 2015). While this plastic *versus* genetic distinction is important when assessing possible reversibility of life-history trends, disentangling the two components in wild populations is notoriously difficult, because realised life-histories will depend on the combined effect of genetic background, maternal effects, and phenotypic plasticity (e.g. Gienapp *et al.* 2008; Kuparinen *et al.* 2009; Hansen *et al.* 2012). Nevertheless, the general trade-offs among life-history traits governing growth, maintenance and reproduction hold regardless of the driver causing the change (Charnov 1993; Jensen 1997), and this allows us to predict the general nature of correlated responses.

All organisms have limited amounts of energy that must be allocated to different body functions; increased energy allocation to reproduction takes energy away from somatic growth, and vice versa.

Recognition of these trade-offs has led to the development of life-history theory (e.g. Roff 1992), which can be used, among other things, to infer age at maturation and energy-based life-history traits from individual growth trajectories (Brunel *et al.* 2013). Assuming that a fixed amount of total energy is available, life-history trade-offs provide some general rules on how changes in one trait will lead to changes in other correlated traits (Fig 1). For example, earlier maturity trades off with later reproductive output by reducing somatic growth and increasing natural mortality through survival costs of reproduction (Bell 1980). These reproductive costs include increased mortality due to reproductive changes in behaviour (competition for mates, nest guarding, spawning migrations), reduced feeding, energetic investment in gonads instead of somatic growth, and less energy allocation to immune functions (Jørgensen and Fiksen 2010; Jørgensen and Holt 2013). The trade-off between growth and maturation in animals with indeterminate growth is typically seen as a negative correlation between intrinsic growth rate against size at maturity and adult body size (Charnov 1993). Accounting for these trade-offs has a large impact on the outcomes of models aiming to predict population level consequences of life-history changes (see below).

The trade-offs in the energy allocation for different functions (growth, reproduction, maintenance) are also at the heart of correlations among species and population life-history parameters observed in broadly defined taxonomic categories. These correlations are known as life-history invariants, and provide some general rules about the expected relationships between asymptotic length (L_{∞}), age and size-at-maturation, growth rate, natural mortality and length-fecundity relationship (e.g. Beverton 1992; Charnov *et al.* 2013). While these relationships encompass large variation, they nevertheless provide useful insights into population parameters in data-poor stocks and are used to inform ecological risk assessments (Patrick *et al.* 2010; Hobday *et al.* 2011a). They are also implemented in global databases, such as FishBase. A promising new approach to reduce variation and improve predictive value of life-history invariants is to derive these life-history ratios separately for species with different life-history strategies (Prince *et al.* 2015).

In this review we focus on life-history and growth parameters and for illustration purposes use the von Bertalanffy (VB) growth model (von Bertalanffy 1957, see also Pardo *et al.* 2013) (Fig. 1). The two VB parameters of interest here are asymptotic length (L_{∞}) and the rate at which growth approaches the asymptote (k), which is often referred to as a growth coefficient. We should note that, as with any model, the VB model has its shortcomings. First, k is difficult to interpret ecologically, and cannot easily be compared across species (see Pauly 2010, who also offered a conversion of k to an ecologically meaningful growth performance index that is comparable across species). Second, in heavily fished stocks where few individuals survive to the older and larger age classes, the estimation of L_{∞} can be problematic (Haddon 2011). Yet, despite these limitations, the model is commonly used in fisheries, due

to its flexibility and generality to summarise somatic fish growth. The model is also used to describe many of the life-history invariants and correlations between L_{∞} and k with natural mortality (M), carrying capacity (K) and per-capita population growth rates (r), i.e. the key parameters defining population productivity and ability to sustain harvesting (Dulvy *et al.* 2004; Hobday 2011b). We nevertheless emphasise that our prediction on faster life-histories does not depend on the VB or any specific growth model, as the two key aspects of this prediction are simply earlier maturation and smaller adult body size.

Drivers and empirical evidence on life-history changes in marine ectotherms

Harvesting: short-term density-dependent growth changes

The effects of harvesting on age structure, population dynamics and density-dependent processes of fished stocks are at the core of basic fisheries models and stock-recruitment relationships, yet a lack of life-history data and poor understanding of interconnected processes remain a limitation of such models (Rose *et al.* 2001; Lorenzen 2016). Complex growth patterns are difficult to model explicitly, and most fisheries models that use growth models to predict body weight-at-length or -age either treat the growth parameters as fixed, or estimate the parameters within the model and assume they do not vary over time (Haddon 2011; Kuparinen *et al.* 2012, but see Whitten *et al.* 2013 for a case where cohort-specific growth is used to account for density dependence on adult growth). Typically, strong density dependence is assumed to operate during the juvenile stages and is modelled through stock-recruitment relationships. In contrast, density effects on the age groups captured by fisheries are usually considered to be weak (Rose *et al.* 2001; Lorenzen 2016).

There seems to be no consensus as to how density-dependent processes might affect life-history parameters (Prince *et al.* 2015). Lorenzen and Enberg (2002) found indications of density dependence on adult growth in many marine stocks and estimated that for species in a 20-45 cm size range, a 100% increase in stock biomass will lead to a 10% decrease in L_{∞} . In the evolutionary and physiological interpretation of the VB growth model presented by Charnov *et al.* (2013), L_{∞} or maximum body size, is determined by both food availability and/or changes in maintenance and reproduction, whereas k , or early growth rate, is affected only by the allocation to maintenance and reproduction. In this formulation, increased food availability is expected to increase L_{∞} , but should not change k . Empirical data provide examples of both situations (Table 1). High abundance of certain age groups had negative effects on juvenile growth rates in sockeye salmon (*Oncorhynchus nerka*, Salmonidae) (Crozier *et al.* 2010) and has been linked to a decrease in L_{∞} from about 100cm to 40 cm in the Baltic Sea cod (*Gadus morhua*, Gadidae) (Svedang and Hornborg 2014). Reduced population sizes and thus lower density were correlated with increased k in southern Bluefin tuna (*Thunnus maccoyii*, Scombridae) (Polacheck *et al.*

2004), but decreased k and increased L_{∞} in southern king weakfish (*Macrodon atricauda*, Sciaenidae) in Southern Brazil (Cardoso and Haimovici 2011).

One limitation of these studies is that they do not statistically test alternative hypotheses that could explain the observed growth patterns, as in addition to density dependence they could include changes in reproductive investment, maturation age and length, maintenance costs and many environmental factors. This underscores the general difficulties in attributing empirically observed life-history changes to a specific driver. We suggest that, for many stocks, in the absence of long-term population trends, density-dependent processes will generally fluctuate over time in response to changes in harvesting levels and recruitment and will not be as important as the long-term trend towards faster life-histories caused by global warming or fisheries-induced evolution.

Global warming: response to productivity change

In this section we focus on the life-history responses to changes in ocean productivity. The effects of global warming on marine organisms are multifaceted (Sarmiento *et al.* 2004; Doney *et al.* 2012), and also include ocean acidification, yet the impacts of ocean acidification on life-history traits remain speculative (Munday *et al.* 2013). We also do not discuss the global warming effects on the distributions of marine organisms, as they have been covered in a number of recent reviews (Poloczanska *et al.* 2013; Bates *et al.* 2014).

Changes in ocean temperature, circulation and mixing all affect nutrient supply and primary productivity (Poloczanska *et al.* 2008; Brown *et al.* 2009; Taucher and Oschlies 2011). A number of modelling studies find that climate change will lead to reduced ectotherm body size through nutrient limitation due to reduced mixing of water layers and smaller size of phytoplankton (e.g. Woodworth-Jefcoats *et al.* 2015). Yet, despite the important implications that productivity shifts are expected to have on marine ecosystem dynamics, our ability to model and forecast the magnitude or even direction of the productivity change is still limited (Polovina and Woodworth 2012; Doney *et al.* 2012). Even greater spatial variability and uncertainty exists in coastal areas. In a semi-global analysis, Barange *et al.* (2014) predicted an overall minor increase in shelf productivity (<10%), with increased productivity at high latitudes and decreased productivity at low and mid latitudes, but considerable regional variation due to local circulation patterns and terrestrial inputs.

Changes in ocean productivity and hence food supply will operate on life-history traits in a broadly similar manner to the density-dependent effects discussed above, except that their combined effect is likely to be opposite – as productivity increases, density dependence is expected to decline, assuming the stock abundance remains at a similar level and there are no large changes in inter-specific interactions.

Increased productivity is likely to amplify the temperature-induced trend towards faster growth and earlier maturation, whereas decreased productivity will lead to slower growth rates. In many cases these effects might be difficult to separate from the response to increased temperature alone, especially in cold regions where higher temperatures often mean higher productivity. As with density dependence, rapid response to e.g. increased productivity through faster growth rates could mask long-term trends due to physiological or evolutionary factors (Fig. 1 in Crozier and Hutchings 2014; Thresher *et al.* 2014; Kokkonen *et al.* 2015). While there is great uncertainty in forecasting productivity changes, its monitoring is considerably easier and is routinely done (e.g. through chlorophyll *a* mapping via satellite images). Hence, up-to-date information on productivity could be a part of routine management (Pethybridge *et al.* 2015) and also incorporated into management decision processes (see Fig. 3).

Global warming: response to temperature change

A decline in average individual body size has been suggested as a “third universal ecological response to global warming” (Daufresne *et al.* 2009; Cheung *et al.* 2013). For ectotherms this ecological response is often based on the temperature-size rule (Atkinson 1994) that describes the tendency for increased juvenile growth (increase in k and hence faster approach towards asymptotic size) and developmental rates combined with the decreased adult body size (L_{∞}) at higher temperatures. Studies that have explicitly tested the temperature-size rule in marine invertebrates indicate that one degree of warming results in a 0.5–4% decrease in body size (Daufresne *et al.* 2009; Irie and Fischer 2009). Similar patterns have been observed in marine fishes with, for example, L_{∞} declining by an average of 16% across eight North Sea fish stocks with around 2°C of warming (Baudron *et al.* 2014). Different physiological and evolutionary mechanisms could produce this theoretically predicted and empirically observed pattern, but more importantly they have similar life-history consequences of earlier maturation and a decrease in L_{∞} (Table 2).

The physiologically-related pathways in Table 2 are broadly grouped into the theory of oxygen- and capacity-limited tolerance, which states that growth and developmental rates are determined by the aerobic performance or oxygen supply to tissue and respiration rates that are strongly affected by temperature (Portner and Knust 2007). Growth rate is therefore expected to increase with temperature from a lower critical value to an optimum determined by aerobic scope, after which temperatures become increasingly too warm and aerobic performance declines (Portner and Knust 2007; Neuheimer *et al.* 2011). The aerobic scope curves are likely to be different for different life stages, and evidence is accumulating that juveniles of ectotherms have higher temperature optima than adults (Ohlberger 2013). Whilst recent work suggests that the optimal temperature for growth is likely to be less than that for aerobic scope due to temperature-dependent costs of locomotion and digestion (Holt and Jørgensen 2015), performance still displays a dome-shaped relationship to temperature and the different effect on

adults and juveniles still holds. Finally, growth and development are governed by different processes and scale differently to temperature, where developmental rates appear to be more sensitive to temperature than growth (Forster *et al.* 2011), and the size-fecundity relationship changes with temperature (Arendt 2011). In combination, these physiological responses lead to the general global-warming-induced trend towards maturation at smaller size and age, and decreased lifetime reproductive output (Table 2), although specific conditions might alter the general trend. Modelling indicates that half of the predicted declines in marine fish body size can be attributed to changes in environmentally-dependent physiology (Cheung *et al.* 2013).

Discussion on climate change effects would be incomplete without mentioning the evolutionary response to the strong selective force that rapid global warming represents (e.g. Hill *et al.* 2011; Pespeni *et al.* 2013). For example, Rummer *et al.* (2014) found that all six studied coral reef fish species had significantly reduced performance at temperatures 3°C above ambient, leading the authors to conclude that many tropical reef species have little capacity for plastic responses to rising water temperatures. At higher temperatures smaller individuals often perform better than larger ones (Portner and Farrell 2008), which results in selection for smaller body size. To some extent, adults might be able to avoid temperature increases by moving into deeper water layers (e.g. Dulvy *et al.* 2008). However, in many cases these movements are limited by topography, suitable habitat and prey, and indeed globally most changes in species distributions are seen as poleward shifts (Poloczanska *et al.* 2013; Cheung *et al.* 2014). A strong correlation between temperature and lifespan of ectotherms, both at inter- and intra-specific level, suggests that a 3°C increase in global temperatures is expected to shorten the lifespan by 8-42% (Munch and Salinas 2009). Shorter lifespan will decrease lifetime reproductive output, and that will lead to selection towards earlier reproduction and higher investment earlier in life, and consequently to smaller adult body sizes. Despite these predictions, empirical evidence for genetic adaptation to climate change is still very limited (Teplitsky and Millien 2014), which, as with fisheries-induced evolution, can mostly be attributed to the fact that convincingly demonstrating evolutionary change in wild populations is extremely difficult (Crozier and Hutchings 2014).

Harvesting: evolutionary response

Most fishing gear is selective with respect to size, age or behaviour of individuals (e.g. Kuparinen *et al.* 2009; Garcia *et al.* 2012). Over time, fishing will disproportionally increase mortality of selected phenotypes, which can lead to fisheries-induced evolution. For many stocks, fishing mortality of individuals recruited to fisheries can be considerably higher than natural mortality, which means that fishing will act as a stronger selective force than natural processes for that life stage (Darimont *et al.* 2009). In some cases these fishery-selection differentials have been calculated from empirical data. For example, the fisheries-imposed selection differential on growth rate of two whitefish species in Swiss

lakes was -7% to -9% per generation (Nussle *et al.* 2011), selection for size-at-age of Alaska sockeye salmon was around -5 to -10 mm per year (Kendall *et al.* 2014), and selection for earlier migration dates of Columbia River sockeye salmon was around -1 day/year between 1950 and 2010 (Crozier *et al.* 2011). In each case, observed direction of change in the life-history trait matched expectations from the selection differentials, providing support for fisheries-induced evolutionary change.

Evidence of evolutionary change in life-history traits as a result of harvesting has been well demonstrated in experimental populations (Conover and Munch 2002; Van Wijk *et al.* 2013; Uusi-Heikkilä *et al.* 2015), and subsequently in wild populations, where changes in frequencies of genes related to life-history traits matched expectations from life-history theory and were best explained by fishing (Jakobsdóttir *et al.* 2012; Therkildsen *et al.* 2013). Nevertheless, evidence correlating life-history changes to fishing intensity still mostly comes from time series analyses of wild stocks. These consistently show decreasing size and age at maturity, smaller adult length-at-age, and to some extent also increasing reproductive investment (Sharpe and Hendry 2009; Devine *et al.* 2012) (Table 1). Rates of these trends vary between 0.1 to 4% per year and depend on fishing intensity, with an average rate of 1% per year for instantaneous long-term fishing mortality (F) of around 0.5 per year (Audzijonyte *et al.* 2013a). Age and size at maturation appear to be among the fastest changing traits, whereas increases in reproductive investment are considerably slower. Empirical phenotypic rates of change in life-history parameters in harvested stocks are on average four times faster than evolutionary rates inferred by simulations (Audzijonyte *et al.* 2013a), suggesting that around a quarter of the observed phenotypic change could be due to evolution; this estimate is consistent with the known heritability of life-history traits (0.2-0.4, Carlson and Seamons 2008).

Summary: climate change and fishing are likely to lead to faster life-histories

The theoretical predictions presented above, as well as empirical data, suggest that current human impacts on marine ectotherms will often lead to synergistic effects on their life-history trends (Table 1, Fig. 1). First, growth rates in early life stages (including k in the VB curve) will tend to increase due to physiological responses to warming. This increase will be amplified in areas where primary productivity increases. The trend towards faster juvenile growth might be counteracted by the fisheries-induced evolution (FIE) towards slower growth, although empirical evidence on how FIE will affect juvenile growth is inconclusive. Second, in many stocks we expect to see decreased maturation age. This is indeed one of the best documented life-history trends (Table 1). Decrease in maturation age will be driven both by physiologically faster juvenile growth due to warming and by evolutionary response to elevated adult mortality caused by higher temperatures (Table 2) and fishing (irrespective of how FIE affects juvenile growth). Third, earlier maturation and costs of reproduction, increased reproductive investment, and susceptibility of large fish to decreased oxygen supply in warming waters can be expected to result in smaller adult body size, including smaller L_{∞} . Of these three predictions, the trends in the juvenile growth

are most uncertain, but they will not have substantial effect on earlier maturation and decreased adult body size, as these are driven by several other mechanisms as well.

The majority of empirical examples reported in the literature confirm this expectation, regardless of the ultimate attribution of the causal factors (Table 1). Exceptions to this trend toward faster life-histories can occur in situations where climate change leads to temporary decrease in temperatures (e.g. some deep sea populations, Thresher *et al.* 2007) or where decreased productivity due to climate change will lead to food limitation in juveniles and hence outweigh the expected increase in early growth rates due to temperature and/or harvesting (e.g. some salmon streams, Crozier *et al.* 2011). Decadal-scale climatic oscillations can also override any warming trend on shorter time scales (e.g. Attrill and Power 2002). Finally, species interactions will alter these general trends through changes in natural mortality and resource availability (Ohlberger 2013). Different species will respond to fishing and climate differently, and the expected reductions in size will vary within the same geographic region. Subsequent shifts in species size composition, abundance and distributions will modify predator-prey interactions, further affecting realised life-histories. We cannot make precise theoretical predictions for these second-order responses, but they should be considered and explored (Audzijonyte *et al.* 2013b).

Implications for population persistence: life-history changes will lead to increased stochasticity

The directional trend towards faster life-histories discussed above will result in changes to natural mortality and population growth rates. However, before we consider management implications, it is important to mention that, in addition to directional trends, climate change and harvesting are also likely to lead to reduced diversity of life-history strategies through both short-term demographic and long-term evolutionary mechanisms (Table 3).

A number of recent studies have demonstrated that the combination of age truncation through fishing and increased environmental variation through climate change will amplify fluctuations in fish stock abundance (Hsieh *et al.* 2008; Shelton and Mangel 2011; Rouyer *et al.* 2012). There is an emerging consensus that the presence of multiple age classes increases the diversity of egg sizes, and the spatial and temporal distribution of spawning events, and hence serves as a bet-hedging strategy to maximise resilience (Secor 2007; Hixon *et al.* 2014; Shelton *et al.* 2015). The trend towards faster life-histories discussed here now means that in addition to the demographic truncation effect we should also consider the consequences of the physiological and/or evolutionary trend towards earlier maturation and smaller adult size, an area that remains largely unexplored. It is possible that earlier maturation may somewhat counteract the demographic truncation effect, by allowing individuals to reproduce earlier. However, the effects of life-history changes on population stability in the presence of fishing and climate change may

have many unexpected feedback mechanisms through changes in behaviour, age-specific mortality and fecundity (Jørgensen and Fiksen 2010; Audzijonyte *et al.* 2013b).

Directional selection from increased temperatures and size-selective fishing also means that we expect to see the narrowing of maturation and temperature reaction norms, and overall decrease in phenotypic and genetic diversity (Table 3). These outcomes potentially have large implications for population persistence. Life-history diversity provides an important “portfolio effect” (Schindler *et al.* 2010; Anderson *et al.* 2015), buffering stocks against environmental fluctuations. A population that is rich in genetic variation for life-history traits will be better able to mount an evolutionary response to changing conditions (Reed *et al.* 2011). The same concept applies to population complexes, such as in the salmon populations spawning in different rivers. The diverse life-history traits expressed by hundreds of sockeye salmon populations in Bristol Bay, Alaska, have possibly reduced the estimated fishery closure events by a factor of 10, compared to what would have occurred if populations had been homogenous (Schindler *et al.* 2010). The short-term buffering due to the “portfolio effect” can occur even in the absence of genetically-based population differences; all that is required is different plastic responses to environmental and habitat heterogeneity. However, longer-term benefits require inter-population genetic heterogeneity. The existence of a diverse portfolio of locally-adapted populations greatly increases the probability that at least some will contain genetic resources that will allow them to persist under radically different future conditions and in this way reduce the biomass fluctuations. Notably, such locally adapted populations can be just as important in widespread marine species (Nielsen *et al.* 2012).

Management implications of changes in life-history characteristics and diversity

Effects of life-history changes on fish stock dynamics

Population parameters that will change with faster life-histories include, among others, natural mortality, steepness of the stock-recruitment relationship, growth model parameters, age at 50% maturity and maximum size. Changes in these parameters will have implications for fisheries stock assessment. Intuitive prediction and broad inter-species comparisons (Jennings *et al.* 1998; Hutchings *et al.* 2012) would suggest that faster juvenile growth, earlier maturation and lower asymptotic size lead to an increased per capita population growth rate (r) and decreased carrying capacity (K) (Heino *et al.* 2013). In terms of fisheries outcomes, this implies that biomass at maximum sustainable yield (B_{MSY}) may decrease ($B_{MSY}=K/2$), fishing mortality (F_{MSY}) needed to achieve the MSY should increase ($F_{MSY}=r/2$), while the actual sustainable yield may not change much, since it is the product of both r and K ($MSY=rK/4$).

Heino *et al.* (2013) described how faster life-histories caused by fisheries-induced evolution (FIE) may affect r , K and fisheries reference points. They argued that if FIE increases a stock’s reproductive

potential through earlier maturation and shorter generation time, it will lead to increased steepness of the stock-recruitment relationship, meaning that higher recruitment occurs at lower biomass. Yet, if FIE reduces pre-recruit survival, through e.g. lower egg quality from smaller females, then the steepness of the stock-recruitment relationship will decrease. Similarly, FIE might change the ratio of spawning stock biomass (SSB) to total biomass. If fish mature earlier, then overestimation of maturation age will underestimate the SSB. However, if FIE reduces the weight of mature fish, then SSB might actually be overestimated if an old age-weight relationship is used (Fig. 2). Earlier maturation means that a cohort's biomass will peak earlier. This would lead to higher fishing mortality on older individuals in cases where selectivity remained unchanged (see the Baltic Sea cod case for a similar result due to density dependence effects; Svedang and Hornborg 2014).

These general predictions should apply regardless of whether faster life-histories are caused by fishing or global warming. However, it is essential that studies exploring the population level consequences of life-history changes account for the trade-offs between growth and reproduction (see section 2). While there is a general consensus that faster life-histories are likely to decrease carrying capacity (a decrease in adult body size of 5-20% led to a 5-30% decrease in K , see Enberg *et al.* 2009, 2010; Audzijonyte *et al.* 2013b; Kuparinen *et al.* 2014), the modelled effect of earlier maturation on r strongly depends on the assumptions about the survival costs of reproduction. As an example, a model of Atlantic cod (*Gadus morhua*, Gadidae) used by Enberg *et al.* (2010) predicted that FIE will increase the ratio of SSB to total biomass, simultaneously increasing the steepness of the stock-recruitment relationship. The authors argued that populations adapted to fishing would produce more recruits for a given level of total biomass, which is likely to increase r . In contrast, modelling studies that included some version of reproduction costs suggest that a combination of earlier maturation (1 to 3 years earlier) and smaller adult body size (ca 20% decrease in L_{∞} and L_{mat}) is likely to decrease r by 20-50% (Hutchings *et al.* 2005; Kuparinen *et al.* 2014; Audzijonyte and Kuparinen 2016). These simulations are also corroborated by empirical studies exploring r among closely related populations of iteroparous species, showing that natural mortality rather than maturation age is the major determinant of population growth rate (squirrels: Mcadam *et al.* 2007; gadoid fishes: Wright 2014). Hence if the benefits of earlier reproduction are outweighed by increased natural mortality (Fig. 1B) the overall effect of faster life-histories on population productivity (r) can be negative.

There are two ways in which the simulation studies presented above may still be too conservative. First, some of them considered the effects of FIE alone, yet our argument is that life-history effects of climate change and fisheries can often be synergistic and amplify each other. Second, these studies generally assumed that egg quality is independent of female age or population context, and that egg production is linearly related to female weight. This assumption means that the total egg production is equivalent to the

SSB. While it is a standard assumption in fisheries, often justified by the absence of time series of reproductive traits (Saborido-Rey and Trippel 2013), it ignores the importance of “Big Old Fat Fecund Female Fish” to the stock’s reproduction (Secor 2007, Hixon *et al.* 2014). If relative egg production and/or quality increases with female age or size, reducing the average age of the spawning stock will even further *decrease* the steepness of the stock-recruitment curves; the effect will be stronger for long-lived species (Venturelli *et al.* 2009). In a number of cases, allowing for time-variable total egg production that takes into account age structure and growth led to different inferences about stock biomasses and recommended fishing rates (Spencer and Dorn 2013). These findings suggest that routine monitoring of reproductive parameters is an important consideration in a precautionary approach to fisheries management.

Impact of life-history trends in single species management

With respect to this review, the main question of interest to fisheries management is – what are the consequences of basing harvesting decisions on assessments that ignore possible trends in life-history parameters? If stocks are monitored and assessed routinely, would the management be self-correcting even without taking into account possible changes in growth and maturation? The answers to these questions all depend on the rate of change in life-history parameters - a change of 0.5% per year will be less risky than a change of 5% per year. Empirical studies (Table 1) often report rates of phenotypic change in the range of 0.5-2% per year, with age-at-maturation among the fastest changing traits. Assuming a conservative estimate of *ca* 0.5% change per year, this would lead to *ca* 5% change over 10 years (see also van Walraven *et al.* 2010). For age-at-maturation, a more realistic value might be *ca* 1% per year (Sharpe and Hendry 2009; Audzijonyte *et al.* 2013a). Given that it is common for stock assessments to use 10-year-old maturation data (e.g. Stewart *et al.* 2011; ICES Advice 2014) and, when growth models are used, an average growth curve for the entire assessment period is assumed, it would be important to ascertain the likely errors in the stock assessment and management recommendations if the actual age-at-maturation is overestimated by 1 (or 2) years and length-at-age of adult individuals is overestimated by *ca* 5% (10%). To answer these questions specific simulations are needed. Some harvest strategies or management plans (combinations of monitoring, assessment and decision rules) may be relatively robust to ignorance about life-history changes, while others may be quite sensitive. Results from a few studies that specifically looked at the consequences of mis-specifying life-history parameters in fisheries models are provided below.

A model with cohort-specific growth rates in an age-structured assessment of southeast Australia’s blue grenadier (*Macruronus novaezelandiae*, Merlucciidae) showed that a 5-15% misspecification of length-at-age led to *ca* 5-20% difference in inferred stock biomass (Whitten *et al.* 2013). A similar result was

found in a study of Baltic Sea herring (*Clupea harengus*, Clupeidae), where growth was modelled as a time-variable parameter in a Bayesian framework, using information on zooplankton abundance, sea surface temperature and small prey fish biomass as priors (Mantyniemi *et al.* 2013). Wells *et al.* (2013) showed that overestimation of L_{∞} (expected in cases of faster life-histories) will overestimate fishing mortality rates, because the absence of large individuals in stocks with smaller adult length-at-age would bias fishery mortality estimates for older age groups upwards. Bias in selectivity estimates can occur if data are pooled from stocks that have different responses to temperature (e.g. increased and decreased growth as temperature increased up to and above the optimum) (Punt *et al.* 2014). A 20% underestimate in VB k will lead to *ca* 20% reduction in the estimation of yield-per-recruit biomass at fishing mortality $F_{0.1}$ (year⁻¹) (the F value where the curve of production against fishing mortality has a gradient of 10%) (Pardo *et al.* 2013). Concerns about changes in growth may be less important for assessment models (such as some cohort analyses) that use annual empirical information about weight at age rather than fitted and assumed growth curves (but data quality and quantity are critical, see Kuriyama *et al.* 2015). However, even for stocks managed using such models and data, projections about future stock dynamics may become biased if possible trends in growth and maturation are not considered. Also the drawback of such empirical fitting is that no insight into the processes governing growth is gained (Lorenzen 2016). Finally, one of the major ways in which faster life-histories will affect stock productivity is through an expected increase in natural mortality (Fig. 1B). Broad empirical and theoretical comparisons suggest that a change in k from 0.1 to 0.2 would increase M from 0.1 to 0.3 (year⁻¹), and a change in maturation age from 5 to 4 years would increase M from 0.2 to 0.3 (e.g. Carruthers *et al.* 2012) (Fig. 1B). Jørgensen and Fiksen (2010) suggest that, due to harvest induced evolution in life-histories and behaviour, natural mortality rates increase proportionally to harvest, such as “fishing two fish kills three”. When studied in an ecosystem modelling framework, a decrease in body length of *ca* 4% led to 10-50% increase in predation mortality in four out of five studied species (Audzijonyte *et al.* 2013b).

Given the evidence for dynamic life-histories in marine species, is it then reasonable to use simplifying assumptions about life-history, such as fixed age or length at maturation, fixed growth curves, fixed *a priori* assumed size independent natural mortality, and fixed steepness of the stock-recruitment relationship, in stock assessment models (see also Lorenzen 2016)? For example, although age at maturation is shown to be one of the most rapidly changing life-history traits (Table 1, Sharpe and Hendry 2009; Devine *et al.* 2012; Audzijonyte *et al.* 2013a), this parameter is often fixed in stock assessments. This is true even in cases where trends in maturation have been demonstrated (e.g. for North Sea plaice (*Pleuronectes platessa*, Pleuronectidae), see van Walraven *et al.* 2010 and ICES Advice 2014). Mangel *et al.* (2013) state that the standard practice of assuming fixed values for steepness of the stock-recruitment relationship and natural mortality will *a priori* determine fisheries reference points in a standard stock assessment that uses two-parameter (Beverton-Holt or Ricker) stock recruitment

relationships. The resulting estimates from a stock assessment “can provide a false sense of precision and information because an important source of uncertainty is hidden”. We believe that acknowledging and explicitly including important sources of uncertainty in fisheries models will reflect the state of affairs about our knowledge and encourage better monitoring (Kuparinen *et al.* 2012; Evans *et al.* 2013).

Life-history trends in multi-species models

The number of parameters and interactions in multispecies models means that the life-history traits of individual species (and hence their temporal change) may not dominate their dynamics in the way they do in population models used in fisheries. Yet, individual size remains one of the key characteristics defining functional groups in most complex ecosystem models (e.g. OSMOSE, Shin and Cury 2004; Atlantis, Fulton 2011) and forms the basis of size- and trait-based models (Andersen and Beyer 2006; Blanchard *et al.* 2009). As tractable complexity is only possible in a few dimensions at a time (e.g. either in food web structure, spatial resolution, or life-history detail), most models do not consider spatial or temporal variation in size or any other life-history parameters beyond what is produced by physiologically mediated responses, or where spatial forage fields are non-uniform and result in changes to realised growth rates. Nevertheless, there is a growing realisation that non-stationary and spatially varying life-histories may be essential to simulate effects of directional pressures such as climate change (e.g. Brännström *et al.* 2012).

Life-history change is recognised mechanistically in some ecological models (single and multispecies) through the use of multiple growth curves per species (e.g. Little *et al.* 2007). Such models allow for changes in aggregate average life-history characteristics if there is differential catchability or gear selectivity. Models that explicitly consider evolution are typically constrained to small species complexes (DeAngelis and Mooij 2005; Miethe *et al.* 2009) or explore evolution of food webs (e.g. Rossberg *et al.* 2005) rather than life-history traits. However, where ecosystem models have been modified to allow for trends in size-at-age, they showed that even small changes in body sizes can get amplified or dampened through species interactions (Audzijonyte *et al.* 2013b) and have large effects on the trophic dynamics, diets and recovery of harvested species (Audzijonyte *et al.* 2014).

Brännström *et al.* (2012) state that further development of models which consider evolution in the context of food webs is critically important for addressing (i) the biological adaptive capacity of ecosystems and (ii) the vulnerability of these systems to stock and food web collapse as a result of secondary extinctions due to trophic cascades. Allowance for plasticity in growth, consumption and reproductive rates is similarly important. Moreover, multi-species and ecosystem models are beginning to provide strategic ecosystem-based management advice, assessing ecosystem and food web implications of management decisions (e.g. Fulton *et al.* 2014). Consequently, just as overlooking the potential for changing life-

history characteristics can undermine management relevant information from single species models, so too the omission of shifting life-history parameters in ecosystem models has the potential to lead to erroneous conclusions about the implications of stressors and management for the long term status of the system and its sub-components (Collie and Gislason 2001; Audzijonyte *et al.* 2013b, 2014).

How to address changes in life-histories of marine organisms - a plan of action

Given the evidence for the changes in life-history of exploited marine species, and the compounding effects of climate change, scientists and marine managers have a range of options in terms of life-history research and subsequent application to management issues (Fig. 3). Scientists involved in stock assessments should acknowledge that trends in life-history traits are likely to be happening and consider what implications this has for the marine populations and ecosystem management arenas in which they work. The framework proposed in Fig. 3 would consider the interaction and relative strength of density dependence, physiological and evolutionary processes. If there is evidence for strong year classes due to relaxed density dependence (top box in Fig. 3) or improved food supply due to increased productivity (second top box in Fig. 3), it is likely that the trend towards decreased adult body size might be masked. It is important to highlight that the underlying physiological and evolutionary processes would still be operational during the periods of faster growth due to improved food supply.

The realisation of the importance of life-history trends is evident in several resource management working groups (Piet *et al.* 2010), but wider investigation is needed. To support management, improved monitoring strategies are required to provide up-to-date life-history characteristics (e.g. Hollowed *et al.* 2013), such as maturation data, growth rate and natural mortality. We acknowledge the difficulties of collecting such data, especially on suitable spatial scales, yet the importance has never been greater. Size-at-age and size-at-maturity could be one of the most important measures determining organism responses to thermal conditions (Ohlberger 2013), and is also one of the main parameters showing rapid response to fisheries-induced population dynamics and productivity (Brander 2010).

Since life-history information for many stocks is missing, we suggest that as the first step, assessment scientists could explore the sensitivity of model outcomes to misspecification of the life-history parameters using simulations (Szuwalski and Punt 2013; Deroba *et al.* 2014). The next step could be to make better use of existing information, such as in otolith collections, to infer life-histories and possible trends. Otolith data have been used to infer individual-level variation of growth (Morrongiello and Thresher 2015), larval growth rates and the age at which juveniles settle into adult habitats (Fowler and Jennings 2003; Hale and Swearer 2008), differential selection on size-at-age imposed by fisheries (Swain *et al.* 2007; Kendall *et al.* 2014), and the age at sexual maturity, either directly through observing a

transition zone (Francis and Horn 1997) or through use of energy-based modelling that recognises the main trade-offs between growth, maintenance and reproduction (Brunel *et al.* 2013). Otolith-based bio-chronologies can provide long-term datasets to aid the investigation of the impacts of environmental variability and fishing pressure on fish stocks (Morrongiello *et al.* 2012). The repeated measures nature of otolith-derived growth data also means that more sophisticated statistical techniques can be employed to decompose population-level patterns into their within and among individual components. This is very important from a climate change perspective as the likelihood of species being able to respond to warming depends on the levels of phenotypic plasticity (Nussey *et al.* 2007).

Finally, molecular analyses of life-history traits also provide an increasingly valuable source of information. Genes and markers responsible for life-history traits have now been identified in some fishes (Van Wijk *et al.* 2013; Therkildsen *et al.* 2013; Uusi-Heikkilä *et al.* 2015). If these markers are consistently correlated to life-history traits in different species, they can be used for targeted genetic monitoring. Even when monitoring data are collected for other purposes, it should be imperative to preserve material in ways that will be suitable for retrospective genomic analyses, particularly given the rapid decrease in analysis costs (Hayden 2014). Currently the main costs typically come from the subsequent bioinformatics analyses, but here also better tools are being developed every year. Nevertheless, careful analytical design will be needed to provide information relevant to the time and spatial scales useful for assessments.

With regard to management decisions underpinning sustainable use of marine resources exposed to life-history changes, a combination of precautionary and learning through adaptive management is recommended to maximise species resilience to climate change. For example, based on evidence that age truncation will lead to increased recruitment variability (Secor 2007), precautionary management might seek to increase resilience by increasing the number of age classes contributing to reproductive output. This might be achieved with lower fishing mortality rates, or by using spatial protection for low mobility species (Plagányi *et al.* 2013), or with upper size limits or dynamic spatial management for more mobile species (Hobday *et al.* 2014). Adaptive management would mean that management strategies would be flexible and could respond rapidly as new information is received (e.g. on the status of the stock, realised mortality rates, or in the context of this review assessments could be continuously updated with new life-history information). Notably, when a positive effect of maternal age is taken into account, reducing harvest rates on older, larger individuals may actually increase catch compared to the *status quo*, as productivity increases and more young fish offset the foregone harvest on older fish (Shelton *et al.* 2015). Spatial management may offer protection to some proportion of a population in no-take reserves, which may also help to maintain genetic diversity and demographic structure of the population, increasing adaptive potential and resilience to fluctuations. Clearly, the effectiveness of any measures will vary

across stocks, and predictions are difficult as processes determining stock resilience are not well understood. Overall, because it is unclear what exploitation rate is sustainable and maintains the adaptive potential and demographic resilience of marine populations to rapid environmental change, the best precautionary management option will be the one that protects life-history diversity and maintains larger stock biomass (Froese 2004). Without further study, we cannot recommend a single biomass reference point, but it is likely to be higher than the biomass leading to maximum sustainable yield. Further work would be necessary to explore whether more conservative target reference points, such as the biomass leading to maximum economic yield (or its proxy of 48% of the unfished level ($B_{0.48}$)), which has been adopted by fisheries management in places such as Australia (Rayns 2007), is sufficient.

As human influence on marine ecosystems is increasing, a return to baseline conditions is unlikely in most habitats and management based on the past will fail to account for new patterns (Merrie *et al.* 2014). Hence, adaptive management informed by monitoring is a suitable pathway to follow. Because resources will always be limited, calls for monitoring should be clear about the specific variables that can provide the information to assess the effectiveness of the management approach - we suggest that up-to-date information on life-history characteristics, such as age-at-maturity and growth rate is essential for implementing adaptive management strategies.

Acknowledgements

This study was supported by the Kone Foundation (AA), Academy of Finland (AK) and CSIRO. We thank two anonymous referees and the editor for comments that sharpened the focus of this manuscript.

References

- Andersen, K.H., Beyer, J.E. (2006) Asymptotic size determines species abundance in the marine size spectrum. *American Naturalist* 168, 54-61.
- Anderson, S.C., Moore, J.W., McClure, M.M., Dulvy, N.K., Cooper, A.B. (2015). Portfolio conservation of metapopulations under climate change. *Ecological Applications* 25, 559-572.
- Arendt, J.D. (2011) Size-fecundity relationships, growth trajectories, and the temperature-size rule for ectotherms. *Evolution* 65, 43-51.
- Arendt, J.D. (2015) Why get big in the cold? Size-fecundity relationships explain the temperature-size rule in a pulmonate snail (*Physa*). *Journal of Evolutionary Biology* 28, 169-178.
- Atkinson, D. (1994) Temperature and Organism Size - a Biological Law for Ectotherms. *Advances in Ecological Research*, 25, 1-58.
- Attrill, M.J., Power, M. (2002) Climatic influence on a marine fish assemblage. *Nature* 417, 275-278.
- Audzijonyte, A., Kuparinen, A. (2016) The role of life histories and trophic interactions in population recovery. *Conservation Biology* (in press)

- 678 Audzijonyte, A., Kuparinen, A., Fulton, E.A. (2013a) How fast is fisheries-induced evolution?
679 Quantitative analysis of modelling and empirical studies. *Evolutionary Applications* 6, 585-595.
- 680 Audzijonyte, A., Kuparinen, A., Gorton, R., Fulton, E.A. (2013b) Ecological consequences of body size
681 decline in harvested fish species: positive feedback loops in trophic interactions amplify human impact.
682 *Biology Letters* 9:1103
- 683 Audzijonyte, A., Kuparinen, A., Fulton, E.A. (2014) Ecosystem effects of contemporary life-history
684 changes are comparable to those of fishing. *Marine Ecology Progress Series* 495, 219-231.
- 685 Audzijonyte, A., Fulton E.A., Kuparinen A. (2015) The impacts of fish body size changes on stock
686 recovery: a case study using an Australian marine ecosystem model. *ICES Journal of Marine Science*
687 72, 782-792
- 688 Barange, M., Merino, G., Blanchard, J.L., *et al.* (2014) Impacts of climate change on marine ecosystem
689 production in societies dependent on fisheries. *Nature Climate Change* 4, 211-216.
- 690 Bates, A.E., Pecl, G.T., Frusher, S., *et al.* (2014) Defining and observing stages of climate-mediated range
691 shifts in marine systems. *Global Environmental Change-Human and Policy Dimensions* 26, 27-38.
- 692 Baudron, A.R., Needle, C.L., Rijnsdorp, A.D., Marshall, C.T. (2014) Warming temperatures and smaller
693 body sizes: synchronous changes in growth of North Sea fishes. *Global Change Biology* 20, 1023-1031.
- 694 Bell, G. (1980) The costs of reproduction and their consequences. *The American Naturalist* 116, 45-76.
- 695 Beverton, R.J.H. (1992) Patterns of Reproductive Strategy Parameters in Some Marine Teleost Fishes.
696 *Journal of Fish Biology* 41, 137-160.
- 697 Blanchard, J.L., Jennings, S., Law, R., *et al.* (2009) How does abundance scale with body size in coupled
698 size-structured food webs? *Journal of Animal Ecology* 78, 270-280.
- 699 Brander, K. (2010) Impacts of climate change on fisheries. *Journal of Marine Systems* 79, 389-402.
- 700 Brander, K. (2013) Climate and current anthropogenic impacts on fisheries. *Climatic Change* 119, 9-21.
- 701 Brännström, Å., Johansson, J., Loeuille, N., *et al.* (2012) Modelling the ecology and evolution of
702 communities: a review of past achievements, current efforts, and future promises. *Evolutionary Ecology*
703 *Research* 14, 601-625.
- 704 Brown, C.J., Fulton, E.A., Hobday, A.J., *et al.* (2009) Effects of climate-driven primary production
705 change on marine food webs: implications for fisheries and conservation. *Global Change Biology* 16,
706 1194-1212.
- 707 Brunel, T., Ernande, B., Mollet, F.M., Rijnsdorp, A.D. (2013). Estimating age at maturation and energy-
708 based life-history traits from individual growth trajectories with nonlinear mixed-effects models.
709 *Oecologia* 172, 631–643.
- 710 Cardoso, L.G., Haimovici, M. (2011) Age and changes in growth of the king weakfish *Macrodon*
711 *atricauda* (Gunther, 1880) between 1977 and 2009 in southern Brazil. *Fisheries Research* 111, 177-187.
- 712 Carlson, S.M., Seamons, T.R. (2008) A review of quantitative genetic components of fitness in
713 salmonids: implications for adaptation to future change. *Evolutionary Applications* 1, 222-238.

714 Carruthers, T.R., Walters, C.J., McAllister, M.K. (2012) Evaluating methods that classify fisheries stock
715 status using only fisheries catch data. *Fisheries Research* 119, 66-79.

716 Charnov, E. (1993) *Life History Invariants: Some Explorations of Symmetry in Evolutionary Ecology*.
717 Oxford University Press.

718 Charnov, E.L., Gislason, H., Pope, J.G. (2013) Evolutionary assembly rules for fish life histories. *Fish
719 and Fisheries* 14, 213-224.

720 Cheung, W.W.L., Sarmiento, J.L., Dunne, J., *et al.* (2013) Shrinking of fishes exacerbates impacts of
721 global ocean changes on marine ecosystems. *Nature Climate Change* 3, 254-258.

722 Cheung, W.W.L., Brodeur, R.D., Okey, T.A., Pauly D. (2014). Projecting future changes in distributions
723 of pelagic fish species of Northeast Pacific shelf seas. *Progress in Oceanography*, 130, 19-31.

724 Chevin, L.M., Lande, R., Mace, G.M. (2010) Adaptation, Plasticity, and Extinction in a Changing
725 Environment: Towards a Predictive Theory. *PloS Biology* 8, e1000357.

726 Christensen, V., Walters, C.J. (2004) Ecopath with Ecosim: methods, capabilities and limitations.
727 *Ecological Modelling* 172, 109-139.

728 Collie, J.S., Gislason, H. (2001) Biological reference points for fish stocks in a multispecies context.
729 *Canadian Journal of Fisheries and Aquatic Sciences* 58, 2167–2176.

730 Conover, D.O., Gilmore, T., Munch, S.B. (2003) Estimating the relative contribution of spring-and
731 summer-spawned cohorts to the Atlantic coast bluefish stock. *Transactions of the American Fisheries
732 Society* 132, 1117-1124.

733 Crozier, L.G., Hutchings, J.A. (2014) Plastic and evolutionary responses to climate change in fish.
734 *Evolutionary Applications* 7, 68-87.

735 Crozier, L.G., Scheuerell, M.D., Zabel, R.W. (2011) Using Time Series Analysis to Characterize
736 Evolutionary and Plastic Responses to Environmental Change: A Case Study of a Shift toward Earlier
737 Migration Date in Sockeye Salmon. *American Naturalist* 178, 755-773.

738 Crozier, L.G., Zabel, R.W., Hockersmith, E.E., Achord, S. (2010) Interacting effects of density and
739 temperature on body size in multiple populations of Chinook salmon. *Journal of Animal Ecology* 79,
740 342-349.

741 Darimont, C.T., Carlson, S.M., Kinnison, M.T., Paquet, P.C., Reimchen, T.E., Wilmer, C.C. (2009)
742 Human predators outpace other agents of trait change in the wild. *Proceedings of the National Academy
743 of Sciences of the United States of America* 106, 952-954.

744 Daufresne, M., Lengfellner, K., Sommer, U. (2009) Global warming benefits the small in aquatic
745 ecosystems. *Proceedings of the National Academy of Sciences of the United States of America* 106,
746 12788-12793.

747 DeAngelis, D.L., Mooij, W.M. (2005) Individual-Based Modeling of Ecological and Evolutionary
748 Processes. *Annual Review of Ecology, Evolution, and Systematics* 36, 147-168.

749 Deroba, J.J., Butterworth, D.S., Methot, R.D., *et al.* (2014). Simulation testing the robustness of stock
750 assessment models to error: some results from the ICES strategic initiative on stock assessment
751 methods. *ICES Journal of Marine Science* doi:10.1093/icesjms/fst237

752 Devine, J.A., Wright, P.J., Pardoe, H., Heino, M. (2012) Comparing rates of contemporary evolution in
753 life-history traits for exploited fish stocks. *Canadian Journal of Fisheries and Aquatic Sciences* 69,
754 1105-1120.

755 Doney, S.C., Ruckelshaus, M., Duffy, J.E., *et al.* (2012) Climate Change Impacts on Marine Ecosystems.
756 *Annual Review of Marine Science*, 4, 11-37.

757 Dulvy, N.K., Ellis, J.R., Goodwin, N.B., Grant, A., Reynolds, J.D., Jennings, S. (2004) Methods of
758 assessing extinction risk in marine fishes. *Fish and Fisheries* 5, 255-276.

759 Dulvy, N. K., Rogers, S. I., Jennings, S., Stelzenmuller, V., Dye, S. R., Skjoldal H. R. (2008). Climate
760 change and deepening of the North Sea fish assemblage: a biotic indicator of warming seas. *Journal of*
761 *Animal Ecology* 45, 1029-1039.

762 Enberg, K., Jørgensen, C., Dunlop, E. S., Heino, M., Dieckmann, U. (2009). Implications of fisheries-
763 induced evolution for stock rebuilding and recovery. *Evolutionary Applications* 2, 394–414.

764 Enberg, K., Jørgensen, C., Mangel, M. (2010) Fishing-induced evolution and changing reproductive
765 ecology of fish: evolution of steepness. *Canadian Journal of Fisheries and Aquatic Sciences* 67, 1708-
766 1719.

767 Enberg, K., Jørgensen, C., Dunlop, E.S., *et al.* (2012) Fishing-induced evolution of growth: Concepts,
768 mechanisms and the empirical evidence. *Marine Ecology* 33, 1-25.

769 Evans, M.R., Grimm, V., Johst, K., *et al.* (2013) Do simple models lead to generality in ecology? *Trends*
770 *in Ecology & Evolution* 28, 578-583.

771 Figueira, W.F., Biro, P., Booth, D.J., Valenzuela, V.C. (2009) Performance of tropical fish recruiting to
772 temperate habitats: role of ambient temperature and implications of climate change. *Marine Ecology*
773 *Progress Series* 384, 231-239.

774 Forster, J., Hirst, A.G., Woodward, G. (2011) Growth and Development Rates Have Different Thermal
775 Responses. *American Naturalist* 178, 668-678.

776 Forster, J., Hirst, A.G., Atkinson, D. (2012) Warming-induced reductions in body size are greater in
777 aquatic than terrestrial species. *Proceedings of the National Academy of Sciences of the United States of*
778 *America* 109, 19310-19314.

779 Fowler, A.J., Jennings, P.R. (2003) Dynamics in 0+ recruitment and early life history for snapper (*Pagrus*
780 *auratus*, Sparidae) in South Australia. *Marine and Freshwater Research* 54, 941-956.

781 Francis, R.I.C.C., Horn, P.L. (1997) Transition zone in otoliths of orange roughy (*Hoplostethus*
782 *atlanticus*) and its relationship to the onset of maturity. *Marine Biology* 129, 681-687.

783 Froese, R. (2004) Keep it simple: three indicators to deal with overfishing. *Fish and Fisheries* 5, 86-91.

784 Fulton, E.A. (2011) Interesting times: winners, losers, and system shifts under climate change around
785 Australia. *ICES Journal of Marine Science* 68, 1329–1342.

786 Fulton, E.A., Smith, A.D.M., Smith, D.C., Johnson, P. (2014) An Integrated Approach Is Needed for
787 Ecosystem Based Fisheries Management: Insights from Ecosystem-Level Management Strategy
788 Evaluation. *Plos One* 9.

789 Garcia, S. M., Kolding, J., Rice, J., *et al.* (2012). Reconsidering the consequences of selective fisheries.
790 *Science*, 335, 1045-1047.

791 Gienapp, P., Teplitsky, C., Alho, J.S., Mills, J.A., Merilä, J. (2008) Climate change and evolution:
792 disentangling environmental and genetic responses. *Molecular Ecology* 17, 167-178.

793 Gislason, H., Daan, N., Rice, J.C., Pope, J.G. (2010) Size, growth, temperature and the natural mortality
794 of marine fish. *Fish and Fisheries* 11, 149-158.

795 Haddon, M. (2011) *Modelling and Quantitative Methods in Fisheries*, Second Edition, 2nd edn Vol.,
796 Chapman and Hall/ CRC.

797 Hale, R., Swearer, S.E. (2008) Otolith microstructural and microchemical changes associated with
798 settlement in the diadromous fish *Galaxias maculatus*. *Marine Ecology Progress Series* 354, 229-234.

799 Halpern, B.S., Longo, C., Hardy, D., *et al.* (2012) An index to assess the health and benefits of the global
800 ocean. *Nature* 488, 615-620.

801 Hansen, M.M., Olivieri, I., Waller, D.M., Nielsen, E.E. (2012) Monitoring adaptive genetic responses to
802 environmental change. *Molecular Ecology* 21, 1311-1329.

803 Hayden, E. C. (2015). The \$1000 genome. *Nature* 507, 294-295.

804 Heino, M., Baulier, L., Boukal, D.S., *et al.* (2013) Can fisheries-induced evolution shift reference points
805 for fisheries management? *ICES Journal of Marine Science* 70, 707-721.

806 Heino, M., Pauli, B.D., Dieckmann, U. (2015) Fisheries-induced evolution. *Annual Review of Ecology,*
807 *Evolution, and Systematics*, 46, 461-480.

808 Helidoniotis, F., Haddon, M. (2014) The effectiveness of broad-scale legal minimum lengths for
809 protecting spawning biomass of *Haliotis rubra* in Tasmania. *New Zealand Journal of Marine and*
810 *Freshwater Research* 48, 70-85.

811 Hill, J.K., Griffiths, H.M., Thomas, C.D. (2011) Climate change and evolutionary adaptations at species'
812 range margins. *Annual Review of Entomology* 56, 143-159.

813 Hixon, M.A., Johnson, D.W., Sogard, S.M. (2014) BOFFFFs: on the importance of conserving old-
814 growth age structure in fishery populations. *ICES Journal of Marine Science* 71, 2171-2185.

815 Hobday, A.J., Smith, A.D.M., Stobutzki, I.C., *et al.* (2011a) Ecological risk assessment for the effects of
816 fishing. *Fisheries Research* 108, 372-384.

817 Hobday, A.J. (2011b) Sliding baselines and shuffling species: implications of climate change for marine
818 conservation. *Marine Ecology-an Evolutionary Perspective* 32, 392-403.

819 Hobday, A. J., Maxwell, S.M., Forgie, J., *et al.* (2014). Dynamic ocean management: Integrating
820 scientific and technological capacity with law, policy and management. *Stanford Environmental Law*
821 *Journal* 33(2), 125-165.

822 Hoffmann, A.A., Sgro, C.M. (2011) Climate change and evolutionary adaptation. *Nature* 470, 479-485.

823 Hollowed, A.B., Barange, M., Beamish, R.J., *et al.* (2013) Projected impacts of climate change on marine
824 fish and fisheries. *ICES Journal of Marine Science* 70, 1023-1037.

825 Holt, R. E., Jørgensen, C. (2015) Climate change in fish: effects of respiratory constraints on optimal life
826 history and behaviour. *Biology Letters* 11, 1032.

827 Hsieh, C.H., Reiss, C.S., Hewitt, R.P., Sugihara, G. (2008) Spatial analysis shows that fishing enhances
828 the climatic sensitivity of marine fishes. *Canadian Journal of Fisheries and Aquatic Sciences* 65, 947-
829 961.

830 Hutchings, J.A., Myers, R.A., Garcia, V.B., Lucifora, L.O., Kuparinen, A. (2012) Life-history correlates
831 of extinction risk and recovery potential. *Ecological Applications* 22, 1061-1067.

832 ICES Advice 2014. Plaice in Subarea IV (North Sea) (updated) [www.ices.dk/sites/pub/.../Advice/2014/](http://www.ices.dk/sites/pub/.../Advice/2014/2014/ple-nsea.pdf)
833 [2014/ple-nsea.pdf](http://www.ices.dk/sites/pub/.../Advice/2014/2014/ple-nsea.pdf) (last accessed on May 5, 2015)

834 IPCC-AR5 (2014) Summary for Policymakers. In *Climate Change 2014: Impacts, Adaptation, and*
835 *Vulnerability*.

836 Irie, T., Fischer, K. (2009) Ectotherms with a calcareous exoskeleton follow the temperature-size rule-
837 evidence from field survey. *Marine Ecology Progress Series* 385, 33-37.

838 Jakobsdóttir, K.B., Pardoe, H., Magnússon, A., *et al.* (2012) Historical changes in genotypic frequencies
839 at the Pantophysin locus in Atlantic cod (*Gadus morhua*) in Icelandic waters: evidence of fisheries-
840 induced selection? *Evolutionary Applications* 4, 562-573.

841 Jensen, A.L. (1997) Origin of the relation between K and Linf and synthesis of relations among life
842 history parameters. *Canadian Journal of Fisheries and Aquatic Sciences* 54, 987-989.

843 Jennings, S., Reynolds, J.D., Mills, S.C. (1998) Life history correlates of responses to fisheries
844 exploitation. *Proceedings of the Royal Society B*, 265, 333-339

845 Jørgensen, C., Ernande, B., Fiksen, Ø., Dieckmann, U. (2006) The logic of skipped spawning in fish.
846 *Canadian Journal of Fisheries and Aquatic Sciences* 63, 200-211.

847 Jørgensen, C., Enberg, K., Dunlop, E.S., *et al.* (2007) Managing evolving fish stocks. *Science* 318, 1247-
848 1248.

849 Jørgensen, C., Fiksen Ø. (2010) Modelling fishing-induced adaptations and consequences for natural
850 mortality. *Canadian Journal of Fisheries and Aquatic Sciences*, 67, 1086-1097.

851 Jørgensen, C., Holt, R.E. (2013) Natural mortality: Its ecology, how it shapes fish life histories, and why
852 it may be increased by fishing. *Journal of Sea Research* 75, 8-18.

853 Kendall, N.W., Dieckmann, U., Heino, M., Punt, A.E., Quinn, T.P. (2014) Evolution of age and length at
854 maturation of Alaskan salmon under size-selective harvest. *Evolutionary Applications* 7, 313-322.

855 Kokita, T. (2003) Potential latitudinal variation in egg size and number of a geographically widespread
856 reef fish, revealed by common-environment experiments. *Marine Biology* 143, 593-601.

857 Kokkonen, E., Vainikka, A., Heikinheimo, O. (2015) Probabilistic maturation reaction norm trends reveal
858 decreased size and age at maturation in an intensively harvested stock of pikeperch *Sander lucioperca*.
859 *Fisheries Research* 167, 1-12.

860 Kozłowski, J., Czarnołęski, M., Dańko, M. (2004) Can optimal resource allocation models explain why
861 ectotherms grow larger in cold? *Integrative and Comparative Biology* 44, 480-493.

862 Kraak, S.B.M. (2007) Does the probabilistic maturation reaction norm approach disentangle phenotypic
863 plasticity from genetic change? *Marine Ecology Progress Series* 335, 295-300.

864 Kuparinen, A., Cano Arias, J.M., Loehr, J., Herczeg, G., Gonda, A., Merilä, J. (2011) Fish age at
865 maturation is influenced by temperature independently of growth. *Oecologia* 167, 435-443.

866 Kuparinen, A., Kuikka, S., Merila, J. (2009) Estimating fisheries-induced selection: traditional gear
867 selectivity research meets fisheries-induced evolution. *Evolutionary Applications* 2, 234-243.

868 Kuparinen, A., Mantyniemi, S., Hutchings, J.A., Kuikka, S. (2012) Increasing biological realism of
869 fisheries stock assessment: towards hierarchical Bayesian methods. *Environmental Reviews* 20, 135-
870 151.

871 Kuparinen, A., Stenseth, N.C., Hutchings, J.A. (2014) Fundamental population-productivity relationships
872 can be modified through density-dependent feedbacks of life-history evolution. *Evolutionary*
873 *Applications*, 7, 1218-1225

874 Kuriyama, P. T., Ono, K., Hurtado-Ferro, F. *et al.* (2015). An empirical weight-at-age approach reduces
875 estimation bias compared to modeling parametric growth in integrated, statistical stock assessment
876 models when growth is time varying. *Fisheries Research*

877 Little, L.R., Punt, A.E., Mapstone, B.D., et al. (2007) ELFSim - A model for evaluating management
878 options for spatially structured reef fish populations: An illustration of the "larval subsidy" effect.
879 *Ecological Modelling* 205, 381-396.

880 Lorenzen, K. (2016) Toward a new paradigm for growth modeling in fisheries stock assessments:
881 Embracing plasticity and its consequences. *Fisheries Research*

882 Lorenzen, K., Enberg, K. (2002) Density-dependent growth as a key mechanism in the regulation of fish
883 populations: evidence from among-population comparisons. *Proceedings of the Royal Society B* 269,
884 49-54.

885 Mangel, M., MacCall, A.D., Brodziak, J., *et al.* (2013) A perspective on steepness, reference points, and
886 stock assessment. *Canadian Journal of Fisheries and Aquatic Sciences* 70, 930-940.

887 Mantyniemi, S., Uusitalo, L., Peltonen, H., Haapasaari, P., Kuikka, S. (2013) Integrated, age-structured,
888 length-based stock assessment model with uncertain process variances, structural uncertainty, and
889 environmental covariates: case of Central Baltic herring. *Canadian Journal of Fisheries and Aquatic*
890 *Sciences* 70, 1317-1326.

891 Manyak-Davis, A., Bell, T.M., Sotka, E.E. (2013) The Relative Importance of Predation Risk and Water
892 Temperature in Maintaining Bergmann's Rule in a Marine Ectotherm. *American Naturalist* 182, 347-
893 358.

894 Mcadam, A.G., Boutin, S., Sykes, A.K., Humphries, M.M. (2007) Life histories of female red squirrels
895 and their contributions to population growth and lifetime fitness. *Ecoscience* 14, 362-369.

896 Melville-Smith, R., De Lestang, S. (2006). Spatial and temporal variation in the size at maturity of the
897 western rock lobster *Panulirus cygnus* George. *Marine Biology*, 150, 183-195.

898 Merrie, A., Dunn, D.C., Metian, M., *et al.* (2014) An ocean of surprises - Trends in human use,
899 unexpected dynamics and governance challenges in areas beyond national jurisdiction. *Global*
900 *Environmental Change-Human and Policy Dimensions* 27, 19-31.

901 Miethe, T., Pitchford, J., Dytham, C. (2009) An individual-based model for reviewing marine reserves in
902 the light of fisheries-induced evolution in mobility and size at maturation. *Journal of Northwest Atlantic*
903 *Fisheries Science* 41, 151-162.

904 Morrongiello, J.R., Thresher, R.E. (2015) A statistical framework to explore ontogenetic growth variation
905 among individuals and populations: a marine fish example. *Ecological Monographs* 85, 93-115.

906 Morrongiello, J.R., Thresher, R.E., Smith, D.C. (2012) Aquatic biochronologies and climate change.
907 *Nature Climate Change* 2, 849-857.

908 Morrongiello, J.R., Walsh, C.T., Gray, C.A., Stocks, J.R., Crook, D.A. (2014) Environmental change
909 drives long-term recruitment and growth variation in an estuarine fish. *Global Change Biology* 20,
910 1844-1860.

911 Munch, S.B., Salinas, S. (2009) Latitudinal variation in lifespan within species is explained by the
912 metabolic theory of ecology. *Proceedings of the National Academy of Sciences of the United States of*
913 *America* 106, 13860-13864.

914 Munday, P.L., Warner, R.R., Monro, K., Pandolfi, J.M., Marshall, D.J. (2013) Predicting evolutionary
915 responses to climate change in the sea. *Ecology Letters* 16, 1488-1500.

916 Neuheimer, A.B., Taggart, C.T. (2010) Can changes in length-at-age and maturation timing in Scotian
917 Shelf haddock (*Melanogrammus aeglefinus*) be explained by fishing? *Canadian Journal of Fisheries*
918 *and Aquatic Sciences* 67, 854-865.

919 Neuheimer, A.B., Thresher, R.E., Lyle, J.M., Semmens, J.M. (2011) Tolerance limit for fish growth
920 exceeded by warming waters. *Nature Climate Change* 1, 110-113.

921 Nielsen, E.E., Cariani, A., Mac Aoidh, E., *et al.* (2012) Gene-associated markers provide tools for
922 tackling illegal fishing and false eco-certification. *Nature Communications* 3.

923 Nussey, D.H., Wilson, A.J., Brommer, J.E. (2007) The evolutionary ecology of individual phenotypic
924 plasticity in wild populations. *Journal of Evolutionary Biology* 20, 831-844.

925 Nusslé, S., Bréchon, A., Wedekind, C. (2011) Change in individual growth rate and its link to gill-net
926 fishing in two sympatric whitefish species. *Evolutionary Ecology*, 25, 681-693.

- 927 Ohlberger, J. (2013) Climate warming and ectotherm body size—from individual physiology to
928 community ecology. *Functional Ecology* 27, 991-1001.
- 929 Olsen, E.M., Heino, M., Lilly, G.R., et al. (2004) Maturation trends indicative of rapid evolution preceded
930 the collapse of northern cod. *Nature* 428, 932-935.
- 931 Palkovacs, E.P., Kinnison, M.T., Correa, C., Dalton, C.M., Hendry, A.P. (2012) Fates beyond traits:
932 Ecological consequences of human-induced trait change. *Evolutionary Applications* 5, 183-191.
- 933 Pardo, S.A., Cooper, A.B., Dulvy, N.K. (2013) Avoiding fishy growth curves. *Methods in Ecology and*
934 *Evolution* 4, 353-360.
- 935 Patrick, W.S., Spencer, P., Link, J., et al. (2010) Using productivity and susceptibility indices to assess
936 the vulnerability of United States fish stocks to overfishing. *Fishery Bulletin* 108, 305-322.
- 937 Peck, L.S., Clark, M.S., Morley, S.A., Massey, A., Rossetti, H. (2009) Animal temperature limits and
938 ecological relevance: effects of size, activity and rates of change. *Functional Ecology* 23, 248-256.
- 939 Perry, R.I., Cury, P., Brander, K., Jennings, S., Mollmann, C., Planque, B. (2010) Sensitivity of marine
940 systems to climate and fishing: Concepts, issues and management responses. *Journal of Marine Systems*
941 79, 427-435.
- 942 Pespeni, M.H., Sanford, E., Gaylord, B., et al. (2013) Evolutionary change during experimental ocean
943 acidification. *Proceedings of the National Academy of Sciences of the United States of America* 110,
944 6937-6942.
- 945 Pethybridge, H. R., Young, J. W., Kuhnert, P. M., Farley, J. H. (2015). Using stable isotopes of albacore
946 tuna and predictive models to characterize bioregions and examine ecological change in the SW Pacific
947 Ocean. *Progress in Oceanography*.
- 948 Piet, G.J., Albella, A.J., Aro, E., et al. 2010. Marine Strategy Framework Directive, Task Group 3 Report:
949 Commercially exploited fish and shellfish. *Luxembourg: Office for Official Publications of the*
950 *European Communities*, 81 pp. doi:10.2788/83073.
- 951 Planque, B., Fromentin, J.M., Cury, P., et al. (2010) How does fishing alter marine populations and
952 ecosystems sensitivity to climate? *Journal of Marine Systems* 79, 403-417.
- 953 Plagányi, E. E., Skewes, T.D., Dowling, N.A., Haddon, M. (2013) Risk management tools for sustainable
954 fisheries management under changing climate: a sea cucumber example. *Climatic Change* 119, 181-197.
- 955 Polacheck, T., Eveson, J.P., Laslett, G.M. (2004) Increase in growth rates of southern bluefin tuna
956 (*Thunnus maccoyii*) over four decades: 1960 to 2000. *Canadian Journal of Fisheries and Aquatic*
957 *Sciences* 61, 307-322.
- 958 Poloczanska, E.S., Brown, C.J., Sydeman, W.J., et al. (2013) Global imprint of climate change on marine
959 life. *Nature Climate Change* 3, 919-925.
- 960 Poloczanska, E.S., Hawkins, S.J., Southward, A.J., Burrows, M.T. (2008) Modeling the response of
961 populations of competing species to climate change. *Ecology* 89, 3138-3149.

Polovina, J.J., Woodworth, P.A. (2012) Declines in phytoplankton cell size in the subtropical oceans estimated from satellite remotely-sensed temperature and chlorophyll, 1998-2007. *Deep-Sea Research Part II-Topical Studies in Oceanography* 77-80, 82-88.

Portner, H.O. (2001) Climate change and temperature-dependent biogeography: oxygen limitation of thermal tolerance in animals. *Naturwissenschaften* 88, 137-146.

Portner, H.O. (2010) Oxygen- and capacity-limitation of thermal tolerance: a matrix for integrating climate-related stressor effects in marine ecosystems. *Journal of Experimental Biology* 213, 881-893.

Portner, H.O., Knust, R. (2007) Climate change affects marine fishes through the oxygen limitation of thermal tolerance. *Science* 315, 95-97.

Portner, H.O., Farrell, A.P. (2008) Physiology and climate change. *Science* 322, 690-692.

Punt, A.E., Hurtado-Ferro, F., Whitten, A.R. (2014) Model selection for selectivity in fisheries stock assessments. *Fisheries Research* 158, 124-134.

Prince, J.D., Hordyk, A., Valencia, S.V., Loneragan, N., Sainsbury K., (2015) Revisiting the concept of Beverton-Holt Life History Invariants with the aim of informing data-poor fisheries assessment. *ICES Journal of Marine Science* 72 (1), 194-203.

Rayns, N. (2007) The Australian government's harvest strategy policy. *ICES Journal of Marine Science* 64, 596-598.

Reed, T.E., Schindler, D.E., Waples, R.S. (2011) Interacting Effects of Phenotypic Plasticity and Evolution on Population Persistence in a Changing Climate. *Conservation Biology* 25, 56-63.

Reznick, D.N., Shaw, F.H., Rodd, F.H., Shaw, R.G. (1997) Evaluation of the rate of evolution in natural populations of guppies (*Poecilia reticulata*). *Science* 275, 1934-1937.

Roff, D.A. (1992) The evolution of life histories; theory and analysis. Chapman and Hall, London.

Rogers, L.A., Stige, L.C., Olsen, E.M., *et al.* (2011). Climate and population density drive changes in cod body size throughout a century on the Norwegian coast. *Proceedings of the National Academy of Sciences of the United States of America*, 108, 1961-1966.

Rose, K.A., Cowan, J.H., Winemiller, K.O., Myers, R.A., Hilborn, R. (2001) Compensatory density dependence in fish populations: importance, controversy, understanding and prognosis. *Fish and Fisheries* 2, 293-327.

Rossberg, A.G., Matsuda, H., Amemiya, T., Itoh, K. (2005) An explanatory model for food-web structure and evolution. *Ecological Complexity* 2, 312-321.

Rountrey, A.N., Coulson, P.G., Meeuwig, J.J., Meekan, M. (2014) Water temperature and fish growth: otoliths predict growth patterns of a marine fish in a changing climate. *Global Change Biology* 20, 2450-2458.

Rouyer, T., Sadykov, A., Ohlberger, J., Stenseth, N.C. (2012) Does increasing mortality change the response of fish populations to environmental fluctuations? *Ecology Letters* 15, 658-665.

997 Rummer J.L., Couturier, C.S., Stecyk, J.A., *et al.* (2014) Life on the edge: thermal optima for aerobic
 998 scope of equatorial reef fishes are close to current day temperatures. *Global Change Biology* 20, 1055-
 999 1066.

1000 Saborido-Rey, F., Trippel, E. A. (2013). Fish reproduction and fisheries. *Fisheries Research*, 138, 1-4.

1001 Sarmiento, J.L., Slater, R., Barber, R., *et al.* (2004) Response of ocean ecosystems to climate warming.
 1002 *Global Biogeochemical Cycles* 18.

1003 Schaefer, M.B. (1954) Some aspects of the dynamics of populations important to the management of
 1004 commercial marine fisheries. *Inter-Am. Trop. Tuna Comm. Bull.* 1, 27-56.

1005 Schindler, D.E., Hilborn, R., Chasco, B., *et al.* (2010) Population diversity and the portfolio effect in an
 1006 exploited species. *Nature* 465, 609-612.

1007 Secor, D.H. (2007) The year-class phenomenon and the storage effect in marine fishes. *Journal of Sea*
 1008 *Research* 57, 91-103.

1009 Sharpe, D., Hendry, A. (2009) Life history change in commercially exploited fish stocks: an analysis of
 1010 trends across studies. *Evolutionary Applications* 2, 260-275.

1011 Shelton, A.O., Mangel, M. (2011) Fluctuations of fish populations and the magnifying effects of fishing.
 1012 *Proceedings of the National Academy of Sciences of the United States of America* 108, 7075-7080.

1013 Shelton, A.O., Munch, S.B., Keith, D., Mangel, M. (2012) Maternal age, fecundity, egg quality, and
 1014 recruitment: linking stock structure to recruitment using an age-structured Ricker model. *Canadian*
 1015 *Journal of Fisheries and Aquatic Sciences* 69, 1631-1641.

1016 Shelton, A. O., Hutchings, J. A., Waples, R. S., Keith, D. M., Akçakaya., H. R. (2015) Maternal age
 1017 effects on Atlantic cod recruitment and implications for future population trajectories. *ICES Journal of*
 1018 *Marine Science*.

1019 Shin, Y.-J., Cury, P. (2004) Using an individual-based model of fish assemblages to study the response of
 1020 size spectra to changes in fishing. *Canadian Journal of Fisheries and Aquatic Sciences* 61, 414-431.

1021 Spencer, P.D., Dorn, M.W. (2013) Incorporation of weight-specific relative fecundity and maternal
 1022 effects in larval survival into stock assessments. *Fisheries Research* 138, 159-167.

1023 Stewart, I.J., Forrest, R.E, Gradin C., et al. (2011) Status of the Pacific Hake (Whiting) stock in U.S. and
 1024 Canadian Waters in 2011. *Joint U.S. and Canadian Hake Technical Working Group*

1025 Svedang, H., Hornborg, S. (2014) Selective fishing induces density-dependent growth. *Nature*
 1026 *Communications* 5.

1027 Swain, D. P., Sinclair, A. F., Hanson, J. M. (2007). Evolutionary response to size-selective mortality in an
 1028 exploited fish population. *Proceedings of the Royal Society B: Biological Sciences* 274, 1015–1022.

1029 Swain, D.P. (2011) Life-history evolution and elevated natural mortality in a population of Atlantic cod
 1030 (*Gadus morhua*). *Evolutionary Applications* 4, 18-29.

- 1031 Szuwalski, C.S., Punt, A.E. (2013) Fisheries management for regime-based ecosystems: a management
1032 strategy evaluation for the snow crab fishery in the eastern Bering Sea. *ICES Journal of Marine Science*
1033 70, 955-967.
- 1034 Taucher, J., Oschlies, A. (2011) Can we predict the direction of marine primary production change under
1035 global warming? *Geophysical Research Letters* 38.
- 1036 Teplitsky, C., Millien, V. (2014) Climate warming and Bergmann's rule through time: is there any
1037 evidence? *Evolutionary Applications* 7, 156-168.
- 1038 Therkildsen, N.O., Hemmer-Hansen, J., Als, T.D., *et al.* (2013) Microevolution in time and space: SNP
1039 analysis of historical DNA reveals dynamic signatures of selection in Atlantic cod. *Molecular Ecology*
1040 22, 2424-2440.
- 1041 Thomas, A.C., Jarman, S.N., Haman, K.H., Trites, A.W., Deagle, B.E. (2014) Improving accuracy of
1042 DNA diet estimates using food tissue control materials and an evaluation of proxies for digestion bias.
1043 *Molecular Ecology* 23, 3706-3718.
- 1044 Thresher, R.E., Koslow, J.A., Morison, A.K., Smith, D.C. (2007) Depth-mediated reversal of the effects
1045 of climate change on long-term growth rates of exploited marine fish. *Proceedings of the National*
1046 *Academy of Sciences of the United States of America* 104, 7461-7465.
- 1047 Thresher, R., Morrongiello, J., Sloyan, B.M., *et al.* (2014) Parallel decadal variability of inferred water
1048 temperatures for Northern and Southern Hemisphere intermediate water masses. *Geophysical Research*
1049 *Letters* 41, 1232-1237.
- 1050 Uusi-Heikkilä, S., Whiteley, A.R., Kuparinen, A., *et al.* (2015) The evolutionary legacy of size-selective
1051 harvest extends from genes to populations. *Evolutionary Applications* (in press).
- 1052 Vainikka, A., Gaðrdmark, A., Bland, B., and Hjelm, J. (2009). Two- and three-dimensional maturation
1053 reaction norms for the eastern Baltic cod, *Gadus morhua*. *ICES Journal of Marine Science* 66, 248–257.
- 1054 Van Wijk, S.J., Taylor, M.I., Creer, S., *et al.* (2013) Experimental harvesting of fish populations drives
1055 genetically based shifts in body size and maturation. *Frontiers in Ecology and the Environment* 11, 181-
1056 187.
- 1057 Venturelli, P.A., Shuter, B.J., Murphy, C.A. (2009) Evidence for harvest-induced maternal influences on
1058 the reproductive rates of fish populations. *Proceedings of the Royal Society B: Biological Sciences* 276,
1059 919-924.
- 1060 van Walraven, L., Mollet, F.M., van Damme, C.J.G. , Rijnsdorp, A.D. (2010). Fisheries-induced
1061 evolution in growth, maturation and reproductive investment of the sexually dimorphic North Sea plaice
1062 (*Pleuronectes platessa* L.). *Journal of Sea Research*, 64, 85–93
- 1063 von Bertalanffy, L. (1957) Quantitative Laws in Metabolism and Growth. *The Quarterly Review of*
1064 *Biology* 32, 217-231.
- 1065 Wells, R.J.D., Kohin, S., Teo, S.L.H., Snodgrass, O.E., Uosaki, K. (2013) Age and growth of North
1066 Pacific albacore (*Thunnus alalunga*): Implications for stock assessment. *Fisheries Research* 147, 55-62.

- Whitten, A.R., Klaer, N.L., Tuck, G.N., Day, R.W. (2013) Accounting for cohort-specific variable growth in fisheries stock assessments: A case study from south-eastern Australia. *Fisheries Research* 142, 27-36.
- Woodworth-Jefcoats, P. A., Polovina, J. J., Howell, E. A., Blanchard, J. L. (2015). Two takes on the ecosystem impacts of climate change and fishing: Comparing a size-based and a species-based ecosystem model in the central North Pacific. *Progress in Oceanography*, 138, 533-545.
- Wright, P.J. (2014) Are there useful life history indicators of stock recovery rate in gadoids? *ICES Journal of Marine Science* 71, 1393-1406.

Table 1. Some literature examples on changes life-history parameters: juvenile growth (JG, often assessed as k in the VB curve), maturation (MAT, assessed as either age at 50% maturation or probabilistic maturation reaction norms) and asymptotic length or adult size (AS, often assessed as L_{∞} in the VB curve). Increase/decrease in parameter values is shown with +/-, no change indicated with 0, parameters that were not assessed are shown with na. Uncertain inferences in parameter change are shown in parentheses. DD – density dependent, T – temperature, FIE – fisheries-induced evolution. Species/stocks included in the review studies (e.g. Sharpe and Hendry 2009) are not listed separately.

Species	Area	MAT	JG	AS	Driver	Reference	Notes
Southern bluefin tuna	South Pacific	na	+	(–)	relaxed DD	Polachek et al. 2004	1960s-1990s, decrease in L_{∞} uncertain
Southern king weakfish	Southern Brazil	–	–	+	FIE, relaxed DD	Cardoso & Haimovici 2011,	1976 to 2011; changes coincided with heavy exploitation
Cod	Baltic Sea	na	na	–	increased DD	Svedang & Hornborg, 2014	k values not reported
Cod	Baltic Sea	–	na	–	FIE, environment	Vainikka et al. 2009	1987-2003, condition corrected trend in maturation found only in females
Pike perch (<i>Sander lucioperca</i> , Percidae)	Baltic Sea	–	na	0	FIE, higher T	Kokkonen et al. 2015	Increased productivity may have increased growth, masking decrease in size due to FIE
The western blue groper (<i>Achoerodus gouldii</i> , Labridae)	Western Australia	na	+	na	higher T	Rountrey et al 2014	1952–2003, changes in growth for ages 1-20y (maturation at ca 17 y)
Six commercial fish species	North Sea	na	na	-	higher T	Baudron et al. 2014	16% change per 2C (ca 0.8% /year)
Western rock lobster (<i>Panulirus cygnus</i> ,	West Australia	–	na	na	FIE, higher T	Melville-Smith & de Lestang, 2006	1975-2005, decrease in the carapace length at maturation

Palinuridea)							
Banded morwong (<i>Cheilodactylus spectabilis</i> , Cheilodactylidae)	Tasman Sea	na	+/-	na	higher T	Neuheimer et al. 2011	<i>k</i> decreased when T is above optimum
Three shallow water fishes (<250m)	SW Pacific	na	+	na	higher T	Thresher et al. 2007,	Redfish, Jackass morwong, Banded morwong
Three deep water fishes (>1000m)	SW Pacific	na	-	na	lower T	Thresher et al. 2007,	Orange roughy, Smooth oreo, Warty oreo; no change at medium depth
Blacklip abalone (<i>Haliotis rubra</i> , Haliotidae)	Tasmania	-	+	-	higher T	Helidoniotis & Haddon 2014	
Cod	E Atlantic	na	+/-	na	higher T	Rogers et al. 2011	Increased T has + effect in spring, but - effect in summer (above optimum)
Chinook salmon (<i>Oncorhynchus tshawytscha</i> , Salmonidae)	W USA rivers	na	+/-	na	higher T	Crozier et al. 2010	Increased T has + effect at low density, but - effect at high density (due to DD)
European plaice	North Sea	-	+	-	FIE	Walraven et al. 2010	L_{∞} change at 1%/year for F=0.4
Haddock (<i>Melanogrammus aeglefinus</i> , Gadidae)	Eastern Canada, Scotian Shelf	-	0	-	FIE	Neuheimer & Taggart 2010	measures of temperature-corrected length-at-age
Pacific hake (<i>Merluccius productus</i> , Merlucciidae)	USA and Canada west costs	na	+	-	unclear	Stewart et al. 2011	Step change in <i>k</i> (0.1 to 0.5) and L_{∞} (80 to 60 m) between 1975-1990 and 1991-2010 inferred
Cod	NW Atlantic	-	na	na	FIE	Swain 2011; Therkildsen et al. 2013	decrease in PMRN correlated to genetic change and fishing mortality

Sockeye salmon	Alaska	–	NA	na	FIE	Kendall et al. 2014	Observed change corresponded to the estimated selection differentials
Tiger flathead (<i>Platycephalus richardsoni</i> , Platycephalidae)	SE Australia	na	+/-	na	higher T	(Morrongiello and Thresher 2015)	<i>k</i> increased with temperature to a presumed optima, then declines
Estuary perch (<i>Macquaria colonorum</i> , Percichthyidae)	Victoria	na	+	na	higher T and increased freshwater flow	(Morrongiello et al. 2014)	Juvenile growth strongly related to temperature with a lesser flow effect, but adult growth relatively unaffected by environmental change
37 commercial fish stocks	mostly North Atlantic	–	na	na	FIE	Sharpe & Hendry 2009; Devine et al. 2012	Rate of change in PMRN correlated to fishing intensity
14 marine and freshwater fish stocks		na	na	–	FIE	Audzijonyte et al. 2013a	0.5-2% /year for F=0.5

Table 2. Pathways explaining the effects of temperature increase on ectotherm body size.

Pathway	Consequences	Examples
Oxygen concentration in water declines with temperature. Different surface area to volume ratios in small vs. large animals affect oxygen uptake	Large individuals with greater oxygen demands are more sensitive to temperature increases due to reduced relative surface area for oxygen diffusion. This results in disproportional mortality of oldest age classes, leading to smaller adult size-at-age and L_{∞}	Negative within-species relationship between size and temperature in aquatic species, especially in large bodied taxa (Forster et al. 2012); increased mortality from acute high temperatures in large individuals of marine invertebrate species (Peck et al. 2009); loss of larger eel pout (<i>Lota lota</i> , Lotidae) in the North Sea due to heat stress (Portner and Knust 2007); assemblage-wide average body weights predicted to shrink in a warmer

future due to oxygen limitation and the energetic costs of a large body size (Cheung et al. 2013)

Different response rate of anabolic and catabolic processes to temperature

Catabolic rates increase faster with temperature (and size, see above) than anabolic rates. As a result, increasing costs of catabolism at higher temperatures have a stronger effect on large individuals, leading to a cessation of growth at smaller sizes/ younger ages, disproportionate mortality of oldest age classes, and to smaller adult size-at-age and L_{∞}

Increased relative catabolic rate in larger and older individuals reduces their growth potential (Kozłowski et al. 2004)

Different responses of growth and reproductive processes to temperature

Changes in temperature may advance maturation irrespective of growth, resulting in small adult body size due to earlier energy allocation to reproduction

Earlier age at maturation caused by higher temperature independent of growth rate and body size in stickleback (*Pungitius pungitius*, Gasterosteidae) males (Kuparinen et al. 2011); smaller body size and size at maturity at higher temperatures despite faster growth rates in blacklip abalone (Helidoniotis and Haddon, 2014)

Temperature-dependent differences in size-fecundity or size-reproductive rate relationships

Fecundity rate may increase with size faster under cool conditions than warm conditions. In such a case, the advantages of delayed maturation decrease as temperatures warm leading to earlier maturation and decrease in L_{∞}

The size-fecundity relationship of a freshwater snail is steeper in cooler environments, meaning that it is maladaptive to be a larger size at maturation in warm environments (Arendt 2015); size-specific clutch size is greater at cooler temperatures in a reef fish (Kokita 2003)

Increased temperature leads to higher predation rates and increased natural mortality	Mortality, regardless of size and age, increases with temperature. Increased natural mortality leads to evolutionary response maximising earlier reproduction and decrease in L_{∞}	Active species have higher survival at elevated temperatures compared to sessile species, resulting in potential increases in predation at higher temperatures (Peck et al. 2009); predation rates on isopods are greater at warmer temperatures, likely causing the observed pattern of smaller size and age at maturity in warmer environments (Manyak-Davis et al. 2013)
---	--	---

Table 3. Mechanisms explaining how fishing and climate change can be expected to decrease life-history diversity in marine ectotherms

Mechanism and potential consequences	How it works	Examples
Climate change: depletion of phenotypic/ genetic variation due to directional selection on temperature reaction norms reduces the ability of populations and species to respond to additional change	Rapid and prolonged warming will result in strong selection against individuals with lower thermal tolerances or limited phenotypic plasticity	Decreased phenotypic variability in individual growth thermal reaction norms across 21 cohorts associated with warming waters in tiger flathead (Morrongiello and Thresher 2015)
Fishing: Depletion of phenotypic/genetic variation due to long-term directional selection of size-selective fisheries reduces the ability of populations and species to respond to additional change	Long-term size-selective fishing will produce strong selection against large-bodied fish.	Decreased phenotypic variability in size-at-age of Atlantic cod (Olsen et al. 2008)
Fishing: Reduced average age of the spawning females increases variability in spawning events	Most fishing selects oldest individuals, leading to demographic population truncation. Young females are more likely to skip spawning if their energy reserves drop	Average individual age of fished populations goes down, increasing the likelihood of skipped spawning and reducing spawning

season length (Jørgensen et al. 2006; Secor 2007; Hixon et al. 2014).

Fishing: Reduced average size and age of a spawning female leads to reduction in egg size and diversity.

Fishing selectively removes larger and older females. Larger fish have more fat content per body weight, higher gonadal mass to somatic mass, and can divert relatively more energy into reproduction. Older individuals often produce larger eggs with higher energy reserves compared to the young females. Larger eggs from older females may perform better under poorer environmental conditions

Lower reproductive rates in 25 marine fish due to age truncation (Venturelli et al. 2009). Importance of large fish for population resilience (Hixon et al. 2014).

Fishing: Reduced diversity of spawning behavior due to truncated age distribution in harvested stocks decreasing buffering capacity against variable environmental conditions and predation mortality in larval stages

Age distribution of fished stocks is usually truncated towards younger ages. Females of different ages often spawn at different places and times.

European plaice, Atlantic herring, cod and others (reviewed in Secor 2007)

Figure legends

Fig. 1. A: Von Bertalanffy growth curves for slow (dashed line) and fast (thick solid line) life-histories. An arrow indicates the suggested general trend towards faster life-histories. Thickened lines indicate the length of maturity for the different life-histories. **B.** Relationship between life-history parameters and natural mortality based on empirical and simulated data (modified from Carruthers et al. 2012).

Fig. 2. Implications of faster life-histories for stock biomass and productivity: four possible scenarios. **A:** Earlier maturation increases spawning stock biomass and productivity, compensating for smaller adult size, total biomass and carrying capacity remain unchanged. **B:** Earlier maturation leads to shorter lifespan as survival costs of reproduction reduce longevity. In combination with reduced adult size this leads to decreased carrying capacity. **C:** Earlier maturation increases the spawning stock biomass, egg production and recruitment, leading to increased productivity. **D:** Decreased size of an average female leads to lower egg production and recruitment, and to lower productivity.

Fig. 3. Suggested framework for precautionary and adaptive management that evaluates different factors that are likely to affect life-history traits and for which monitoring could be established. A trend towards faster life-histories (from dashed to thick solid line) is shown with a red arrow.

Fig. 1

A

B

Fig. 2

Fig. 3

Expected change in growth and maturation age leading to faster life-histories

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Audzijonyte, A;Fulton, E;Haddon, M;Helidoniotis, F;Hobday, AJ;Kuparinen, A;Morrongiello, J;Smith, ADM;Upston, J;Waples, RS

Title:

Trends and management implications of human-influenced life-history changes in marine ectotherms

Date:

2016-12-01

Citation:

Audzijonyte, A., Fulton, E., Haddon, M., Helidoniotis, F., Hobday, A. J., Kuparinen, A., Morrongiello, J., Smith, A. D. M., Upston, J. & Waples, R. S. (2016). Trends and management implications of human-influenced life-history changes in marine ectotherms. FISH AND FISHERIES, 17 (4), pp.1005-1028. <https://doi.org/10.1111/faf.12156>.

Persistent Link:

<http://hdl.handle.net/11343/291038>