Youth mental health services in Italy: An achievable dream?

Authors

Alice Masillo^{1*} & Martina Brandizzi^{1*}, Barnaby Nelson², Nella Lo Cascio^{1,3}, Riccardo Saba¹,
Juliana Fortes Lindau⁴, Ludovica Telesforo⁴, Dori Montanaro⁵, Marco D'Alema⁵, Paolo Girardi⁴,
Patrick McGorry², Paolo Fiori Nastro¹.

Affiliations:

1: Department of Neurology and Psychiatry, Sapienza University of Rome, Faculty of Medicine and Odontology, Rome 00156, Italy;

2: Orygen, The National Centre of Excellence in Youth Mental Health, University of Melbourne, 35 Poplar Rd (Locked Bag 10), Parkville, Victoria 3052, Australia;

3: Department of Neurosciences and Neuro-rehabilitation, Bambino Gesù Children's Hospital Rome, Italy

4: Neurosciences, Mental Health and Sensory Functions (NESMOS) Department, Sapienza University of Rome, Faculty of Medicine and Psychology, Sant'Andrea Hospital, Via di Grottarossa 1035, 00189 Rome, Italy;

5: Community Mental Health Service, ASL Rome H, 00041 Rome, Italy.

*Joint first authors

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/eip.12328](https://doi.org/10.1111/eip.12328)

Author Manuscript

Youth mental health services in Italy: An achievable dream?

Abstract

Aim

“Liberiamo il futuro” (LIF) project was designed to assess psychological problems of adolescents and young adults and to identify individuals at high-risk for developing a psychosis through a collaboration between a University team, Child and Adolescent Mental Health Services and Adult Mental Health Services. This paper presents the baseline demographic and clinical characteristics of the cohort, particularly the nature and severity of psychopathology.

Method

All help-seeking young people aged 12-35 years residing in the health district involved in LIF were invited to participate in the study and completed a battery of self-report and interviewer-administered measures of psychopathology and functioning at baseline.

Results

A total of 338 adolescents and young people (mean age 17.42) participated in the study. The majority of the sample (N=107, 35%) had an anxiety disorder, followed by mood disorders (N=62, 21%). Only 35 (12%) participants had no psychiatric diagnosis. After a screening phase, 166 (52%) individuals were assessed to detect the presence of an Ultra High Risk (UHR) state. Of these, 38.60% (N=64) met UHR criteria. Overall, the majority of the sample resulted moderately functionally impaired at baseline.

Conclusions

LIF project showed that psychological problems, associated with impaired psychosocial functioning, are very common among help-seeking young people. The help-seeking behaviour of young people is in contrast with the barriers presented by the Italian community mental health system that is modelled around adults' requirements. A need of a strong, stigma-free, young oriented system of care for young people up to the mid-20s emerged.

Introduction

Worldwide mental disorders are the fifth leading disorder category of global DALYs (disability-adjusted life years). Together with substance use disorders they accounted for 7.4% of total disease burden in 2010 (1). Despite the apparently small contribution of YLLs (years of life lost to premature mortality), they represent a striking and growing challenge for health systems in developed and developing regions (2), (3). The peak of onset of mental disease is from the early teens to the mid-twenties and the incidence and prevalence of mental illness in adolescents and young adults is the highest of any age group (4) (5).

The early intervention paradigm advocates detecting and treating mental illness at their earliest stages, during adolescence and early adulthood, with the goal of influencing the trajectory of illness evolution (preventing illness onset or improving prognosis; (6), (7)). This is not a simple task. The clinical profile of mental ill-health in young people is characterized by mixed sub-threshold symptom patterns (8). Adolescence is a fundamental transition life stage that leads to many changes both physically and mentally. The human body matures into adulthood and becomes ready for sexual and reproductive activities. Concurrently, adolescents struggle with the process of individuation, achieving their own identity distinct from their parents and relatives. A physiological crisis often occurs during this phase: adolescents may react to these fast changes in their lives with a kaleidoscope of behavioural changes and mood states. For example, becoming depressed, sleep

disturbances, appetite change, emotional swings, behavioural withdrawal, and apathy are common experiences after disturbances to interpersonal relationships (e.g., relationship break-ups) in this developmental stage(9),(8). Usually, these symptoms are short lived and much of the associated distress resolves quickly. However, they should not be dismissed as trivial purely because they are so common. Physiological difficulties sometimes-may hide a more severe inner-self fragility that struggles to cope with external pressures of maturation (i.e. more competitive education stages, more complex and intimate relationships, more frequent fights with parents/relatives etc.)(10), (11). This mixture of inner fragility and increasing outer changes may represent fertile ground for the development of a mental disease (12), (13). Early detection of these impaired mental states represents a challenge for clinicians and may be the gold standard for prevention. Despite this, access to, and use of, health services by young people to improve mental health outcomes is low (14), (15), (16), (17) . In general only a few young people (less than one in six) with mental disorders access services or receive appropriate care (18), (19). This confirms a paucity of service utilization just at the time when serious mental health problems are beginning to emerge (15). The current main service configuration of distinct Child and Adolescent Mental Health Services (CAMHS) and Adult Mental Health Services (AMHS) is a barrier to providing continuity of care, not restricted to age boundaries alone (20), (15), (21). Research findings indicate conceptual, clinical, and ideological differences between CAMHS and AMHS in relation to thresholds regarding acceptance criterion, professional differences and service structures/configurations which have been found to be an impediment to continuity of care for young people, especially those who make a transition from one system of care to another (15). For these reasons, the current main service configuration of CAMHS and AMHS has been described as the “weakest link in a system where it should be most robust” (22).

These issues of the lack in continuity of care and failure to tailor services specifically to young people are paramount in Italy. Data suggest that only a small proportion (19%) of adolescents

treated by CAMHS move to AMHS (23) and programs addressing this problem for various reasons struggle to be established in this country. With the aim of deeper assessing psychological problems of adolescents and young adults and identifying patients at high-risk for developing a psychosis through an extended collaboration between CAMHS and AMHS services, the “Liberiamo il futuro” (LIF) project was developed. LIF is a multicentre project carried out by the contribution of Sapienza University of Rome and six AMHS and six CAMHS located in one of the eight Local Health Districts of Rome, Italy, i.e. the Rome H area. Briefly, the main aim of LIF is to identify subjects at high-risk for developing a psychosis according to the at risk mental state for psychosis (UHR) (6) and the Basic Symptoms criteria (24) among help-seeking adolescents (age range: 12–18 years) and young adults (18–35 years). The inclusion criteria were: age between 12 and 35 years; IQ ≥ 70 ; sufficient knowledge of the Italian language; the disorder is not secondary to or correlated with a general medical condition; willingness and ability to provide free written informed consent (the informed consent was provided by parents or guardian in case of minors). The aim of the present study was to describe the baseline demographic and clinical characteristics of participants in the LIF project.

Method

Setting

Data were collected in six CAMHS and six AMHS located in the area of Roma H and involved in the LIF project between January 2012 and June 2015.

Participants

The study included all help-seeking young people since January 2012 aged 12–35 years old (25, 26) who resided in the Rome H area. Exclusion criteria for the LIF project include psychiatric disorders or symptoms due to a somatic factor or psychotropic substances, IQ below 70, and insufficient knowledge of Italian or other communication problems severely impairing clinical assessments.

338 individuals and/or their parents or guardians gave written informed consent to participate. Ethical approval was obtained from the local research ethics committee. The mean age was 17.42 years (SD: 5.62; Median: 16, Mode: 12). One hundred-seventy-three participants were female (51.2%) and 165 (48.8%) were male. Full details of the study are provided in Brandizzi et al. 2014 (27).

Instruments

Case-history, socio-demographic and psychosocial variables were recorded during a baseline clinical assessment using a non-standardized questionnaire.

Prodromal Questionnaire. Prodromal symptoms of psychosis were assessed with the Prodromal Questionnaire-92 (PQ92; (28)), a self-report screening questionnaire that aims to identify individuals who may benefit from a clinical diagnostic interview. The 92 true/false items can be divided into four major subscales: (1) positive symptoms; (2) negative symptoms; (3) disorganized symptoms; and (4) general symptoms. Patients who score 18 or more on the PQ-92 positive symptoms subscale (29) or evaluated as possibly at risk according to clinical impression were examined with the Structured Interview for Psychosis-risk Syndromes (SIPS) and Schizophrenia Proneness Instrument- Adult (SPI-A (30)) and Child and Adolescents (SPI-CY (31)) to detect if they are at risk for developing a psychosis according to the ultra-high risk (UHR) (32) and basic symptom (BS) criteria (24).

Global Functioning: Social and Role Scales. Social and role functioning was assessed using the Global Functioning: Social (GFSS) and Global Functioning: Role scales (GFRS) (33). The GFSS assesses peer relationships, peer conflict, age-appropriate intimate relationships, and involvement with family members. The GFRS rates performance and amount of support needed in one's specific role (i.e., school or work). For both scales, scores range from 1 to 10 (10 = superior functioning to 1

= extreme dysfunction). Ratings for each of the two GF scales were based on available clinical information.

Interpersonal Sensitivity Measure (IPSM). To measure interpersonal sensitivity, we used the Italian version (34) of Interpersonal Sensitivity Measure (IPSM) (35), a 36-item self-report questionnaire that identifies people with an undue and excessive awareness of and sensitivity to the behaviour and mood of others.

Structured Interview for Psychosis-risk Syndromes (SIPS). The SIPS (36) is a clinician-administered, semi-structured interview specifically designed to establish risk of psychosis. There are four major symptom dimensions on the Scale of Psychosis-Risk Symptoms (SOPS): positive, negative, disorganized and general symptoms.

Schizophrenia Proneness Instrument- Adult (SPI-A) and Child and Adolescents (SPI-CY) (30), (31). SPI-A and SPI-CY are two semi-structured interviews derived from Bonn Scale for the Assessment of Basic Symptoms (BSABS; Gross (37, 38)). Basic Symptoms (BS) are subjectively experienced sub-clinical disturbances in drive, affect, thinking, speech, perception, motor action, central vegetative functions and stress tolerance. Among all BS, COPER criteria (10 cognitive-perceptive BS) and the COGDIS criteria (the 9 cognitive BS that are the most predictive of later psychosis (39)) were assessed.

Structured Clinical Interview (SCID-I) for DSM-IV and Kiddie-Schedule for Affective Disorders and Schizophrenia Present and Lifetime (K-SADS-PL): Axis I diagnoses were evaluated with the SCID-I (40) (age 19-35) and with the K-SADS-PL (41) (age 12-18 years).

Statistical analysis

Analyses were conducted using SPSS 20.0.0. Descriptive analysis including mean and SD for continuous variables and absolute and relative frequencies for categorical variables were performed. Chi-square test and Fisher's Exact Test were employed for categorical variables. One-way ANOVA was used for comparisons between groups of subjects.

Results

Demographic characteristics

A summary of the participant characteristics is provided in Table 1. Three hundred thirty-eight individuals were involved in the study. About half of participants were female (51%; $n = 173$) and the mean age of participants was 17.42 years (SD: 5.6), with 36% ($N=122$) aged 12–14 years, 37% ($N=125$) aged 15–18, and 15% ($N=52$) aged 19–24 years, and 12% ($N=39$) aged 25–35. There were no significant differences between male (18.4 years, SD: 5.63) and female (18.2 years, SD: 5.60) with respect to mean age at baseline assessment ($F = 4.635$; $P = 0.201$). The majority of participants indicated that their parents were Italian born (89.4% of fathers and 89.8% of mothers) and to currently be single ($N=328$; 97%) and residing in a semi-urban area ($N=208$; 63%). Seventy-nine per cent ($N=262$) of the sample indicated being enrolled in some form of education, 12% ($N=38$) were unemployed and 10% ($N=33$) were currently employed. According with the mean age of the sample the majority of them had completed primary or a lower secondary school ($N=253$, 68%). Not unexpectedly, education was associated with age, with 63.1% of 12–14-years-olds completed primary school, compared with 78.4% of 15–18-year-olds who completed lower secondary school and 50.8% of 19–24-year-olds who completed upper secondary school. Female participants were significantly more likely to have a higher level of education than male participants ($\chi^2 = 22.22$; $P < 0.001$).

Clinical characteristics

General clinical characteristics are presented in Table 2. Of all young people entering our services during the study period, 6% (N=18) were referred by a psychiatrist, 19% (N=62) were self-referrals, 33% (N=107) were arranged by family members, 22% (N=71) were referred by standard health-referral pathways, such as GPs, other health professionals or contact with public mental health services, and the remainder (19%, N= 63) were referred from schools and legal agencies. The main reason for referral for the majority of the sample was relational or school problems (33%, N=110) and anxiety symptoms (29%, N= 96). There were gender related significant differences regarding reason for referral ($\chi^2=16.3$, $P<0.005$), with females more commonly being referred to the services for mood-related symptoms than males (46 vs 21). Seventy-eight individuals (23%) admitted having used alcohol or illicit substances at least once during their lifetime. Only 35 (12%) participants had no psychiatric diagnosis. The majority of the sample (N=107, 35%) had an anxiety disorder, followed by mood disorders (N=62, 21%). There were between gender differences for psychiatric diagnosis ($\chi^2=18.3$; $p<0.05$), with females diagnosed more frequently with anxiety and mood disorder than males. More than half of participants (N=180, 54%) have never had psychiatric treatment before asking help of the services involved, whereas 32% (N=107) had been on psychotherapeutic treatment and 8% (N=28) on psychiatric medications during their lifetime.

Mental health characteristics

Participant scores on interviews and self-rated measures of psychopathology are presented in Table 3.

The prodromal symptoms questionnaire:

Among the whole sample 96 participants (29%) scored 18 or more on the positive symptoms scale of the PQ, with a mean score of 14.22 (SD: 9.00). Female subjects tended to have more positive

($F=5.27$; $P<0.05$), negative ($F=8.65$; $P<0.005$), general ($F=14.24$; $P<0.005$) self-reported symptoms compared with males, as well as higher total scores on the PQ-92 ($F=8.48$; $P<0.005$).

Global, social and role functioning:

The mean GAF score was 64.44 (SD: 11.92), indicating some mild psychiatric symptoms or some difficulty in social, occupational, or school functioning. The mean role functioning (measured by GFRS) was 6.47 (SD: 1.54) indicating moderately/mildly impaired functioning. The mean social functioning (measured by GFSS) was 6.69 (SD: 1.27) referring to a moderate/ mild impairment in social functioning that is beyond an expectable reaction to a psychosocial stressor. There were between gender differences in functioning with males reporting lower global ($F=7.422$; $p<0.05$), role ($F=12.807$; $p<0.005$) and social ($F=11.316$; $p<0.005$) functioning levels than females.

Interpersonal sensitivity:

The mean score of the IPSM scale was 91.04 (SD: 19.04), with females reporting higher scores than males in total scores ($F=13.72$; $P<0.005$), in the interpersonal awareness subscale ($F=15.085$; $P<0.005$) and in the separation anxiety subscale ($F=22.80$; $P<0.005$). These scores indicated difficulties in interpersonal relationships, due to an undue awareness of behaviour and mood of others and to a fragility of inner-self.

Psychotic prodromal symptoms:

Based on a combination of PQ-92 positive symptoms scale scores (cut-off of 18) and clinical impression, of all participants ($N=338$), 166 (52%) individuals were assessed with the SIPS and with the COPER and COGDIS criteria. Of these 166 participants, 55% ($N=92$) were not UHR or experiencing First Episode Psychosis (FEP) according to the SIPS criteria; 38.60% ($N=64$) met UHR criteria, mainly the Attenuated Psychosis Syndrome (APS) criteria ($N=56$, 34%). 92 subjects (58.2%) reported COPER and/or COGDIS criteria (the BS criteria). Sixty subjects (38%) of subjects reported only COGDIS criteria. Twenty-six subjects (16%) reported both UHR and COGDIS criteria. There were not significant gender differences in psychosis risk criteria (see table 4). Among the fully-assessed participants, 10 (6%) were suffering from FEP.

Discussion

The purpose of this paper was to describe the baseline demographic and clinical characteristics of the LIF cohort, particularly the nature and severity of psychopathology in this population. One of the main aims of LIF was to answer the needs of young people with psychological problems within the area of Rome where to date services for youth mental health have not been available. In order to conduct this study collaboration between CAMHS and AMHS services was necessary. The 338 young people recruited at baseline represent a relatively heterogeneous cohort in relation to the nature of their mental health problems and levels of functional impairment. In our sample half of participants were male (49%; $n = 165$), indicating that there were not differences in service utilisation between gender. This is in contrast with evidence in most high-income countries, indicating a consistent pattern in services utilisation with women and older people using specialist services more than men and younger people (42). Also along young people females seem to use more often mental health services than male (43). One possible explanation of this result is that most participants were very young and were more often accompanied by parents or relatives. In fact, the main source of referral of our study were family members (33%). This result was in line with recent studies showing that in Italy family members are often involved in the care of mentally ill patients (44), (45). Relatives played different roles in each phase of the disorder. In particular, at the onset of psychosis, relatives could promote or delay referral to mental health professionals (46). For these reasons the improvement of relatives' knowledge about mental disorders should be prioritized in mental health research and practice, mainly to reduce the delay in the detection and treatment of mental disorders among young people. According to studies about accessibility and pathways to psychiatric care in Italian community-based mental health system (47), (48), the second most common source of referral were GPs and other health professionals in contact with public mental health services (22%). However, these percentages were not high; these results underlined

that a stronger collaboration of psychiatrists and mental health professionals with general practitioners is still needed.

The majority of the sample was less than 18 years old (mean age of 17.42, SD: 5.6). The current service configuration is of distinct CAMHS and AMHS services with 18 years old as the age of transfer from a service to another. As McGorry (22) has highlighted, our results suggest that where the system should be the most robust, indeed it shows its weakest part. Even though the main reason of referral to services was relational and school difficulties (33%), our findings showed that only 12% of young people help-seekers for psychological problems had no psychiatric diagnosis. The remainder were diagnosed with anxiety (35%) and mood disorders (21%), in line with epidemiological studies that consistently indicate that anxiety disorders are the most prevalent mental disorders among young people (49). Moreover, according to previous findings (50), (49), anxiety disorders were significantly more prevalent in females (57% vs 43%). These data supported the importance of a meticulous clinical assessment even if the main complaint appears trivial. Among the whole sample, following the screening phase, 166 (52%) individuals were assessed with the SIPS. Of these, 64 (38.6%) were identified as being at ultra high risk for developing psychosis. These results are remarkably consistent with the UHR prevalence amongst referrals to youth-focused mental health services in Australia, called Headspace services (43). The importance of early detection of psychosis has been confirmed by a recent meta-analysis showing that the UHR state is associated with 36% transition risk of developing psychosis within the first 3 years of clinical presentation, and the risk progressively increases across this period (51). The high rate of UHR individuals confirmed that a screening method identifies more UHR+ patients at entry to mental health services, which would otherwise go undetected and untreated (52). Furthermore 58% of the sample reported the presence of COPER and/or COGDIS criteria; while only 16% reported both UHR and COGDIS criteria. As emerged from a recent study a combined UHR and BS approach may improve the prediction of psychosis and the diagnostic/outcome-related

homogeneity of risk groups (39). Following these results we may hypothesize that this sub-group may be in particular need of clinical attention. Follow-up assessments among this group are currently being conducted to evaluate transition rates and other clinical outcomes.

In contrast with recent literature (53), (43), only 23% of the total sample reported use of licit and illicit substances (e.g. alcohol, cannabis, cocaine, opioids, etc.). One possible explanation may be that adolescents and young adults involved in the study underreported their use of drugs and alcohol, even though professionals performed the interview.

Moreover, having a psychiatric disorder, or symptoms due to psychotropic substances was an exclusion criteria of LIF. Evaluating if a substance use is a primary or secondary problem in adolescence is not a simple task; drug use may be considered as a self-treatment of emerging psychiatric problems(54), (55), (56). Subjects who presented with symptoms (such as panic attacks, sub-threshold psychotic symptoms etc.) only in the context of substance use (intoxication/withdrawal) were not involved in LIF, with the awareness of excluding many “limit” cases.

In line with Scott (57), although most participants were in the early phases of illness, disability levels were already pronounced. The mean score of social and role functioning showed a mild/moderate impairment in peers’ relationships and age-appropriate role functioning. Interestingly, of the whole sample, 36% of participants reported a decreasing role functioning and 37% of social functioning during the year before the first assessment. Moreover, our sample reported high level of interpersonal sensitivity. This personality trait was found to be associated with avoidant coping strategies, persecutory ideations and depressive symptoms among non-clinical and ultra-high-risk samples (58), (59), (60), (61). Highly interpersonal sensitive individuals also report low social functioning levels (62). In line with previous findings (58), (60), (62), UHR individuals showed higher level of interpersonal sensitivity compared to other help-seekers and this personality feature could be associated with worse functional outcome. To confirm this data, follow up assessments are in progress.

The psychosocial functioning impairment of our sample has clinical relevance. Recent studies indicated that UHR individuals with poor baseline functioning continue to show poor functioning after several years, even when their positive symptoms never cross the threshold into full psychotic severity (63), (64), (65). Moreover, these individuals are often still in need of clinical intervention (66), (67), (68).

Limitations

Our results should be interpreted in view of the limitations of this study. The main limitation is the lack of follow-up data to rate the continuity of care between CAMHS and AMHS. Another limitation is that axis I diagnoses were evaluated with the SCID-I according to DSM-IV and not to the last version of the manual (DSM-5(69)). Moreover substance use was evaluated just by a clinical interview and not using a specific questionnaire. Data on interrater reliability for structured and semi-structured interviews (SCID, K-SADS, SPI-CY, SPI-A, SIPS) are not available. However, these instruments were performed for trained researchers. Furthermore data about how many people did not agree to participate in the study are not available. It is therefore possible that there was a sampling bias of those who were most unwell not agreeing to participate.

Conclusions

Our results showed that psychological problems are very common among help-seeking young people. Psychiatric symptoms are often associated with impaired psychosocial functioning that is a negative predictor of outcome and needs to be targeted with focused treatment. The mean age of our sample highlights the problem of transferring from a child and adolescent service to an adult service at the age of 18. Nevertheless, the help-seeking behaviour of young people is in contrast with the barriers presented by the Italian community mental health system that is modelled around adults'

needs. As many researchers have argued, to try to go beyond these barriers is necessary in order to build a strong, stigma-free and effective system of care for young people up to the mid-20s (22). This reform is gaining significant ground in Australia (5, 43). In 2006, the Australian Federal Government established Headspace, the National Youth Mental Health Foundation, to promote and support early intervention in 12–25-year-olds. From this initiative, 55 Headspace clinical services currently operate throughout Australia. They are specialist-, youth- and carer-friendly youth services that ensure that young people's needs are met with a coordinated and integrated response. They include mental health services (including alcohol and drug services), primary health-care services (including sexual health consultations) and vocational assistance. Headspace clinical services provide a youth-friendly, stigma-free environment and offer a soft-entry point more appealing for young people. Similar programmes have since been established in other countries (UK, Ireland and Denmark, (43), (70)). For the Italian psychiatric system, this model should be a way forward to build an effective mental disorder prevention service. Investing in this direction is an important challenge for the future of Italian psychiatry.

References

1. Whiteford HA, Degenhardt L, Rehm J, Baxter AJ, Ferrari AJ, Erskine HE, et al. Global burden of disease attributable to mental and substance use disorders: findings from the Global Burden of Disease Study 2010. *Lancet*. 2013;382(9904):1575-86.
2. Wang PS, Aguilar-Gaxiola S, Alonso J, Angermeyer MC, Borges G, Bromet EJ, et al. Use of mental health services for anxiety, mood, and substance disorders in 17 countries in the WHO world mental health surveys. *Lancet*. 2007;370(9590):841-50.
3. Wang PS, Angermeyer M, Borges G, Bruffaerts R, Tat Chiu W, G DEG, et al. Delay and failure in treatment seeking after first onset of mental disorders in the World Health Organization's World Mental Health Survey Initiative. *World psychiatry : official journal of the World Psychiatric Association (WPA)*. 2007;6(3):177-85.
4. Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of general psychiatry*. 2005;62(6):593-602.
5. Patrick D McGorry RP, Ian B Hickie and Anthony F Jorm. Investing in youth mental health is a best buy. *Medical Journal of Australia*. 2007;187(7):5.
6. Fusar-Poli P, Borgwardt S, Bechdolf A, Addington J, Riecher-Rossler A, Schultze-Lutter F, et al. The psychosis high-risk state: a comprehensive state-of-the-art review. *JAMA Psychiatry*. 2013;70(1):107-20.
7. Nelson B, Yuen HP, Wood SJ, Lin A, Spiliotacopoulos D, Bruxner A, et al. Long-term follow-up of a group at ultra high risk ("prodromal") for psychosis: the PACE 400 study. *JAMA Psychiatry*. 2013;70(8):793-802.
8. McGorry PD, Goldstone S. Is this normal? Assessing mental health in young people. *Australian family physician*. 2011;40(3):94-7.

9. Montgomery MJ. Psychosocial intimacy and identity from early adolescence to emerging adulthood. *Journal of Adolescent Research*. 2005;20(3):346-74.
10. Herres J. Adolescent coping profiles differentiate reports of depression and anxiety symptoms. *Journal of affective disorders*. 2015;186:312-9.
11. Tandon SD, Dariotis JK, Tucker MG, Sonenstein FL. Coping, Stress, and Social Support Associations With Internalizing and Externalizing Behavior Among Urban Adolescents and Young Adults: Revelations From a Cluster Analysis. *Journal of Adolescent Health*. 2013;52(5):627-33.
12. Fagioli M. Teoria della nascita e castrazione umana: L'Asino d'Oro; 2012.
13. van Os J, Linscott RJ, Myin-Germeys I, Delespaul P, Krabbendam L. A systematic review and meta-analysis of the psychosis continuum: evidence for a psychosis proneness-persistence-impairment model of psychotic disorder. *Psychological medicine*. 2009;39(2):179-95.
14. de Girolamo G, Dagani J, Purcell R, Cocchi A, McGorry PD. Age of onset of mental disorders and use of mental health services: needs, opportunities and obstacles. *Epidemiology and psychiatric sciences*. 2012;21(1):47-57.
15. Singh SP. Transition of care from child to adult mental health services: the great divide. *Current opinion in psychiatry*. 2009;22(4):386-90.
16. Burgess PM, Pirkis JE, Slade TN, Johnston AK, Meadows GN, Gunn JM. Service use for mental health problems: findings from the 2007 National Survey of Mental Health and Wellbeing. *The Australian and New Zealand journal of psychiatry*. 2009;43(7):615-23.
17. Costello EJ, Egger H, Angold A. 10-year research update review: the epidemiology of child and adolescent psychiatric disorders: I. Methods and public health burden. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2005;44(10):972-86.
18. Kataoka SH, Zhang L, Wells KB. Unmet need for mental health care among U.S. children: variation by ethnicity and insurance status. *The American journal of psychiatry*. 2002;159(9):1548-55.

19. Pottick KJ, Bilder S, Vander Stoep A, Warner LA, Alvarez MF. US patterns of mental health service utilization for transition-age youth and young adults. *The journal of behavioral health services & research*. 2008;35(4):373-89.
20. Davis M, Geller JL, Hunt B. Within-state availability of transition-to-adulthood services for youths with serious mental health conditions. *Psychiatric services (Washington, DC)*. 2006;57(11):1594-9.
21. Singh SP, Paul M, Ford T, Kramer T, Weaver T, McLaren S, et al. Process, outcome and experience of transition from child to adult mental healthcare: multiperspective study. *The British journal of psychiatry : the journal of mental science*. 2010;197(4):305-12.
22. McGorry PD. The specialist youth mental health model: strengthening the weakest link in the public mental health system. *The Medical journal of Australia*. 2007;187(7 Suppl):S53-6.
23. Stagi P, Galeotti S, Mimmi S, Starace F, Castagnini AC. Continuity of care from child and adolescent to adult mental health services: evidence from a regional survey in Northern Italy. *European child & adolescent psychiatry*. 2015.
24. Schultze-Lutter F. Subjective symptoms of schizophrenia in research and the clinic: the basic symptom concept. *Schizophrenia bulletin*. 2009;35(1):5-8.
25. Kessler RC, Amminger GP, Aguilar-Gaxiola S, Alonso J, Lee S, Ustun TB. Age of onset of mental disorders: a review of recent literature. *Current opinion in psychiatry*. 2007;20(4):359-64.
26. Kessler RC, Angermeyer M, Anthony JC, R DEG, Demyttenaere K, Gasquet I, et al. Lifetime prevalence and age-of-onset distributions of mental disorders in the World Health Organization's World Mental Health Survey Initiative. *World psychiatry : official journal of the World Psychiatric Association (WPA)*. 2007;6(3):168-76.
27. Brandizzi M, Schultze-Lutter F, Masillo A, Lanna A, Curto M, Lindau JF, et al. Self-reported attenuated psychotic-like experiences in help-seeking adolescents and their association with age, functioning and psychopathology. *Schizophr Res*. 2014;160(1-3):110-7.

28. Loewy RL, Bearden CE, Johnson JK, Raine A, Cannon TD. The prodromal questionnaire (PQ): preliminary validation of a self-report screening measure for prodromal and psychotic syndromes. *Schizophr Res.* 2005;77(2-3):141-9.
29. Ising HK, Veling W, Loewy RL, Rietveld MW, Rietdijk J, Dragt S, et al. The validity of the 16-item version of the Prodromal Questionnaire (PQ-16) to screen for ultra high risk of developing psychosis in the general help-seeking population. *Schizophrenia bulletin.* 2012;38(6):1288-96.
30. Schultze-Lutter F, Addington J, Ruhrmann S, Klosterkötter J. Schizophrenia Proneness Instrument, Adult version (SPI-A). . Editore. GF, editor. Rome2007.
31. Fux L, Walger P, Schimmelmann BG, Schultze-Lutter F. The Schizophrenia Proneness Instrument, Child and Youth version (SPI-CY): Practicability and discriminative validity. *Schizophrenia Research.* 2013;146(1–3):69-78.
32. Miller TJ, McGlashan TH, Rosen JL, Somjee L, Markovich PJ, Stein K, et al. Prospective diagnosis of the initial prodrome for schizophrenia based on the Structured Interview for Prodromal Syndromes: preliminary evidence of interrater reliability and predictive validity. *The American journal of psychiatry.* 2002;159(5):863-5.
33. Cornblatt BA, Auther AM, Niendam T, Smith CW, Zinberg J, Bearden CE, et al. Preliminary findings for two new measures of social and role functioning in the prodromal phase of schizophrenia. *Schizophrenia bulletin.* 2007;33(3):688-702.
34. Masillo A, Valmaggia LR, Lanna A, Brandizzi M, Lindau JF, Curto M, et al. Validation of the Italian version of interpersonal sensitivity measure (IPSM) in adolescents and young adults. *Journal of affective disorders.* 2014;156:164-70.
35. Boyce P, Parker G. DEVELOPMENT OF A SCALE TO MEASURE INTERPERSONAL SENSITIVITY. *Australian and New Zealand Journal of Psychiatry.* 1989;23(3):341-51.
36. McGlashan T, Walsh B, Woods S. *The Psychosis-Risk Syndrome: Handbook for Diagnosis and Follow-Up*: Oxford University Press, USA; 2010.

37. Gross G. The 'basic' symptoms of schizophrenia. *The British journal of psychiatry Supplement*. 1989(7):21-5; discussion 37-40.
38. Huber G, Gross G. The concept of basic symptoms in schizophrenic and schizoaffective psychoses. *Recenti progressi in medicina*. 1989;80(12):646-52.
39. Schultze-Lutter F, Klosterkötter J, Ruhrmann S. Improving the clinical prediction of psychosis by combining ultra-high risk criteria and cognitive basic symptoms. *Schizophr Res*. 2014;154(1-3):100-6.
40. First MB, Spitzer RL, Gibbon M, Williams JBW. Structured Clinical Interview for DSM-IV® Axis I Disorders (SCID-I), Clinician Version, Administration Booklet: American Psychiatric Publishing; 2012.
41. Kaufman B, Brent, Rao & Ryan. Kiddie-Sads-Present and Lifetime Version (K-SADS-PL): Kaufman, Birmaher, Brent, Rao & Ryan; 1996.
42. Hansen AH, Høy A. Gender differences in the use of psychiatric outpatient specialist services in Tromsø, Norway are dependent on age: a population-based cross-sectional survey. *BMC health services research*. 2015;15(1):477.
43. Purcell R, Jorm AF, Hickie IB, Yung AR, Pantelis C, Amminger GP, et al. Demographic and clinical characteristics of young people seeking help at youth mental health services: baseline findings of the Transitions Study. *Early intervention in psychiatry*. 2014:n/a-n/a.
44. Del Vecchio V, Luciano M, Sampogna G, De Rosa C, Giacco D, Tarricone I, et al. The role of relatives in pathways to care of patients with a first episode of psychosis. *The International journal of social psychiatry*. 2015;61(7):631-7.
45. Luciano M, Sampogna G, del Vecchio V, Giacco D, Mule A, de Rosa C, et al. The family in Italy: cultural changes and implications for treatment. *International review of psychiatry (Abingdon, England)*. 2012;24(2):149-56.

46. Magliano L, Fiorillo A, Malangone C, De Rosa C, Maj M. Patient functioning and family burden in a controlled, real-world trial of family psychoeducation for schizophrenia. *Psychiatric services* (Washington, DC). 2006;57(12):1784-91.
47. Amaddeo F, Zambello F, Tansella M, Thornicroft G. Accessibility and pathways to psychiatric care in a community-based mental health system. *Social psychiatry and psychiatric epidemiology*. 2001;36(10):500-7.
48. Volpe U, Fiorillo A, Luciano M, Del Vecchio V, Palumbo C, Calo S, et al. Pathways to mental health care in Italy: results from a multicenter study. *The International journal of social psychiatry*. 2014;60(5):508-13.
49. Rockhill C, Kodish I, DiBattisto C, Macias M, Varley C, Ryan S. Anxiety disorders in children and adolescents. *Current problems in pediatric and adolescent health care*. 2010;40(4):66-99.
50. Costello EJ, Mustillo S, Erkanli A, Keeler G, Angold A. Prevalence and development of psychiatric disorders in childhood and adolescence. *Archives of general psychiatry*. 2003;60(8):837-44.
51. Fusar-Poli P, Bonoldi I, Yung AR, Borgwardt S, Kempton M, Barale F, et al. Predicting psychosis: a meta-analysis of transition outcomes in individuals at high clinical risk. *Archives of general psychiatry*. 2012;69(3):1-10.
52. Rietdijk J, Klaassen R, Ising H, Dragt S, Nieman D, van de Kamp J, et al. Detection of people at risk of developing a first psychosis: comparison of two recruitment strategies. *Acta psychiatrica Scandinavica*. 2012;126(1):21-30.
53. Hermens DF, Scott EM, White D, Lynch M, Lagopoulos J, Whitwell BG, et al. Frequent alcohol, nicotine or cannabis use is common in young persons presenting for mental healthcare: a cross-sectional study. *BMJ open*. 2013;3(2).
54. Khantzian EJ. The self-medication hypothesis of substance use disorders: a reconsideration and recent applications. *Harvard review of psychiatry*. 1997;4(5):231-44.

55. Mariani JJ, Khantzian EJ, Levin FR. The self-medication hypothesis and psychostimulant treatment of cocaine dependence: an update. *The American journal on addictions / American Academy of Psychiatrists in Alcoholism and Addictions*. 2014;23(2):189-93.
56. Griffin KW, Lowe SR, Acevedo BP, Botvin GJ. Affective Self-Regulation Trajectories During Secondary School Predict Substance Use Among Urban Minority Young Adults. *Journal of child & adolescent substance abuse*. 2015;24(4):228-34.
57. Scott EM, Hermens DF, Glozier N, Naismith SL, Guastella AJ, Hickie IB. Targeted primary care-based mental health services for young Australians. *The Medical journal of Australia*. 2012;196:136-40.
58. Valmaggia LR, Freeman D, Green C, Garety P, Swapp D, Antley A, et al. Virtual reality and paranoid ideations in people with an 'at-risk mental state' for psychosis. *The British journal of psychiatry Supplement*. 2007;51:s63-8.
59. Green CEL, Freeman D, Kuipers E, Bebbington P, Fowler D, Dunn G, et al. Paranoid Explanations of Experience: A Novel Experimental Study. *Behavioural and Cognitive Psychotherapy*. 2011;39(1):21-34.
60. Masillo A, Day F, Laing J, Howes O, Fusar-Poli P, Byrne M, et al. Interpersonal sensitivity in the at-risk mental state for psychosis. *Psychological medicine*. 2012;42(9):1835-45.
61. Joo HJ, Yeon B, Lee KU. The impact of personality traits on emotional responses to interpersonal stress. *Clinical psychopharmacology and neuroscience : the official scientific journal of the Korean College of Neuropsychopharmacology*. 2012;10(1):54-8.
62. Masillo A, Valmaggia LR, Saba R, Brandizzi M, Lindau JF, Solfanelli A, et al. Interpersonal sensitivity and functioning impairment in youth at ultra-high risk for psychosis. *European child & adolescent psychiatry*. 2015.
63. Yung AR, Yuen HP, Berger G, Francey S, Hung TC, Nelson B, et al. Declining transition rate in ultra high risk (prodromal) services: dilution or reduction of risk? *Schizophrenia bulletin*. 2007;33(3):673-81.

64. Addington J, Cornblatt BA, Cadenhead KS, Cannon TD, McGlashan TH, Perkins DO, et al. At clinical high risk for psychosis: outcome for nonconverters. *The American journal of psychiatry*. 2011;168(8):800-5.
65. Schlosser DA, Jacobson S, Chen Q, Sugar CA, Niendam TA, Li G, et al. Recovery from an at-risk state: clinical and functional outcomes of putatively prodromal youth who do not develop psychosis. *Schizophrenia bulletin*. 2012;38(6):1225-33.
66. Fulford D, Niendam TA, Floyd EG, Carter CS, Mathalon DH, Vinogradov S, et al. Symptom dimensions and functional impairment in early psychosis: More to the story than just negative symptoms. *Schizophrenia Research*. 2013;147(1):125-31.
67. Simon AE, Borgwardt S, Riecher-Rossler A, Velthorst E, de Haan L, Fusar-Poli P. Moving beyond transition outcomes: meta-analysis of remission rates in individuals at high clinical risk for psychosis. *Psychiatry Res*. 2013;209(3):266-72.
68. Brandizzi M, Valmaggia L, Byrne M, Jones C, Iwegbu N, Badger S, et al. Predictors of functional outcome in individuals at high clinical risk for psychosis at six years follow-up. *Journal of psychiatric research*. 2015;65:115-23.
69. APA. *Diagnostic and Statistical Manual of Mental Disorders: Dsm-5*: Amer Psychiatric Pub Incorporated; 2013.
70. Illback RJ, Bates T. Transforming youth mental health services and supports in Ireland. *Early intervention in psychiatry*. 2011;5 Suppl 1:22-7.

		N	Mean	SD	95% CI		Statistics		
					L	U	F	Sig.	
Age	M	165	17.06	5.63	16.20	17.93	1.298	0.255	
	F	173	17.76	5.60	16.92	18.60			
	Total	338	17.42	5.62	16.82	18.02			
		Tot		M (N=165)		F(N=173)		Statistics	
		N	%	N	%	N	%	χ ²	P
Age groups	12-14	122	36	69	56.60	53	43.40	4.635	0.201
	15-18	125	37	55	44.00	70	56.00		
	19-25	52	15	23	44.20	29	55.80		
	26-35	39	12	18	46.20	21	53.80		
Father nationality	Italian	295	89	147	49.8	148	50.2	4.35	0.478
	Not Italian	35	11	15	42.9	20	57.1		
Mother nationality	Italian	298	90	149	50.0	149	50.0	0.33	0.369
	Not Italian	34	10	14	41.2	20	58.8		
Marital status	Single	328	97	161	49.10	167	50.90	2.014	0.365
	Married	7	2	3	42.90	4	57.10		
	Divorced	2	1	0	0.00	2	100.00		
Education	Primary school	88	26	57	64.80	31	35.20	22.225	0.000*
	Lower secondary school	165	49	64	38.80	101	61.20		
	Upper general secondary school	63	19	27	42.90	36	57.10		
	Upper vocational secondary school	14	4	11	78.60	3	21.40		
	Graduated	6	2	4	66.70	2	33.30		
Occupation	Student	262	79	131	50.00	131	50.00	3.125	0.537
	Unemployed	38	12	14	41.20	22	58.80		
	Employed	33	10	16	52.90	17	47.10		
Living area	Suburb	80	24	37	48.50	43	51.50	2.139	0.544
	Semi-suburb area	208	63	105	50.50	103	49.50		
	City-centre	45	14	19	42.20	26	57.80		

Table 1: summary of demographic characteristics of LIF participants (N=338).

		Tot		M		F		Statistics	
		N	%	N	%	N	%	χ^2	P
Source of referral	Psychiatrist	18	6	10	55.60	8	44.40	8.057	0.09
	Other health professionals	71	22	30	42.30	41	57.70		
	School or legal agencies	63	19	40	63.50	23	36.50		
	Family	107	33	49	45.80	58	54.20		
	Self-referrals	62	19	27	43.50	35	56.50		
Reason for referral	Anxiety	96	29	43	44.80	53	55.20	16.352	0.003*
	Depression /mood	67	20	21	31.30	46	68.70		
	Psychotic symptoms	16	5	12	75.00	4	25.00		
	Relational or school difficulties	110	33	63	57.30	47	42.70		
	Other	44	13	22	50.00	22	50.00		
Psychotropic substance use	No	260	77	128	49.20	132	50.80	0.077	0.781
	Yes	78	23	37	47.40	41	52.60		
Psychiatric diagnosis	No diagnosis	35	12	21	60.00	14	40.00	18.344	0.011*
	Anxiety	107	35	46	43.00	61	57.00		
	Mood	62	21	22	35.50	40	64.50		
	Psychosis	10	3	6	60.00	4	40.00		
	PD	12	4	5	41.70	7	58.30		
	Adjustment	26	9	18	69.20	8	30.80		
	Childhood (ADHD, conduct disorder)	34	11	23	67.60	11	32.40		
	other	16	5	6	37.50	10	62.50		
Previous psychiatric treatment	Nothing	180	54	87	48.30	93	51.70	3.645	0.456
	Psychotherapy	107	32	49	45.80	58	54.20		
	Drugs	28	8	15	53.60	13	46.40		
	Combined	15	5	8	53.30	7	46.70		
	Other	6	2	5	83.30	1	16.70		

Table 2: summary of clinical characteristics of LIF participants (N=338)

		N	Mean	SD	95% CI		Statistics		
					L	U	F	Sig.	
PQ total	M	162	29.9	17.74	27.14	32.65	8.481	0.004*	
	F	169	35.49	17.21	32.88	38.11			
	Total	331	32.75	17.67	30.84	34.66			
PQ positive	M	164	13.08	9.08	11.68	14.48	5.275	0.022*	
	F	170	15.33	8.82	13.99	16.67			
	Total	334	14.22	9.00	13.25	15.19			
PQ negative	M	162	7.14	4.77	6.40	7.88	8.651	0.004*	
	F	169	8.65	4.56	7.96	9.34			
	Total	331	7.91	4.72	7.40	8.42			
PQ general	M	163	5.75	3.62	5.19	6.32	14.242	0.000*	
	F	169	7.27	3.67	6.71	7.82			
	Total	332	6.52	3.72	6.12	6.93			
PQ disorganized	M	164	3.87	2.76	3.44	4.29	1.771	0.184	
	F	169	4.3	3.11	3.82	4.77			
	Total	333	4.08	2.95	3.77	4.4			
GFRS	M	163	6.17	1.53	5.93	6.41	12.807	0.000*	
	F	168	6.77	1.49	6.54	7.00			
	Total	331	6.47	1.54	6.31	6.64			
GFSS	M	163	6.46	1.33	6.25	6.67	11.316	0.001*	
	F	168	6.92	1.16	6.75	7.10			
	Total	331	6.69	1.27	6.56	6.83			
GAF	M	155	62.58	12.40	60.61	64.55	7.422	0.007*	
	F	166	66.17	11.22	64.45	67.90			
	Total	321	64.44	11.92	63.13	65.75			
IPSM: total scores	M	154	87.02	20.48	83.76	90.28	13.727	0.000*	
	F	164	94.82	17.00	92.2	97.45			
	Total	318	91.04	19.14	88.93	93.16			
IPSM: need of approval	M	154	22.61	4.85	21.84	23.38	2.951	0.087	
	F	164	23.46	3.98	22.85	24.08			
	Total	318	23.05	4.43	22.56	23.54			
IPSM: interpersonal awareness	M	154	17.61	5.21	16.78	18.44	15.085	0.000*	
	F	164	19.82	4.91	19.06	20.58			
	Total	318	18.75	5.17	18.18	19.32			
IPSM: separation anxiety	M	154	19.05	5.83	18.12	19.98	22.80	0.000*	
	F	164	22.05	5.37	21.23	22.88			
	Total	318	20.6	5.79	19.96	21.24			
IPSM: timidity	M	154	18.25	5.20	17.43	19.08	0.885	0.347	
	F	164	18.79	5.02	18.02	19.57			
	Total	318	18.53	5.10	17.97	19.1			
IPSM: fragile inner self	M	154	9.86	3.82	9.25	10.47	1.538	0.216	
	F	164	10.41	4.08	9.78	11.04			
	Total	318	10.14	3.96	9.70	10.58			
		Total		Male		Female		Statistics	
		N	%	N	%	N	%	χ^2	P
PQ $\geq$ 18	no	238	71	120	50.40	118	49.60	0.576	0.448

	yes	96	29	44	45.80	52	54.20		
GFRS reduction	same	203	61	99	48.80	104	51.20	0.65	0.722
	reduction	119	36	61	51.30	58	48.70		
	increment	8	3	3	37.50	5	62.50		
GFSS reduction	same	206	62	101	49.00	105	51.00	0.336	0.846
	reduction	122	37	61	50.00	61	50.00		
	increment	3	1	1	33.30	2	66.70		

Table 3: summary of psychopathological characteristics of LIF participants(N=338). PQ: Prodromal Questionnaire; GFRS: Global Functioning Role Scale; GFSS: Global Functioning Role Scale; GAF: Global assessment of functioning; IPSM: Interpersonal Sensitivity Measure.

		Tot		M		F		Statistics	
		N	%	N	%	N	%	χ^2	P
Study:	not completed	157	49	75	47.80	82	52.20	0.526	0.468
Study:	completed	166	52	86	51.80	80	48.20		
UHR	yes	64	38.60	33	51.60	31	48.40	0.002	0.96
BS	present	92	58.20	49	47.60	54	52.40	1.742	0.187
COGDIS	present	60	38.00	34	49.30	35	50.70	0.275	0.600
COPER	present	92	58.20	49	47.60	54	52.40	1.742	0.187

Table 4: summary of psychotic prodromal symptoms of full-assessed LIF participants (N=166). UHR: Ultra High Risk; COGDIS: Cognitive Symptoms; COPER: Cognitive Perceptive Symptoms; BS: Basic Symptoms.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Masillo, A;Brandizzi, M;Nelson, B;Lo Cascio, N;Saba, R;Lindau, JF;Telesforo, L;Montanaro, D;D'Alema, M;Girardi, P;McGorry, P;Nastro, PF

Title:

Youth mental health services in Italy: An achievable dream?

Date:

2018-06-01

Citation:

Masillo, A., Brandizzi, M., Nelson, B., Lo Cascio, N., Saba, R., Lindau, J. F., Telesforo, L., Montanaro, D., D'Alema, M., Girardi, P., McGorry, P. & Nastro, P. F. (2018). Youth mental health services in Italy: An achievable dream?. *EARLY INTERVENTION IN PSYCHIATRY*, 12 (3), pp.433-443. <https://doi.org/10.1111/eip.12328>.

Persistent Link:

<http://hdl.handle.net/11343/291156>