Received Date : 11-Mar-2016

Accepted Date : 12-Mar-2016

Article type : Correspondence

Supporting Information : stephen.fox@petermac.org

Dear Sir/Madam

We thank Dr Van Bockstal and colleagues for their letter, titled 'The Baader-Meinhof phenomenon in ductal carcinoma *in situ* of the breast' in response to our review 'Ductal carcinoma *in situ*- update on risk assessment and management'¹. Van Bockstal *et al* raise the issue of the often ignored ductal carcinoma *in situ* (DCIS) microenvironment, in particular myxoid-appearing stroma, and its role in modulating disease outcome in DCIS.

As Van Bockstal *et al* acknowledge, detailed appraisal of the prognostic significance of the DCIS microenvironment was beyond the scope of our review. While we did not specifically draw attention to the changes in the DCIS stroma seen in Figure 1 of our review, we acknowledge the importance of the DCIS microenvironment in determining disease outcome. This has been demonstrated by the work of Van Bockstal *et al*², which we cited in our review, and others investigating the role of myoepithelial cells^{3,4}, stromal proteins^{5,6}, and the immune response in DCIS⁷. Indeed, we have published extensively on the role of the tumour microenvironment in DCIS, including angiogenic patterns surrounding DCIS^{8,9} and the significance of the immune infiltrate to prognosis⁷.

The microenvironment is clearly a major factor in DCIS biology which in the past has been overshadowed by an emphasis on the tumour epithelium. As Van Bockstal *et al* state, it is hoped that increased awareness of and research into the non-epithelial components of a DCIS lesion will lead to the development of improved DCIS biomarkers to optimise patient management.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/HIS.12968](https://doi.org/10.1111/HIS.12968)

This article is protected by copyright. All rights reserved

Yours faithfully

Jia-Min B. Pang

Kylie L. Gorringer

Stephen B. Fox

References

1. Pang JM, Gorringer KL, Fox SB. Ductal carcinoma in situ - update on risk assessment and management. *Histopathology* 2016;68:96-109.
2. Van Bockstal M, Lambein K, Gevaert O, et al. Stromal architecture and periductal decorin are potential prognostic markers for ipsilateral locoregional recurrence in ductal carcinoma in situ of the breast. *Histopathology* 2013;63:520-33.
3. Russell TD, Jindal S, Agunbiade S, et al. Myoepithelial Cell Differentiation Markers in Ductal Carcinoma in Situ Progression. *The American journal of pathology* 2015;185:3076-89.
4. Allen MD, Thomas GJ, Clark S, et al. Altered microenvironment promotes progression of preinvasive breast cancer: myoepithelial expression of α v β 6 integrin in DCIS identifies high-risk patients and predicts recurrence. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2014;20:344-57.
5. Witkiewicz AK, Dasgupta A, Nguyen KH, et al. Stromal caveolin-1 levels predict early DCIS progression to invasive breast cancer. *Cancer biology & therapy* 2009;8:1071-9.
6. Witkiewicz AK, Freydin B, Chervoneva I, et al. Stromal CD10 and SPARC expression in ductal carcinoma in situ (DCIS) patients predicts disease recurrence. *Cancer biology & therapy* 2010;10:391-6.
7. Bates GJ, Fox SB, Han C, et al. Quantification of regulatory T cells enables the identification of high-risk breast cancer patients and those at risk of late relapse. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2006;24:5373-80.
8. Engels K, Fox SB, Whitehouse RM, Gatter KC, Harris AL. Up-regulation of thymidine phosphorylase expression is associated with a discrete pattern of angiogenesis in ductal carcinomas in situ of the breast. *Journal of Pathology* 1997;182:414-20.

9. Engels K, Fox SB, Whitehouse RM, Gatter KC, Harris AL. Distinct angiogenic patterns are associated with high-grade in situ ductal carcinomas of the breast. *Journal of Pathology* 1997;181:207-12.

Author Manuscript

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Pang, J-MB;Gorringe, KL;Fox, SB

Title:

Reply to the Baader-Meinhof phenomenon in ductal carcinoma in situ of the breast

Date:

2016-09-01

Citation:

Pang, J. -M. B., Gorringe, K. L. & Fox, S. B. (2016). Reply to the Baader-Meinhof phenomenon in ductal carcinoma in situ of the breast. HISTOPATHOLOGY, 69 (3), pp.523-524. <https://doi.org/10.1111/his.12968>.

Persistent Link:

<http://hdl.handle.net/11343/291281>