Received Date : 09-Oct-2015

Revised Date : 23-Feb-2016

Accepted Date : 11-Apr-2016

Article type : Original Article: Airway Diseases

Editor: Douglas Robinson

Residential greenness is differentially associated with childhood allergic rhinitis and aeroallergen sensitization in seven birth cohorts

Elaine Fuertes^a, Iana Markevych^{a,b}, Gayan Bowatte^c, Olena Gruzieva^d, Ulrike Gehring^e, Allan Becker^f, Dietrich Berdel^g, Andrea von Berg^g, Anna Bergström^d, Michael Brauer^{h,i}, Bert Brunekreef^e, Irene Brüske^a, Chris Carlsten^{h,i}, Moira Chan-Yeungⁱ, Shyamali C Dharmage^c, Barbara Hoffmann^{j,k}, Claudia Klümper^j, Gerard H. Koppelman^{l,m}, Anita Kozyrskyj^{n,o}, Michal Korek^d, Inger Kull^{d,p,q}, Caroline Lodge^c, Adrian Lowe^c, Elaina MacIntyre^r, Göran Pershagen^d, Marie Standl^a, Dorothea Sugiri^j, Alet Wijga^s, MACS, Joachim Heinrich^{a,t}

a) Institute of Epidemiology I, Helmholtz Zentrum München - German Research Center for Environmental Health, Neuherberg, Germany

b) Division of Metabolic and Nutritional Medicine, Dr. von Hauner Children's Hospital, Ludwig-Maximilians-University of Munich, Munich, Germany

c) Allergy and Lung Health Unit, Melbourne School of Population and Global Health, The University of Melbourne, Australia

d) Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

e) Institute for Risk Assessment Sciences, Utrecht University, Utrecht, The Netherlands

f) Department of Pediatrics and Child Health, University of Manitoba, Winnipeg, Canada

g) Research Institute, Department of Pediatrics, Marien-Hospital Wesel, Germany

h) School of Population and Public Health, University of British Columbia, Vancouver, Canada

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/all.12915](#)

This article is protected by copyright. All rights reserved

- i) Department of Medicine, University of British Columbia, Vancouver, Canada
- j) IUF – Leibniz Research Institute for Environmental Medicine, Heinrich-Heine University of Düsseldorf, Germany
- k) Medical Faculty, Deanery of Medicine, Heinrich-Heine University of Düsseldorf, Germany
- l) University of Groningen, University Medical Center Groningen, Groningen, Department of Pediatric Pulmonology and Pediatric Allergology, Beatrix Children's Hospital, The Netherlands
- m) Groningen Research Institute for Asthma and COPD, University of Groningen, Groningen, The Netherlands
- n) Department of Pediatrics, Faculty of Medicine & Dentistry, Women and Children's Health Research Institute, Edmonton, Canada
- o) School of Public Health, University of Alberta, Edmonton, Canada
- p) Department of Clinical Science and Education, Karolinska Institutet, Stockholm, Sweden
- q) Sachs' Children and Youth Hospital, Stockholm, Sweden
- r) Department of Environmental Health, Public Health Ontario, Toronto, Canada
- s) Center for Nutrition, Prevention and Health Services, National Institute of Public Health and the Environment, Bilthoven, The Netherlands
- t) Institute and Outpatient Clinic for Occupational, Social and Environmental Medicine, University Hospital Munich, Ludwig-Maximilians-University of Munich, Munich, Germany

Corresponding author:

Elaine Fuertes, PhD

Institute of Epidemiology I, Helmholtz Zentrum München - German Research Center for Environmental Health, Ingolstädter Landstraße 1, 85764 Neuherberg, Germany

Phone: +49 89 3187 4235; Fax: +49 89 3187 3380

elaine.fuertes@helmholtz-muenchen.de

Short title: Residential greenness and allergic rhinitis

Key Words: allergic rhinitis; birth cohorts; greenness; NDVI; sensitization

Abstract word count: 248

Text word count (introduction to discussion): 3634

59 FUNDING

60 The BAMSE study was supported by the Swedish Research Council, the Swedish Heart–Lung
61 Foundation, Stiftelsen Frimurare Barnhuset i Stockholm, Matsumura’s donation, the Stockholm County
62 Council, the Swedish Environmental Protection Agency, the Swedish Society for Medical Research, the
63 Swedish Foundation for Strategic Research and the Swedish Research Council for Health Working Life
64 and Welfare. The PIAMA study is supported by The Netherlands Organization for Health Research and
65 Development, The Netherlands Organization for Scientific Research, The Netherlands Asthma Fund,
66 The Netherlands Ministry of Spatial Planning, Housing, and the Environment, and The Netherlands
67 Ministry of Health, Welfare and Sport. The GINIplus study was mainly supported for the first 3 years
68 by the Federal Ministry for Education, Science, Research and Technology (interventional arm) and
69 Helmholtz Zentrum Munich (former GSF) (observational arm). The 4 year, 6 year and 10 year follow-
70 up examinations of the GINIplus study were covered from the respective budgets of the 5 study centres
71 (Helmholtz Zentrum Munich (former GSF), Marien-Hospital Wesel, LMU Munich, TU Munich and
72 from 6 years onward also from IUF - Leibniz Research-Institute for Environmental Medicine) and a
73 grant from the Federal Ministry for Environment (IUF, FKZ 20462296). The LISApplus study was
74 mainly supported by grants from the Federal Ministry for Education, Science, Research and
75 Technology and in addition from Helmholtz Zentrum Munich (former GSF), Helmholtz Centre for
76 Environmental Research - UFZ, Leipzig, Marien-Hospital Wesel, Pediatric Practice, Bad Honnef for
77 the first 2 years. The 4 year, 6 year and 10 year follow-up examinations of the LISApplus study were
78 covered from the respective budgets of the involved partners (Helmholtz Zentrum Munich (former
79 GSF), Helmholtz Centre for Environmental Research - UFZ, Leipzig, Marien-Hospital Wesel, Pediatric
80 Practice, Bad Honnef, IUF – Leibniz-Research Institute for Environmental Medicine) and in addition
81 by a grant from the Federal Ministry for Environment (IUF, FKZ 20462296). The CAPPS study was
82 supported by the Canadian Institutes of Health Research, the British Columbia Lung Association and
83 the Manitoba Medical Service Foundation. The SAGE study was supported by the Canadian Institutes
84 of Health Research. The first 6 years of the MACS study were funded by Nestec Ltd, a subsidiary of
85 Nestlé Australia. The 12 year follow-up was funded by a project grant from the Asthma Foundation of
86 Victoria. The NHMRC funded Centre for Air Quality and Health Research and evaluation (CAR)
87 funded geocoding of participants’ addresses. The “Traffic Asthma and Genetics” collaboration was
88 supported by the AllerGen Networks of Centres of Excellence. The ESCAPE (grant agreement number:
89 211250) research received funding from the European Community’s Seventh Framework Program

90 (FP7/2007-2011). The aforementioned funding sources had no involvement in the study design, in the
91 collection, analysis and interpretation of data, in the writing of the report and in the decision to submit
92 the article for publication.

Author Manuscript

ABSTRACT

Background: The prevalence of allergic rhinitis is high but the role of environmental factors remains unclear. We examined cohort-specific and combined associations of residential greenness with allergic rhinitis and aeroallergen sensitization based on individual data from Swedish (BAMSE), Australian (MACS), Dutch (PIAMA), Canadian (CAPPS and SAGE) and German (GINIplus and LISApplus) birth cohorts (N=13,016).

Methods: Allergic rhinitis (doctor diagnosis/symptoms) and aeroallergen sensitization were assessed in children aged 6-8 years in six cohorts and 10-12 years in five cohorts. Residential greenness was defined as the mean Normalized Difference Vegetation Index (NDVI) in a 500m buffer around the home address at the time of health assessment. Cohort-specific associations per 0.2 unit increase in NDVI were assessed using logistic regression models and combined in a random-effects meta-analysis.

Results: Greenness in a 500m buffer was positively associated with allergic rhinitis at 6-8 years in BAMSE (odds ratio=1.42, 95% confidence interval [1.13, 1.79]) and GINI/LISA South (1.69 [1.19, 2.41]) but inversely associated in GINI/LISA North (0.61 [0.36, 1.01]) and PIAMA (0.67 [0.47, 0.95]). Effect estimates in CAPPS and SAGE were also conflicting but not significant (0.63 [0.32, 1.24] and 1.31 [0.81, 2.12], respectively). All meta-analyses were non-significant. Results were similar for aeroallergen sensitization at 6-8 years and both outcomes at 10-12 years. Stratification by NO₂ concentrations, population density, an urban versus rural marker and moving did not reveal consistent trends within subgroups.

Conclusion: Although residential greenness appears to be associated with childhood allergic rhinitis and aeroallergen sensitization, the effect direction varies by location.

INTRODUCTION

Green environments are thought to impart beneficial effects on health by increasing physical activity and stress relief, and by facilitating social interactions. They are also associated with reduced noise, air pollution and heat exposures (1). However, surrounding greenness may play a more complex role on allergic health outcomes. Although a causal relationship remains to be established, studies suggest that children who spend more time in outdoor green environments during early-life may benefit from exposure to a greater number and diversity of beneficial microbes (2,3). A similar protective effect has also been documented between sensitization and a diverse early-life exposure to indoor allergens and microbes (4). However, among those sensitized, exposure to pollen-releasing plants and outdoor fungi may exacerbate allergic symptoms in later childhood (5).

The few epidemiological studies that have examined associations between residing in/near green places and allergic health outcomes have yielded inconsistent results. Studies report increased (6), no (7), protective (2,8), or conflicting (9) effects, and a recent study concluded that associations appear to depend on the type of greenness evaluated (for example, parks versus forests (10)). These studies differ with respect to their designs, outcomes, populations and green exposure assessment strategies, which may in part explain some of these discrepant findings. For example, the aforementioned studies defined vegetation level using data on tree canopy cover (6), vegetation or land-use types (2,8), the Normalized Difference Vegetation Index (NDVI) (7,9) or several of these measures (10). It is currently unclear which of these exposure metrics may be best. While some more specific measures are able to classify large green areas into land use types (such as the CORINE land use European data), they are not commonly available on a global scale and do not include small green areas. Further, it is possible that different metrics may be more or less relevant to specific pathways. For example, land use data may be very useful for studying physical activity levels, but this is unlikely to represent the main pathway by which greenness might affect allergic diseases.

As a general measure of vegetation presence, the NDVI index captures vegetation of all sizes using a globally harmonized method, and we chose to use this index to examine cross-sectional associations between residential greenness and allergic rhinitis and aeroallergen sensitization during childhood and early adolescence in seven birth cohorts from Australia, Canada, Germany, the Netherlands and Sweden. As suggestive evidence exists that air pollutants and urbanization may act as confounders or

145 effect modifiers in greenness-health relationships (11,12), we tested interactions between nitrogen
146 dioxide (NO₂) concentrations, population density and a rural/urban indicator with residential
147 greenness, and also adjusted for these factors.

METHODS

Data sources

Seven birth cohorts participated: BAMSE (13), CAPPS (14), GINIplus (15), LISApplus (16,17), MACS (18), PIAMA (19) and SAGE (20). Data on several health outcomes, environmental exposures and covariates from all cohorts except MACS had already been harmonized as part of the Traffic, Asthma and Genetics (21) and European Study of Cohorts for Air Pollution Effects (22) collaborations. MACS is here included as this Australian birth cohort adds additional vegetation and geography heterogeneity. Each cohort received ethical approval from their local authorized Institutional Review Boards.

Outcome assessment

We focused on health outcomes during childhood (6-8 years) and early adolescence (10-12 years). Information on the cohort-specific study designs and outcome definitions, which varied slightly by cohort, are provided in the Supplemental Information, Table S1. Allergic rhinitis was defined based on a diagnosis during a physician assessment at a follow-up visit in CAPPS and SAGE, parental report of a doctor's diagnosis in GINIplus and LISApplus, parental symptom report in PIAMA and BAMSE and parental symptom or treatment report in MACS.

Sensitization was assessed by skin prick testing for CAPPS, MACS and SAGE, with a positive reaction defined as having a wheal diameter of ≥ 3 mm. For all other cohorts, sensitization was assessed by measuring allergen specific IgE levels, with a positive reaction defined as any value ≥ 0.35 kU/L, the lower detection limit of the assay. Birch, *Dactylis*, mugwort, ragweed, rye, timothy grass, trees and weeds were considered as outdoor aeroallergens. *Alternaria alternata*, cats, *Cladosporium herbarum*, cockroaches, dogs, feathers, house dust mites and molds were considered as indoor aeroallergens. All available aeroallergens were included in the sensitization analyses. Not all cohorts had information on all aeroallergens or health data at both time points (Supplemental Information, Table S1).

Greenness assessment

The NDVI, a green biomass density indicator, was used as a surrogate for surrounding greenness. Its calculation is based on the difference of surface reflectance in visible (0.4–0.7 μ m) and near-infrared (0.7–1.1 μ m) wavelengths. Values range from negative one (water) through zero (rock, sand, snow) to positive one (dense green vegetation) (23). The assignment of NDVI to the home addresses of all

cohort participants was done using a harmonized method previously described (24). First, to achieve maximum exposure contrasts, cloud-free satellite images corresponding as close as possible to the spring and summer months during the year of birth of the participants were centrally selected for all cohorts and used to calculate NDVI maps. Negative NDVI pixels were set to zero (replication of analyses with negative NDVI values left as is or set to missing yielded the same results). Second, these images were used to calculate mean greenness in 500m and 1000m circular buffers around the home addresses of participants at 6-8 and 10-12 years of age in order to assess current greenness exposure effects. The 500m buffer was *a priori* selected as the main buffer as it is a proximal measure of a child's neighborhood, may be less prone to exposure misclassification and has been used in previous studies on children (e.g. (25,26)). The 1000m buffer captures a larger area around an individual's neighborhood and was used as a sensitivity analysis.

The NDVI values used in all main analyses were derived from satellite maps taken at the time of birth of the participants and assigned to their 6-8 and 10-12 year addresses under the assumption that the spatial distribution of greenness would remain stable between these time points. To test this assumption, a second set of NDVI values was created based on satellite maps selected approximately a decade after the birth of the participants and assigned to these same 6-8 and 10-12 year addresses. All main analyses were replicated with this second set of NDVI values. Details of the months and years used for the NDVI assignments for each cohort are provided in the Supplemental Information, Table S1.

Statistical analysis

Cohort-specific associations were analyzed using logistic regression. Odds ratios are reported per 0.2 unit increase in NDVI (approximately two times the standard deviation in the total population) with corresponding 95% confidence intervals. The GINIplus and LISApplus cohorts were pooled as the study designs are nearly identical and associations are presented per geographical area instead (the rural GINI/LISA North area and GINI/LISA South, which covers the urban city of Munich and its surroundings). Random-effects meta-analysis was used to calculate combined estimates to allow for potential within-and between-cohort heterogeneity (27). The I^2 statistic was used to examine statistical heterogeneity among cohort-specific effect estimates and can be interpreted as the percentage of the variability in effect sizes attributable to the between-study variability rather than sampling error (28). I^2

values between 50-90% and 75%-100% represent substantial and considerable heterogeneity, respectively (29). Cochran's Q test was used to test for significant heterogeneity. Analyses for CAPPS, GINI/LISA North, GINI/LISA South, PIAMA, SAGE and the combined meta-analyses (using package "meta" (30)) were conducted centrally using the statistical program R, version 3.1.1 (31). Analyses for BAMSE and MACS were done locally using STATA, version 13 and 13.1 (32), respectively, following the same analysis plan.

Minimally adjusted models were adjusted for sex and age. Main models were additionally adjusted for parental atopy (not included for MACS as 97% of participants had a history of parental atopy), older siblings, maternal smoking during pregnancy, secondhand smoke exposure concomitant with the time of health outcome assessment (not available for MACS), socioeconomic status (defined as the highest education attained by either parent for BASME, GINI/LISA North, GINI/LISA South, MACS and PIAMA, and maternal age at birth for CAPPS and SAGE), group (intervention for CAPPS, GINI/LISA North, GINI/LISA South, PIAMA and MACS), region (CAPPS and PIAMA only) and cohort (GINI/LISA North and GINI/LISA South only). The influence of additional adjustments for birth weight and exposure to furry pets and mold/dampness in the home at the time of health outcome assessment was examined in sensitivity analyses (MACS not included as these data were generally not available). Covariates were defined as similarly as possible across cohorts using questionnaire-derived information and their selection is based on previous combined analyses of these cohorts with regard to allergic rhinitis and sensitization (9,22,33).

Effect modification

To assess effect modification by sex, regression analyses were run including an interaction term between NDVI and sex. In a separate analysis, regression analyses were also run separately for males and females. Effect modification by cohort-specific tertiles of NO₂ concentrations and population density in a 1000m buffer around the home address was also assessed, and models were run stratified by whether participants lived in urban or rural surroundings (data sources and methodology described in the Supplemental Information, page 3). Models were also stratified by whether a child had moved between 1) birth and 6-8 years when considering the childhood health outcomes and between 2) birth and 10-12 years when considering the adolescent health outcomes (CAPPS and SAGE not included as data on moving behavior were unavailable).

RESULTS

Study population

In total, 13016 children had available information on NDVI exposure and at least one outcome of interest at one time point. The included cohorts varied in size from 3339 children in PIAMA to 327 children in MACS (Table 1). Of those with available data, 9.8% (1182/12007) had allergic rhinitis and 30.3% (2246/7408) were sensitized to at least one aeroallergen at the age of 6-8 years (13.6% (1346/9885) and 42.1% (1650/3922) are the respective values for 10-12 years). Allergic rhinitis prevalence was lowest in GINI/LISA North and highest among cohorts recruited on the basis of family history (MACS and CAPPS) and SAGE.

Distribution of NDVI values

The mean and range of NDVI values in a 500m buffer were similar across cohorts (Figure 1). NDVI estimates in a 500m buffer were highly correlated with those in a 1000m buffer (Pearson's $r > 0.88$). NDVI estimates in the 500m buffer assessed to the childhood and early adolescence addresses were weak to moderately correlated across cohorts for those who moved between these two time points (range of $r = 0.26$ in PIAMA to $r = 0.55$ in BAMSE). NDVI estimates derived using satellite maps obtained for the year of birth and approximately 10 years later ($r > 0.73$) were highly correlated. As it was not possible to obtain cloud-free images for the same months for all cohorts and given that months have different meanings in the different cohorts (for example, when contrasting European and Australian seasons), comparing NDVI distributions across cohorts is not appropriate. Cohort locations and the distribution of NDVI values per cohort are depicted in the Supplemental Information, Figure S1.

Associations between health outcomes and NDVI

The adjusted cohort-specific associations per 0.2 increase in NDVI for the main models are presented in Figures 2 and 3 for outcomes assessed during childhood (6-8 years) and early adolescence (10-12 years), respectively (results per cohort-specific interquartile range increase in NDVI presented in the Supplemental Information, Figure S2). The minimally adjusted models (for age and sex only) were similar (not shown). Greenness in a 500m buffer was positively associated with allergic rhinitis at 6-8 years in BAMSE (1.42 [1.13, 1.79]) and GINI/LISA South (1.69, [1.19, 2.41]) but inversely associated

in GINI/LISA North (0.61 [0.36, 1.01]) and PIAMA (0.67 [0.47, 0.95]). The effect estimates in the Canadian cohorts were also conflicting but not significant (0.63 [0.32, 1.24] and 1.31 [0.81, 2.12] for CAPPS and SAGE, respectively). The pattern of associations within each cohort for aeroallergen sensitization was similar to those with allergic rhinitis. The pattern also did not differ when associations were stratified into categories of indoor and outdoor allergens, with the exception of SAGE for which the direction of effect estimates varied across outcomes. This suggests that the observed associations with aeroallergen sensitization are not attributable to a single allergen.

Similar results were obtained for both health outcomes at 10-12 years for the four cohorts with available data at both time points. Associations in the seventh cohort MACS, for which health data were only available at this latter age, were non-significant for allergic rhinitis (0.96 [0.59, 1.57]) and inverse for aeroallergen sensitization (0.57 [0.34, 0.96]).

Effect estimates were consistent when NDVI was assessed in a 1000m buffer and when models were further adjusted for birth weight and exposure to furry pets and mold/dampness at the time of health outcome assessment (not shown). There was no good indication of non-linearity between NDVI exposures and the health outcomes when associations were examined using generalized additive models, suggesting that at least for these outcomes, a threshold value for NDVI was not apparent.

Given the substantial/considerable heterogeneity between the cohort-specific associations ($I^2 > 0.7$ for seven of the eight adjusted associations), all meta-analytic results were non-significant (Supplemental Information, Table S2).

Effect modification

Although at least one interaction term between NDVI in a 500m buffer and each potential effect modifier considered was significant for at least one cohort, results were not consistent across cohorts and all interaction terms in the combined analyses were non-significant (Supplemental Information, Table S3). In line with this, associations stratified by sex (Supplemental Information, Figure S3) as well as NO₂ (Supplemental Information, Figure S4) and population density (Supplemental Information, Figure S5) tertiles did not reveal consistent patterns within or between cohorts. Stratification by whether participants' lived in urban or rural surroundings yielded weak evidence for stronger positive

effects in urban settings in the cohorts for which greenness was positively associated with the health outcomes (BAMSE and GINI/LISA South; Supplemental Information, Figure S6), but confidence intervals overlapped. Independently adjusting the main models for NO₂, population density and urban versus rural categorical variables did not change the results, although the effect estimates for BAMSE were attenuated after adjustment for population density and urban versus rural surroundings (for example, 1.18 [0.81, 1.72] and 1.10 [0.78, 1.54], respectively, compared to 1.42 [1.13, 1.79], for the association between childhood allergic rhinitis and NDVI in a 500m buffer). Finally, models stratified by moving behaviour did not yield consistent differences between groups (Supplemental Information, Figure S7).

DISCUSSION

Mean NDVI in a 500m buffer was differentially associated with allergic rhinitis and aeroallergen sensitization in this analysis of seven birth cohorts, resulting in an overall non-significant combined finding. Evaluating sex, NO₂ exposure, population density and an urban/rural marker as effect modifiers did not clarify these trends. Confounding by an unknown factor that varies between-study areas or by several region-specific confounders may be a possible explanation. Alternatively, our results may be simply driven by chance.

It may be worth asking whether a combined meta-analysis is appropriate in this study, given the considerable/substantial heterogeneity observed in the cohort-specific results. We chose to present the meta-analytic results as they answer our original research question. However, the most important lesson from this study may not lie in the direction of the effect estimates but rather upon the use of the NDVI in allergic health research. Although the NDVI is able to capture small-scale greenness in a standardized and objective manner, it does not allow particular types of vegetation to be distinguished, nor are we able to derive individual-level measures of exposure to pollen or other allergenic tree species. The duration and character of potential exposures can also not be assessed. For example, the extent to which NDVI serves as a proxy for exposure to pollen or microbial diversity, or an indicator of areas conducive to physical activity or social interactions, or a proxy for visual impacts related to stress reduction is unclear. We are thus not able to identify which, if any, particular vegetation types, exposure pathway(s) or duration of exposures may drive the observed associations. Consequently, although the use of the NDVI to assess vegetation may be well justified for the evaluation of potential pathways related to stress and for certain health outcomes (for example, birth weight, physical activity and mental health), it appears to be too general of a measure to completely capture the full structure and potential role of the green environment with respect to allergic diseases. We thus caution against its further use in the allergic field and rather recommend that future studies use more detailed data on local tree and herbaceous species and on interactions between people and various measures of vegetation when exploring the role of the residential green environment and the overall living environment on allergic health outcomes. Such measures naturally are more focused on pathways related to pollen dispersion and microbial diversity.

The current study nevertheless has several strengths. It is the largest analysis of residential greenness

on childhood allergic health outcomes to date and the first to include individual-level data from more than one continent. The majority of the health and covariate data had been previously harmonized for these cohorts (21,22), although the allergic rhinitis definitions differed slightly as did the number of objectively measured aeroallergens tested. Also, two cohorts were high allergy-risk by design (MACS and SAGE). These factors could have affected the cohort-specific outcome prevalences, but not necessarily the associations. The high outcome prevalences for some of the cohorts may also have resulted in odd ratios that overestimate the true relative risks, although the overall conclusions of this study would not be affected (34). Several covariates were adjusted for in this analysis, but residual confounding is always possible in observational studies. For example, although models were adjusted for a marker of individual-level socioeconomic status and consistent evidence of effect modification by this factor was not detected (not shown), our measures of individual-level socioeconomic status may not be optimal. It is also possible that area-level factors may play a role.

Data were prospectively collected for all cohorts except SAGE. Thus, we anticipate that recall bias should be minimal, but remains possible, as does selection bias due to loss of follow-up. Given the cross-sectional design of the analyses, bias related to moving or the effect of timing of exposures (current versus early) was not directly assessed. Findings from a previous study indicate that the green environment around the home at birth may be more strongly associated with allergies later in life than the current home green environment for children that have moved (8). In our study, models stratified by whether a child had moved between birth and the time of outcome assessment did not yield consistent trends. Further, it is unlikely that any bias related to the length of residence at the current address would differentially affect the results across cohorts.

The harmonized greenness assignment across studies is also an important strength of this study, but is not without limitations. First, it was not possible to obtain cloud-free images for the same months and years for all cohorts. NDVI estimates were derived from images as close in time as possible during spring and summer months to achieve maximum exposure contrasts between areas of low and high greenness. Second, we related NDVI values derived from maps taken at the time of birth to health outcomes 6-12 years later assuming that the spatial variability in greenness exposures would not have changed during this time, an approach often used in air pollution research (22). This assumption is supported by the fact that a second set of NDVI values derived from satellite images taken ten years

after the birth of the participants were highly correlated with the main NDVI estimates and yielded no differences in the results. This finding suggests that the spatial distribution of residential greenness was temporally stable during the time frame covered in this study (early/mid 1990s to middle/late 2000s) in the areas investigated. Further studies are needed to confirm whether this finding is also valid in other parts of the world, particularly in developing countries where land use patterns might change more rapidly. Third, our decision to assess associations with greenness in 500m and 1000m buffers around the home address did not allow the study of the effect of greenness on a very small (in a 100m buffer) or large scale (for example, 3000m buffer or even at the city-level). The 500m buffer around the home address was *a priori* selected as the main buffer of interest as it is a proximal measure of a child's neighbourhood and is likely to incorporate less exposure misclassification than larger buffers, although it is well-known that pollen can travel much larger distances (35). The optimal buffer size to use when studying similar associations remains to be determined. Fourth, we chose to limit our analysis to vegetation levels around the home address and did not assess associations with types of green space or land use classifications (e.g. presence or percentage of parks, forest and agriculture) as standard data of this type (e.g. the CORINE data) were only available for the European cohorts and, like the NDVI, do not provide information on vegetation types.

Although the evidence supporting a beneficial effect of greenness on several health measures is increasing, studies on allergic health outcomes remain inconsistent. In this harmonized analysis of seven birth cohorts from three continents, the direction of the association between mean NDVI in a 500m buffer and allergic rhinitis and aeroallergen sensitization varied by region, resulting in a non-significant combined finding. Our results thus suggest that using the NDVI as a marker for residential greenness may only have local interpretations. Alternatively, it is possible that there is no real association between residential greenness and allergic health, and that the observed effects are driven by chance or unknown confounding (region-specific) factors.

399 **ACKNOWLEDGEMENTS**

400 We thank all children and parents for their cooperation, and all technical and administrative support
401 staff and medical and field work teams. We also thank all BAMSE, CAPPS, GINIplus, LISApplus,
402 MACS, PIAMA and SAGE investigators.

403

404 **CONFLICT OF INTEREST STATEMENT**

405 All co-authors have no conflicts of interest.

406

407 **AUTHOR CONTRIBUTIONS**

408 EF, IM and JH designed the study. EF wrote the initial draft and had final responsibility for the decision
409 to submit for publication. EF, GB and OG conducted the statistical analyses. IM, GB, MK, UG, DS,
410 MB and CC contributed to the greenness exposure assignment. ABecker, DB, AvB, ABergström, BB,
411 IB, MC-Y, SCD, UG, BH, CK, GHK, AK, IK, CL, AL, EM, GP, MS and AW contributed to the
412 collection and/or provided the health and covariate data. All authors provided substantial contributions
413 to the conception or design of the work, or the acquisition, analysis, or interpretation of data for the
414 work, revised the manuscript for important intellectual content, approved the final version and agreed
415 to be accountable for all aspects of the work.

1. Hartig T, Mitchell R, de Vries S, Frumkin H. Nature and Health. *Annu Rev Public Health*. 2014;**35**(1):21.1–21.22. DOI:10.1146/annurev-publhealth-032013-182443
2. Hanski I, Herten L von, Fyhrquist N, Koskinen K, Torppa K, Laatikainen T, et al. Environmental biodiversity, human microbiota, and allergy are interrelated. *Proc Natl Acad Sci*. 2012;**109**(21):8334–9. DOI:10.1073/pnas.1205624109
3. Von Hertzen L, Hanski I, Haahtela T. Natural immunity: biodiversity loss and inflammatory diseases are two global megatrends that might be related. *EMBO Rep*. 2011;**12**(11):1089–93. DOI:10.1038/embor.2011.195
4. Lynch SV, Wood RA, Boushey H, Bacharier LB, Bloomberg GR, Kattan M, et al. Effects of early-life exposure to allergens and bacteria on recurrent wheeze and atopy in urban children. *J Allergy Clin Immunol*. 2014 Sep 1;**134**(3):593–601.e12.
5. Cakmak S, Dales RE, Burnett RT, Judek S, Coates F, Brook J. Effect of airborne allergens on emergency visits by children for conjunctivitis and rhinitis. *Lancet*. 2002;**359**(9310):947–948. DOI:10.1016/S0140-6736(02)08045-5
6. Lovasi GS, O’Neil-Dunne JPM, Lu JWT, Sheehan D, Perzanowski MS, MacFaden SW, et al. Urban tree canopy and asthma, wheeze, rhinitis, and allergic sensitization to tree pollen in a New York City birth cohort. *Environ Health Perspect*. 2013;**121**(4):494–500. DOI:10.1289/ehp.1205513
7. Fuertes E, Butland BK, Ross Anderson H, Carlsten C, Strachan DP, Brauer M. Childhood intermittent and persistent rhinitis prevalence and climate and vegetation: a global ecologic analysis. *Ann Allergy Asthma Immunol*. 2014;**113**(4):386–92.e9. DOI:10.1016/j.jaci.2013.03.007
8. Ruokolainen L, von Hertzen L, Fyhrquist N, Laatikainen T, Lehtomäki J, Auvinen P, et al. Green areas around homes reduce atopic sensitization in children. *Allergy*. 2015;**70**(2):195–202. DOI:10.1111/all.12545
9. Fuertes E, Markevych I, Berg A von, Bauer C-P, Berdel D, Koletzko S, et al. Greenness and allergies: evidence of differential associations in two areas in Germany. *J Epidemiol Community*

Health. 2014;**68**(8):787-790. 8. DOI:10.1136/jech-2014-203903

10. Dadvand P, Villanueva CM, Font-Ribera L, Martinez D, Basagaña X, Belmonte J, et al. Risks and benefits of green spaces for children: a cross-sectional study of associations with sedentary behavior, obesity, asthma, and allergy. *Environ Health Perspect*. 2014;**122**(12):1329–35. DOI:10.1289/ehp.1308038.
11. Richardson EA, Mitchell R, Hartig T, Vries S de, Astell-Burt T, Frumkin H. Green cities and health: a question of scale? *J Epidemiol Community Health*. 2012;**66**(2):160–5. DOI:10.1136/jech.2011.137240
12. Maas J, Verheij RA, Vries S de, Spreeuwenberg P, Schellevis FG, Groenewegen PP. Morbidity is related to a green living environment. *J Epidemiol Community Health*. 2009;**63**(12):967–73. DOI:10.1136/jech.2008.079038
13. Wickman M, Pershagen G, Nordvall SL. The BAMSE Project: presentation of a prospective longitudinal birth cohort study. *Pediatr Allergy Immunol*. 2002;**13**(Suppl. 15):11–3. DOI:10.1034/j.1399-3038.13.s.15.10.x
14. Chan-Yeung M, Manfreda J, Dimich-Ward H, Ferguson A, Watson W, Becker A. A randomized controlled study on the effectiveness of a multifaceted intervention program in the primary prevention of asthma in high-risk infants. *Arch Pediatr Adolesc Med*. 2000;**154**(7):657–63. DOI:10.1001/archpedi.154.7.657
15. von Berg A, Krämer U, Link E, Bollrath C, Heinrich J, Brockow I, et al. Impact of early feeding on childhood eczema: development after nutritional intervention compared with the natural course – the GINIplus study up to the age of 6 years. *Clin Exp Allergy*. 2010;**40**(4):627–36. DOI:10.1111/j.1365-2222.2009.03444.x
16. Heinrich J, Bolte G, Hölscher B, Douwes J, Lehmann I, Fahlbusch B, et al. Allergens and endotoxin on mothers' mattresses and total immunoglobulin E in cord blood of neonates. *Eur Respir J*. 2002;**20**(3):617–23. DOI:10.1183/09031936.02.02322001

17. Zutavern A, Brockow I, Schaaf B, Bolte G, von Berg A, Diez U, et al. Timing of solid food introduction in relation to atopic dermatitis and atopic sensitization: results from a prospective birth cohort study. *Pediatrics*. 2006;**117**(2):401–11. DOI:10.1542/peds.2004-2521
18. Lowe AJ, Carlin JB, Bennett CM, Abramson MJ, Hosking CS, Hill DJ, et al. Atopic disease and breast-feeding—cause or consequence? *J Allergy Clin Immunol*. 2006;**117**(3):682–7. DOI:10.1016/j.jaci.2005.10.027
19. Brunekreef B, Smit J, de Jongste J, Neijens H, Gerritsen J, Postma D, et al. The prevention and incidence of asthma and mite allergy (PIAMA) birth cohort study: design and first results. *Pediatr Allergy Immunol*. 2002;**13**(15):55–60. DOI:10.1034/j.1399-3038.13.s.15.1.x
20. Kozyrskyj AL, HayGlass KT, Sandford A, Pare PD, Chan-Yeung M, Becker AB. A novel study design to investigate the early-life origins of asthma in children (SAGE study). *Allergy*. 2009;**64**(8):1185–93. DOI:10.1111/j.1398-9995.2009.02033.x
21. MacIntyre EA, Carlsten C, MacNutt M, Fuertes E, Melén E, Tiesler CMT, et al. Traffic, asthma and genetics: combining international birth cohort data to examine genetics as a mediator of traffic-related air pollution's impact on childhood asthma. *Eur J Epidemiol*. 2013;**28**(7):597–606. DOI:10.1007/s10654-013-9828-5
22. Gruzieva O, Gehring U, Aalberse R, Agius R, Beelen R, Behrendt H, et al. Meta-analysis of air pollution exposure association with allergic sensitization in European birth cohorts. *J Allergy Clin Immunol*. 2014;**133**(3):767–76.e7. DOI:10.1016/j.jaci.2013.07.048
23. Weier J, Herring D. Measuring vegetation (NDVI & EVI) [Internet]. 2000 [cited 2013 Dec 30]. Available from: <http://earthobservatory.nasa.gov/Features/MeasuringVegetation/printall.php>
24. Markevych I, Fuertes E, Tiesler CMT, Birk M, Bauer C-P, Koletzko S, et al. Surrounding greenness and birth weight: Results from the GINIplus and LISAplus birth cohorts in Munich. *Health Place*. 2014;**26**:39–46. DOI:10.1016/j.healthplace.2013.12.001
25. Markevych I, Thiering E, Fuertes E, Sugiri D, Berdel D, Koletzko S, et al. A cross-sectional analysis of the effects of residential greenness on blood pressure in 10-year old children: results from the GINIplus and LISAplus studies. *BMC Public Health*. 2014;**14**(1):477.

26. McMorris O, Villeneuve PJ, Su J, Jerrett M. Urban greenness and physical activity in a national survey of Canadians. *Environ Res.* 2015;**137**:94–100. DOI:10.1016/j.envres.2014.11.010
27. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials.* 1986;**7**(3):177–88. DOI:10.1016/0197-2456(86)90046-2
28. Huedo-Medina TB, Sánchez-Meca J, Marín-Martínez F, Botella J. Assessing heterogeneity in meta-analysis: Q statistic or I^2 index? *Psychol Methods.* 2006;**11**(2):193–206. DOI:10.1037/1082-989X.11.2.193
29. Higgins JP, Green S. Cochrane Handbook for Systematic Reviews of Interventions. Chapter 9.5.2: Identifying and measuring heterogeneity. Version 5.1.0 [updated 2011]. Available from: http://handbook.cochrane.org/chapter_9/9_5_2_identifying_and_measuring_heterogeneity.htm.
30. Schwarzer G. meta: Meta-Analysis with R. R package version 3.7-0. <http://CRAN.R-project.org/package=meta>
31. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, www.R-project.org/. 2012.
32. StataCorp. Stata Statistical Software: Release 13. College Station, TX: StataCorp LP. 2013.
33. Fuertes E, Brauer M, MacIntyre E, Bauer M, Bellander T, von Berg A, et al. Childhood allergic rhinitis, traffic-related air pollution, and variability in the GSTP1, TNF, TLR2, and TLR4 genes: Results from the TAG Study. *J Allergy Clin Immunol.* 2013;**132**(2):342–52.e2. DOI:10.1016/j.jaci.2013.03.007
34. Davies HTO, Crombie IK, Tavakoli M. When can odds ratios mislead. *BMJ.* 1998;**316**(7136):989–991. DOI:10.1136/bmj.316.7136.989
35. D’Amato G, Cecchi L, Bonini S, Nunes C, Annesi-Maesano I, Behrendt H, et al. Allergenic pollen and pollen allergy in Europe. *Allergy.* 2007;**62**(9):976–90. DOI:10.1111/j.1398-9995.2007.01393.x

Table 1: Summary statistics of the study population

	BAMSE		CAPPS		GINI/LISA		GINI/LISA		MACS		PIAMA		SAGE	
	N _{total} =3304		N _{total} =357		North N _{total} =2152		South N _{total} =2855		N _{total} =327		N _{total} =3339		N _{total} =682	
	n	%	n	%	n	%	n	%	n	%	n	%	n	%
Outcomes														
Childhood (6-8 yrs)														
Allergic rhinitis	422	13.4	105	29.4	96	4.8	174	6.3	-	-	211	6.6	174	33.2
Aeroallergen sensitization	623	28.5	154	44.8	256	26.1	481	31.1	-	-	543	32.5	189	27.9
Indoor aeroallergen sensitization	413	20.9	126	36.6	174	17.8	276	17.9	-	-	432	25.8	127	18.7
Outdoor aeroallergen sensitization	503	24.3	73	21.3	183	18.7	338	21.9	-	-	305	18.3	125	18.4
Early adolescence (10-12 yrs)														
Allergic rhinitis	587	19.2	-	-	132	8.0	249	10.9	118	37.0	260	10.1	-	-
Aeroallergen sensitization	-	-	-	-	300	34.8	626	43.0	180	55.1	544	42.6	-	-
Indoor aeroallergen sensitization	-	-	-	-	211	24.5	407	28.0	166	50.8	437	34.2	-	-
Outdoor aeroallergen sensitization	-	-	-	-	223	25.9	478	32.8	116	35.5	356	27.9	-	-
Covariates														
Age at childhood ¹	8.2	(0.5)	7.2	(0.2)	6.1	(0.3)	6.0	(0.1)	-	-	8.1	(0.2)	9.1	(0.5)
Age at early adolescence ¹	13.0	(0.8)	-	-	10.1	(0.2)	10.1	(0.2)	11.2	(2.1)	11.4	(0.3)	-	-
Male sex	1668	50.5	194	54.3	1094	50.8	1469	51.5	172	52.6	1720	51.5	379	55.6
Birth weight (grams) ¹	3528.7	(557.3)	3482.1	(650.6)	3536.8	(478.4)	3415.1	(433.7)	-	-	3521.3	(540)	3378.9	(636.7)
Parental atopy	1007	30.8	331	92.7	1005	47.0	1875	66.1	309	94.8	1666	49.9	395	58.7
Older siblings	1602	48.5	198	55.5	1174	54.8	1231	43.2	204	62.4	1680	50.3	433	73.3

Maternal smoking during pregnancy		415	12.6	29	8.2	321	15.1	375	13.4	13	4.00	537	16.2	131	20.0
Parental education ²	Low	64	2.0	-	-	272	12.7	144	5.1	83	25.4	400	12.0	-	-
	Med	1410	43.9	-	-	875	40.8	513	18.0	-	-	1210	36.4	-	-
	High	1740	54.1	-	-	999	46.6	2188	76.9	244	74.6	1716	51.6	-	-
Maternal age (years) ¹		30.8	(4.5)	31.9	(5.0)	30.8	(3.8)	32.4	(4.1)	32.2	(4.1)	30.5	(3.8)	28.9	(5.3)
Intervention	Active	-	-	167	46.8	727	33.8	852	29.8	109	33.3	309	9.3	-	-
	Placebo	-	-	-	-	-	-	-	-	-	-	272	8.1	-	-
Childhood (6-8 yrs)															
Tobacco smoke at home		579	18.6	67	18.8	795	38.4	545	19.8	-	-	494	15.6	182	27.5
Furry pets at home		828	26.2	34	9.5	583	28.1	673	24.0	-	-	1697	54.5	424	62.9
Mold/dampness at home		250	7.9	175	49.0	306	15.0	590	21.9	-	-	913	29.0	475	69.9
NO ₂ concentration ¹ (µg/m ³)		11.9	(5.0)	19.5	(11.3)	23.5	(3.1)	20.1	(5.3)	-	-	22.0	(6.1)	8.1	(2.1)
Population density ³ (1000m buffer)		9341	(15602)	-	-	1218	(1678)	2829	(3389)	-	-	7359	(8395)	-	-
Living in an urban surrounding ⁴		1117	33.8	-	-	24	1.1	1452	51.1	-	-	661	20.9	-	-
Moved since birth		2161	66.7	-	-	713	34.1	1378	48.5	-	-	1611	50.8	-	-
Early adolescence (10-12 yrs)															
Tobacco smoke at home		435	16.1	-	-	464	27.8	309	13.2	-	-	299	11.6	-	-
Furry pets at home		709	22.9	-	-	596	36.0	822	35.5	-	-	1541	59.8	-	-
Mold/dampness at home		261	9.9	-	-	317	19.6	504	22.3	-	-	841	32.6	-	-
NO ₂ concentration ¹ (µg/m ³)		11.5	(5.6)	-	-	23.7	(3.4)	19.8	(5.2)	242 ⁵	(293) ⁵	21.8	(6.1)	-	-
Population density ³ (1000m buffer)		8315	(12778)	-	-	1309	(1852)	2673	(3258)	5131	(5488)	7076	(8677)	-	-
Living in an urban surrounding ⁴		893	27.0	-	-	26	1.2	1333	48.8	-	-	515	19.9	-	-
Moved since birth		2680	82.3	-	-	811	47.2	1546	64.1	173	53.0	1559	60.2	-	-

420

421 ¹ Mean (standard deviation)

422 ² Defined as the highest education attained by either parent

423 ³ Medium (interquartile range) reported

424 ⁴ Defined as $\geq 25\%$ of sealed soil in a 5000m buffer around the home address for BAMSE, GINI/LISA North, GINI/LISA South and PIAMA. Data only available for
425 the European cohorts.

426 ⁵ Minimum distance to a major road in meters (medium (interquartile range)) reported instead as NO2 concentration data were not available for MACS

427 - : not available/not applicable

428 **FIGURE LEGENDS:**

429 **Figure 1:** Cohort-specific distribution of mean NDVI in a 500m buffer around the home addresses in
430 childhood (6-8 years) and early adolescence (10-12 years). Comparisons across cohorts are not
431 appropriate as it was not possible to obtain cloud-free images at the same time for all cohorts. na = not
432 available

433 **Figure 2:** Adjusted associations between allergic rhinitis and overall, indoor and outdoor aeroallergen
434 sensitization assessed during childhood (6-8 years) with mean NDVI in a 500m buffer.

435 **Figure 3:** Adjusted associations between allergic rhinitis and overall, indoor and outdoor aeroallergen
436 aeroallergen sensitization assessed during early adolescence (10-12 years) with mean NDVI in a 500m
437 buffer.

all_12915_f1.tiff

all_12915_f2.tiff

all_12915_f3.tiff

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Fuertes, E;Markevysh, I;Bowatte, G;Gruzieva, O;Gehring, U;Becker, A;Berdel, D;von Berg, A;Bergstrom, A;Brauer, M;Brunekreef, B;Brueske, I;Carlsten, C;Chan-Yeung, M;Dharmage, SC;Hoffmann, B;Kluemper, C;Koppelman, GH;Kozyrskyj, A;Korek, M;Kull, I;Lodge, C;Lowe, A;MacIntyre, E;Pershagen, G;Standl, M;Sugiri, D;Wijga, A;Heinrich, J

Title:

Residential greenness is differentially associated with childhood allergic rhinitis and aeroallergen sensitization in seven birth cohorts

Date:

2016-10-01

Citation:

Fuertes, E., Markevysh, I., Bowatte, G., Gruzieva, O., Gehring, U., Becker, A., Berdel, D., von Berg, A., Bergstrom, A., Brauer, M., Brunekreef, B., Brueske, I., Carlsten, C., Chan-Yeung, M., Dharmage, S. C., Hoffmann, B., Kluemper, C., Koppelman, G. H., Kozyrskyj, A., ... Heinrich, J. (2016). Residential greenness is differentially associated with childhood allergic rhinitis and aeroallergen sensitization in seven birth cohorts. *ALLERGY*, 71 (10), pp.1461-1471. <https://doi.org/10.1111/all.12915>.

Persistent Link:

<http://hdl.handle.net/11343/291285>