Received Date : 17-Jun-2015

Revised Date : 28-Mar-2016

Accepted Date : 12-Apr-2016

Article type : Original Article

Geographic variation in hybridisation and ecological differentiation between three syntopic, morphologically similar species of montane lizards

M L Haines^{a,b,*}, J Melville^a, J Sumner^a, N Clemann^{a,c}, D G Chapple^d, D Stuart-Fox^b

^aSciences Department, Museum Victoria, GPO Box 666, Melbourne, VIC 3001, Australia

^bBiosciences, University of Melbourne, Parkville, VIC 3010, Australia

^cArthur Rylah Institute for Environmental Research, Department of Environment, Land, Water, and Planning, PO Box 137, Heidelberg, VIC 3084, Australia

^dSchool of Biological Sciences, Monash University, Clayton, Victoria 3800, Australia

Key words: Australia; conservation genetics; hybridisation; microsatellites; Scincidae

*Corresponding author. Current address: Department of Biological Sciences, University of Wisconsin-Milwaukee, 3209 N. Maryland Ave., Milwaukee, WI 53211, USA. Email address: hainesm@uwm.edu

Running title: Hybridisation between three lizard species

Abstract

To understand factors shaping species boundaries in closely related taxa, a powerful approach is to compare levels of genetic admixture at multiple points of contact, and determine how this relates to intrinsic and extrinsic factors, such as genetic, morphological and ecological differentiation. In the Australian Alps, the threatened alpine bog skink

Pseudemoia cryodroma co-occurs with two morphologically and ecologically similar

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/mec.13652](https://doi.org/10.1111/mec.13652)

congeners, *P. entrecasteauxii* and *P. pagenstecheri*, and all three species are suspected to hybridise. We predicted that the frequency of hybridisation should be negatively correlated with genetic divergence, morphological differentiation, and microhabitat separation. We tested this hypothesis using a mitochondrial locus, 13 microsatellite loci, morphological and microhabitat data, and compared results across three geographically isolated sites. Despite strong genetic structure between species, we detected hybridisation between all species pairs, including evidence of backcrossed individuals at the two sites where all three species are syntopic. Hybridisation frequencies were not consistently associated with genetic, morphological or ecological differentiation. Furthermore, *P. entrecasteauxii* and *P. pagenstecheri* only hybridised at the two sites where they are syntopic with *P. cryodroma*, but not at the largest site where *P. cryodroma* was not recorded, suggesting that *P. cryodroma* may serve as a bridging species. This study reveals the complex dynamics within a three species hybrid zone and provides a baseline for assessing the impact of climate change and anthropogenic habitat modification on future hybridisation frequencies.

Introduction

Contemporary contact between closely related species provides an opportunity to investigate the mechanisms maintaining or eroding species boundaries. The degree of genetic admixture between lineages at points of contact can range from no admixture to complete admixture, depending on numerous intrinsic and extrinsic factors, and their interaction (Singhal & Moritz 2013). Intrinsic barriers include genetic incompatibilities, which can prevent the development of viable or fertile offspring. Such incompatibilities are expected to accumulate over time due to genetic drift and/or divergent selection (Coyne & Orr 1989; Hewitt 2011; Mallet 2007), a concept central to the Bateson-Dobzhansky-Muller model of hybrid incompatibility (Turelli *et al.* 2001). To minimize energy wasted on the production of unfit offspring, post-zygotic barriers should select for the formation of pre-zygotic barriers, particularly in hybrid zones, and a large literature documents such a process of reinforcement (Liou & Price 1994; Noor 1999; Servedio & Noor 2003; Turelli *et al.* 2001). Common examples of such barriers include timing of reproduction, divergence in mating preferences, and mechanical incompatibility (Dobzhansky 1937). In contrast, extrinsic factors necessarily entail pre-zygotic barriers, comprising geographic barriers (e.g. rivers, mountains) and ecological separation (e.g. aboreal vs fossorial; Mayr 1942).

A powerful approach to understanding factors shaping species boundaries is to compare levels of genetic admixture between species at multiple points of contact, and how this relates to intrinsic and extrinsic factors, such as genetic, morphological and ecological

66 differentiation; yet studies that do so are relatively rare (but see Chatfield *et al.* 2010; Nice *et*
67 *al.* 2013). Morphological divergence is often correlated with ecological divergence (Losos *et*
68 *al.* 1998), and both may be inversely related to the likelihood of admixture. Specifically,
69 morphological differentiation can serve as a pre-zygotic barrier through female preference for
70 specific male secondary sexual characteristics and/or physical incompatibilities between taxa
71 (Coyne & Orr 2004). Hybrid offspring that are morphologically intermediate may be unsuited
72 to either of the parental species' ecological niches, reinforcing existing pre-zygotic barriers.
73 Furthermore, differences in ecological requirements may reduce direct contact between
74 species, decreasing opportunities for hybridisation (Mayr 1942) and/or result in hybrid
75 offspring with lower fitness (Arnold & Hodges 1995; de Leon *et al.* 2010; Harrison & Rand
76 1989). Consequently, species that have multiple, spatially isolated areas of geographic
77 overlap provide an ideal system in which to examine potential factors influencing the degree
78 of genetic admixture between species.

79 Although closely related lineages generally evolve and exist in allopatry (Turelli *et al.*
80 2001), changes in climate and habitat distribution may result in secondary contact and
81 admixture (Rhymer & Simberloff 1996). Taxa that have recently speciated are expected to
82 have accumulated the fewest reproductive barriers, and are consequently more likely to
83 produce viable hybrid offspring (Orr 1995) when environmental changes bring previously
84 isolated species into contact. When hybrids are equally fit compared to the parental species,
85 ongoing admixture can produce a variety of outcomes. For example, hybridisation can result
86 in the formation of hybrid swarms, which have been observed primarily in fishes (e.g. Avise
87 *et al.* 1984; Hasselman *et al.* 2014; Seehausen *et al.* 1997), but also mammals (Latch *et al.*
88 2011; McDevitt *et al.* 2009) and herpetofauna (Pritchard & Edmands 2013; Schulte *et al.*
89 2012). In rare cases, widespread hybridisation can lead to the extinction of pure parentals,
90 resulting in species collapse, as documented in three-spined sticklebacks (Taylor *et al.* 2006)
91 and Darwin's finches (Kleindorfer *et al.* 2014). Hybridisation can also be unidirectional and
92 lead to genetic swamping of the introgressed species. This often occurs when invasive
93 species introgress into native species (Rhymer & Simberloff 1996), and represents a
94 significant conservation threat for numerous taxa (e.g. Abernathy 1994; Allendorf & Leary
95 1988; Dowling & Childs 1992; Johnson *et al.* 2010; Leary *et al.* 1993).

96 Here we investigate contemporary genetic admixture and assess the relationships
97 between the frequency of hybridisation and ecological and morphological differentiation in
98 syntopic alpine skinks (genus *Pseudemoia*). While studies on hybrid systems generally focus
99 on two species and/or a single point of contact (but see Bogdanowicz *et al.* 2012; Fisch *et al.*

2013; Marino *et al.* 2013), we examine three species that exhibit genetic signatures of historical mitochondrial introgression (Haines *et al.* 2014) at three geographically isolated locations. All three species, *P. cryodroma*, *P. entrecasteauxii*, and *P. pagenstecheri*, are morphologically similar and occupy overlapping ecological niches in the montane and sub-alpine regions (> 1100 m above sea level) of south-eastern Australia (Wilson & Swan 2013). *Pseudemoia cryodroma* is a threatened alpine specialist (DSE 2013), restricted to habitats higher than 1200 m above sea level, while *P. entrecasteauxii* and *P. pagenstecheri sensu lato* are widespread generalists that are also sympatric in lowland areas (Wilson & Swan 2013). These three species form a clade within the genus, and divergence between *P. pagenstecheri* and *P. cryodroma* plus *P. entrecasteauxii* occurred as late as 4 mya, with more recent divergence occurring between *P. cryodroma* and *P. entrecasteauxii* (Haines *et al.* 2014). We have previously provided evidence of historic hybridisation between all three species pairs, with probable mitochondrial introgression from *P. pagenstecheri* into *P. entrecasteauxii* and *P. cryodroma*, as well as *P. entrecasteauxii* into its sister species *P. cryodroma* (Haines *et al.* 2014). As the previous study was based on mtDNA and multiple nuclear gene regions, it remains unclear whether contemporary hybridisation occurs among these species (Haines *et al.* 2014).

In the present study, we not only test for recent admixture between species, but also explore potential drivers of hybridisation. Specifically, we assess whether levels of interbreeding are consistent across three geographically isolated sites, and how genetic admixture relates to the degree of morphological and ecological differentiation between species. We predict that, if present, hybridisation will be highest between the recently diverged sister species *P. cryodroma* and *P. entrecasteauxii*, and that the level of admixture between species will be negatively correlated with morphological and ecological differentiation. By testing these predictions, this study provides rare insight into the factors influencing contemporary hybridisation dynamics of three historically interbreeding species.

Materials and methods

Study sites and sampling

Pseudemoia cryodroma, *P. entrecasteauxii*, and *P. pagenstecheri* are sympatric in the Australian Alps in north-eastern Victoria, Australia (Hutchinson & Donnellan 1992) and researchers in the field have observed them syntopically (N Clemann, pers. obs.). The three species were sampled from neighbouring mountain plateaux (Fig. 1; see Table S1, Supporting Information for sampling details) with a mosaic of snow gum (*Eucalyptus*

pauciflora) woodland, alpine heathland, and alpine grassland, isolated from one another by valleys of sclerophyll forests. The distance between sites ranged from 11 to 35 km. All three species were syntopic at Mount Higginbotham (HT) and the Bogong High Plains (BHP); whereas, only *P. entrecasteauxii* and *P. pagenstecheri* were observed on the Dargo High Plains (DHP). Located on the border of Mount Hotham Alpine Resort and the Alpine National Park, HT comprised an area 500 m by 500 m. The BHP site (500 m by 1750 m) was in the Alpine National Park, approximately 500 m south of Mt Nelse, and on the DHP, individuals were sampled along a 10 km section of the Dargo High Plains Road and along Long Spur Track. At DHP, individuals were sampled at multiple intersections of snow gum woodland and alpine grassland, where *P. entrecasteauxii* and *P. pagenstecheri* were syntopic. Large patches of tussock grasslands, where only *P. pagenstecheri* was detected, separated these small areas of syntopy. We focused our sampling on areas of syntopy in order to maximise the likelihood of detecting hybrids. Therefore, we had to cover a greater overall sampling area at DHP to obtain enough samples (>20) for genetic analysis.

During the 2010-2013 breeding seasons (December to March), we collected 111, 42, and 70 *Pseudemoia* spp. from BHP, DHP, and HT, respectively, through noosing and hand capture. Lizards were measured, and then either a tail tip was collected as a genetic sample and the lizard then released at point of capture, or they were retained as museum voucher specimens and liver tissue used as the genetic sample. All tissues and voucher specimens were registered at Museum Victoria (Table S1). Species were identified in the field using current taxonomic identifiers: the presence/absence of vertebral stripes and paravertebral stripes, male breeding colouration, and whether the mid-lateral stripe was well defined (Hutchinson & Donnellan 1992). The 30 individuals that could not be identified morphologically were classified as *Pseudemoia* sp. In the field, we recorded the following standard morphometric measurements: snout-vent length, distance from tip of the snout to the cloaca (SVL); head length, distance from posterior of skull to snout (HL); eye width, distance between eyes (HWE); head width between the widest part of jaws (HWJ); head depth, length at deepest part of the head (HD); interlimb length, length of the body from insertion of forelimb to insertion of hindlimb (ILL); pelvic width, width of the body at insertion of hindlimb (PW); upper forelimb, length from limb insertion to elbow (UFL); lower forelimb, length from elbow to wrist (LFL); forefoot, length from wrist to tip of fourth toe (FFL); upper hindlimb, length from limb insertion to knee (UHLL); lower hindlimb, length from knee to ankle (LHLL); and hind foot, length from ankle to tip of forth toe (HFL; Table S2, Supporting Information). A standard ruler was used to measure SVL to the nearest 1 mm, and

digital calipers were used to measure the remaining morphological variables to the nearest 0.01 mm.

Additionally, we recorded a number of microhabitat features. We focussed specifically on microhabitat variation rather than broader environmental variables (e.g. climate, vegetation, geology) because differences between sites in broad environmental conditions, such as temperature, rainfall, elevation and vegetation, are negligible and because microhabitat variation is most likely to capture the fine-scale variation relevant to potential for admixture in these morphologically and ecologically similar species. We recorded plant litter depth to the nearest 5 cm at the point of capture, and estimated by eye the percentage of the area within a 2 m radius of the point of capture that was dominated by ground cover, shrub (vegetation < 1 m high), tree (> 1 m high), and rock to the nearest 5%. Tree refers to the tree trunk and low-lying branches. These ecological attributes were based on previous studies on lizard habitat use (e.g. Goodman *et al.* 2008; Melville & Swain 2003; Quirt *et al.* 2006; Teasdale *et al.* 2013). All morphometric and habitat measurements were taken by the same researcher (M. Haines).

Mitochondrial DNA sequence analysis

Genomic DNA was extracted from liver and tail tip samples using a Qiagen DNeasy Blood and Tissue Extraction Kit (Qiagen, Hilden, Germany), a GenCatch Blood and Tissue Genomic DNA Miniprep Kit (Epoch Life Sciences, Sugar Land, Texas, USA) or a modified high-salt method (Aljanabi & Martinez 1997). To initially assign each individual to a given species, samples were sequenced for a 794 bp mitochondrial fragment NADH subunit 4 (ND4) and partial tRNAs using the protocol detailed in Haines *et al.* (2014). Using Geneious 6.1.2 (Biomatters, Auckland, New Zealand, available at: <http://www.geneious.com>), sequences were aligned using the default clustering algorithm, edited, and translated to amino acids. No premature stop codons were observed. Fourteen sequences had been published in Haines *et al.* (2014) and previously unpublished sequences were deposited in GenBank (Table S1, Supporting Information).

We conducted Bayesian phylogenetic analyses using MrBayes 3.2 (Ronquist & Huelsenbeck 2003) on CIPRES Science Gateway (Miller *et al.* 2010) and Maximum Likelihood (ML) analyses using PhyML 3.0 (Guindon *et al.* 2010) at <http://www.atgc-montpellier.fr/phyml/>. We assessed partitioning schemes and models of best-fit based on

Akaike Information Criterion (AIC) using MrModeltest 2.3 (Nylander 2004) and selected a GTR + Γ + I model. We included several sequences previously published in Haines et al. (2014), including 14 *P. cryodroma-pagenstecheri-entrecasteauxii* sequences (Table S1), and sequences from the three other species within the genus *Pseudemoia* (*P. baudini*: KM263203; *P. rawlinsoni*: KM263321; *P. spenceri*: KM263326) and a closely related genus (*Niveoscincus metallicus*: KM263269) as outgroups.. The Bayesian analysis consisted of two independent runs, each with four chains of Markov Chain Monte Carlo (MCMC). Chains were sampled every 500 generations and chain convergence was confirmed using an assessment of average standard deviation of split frequencies (<0.01). For the maximum likelihood analysis, the topology was determined using NNI and approximate ratio-likelihood test (aLRT).

Primer screening and development

Fourteen microsatellite loci developed for *P. entrecasteauxii* by Stapley *et al.* (2003) were tested for cross amplification in *P. cryodroma* and *P. pagenstecheri*. Only seven loci amplified cleanly and consistently for all three species (Table S3, Supporting Information). For these markers, fluorescently labelled dyes were attached to forward primers and a GTTTCT ‘pig-tail’ sequence was added to the reverse primers to reduce stuttering when scoring microsatellites (Brownstein *et al.* 1996).

As part of the current study additional microsatellite markers were developed using 454-sequencing to supplement these published markers. One *P. entrecasteauxii* sample was sent to the Australian Genome Research Facility (AGRF, Queensland, Australia) for high throughput DNA sequencing on 1/16 of a plate using the Roche GS FLX (454) system. A detailed sequencing protocol can be found in Gardner *et al.* (2011). Using Geneious 6.1.6 (Biomatters), we searched for sequences with a minimum of 8 tetra-, penta-, or hexa- repeats. Using the default parameters in the program Primer3 (Untergasser *et al.* 2012), primers were designed for potential microsatellite loci. Following the approach by James *et al.* (2011), forward primers were tailed with a 454A adapter primer sequence (5’ GCCTCCCTCGCGCCATCAG 3’; Margulies *et al.* 2005), using a modified protocol of Schuelke (2000). Thirty-seven candidate loci were identified and primers for 17 loci were tested with DNA from each of the three *Pseudemoia* spp. Loci that amplified successfully were further optimized (see below for PCR details). The seven loci that were polymorphic for

the three *Pseudemoia* spp. were used, with seven of the published markers, to genotype all individuals (Table S3, Supporting Information).

Microsatellite genotyping and allelic diversity

All 223 individuals were genotyped at 14 microsatellite loci. For the loci identified by Stapley *et al.* (2003), PCRs were performed in 20 μ L reactions containing 0.5 μ L of each primer (10 μ M), 10 μ L GoTaq Hot Start Master Mix (Promega, Madison, Wisconsin, USA), and 2 μ L genomic DNA. The PCRs using primers designed for this study were performed in 20 μ L reactions containing 0.25 μ L forward primer with 454A tail (10 μ M), 0.15 μ L fluorescent dye with corresponding tails (6-FAM, VIC, NED, or PET; 10 μ M), 0.5 μ L reverse primer (10 μ M), 10 μ L GoTaq Hot Start Master Mix (Promega, Madison, Wisconsin, USA), and 2 μ L genomic DNA. All PCR protocols started with an initial denaturation at 95 °C for 5 min, followed by 40 cycles consisting of denaturation at 95 °C for 30 s, annealing at temperatures ranging from 50-65 °C (for details see Table S4, Supporting Information) for 30 s, and extension at 72 °C for 45 s, followed by a final 5 min extension at 72 °C. The PCR products were sent to Macrogen, Inc (Seoul, South Korea) and analysed on an AB 3730 platform using a LIZ-500 size standard. Chromatograms were scored in Geneious 6.1.6 (Biomatters) and visually checked for accuracy. We checked samples for identical genotypes using Microsoft Excel. In Microchecker version 2.3.3 (Van Oosterhout *et al.* 2004), we checked for evidence of stutter products, large allele dropout, and null alleles.

Allelic diversity was quantified for each species within sites, using only individuals with genetically pure ancestry (see Results). The number of alleles and private alleles were calculated in GenAlEx 6.5 (Peakall & Smouse 2012). We determined allelic richness and private allelic richness, correcting for sample size, using HP-Rare (Kalinowski 2005). We calculated expected and observed heterozygosity and tested for Hardy-Weinberg equilibrium and linkage disequilibrium in Arlequin 3.5.1.3 (Excoffier & Lischer 2010). To assess statistical significance of Hardy-Weinberg equilibrium and linkage disequilibrium results, a Bonferroni correction was used to account for multiple comparisons (Rice 1989). For each site, genetic distance between species was estimated by calculating Jost's *D* in GenAlEx 6.5 and pairwise *F_{ST}* in Genodive 2.0b25 (Meirmans 2006).

Admixture analysis

To determine the presence of contemporary hybridisation at each site, we analysed the microsatellite data from the three sites separately in the programs STRUCTURE 2.3.3

(Pritchard *et al.* 2000) and NewHybrids 1.1 (Anderson & Thompson 2002). STRUCTURE uses a Bayesian clustering algorithm to determine the most likely number of genetic clusters (K) in the dataset, and calculates the individual proportion of membership (Q) of each individual to each of the clusters. Individuals with a Q value close to 1 are considered pure, while those with Q values of ~0.5 for two separate clusters are likely to be F1 hybrids. In STRUCTURE, the parameters were set to allow for admixture between clusters and implemented the correlated allele frequency model. Analyses were run with a burn-in of 10^4 iterations followed by 10^6 iterations. The number of clusters (K) was set from 1-8 and executed five runs for each K. Results were combined in STRUCTURE HARVESTER (Earl & von Holdt 2012) and the most likely K was estimated based on where the Ln(K) plateaued (Pritchard *et al.* 2009) and the highest value of ΔK (Evanno *et al.* 2005). Outputs from STRUCTURE HARVESTER were combined in CLUMPP (Jacobsson & Rosenberg 2007).

In addition to STRUCTURE, the data was analysed with NewHybrids, which specifically calculates the probability of an individual belonging to either of the parental species and one of four hybrid classes (F1, F2, and backcrosses). Since NewHybrids can only accommodate two species, for sites with three species (BHP and HT), we separately analysed individuals from each pair of parental species. Individuals were assigned to a species based on the mtDNA analysis. A model with a Jeffrey's-like prior was applied for allele frequencies and mixing proportions, and implemented burn-in of 10^4 sweeps, followed by 10^6 sweeps. Runs were executed five times and repeated using Uniform priors.

Since the appropriate threshold for identifying pure individuals can vary depending on factors such as allele size convergence, overall proportion of hybrids, and number of loci analysed (Vähä & Primmer 2006), simulations were run to compare how accurately STRUCTURE and NewHybrids could identify both pure and admixed individuals at the commonly used thresholds of $Q \geq 0.90$ and $Q \geq 0.95$ (Bogdanowicz *et al.* 2012; Burgarella *et al.* 2009; Marino *et al.* 2013). Because all three species are sympatric throughout the Australian Alps, pure individuals could not be simulated using genotypes from nearby allopatric populations. Instead a subset of the empirical data was used for the simulations, which has been shown to produce almost identical results (Vähä & Primmer 2006). Preliminary STRUCTURE analyses indicated that the most likely number of clusters corresponded to the number of species morphologically identified at each site (see Results); therefore, we used individuals that met the stringent threshold of $Q \geq 0.95$ for the cluster corresponding to their mitochondrial lineage to simulate pure individuals (Garroway *et al.* 2010; Tsy *et al.* 2013). Using the individuals classified as pure, we simulated genotypes for

500 individuals for each species in HybridLab (Nielsen *et al.* 2006). These simulated genotypes were used to simulate another 500 individuals for each combination of F1s and the simulated pure individuals and F1s were then used to simulate F2s, and backcrosses. To ensure that the overall proportion of simulated hybrids was comparable to that predicted for the empirical data, the number of simulated hybrids was limited to 20 randomly selected individuals from each hybrid class. The resulting dataset was run through STRUCTURE using the same parameters as the original analyses, except K was set to the number of species at that site. Results were summarized in CLUMPP and visualized in MS Excel. The simulated data was also run in NewHybrids using the same settings as for the empirical data to compare the accuracy of applying cut-off values of 0.90 and 0.95 for membership to either parental group or a specific hybrid class. Individuals with $Q \geq 0.90$, or $Q \geq 0.95$ for a parental category were considered to be pure for that species and all other individuals were classified as hybrids for subsequent comparisons of hybridisation frequencies between sites, and in relation to ecology and morphology.

Morphological analysis

To evaluate morphological differentiation, we first tested for differences between the sexes and between sites for each species for each morphological variable. Only adults (SVL > 40 mm; Hutchinson & Donnellan 1992) were used for analyses to eliminate possible ontogenetic effects on morphology. To account for body size, measurements were regressed against SVL and the residuals were used for further analyses. Individuals were assigned to a species or classified as hybrids based on the genetic analyses. Due to small sample sizes, all hybrid classes from possible parental combinations were pooled for subsequent analyses. Univariate analyses were performed in the program R 3.0.2 (R Development Core Team 2013), applying false discovery rate corrections for multiple tests (Benjamini & Hochberg 1995). These initial univariate analyses revealed significant differences between the sexes in all species and hybrids for HL, HWJ, UFL, LHLL, and HFL ($\alpha = 0.05$); therefore, we analysed the sexes separately. As neither sex differed significantly between sites, individuals were pooled across sites for subsequent multivariate tests. A multivariate discriminant function analysis with cross-validation (PROC DISCRIM, SAS 9.3) was implemented to assess how accurately individuals could be assigned to their genetic species based on morphology.

Microhabitat analysis

Using the genetic species identifications, we assessed the degree of habitat differentiation between species at each site. Due to excess zero values in the dataset (i.e. numerous individuals had 0% for one or more of litter depth, shrub, tree and rock cover), only one habitat variable (ground cover) met assumptions of normality and homogeneity of variance, even after data transformation; therefore, non-parametric analyses were performed. As initial plots of the raw data showed clear differences between sites, differences between species were tested for each microhabitat variable separately at each site using Mann-Whitney-Wilcoxon tests in *R*. For sites with more than two species, *p*-values were adjusted for multiple comparisons using false discovery rate. For each site, a non-parametric discriminant analysis with cross-validation (PROC DISCRIM, SAS 9.3) was also performed to assess how accurately individuals could be assigned to their genetic species based on microhabitat characteristics.

Results

Verification of species assignments

The Bayesian and maximum likelihood analyses of the mitochondrial locus ND4 separated individuals into three well-supported clades, each representing one of the three species (Fig. S1, Supporting Information). This mitochondrial species assignment was used as an *a priori* hypothesis of true species identity for all subsequent analyses. The majority of mitochondrial and microsatellite identifications were congruent, and hybrids generally grouped with the mitochondrial clade corresponding to the microsatellite cluster ($Q > 0.5$). Putative F1 hybrids were randomly distributed among mitochondrial clades, suggesting bi-directional hybridisation. There were four mismatches between mitochondrial and microsatellite species assignments, indicating mitochondrial introgression of *P. cryodroma* into *P. pagenstecheri*, *P. pagenstecheri* into *P. cryodroma*, and two instances of *P. entrecasteauxii* into *P. cryodroma*.

Microsatellite diversity

All microsatellites were polymorphic in each species. Locus Pe24 was removed from subsequent analyses because Microchecker consistently showed an excess of homozygotes and deviation from Hardy-Weinberg Equilibrium, following a Bonferroni correction, for species at the sampling location level. This result indicates the presence of null alleles, which can lead to the underestimation of true allelic diversity (Chapuis & Estoup 2007). While other loci (Pe31, Pe124, Pe197, Pe303, Pe304, Pe305, and Pe306) exhibited an excess of

homozygotes and deviated from Hardy-Weinberg Equilibrium, this was randomly distributed across species and localities (Table S5, Supporting Information); consequently, these loci were retained in subsequent analyses. For these 13 loci, 97.6% of alleles were successfully scored. For each species, between 1.9% and 4.5% of all possible pairwise combinations of loci were found to be in linkage disequilibrium, following a Bonferroni correction ($\alpha = 0.05$). However, linkage disequilibrium was not observed consistently between the same loci across species; therefore, no additional loci were excluded. At DHP and HT, *P. entrecasteauxii* had the greatest number of alleles and private alleles, whereas at BHP *P. cryodroma* had the highest values in these two categories (Table S5, Supporting Information). *Pseudemoia entrecasteauxii* exhibited the highest allelic richness and private allelic richness, correcting for sample size, at each locality. Average observed heterozygosity ranged from 0.61 to 0.69 but was not consistently higher or lower for a given species or location. At each locality, both Jost's D and F_{ST} were statistically significant ($p < 0.01$) and were qualitatively the same; therefore, subsequent use of the term 'genetic distance' refers to both measures unless otherwise stated (Table 1). While genetic distance between *P. cryodroma* and *P. entrecasteauxii* was almost identical at BHP and HT, genetic distance for this pair was lower than between the other species pairs at BHP, but comparable to the genetic distance between *P. entrecasteauxii* and *P. pagenstecheri* at HT. In addition, genetic distance was always greater between *P. cryodroma* and *P. pagenstecheri* than *P. cryodroma* and *P. entrecasteauxii*.

Population structure and hybridisation

STRUCTURE results indicated that the most likely number of clusters (K) was equal to the number of species recorded at that locality, with each cluster corresponding to a different species (Fig. S2). Although at HT there was greater support for $K=2$ with *P. cryodroma* and *P. entrecasteauxii* as one cluster and *P. pagenstecheri* as another cluster, when the *P. cryodroma* and *P. entrecasteauxii* cluster was analysed separately, there was highest support for splitting individuals by species into two clusters. The 95% confidence intervals (CIs) for individuals classified as pure individuals were generally between 1.00 and 0.90 and did not overlap with the CI for another cluster. For the simulated datasets, we calculated the number of pure and admixed individuals that were accurately identified at $Q \geq 0.95$ and $Q \geq 0.90$, and found negligible differences (Fig. S3, Supporting Information, Table S6, Supporting Information). In order to maximize the likelihood that individuals classified as hybrids were truly hybrids, the lower threshold was applied (Vähä & Primmer 2006).

Individuals were classified as admixed between two or more species when $0.90 > Q \geq 0.10$ for multiple clusters. A lower limit of $Q \geq 0.10$ was applied to minimize the number of individuals incorrectly classified as having ancestry from all three species. Using this criterion, only 1.9% and 0.6% of simulated hybrid (F1, F2, backcrosses) individuals from BHP and HT, respectively, were mis-classified.

Based on the threshold of $Q \geq 0.90$ for pure individuals, there was evidence of admixture at BHP and HT but not at DHP (Fig. S2; Table S1, Supporting Information). The overall proportion of putative genetic hybrids was higher at HT (22.9%) than BHP (18.0%; Fig. 2), with hybridisation detected between all pairs of species. The majority of hybrid individuals had Q values > 0.70 for one species and < 0.30 for a second species, indicative of backcrossing (Fig. 3A). Notably, *P. cryodroma* – *P. entrecasteauxii* hybrids were either as common or more common as the other hybrid combinations. However, the relative proportion of other hybrid combinations was not consistent across sites. Specifically, the rarest hybrid combination at BHP was *P. entrecasteauxii* – *P. pagenstecheri*, which was detected in less than 2% of individuals. Nevertheless, this type of hybrid at HT was as common as *P. cryodroma* – *P. pagenstecheri* hybrids, and comprised 4% of the dataset. At both BHP and HT, one individual was morphologically identified as *P. cryodroma* yet had $Q \geq 0.10$ for all three clusters. One of the individuals was placed in the *P. cryodroma* mtDNA clade and the other in the *P. entrecasteauxii* clade. This indicates possible hybridisation between two species followed by back-crossing with the third species. Additionally, four individuals identified morphologically from HT and one from BHP were assigned as genetically pure individuals ($Q \geq 0.90$) of a species that did not correspond to their morphologically assigned species. In one instance, an individual that morphologically resembled *P. pagenstecheri* but was classified as *P. cryodroma* ($Q = 0.97$) was assigned to the *P. entrecasteauxii* mtDNA clade. Notably, 17 males from BHP had a single dorsal stripe and a defined lateral stripe, which is characteristic of *P. cryodroma*, yet they also exhibited orange/red ventral breeding colouration, which has only been reported previously in male *P. entrecasteauxii*. Fifteen of these individuals were genetically classed as pure *P. cryodroma* based on both the mtDNA and microsatellite analyses, indicating that ventral breeding colouration is not exclusive to *P. entrecasteauxii*.

The results from the NewHybrids analyses indicate an overall higher proportion of hybrids compared to the STRUCTURE outputs. As with STRUCTURE, there was little difference between applying the cut-offs of $Q \geq 0.90$ and $Q \geq 0.95$ to correctly distinguish

between simulated pure and hybrid individuals (Table S6, Supplementary Information). Consequently, a more conservative $Q \geq 0.90$ was applied to maximize the probability of identifying true hybrids. Using uniform priors, NewHybrids identified slightly more hybrid individuals than did STRUCTURE, and considerably more using Jeffrey's-like priors. Although the NewHybrids results were broadly consistent with the results from the STRUCTURE analysis, we conservatively based the assignment of pure and hybrid individuals on the results from the STRUCTURE analyses. Notably, the NewHybrids analysis revealed stronger evidence for F2 and backcrossed individuals than F1 hybrids; however, individuals were only assigned as F2s with more than 90% probability (Fig. 3B).

The relationship between the proportions of hybrids for each species pair was inconsistently associated with their respective genetic relatedness across sampling sites. Although genetic distance at HT was approximately 1.5 times greater between *P. cryodroma* and *P. pagenstecheri* compared to *P. entrecasteauxii* and *P. pagenstecheri*, we observed an equal number of hybrids for both pairings. However, we observed the opposite pattern at BHP, with approximately equal genetic distances between *P. cryodroma* and *P. pagenstecheri* versus *P. entrecasteauxii* and *P. pagenstecheri* but six *P. cryodroma* – *P. pagenstecheri* hybrids compared to one *P. entrecasteauxii* – *P. pagenstecheri* hybrid. The only pattern consistent with predictions was that as genetic distance increased between *P. entrecasteauxii* and *P. pagenstecheri*, the number of hybrids for this species pair decreased. While this pattern holds for the number of *P. cryodroma* – *P. pagenstecheri* hybrids, the proportion of *P. cryodroma* – *P. pagenstecheri* hybrids at BHP and HT is only one percent different. Therefore, there appears to be little general concordance between genetic distance estimates between species and propensity to hybridise.

Morphological differentiation

The univariate analyses revealed significant differences in HL, HWE, and all limb measurements for both sexes, and males also showed differences in HWJ and PW (Table S7, Supporting Information). In males, *P. entrecasteauxii* had larger head proportions but narrower PW compared to *P. cryodroma*. While *P. cryodroma* had longer UFL and LFL than *P. pagenstecheri*, both species had shorter limb measurements for five of the six limb variables compared to *P. entrecasteauxii*. Pairwise tests for females revealed similar patterns, with *P. entrecasteauxii* having relatively larger heads and longer limbs than *P. cryodroma* and *P. pagenstecheri*.

Accordingly, the discriminant function analyses revealed significant discrimination between species for both males (Wilks' $\lambda = 0.177$, $F_{39, 264} = 5.39$, $p < 0.001$) and females (Wilks' $\lambda = 0.284$, $F_{39, 189} = 2.47$, $p < 0.001$). Canonical variables (CV) 1 and 2 together explained 90.4% and 94.5% of the variation for males and females, respectively (Table 2). For males, HL, LFL, UFL and LHLL contributed most strongly to CV 1, while FFL contributed most strongly to CV 2, followed by the forelimb measurements. Similarly, LFL and FFL contributed most strongly to CV 1 in females (CV 2 was not significant). In both sexes, *P. entrecasteauxii* had longer limbs compared to the other two taxa (Fig. 4). However, there was substantial morphological overlap with correct assignment of males and females to their genetic species being only 61.1% and 33.3% for *P. cryodroma*, 67.9% and 78.3% for *P. entrecasteauxii*, and 85.2% and 50.0% for *P. pagenstecheri*, respectively (Table S8, Supporting Information). Male and female hybrids were most likely to be mis-classified as *P. cryodroma*.

Overall, morphological similarities between species did not correspond consistently with genetic distance or the overall proportion of different hybrid categories. The most common hybrids were between the closely related *P. cryodroma* and *P. entrecasteauxii*, but *P. cryodroma* and *P. pagenstecheri* showed least morphological differentiation. However, *P. entrecasteauxii* and *P. pagenstecheri* showed the greatest morphological differentiation in both sexes, and hybrids between this pair of species were absent at DHP, and uncommon at BHP and HT. This was most prominent in females, where no *P. entrecasteauxii* was mis-classified as *P. pagenstecheri* and only 2 *P. pagenstecheri* were mis-classified as *P. entrecasteauxii*.

Microhabitat differentiation

We assessed the degree of ecological separation between species based on five microhabitat characteristics: litter depth, ground cover, shrub cover, tree cover and rock cover. At all sites, univariate analyses showed significant differences between species in at least one microhabitat variable; however, differences between species were greatest at DHP and HT (Table S9, Supporting Information). Nonparametric discriminant analyses accordingly revealed the greatest habitat separation at DHP. At DHP, CV 1 explained 100% of the variation and correlated positively with ground cover and negatively with rock cover. *Pseudemoia entrecasteauxii* generally occupied areas with less ground cover and greater rock cover than *P. pagenstecheri* (Fig. 5; Table 3). However, these species did exhibit some ecological overlap, and 21.7% of *P. entrecasteauxii* and 15% of *P. pagenstecheri* were mis-

classified as each other based on habitat variables (Table S10, Supporting Information). At BHP, CV 1 and 2 collectively explained 92.1% of the variation, though only the former was significant. Higher values of CV 1 corresponded to shallower litter depth, with *P. cryodroma* occupying habitats with shallower litter depth than *P. entrecasteauxii*. Overall, discrimination was very poor at BHP and correct classification was low (*P. cryodroma* 29.0 %; *P. entrecasteauxii* 47.6%; hybrids 27.3%; *P. pagenstecheri* 14.1%). By contrast, discrimination between species was much greater at HT. Both CV 1 and 2 were significant and explained a combined 99.0% of the variation. CV 1 correlated positively with litter depth and ground cover, and negatively with tree cover, whereas CV 2 correlated positively with tree cover and negatively with rock cover. The most noticeable separation was between *P. entrecasteauxii* and *P. pagenstecheri*, with the former occupying areas with lower litter depth, lower ground cover and more tree cover than the latter. Based on nonparametric discriminant analyses, hybrid individuals at BHP were most likely to be classified as *P. pagenstecheri*. Classification of hybrids was poorest at HT, where hybrids were more likely to be assigned to any of the parental species than the hybrid category.

As with morphology, mis-classification rates based on ecological variables did not show a consistent correlation with genetic distance estimates or proportion of hybrids. The greatest ecological separation was between *P. entrecasteauxii* and *P. pagenstecheri* at DHP, where no hybrids were detected. By contrast, even though ecological differentiation between species was much greater at HT than at BHP, there was a higher proportion of hybrids at HT. Furthermore, the species pair with the most hybrids, *P. cryodroma* and *P. entrecasteauxii*, showed the least ecological differentiation at BHP but the highest ecological differentiation at HT. Genetic distance estimates between *P. entrecasteauxii* and *P. pagenstecheri* were highest at DHP and lowest at HT, yet mis-classification rates were not substantially different between these sites.

Discussion

Our results provide evidence of strong genetic differentiation between three morphologically and ecologically similar taxa that occur syntopically within alpine and subalpine regions of south-eastern Australia and have an evolutionary history of introgression (Haines *et al.* 2014). Despite strong genetic differentiation, we detected evidence of hybridisation between all three species at two sites, BHP and HT, including evidence of backcrossed individuals. At the third site, DHP, only *P. entrecasteauxii* and *P. pagenstecheri* were collected, and there were no hybrids despite admixture between these two taxa at both

BHP and HT. Consistent with the absence of hybrid individuals at DHP, *P. entrecasteauxii* and *P. pagenstecheri* showed the greatest morphological differentiation (all three sites combined) and the greatest genetic and ecological differentiation at DHP of any pair of species at a given site. However, contrary to predictions, genetic, morphological and ecological differentiation did not consistently correspond to the proportion of the other hybrid-crosses at BHP and HT. Below, we first discuss processes that may generate the observed patterns of hybridisation, and then suggest alternative explanations for geographic variation in hybridisation.

Genetic differentiation despite hybridisation

We detected evidence of admixture between *P. cryodroma*, *P. entrecasteauxii* and *P. pagenstecheri* at two of the three localities, yet the proportion of hybrids was generally small, particularly at BHP, and we found no evidence of hybridisation between *P. entrecasteauxii* and *P. pagenstecheri* at DHP. Thus, reproductive barriers must exist; however, they are likely incomplete, as there was strong evidence of backcrossed individuals. Specifically, 19% hybrid individuals were assigned Q values of approximately 0.75 and 0.25 to two different populations (species) in STRUCTURE, indicative of first generation backcrosses. Although the number of loci used was insufficient to confidently assign individuals to specific hybrid classes in NewHybrids (Vähä & Primmer 2006), there was strong evidence ($Q > 0.50$) that 23% of putative hybrids were either F2s or backcrosses. In contrast, the NewHybrids analysis showed a maximum of $Q = 0.17$ for the F1 hybrid category. A similar pattern of fewer F1s compared to backcrossed individuals has been documented in other taxa, such as the *Ensatina eschscholtzii* salamander species complex (Alexandrino *et al.* 2005), wall lizards *Podarcis* spp. (Pinho *et al.* 2009), and spiny lizards *Sceloporus* spp. (Robbins *et al.* 2014). The low observed proportion of F1 individuals suggests potential selection against hybrids.

There is a notable absence of hybrids between *P. entrecasteauxii* and *P. pagenstecheri* at DHP, despite evidence of admixture between these species at both BHP and HT. One explanation for the disparity in hybridisation between sites is that *P. entrecasteauxii* and *P. pagenstecheri* do not hybridise, or rather they do not hybridise directly. An intriguing possibility is that *P. cryodroma* serves as a bridging species. Specifically, hybrids of *P. cryodroma* and *P. entrecasteauxii* (or *P. pagenstecheri*) may hybridise with *P. pagenstecheri* (or *P. entrecasteauxii*), resulting in offspring that exhibit alleles from both *P. entrecasteauxii* and *P. pagenstecheri* despite no direct mating between pure *P. entrecasteauxii* and *P. pagenstecheri*. We did detect an individual with $Q > 0.10$ for all three species at BHP,

suggesting that hybrids involving all three species do occur. Moreover, a similar case has been documented in fishes from the Colorado River Basin where direct hybridisation was not detected between two of the three study species, but there was evidence of hybrids with ancestry from all three species (McDonald *et al.* 2008). Alternatively, it is possible that *P. entrecasteauxii* – *P. pagenstecheri* hybrid-crosses exist at DHP at very low frequencies because these species are further along the speciation continuum at DHP compared to BHP and HT. This is consistent with the greater genetic and ecological differentiation between this species pair at DHP compared to the other two sites.

Geographic variation in hybridisation: relationships between genetic, ecological and morphological differentiation

According to the Bateson-Dobzhansky-Muller model of genetic incompatibility, recently diverged species will have accumulated relatively fewer genetic incompatibilities compared to more distantly related species and are therefore more likely to hybridise. This pattern has been observed in numerous other vertebrates including lizards (Singhal & Moritz 2013), snakes (Tarroso *et al.* 2014), and bats (Bogdanowicz *et al.* 2012). Accordingly, we predicted that hybridisation should be highest between the recently diverged sister species *P. cryodroma* and *P. entrecasteauxii* and indeed this was the case at BHP and HT. However, the total percentage of hybrids as well as the relative proportion of each hybrid combination differed between sites. Hybridisation was only observed at BHP and HT, where all three species were recorded, and was substantially higher at HT (22.9%) than at BHP (18.0%). At BHP, genetic distance between species pairs was negatively associated with the frequency of hybrids but this was not the case at HT. Specifically, *P. cryodroma* – *P. entrecasteauxii* hybrids were more common than *P. entrecasteauxii* and *P. pagenstecheri* hybrids at HT, despite the two species pairs having a similar genetic distance. Thus, genetic distance between species may more closely correspond to the relative strength of pre-zygotic than post-zygotic barriers (Coyne & Orr 1989; Turelli *et al.* 2001), particularly at HT.

We predicted that greater morphological and ecological differentiation between species would correspond to lower levels of admixture because morphological and ecological differentiation are often associated with pre-zygotic barriers to reproduction (Coyne & Orr 2004; Mayr 1942; Turelli *et al.* 2001). Consistent with this prediction, *P. entrecasteauxii* and *P. pagenstecheri* showed the greatest morphological differentiation in both sexes and differences in stripe patterning are most pronounced between this species pair. Furthermore, at DHP, these species exhibited the greatest ecological differentiation between any species

pair at any site. Thus, ecological and morphological differentiation between *P. entrecasteauxii* and *P. pagenstecheri* may generate selection against intermediates and account for the absence of hybrids between the two species at DHP and their rarity at BHP and HT. However, in contrast to predictions, *P. cryodroma* and *P. entrecasteauxii* showed the greatest admixture, yet showed intermediate morphological differentiation across localities, and the highest ecological differentiation of any species pair at HT. Furthermore, overall microhabitat differentiation between species was substantially lower at BHP than HT; yet a higher proportion of hybrids were observed at HT. Lastly, we observed comparable ecological differentiation between *P. entrecasteauxii* and *P. pagenstecheri* at HT and DHP; however, we detected *P. entrecasteauxii* – *P. pagenstecheri* hybrids at HT and none at DHP. Overall, therefore, our results do not support the general prediction that hybridisation should be inversely related to the degree of morphological or ecological differentiation (Mayr 1942; Turelli *et al.* 2001).

Previous work on other morphologically conserved lizards similarly found no consistent patterns between morphological differentiation and reproductive isolation (Singhal & Moritz 2013). Moreover, hybridisation was roughly symmetric for all species pairs based on the mtDNA in contrast to the asymmetric introgression commonly observed between morphologically divergent lizards (Jezkova *et al.* 2013; Olave *et al.* 2011; Robbins *et al.* 2014; Schulte *et al.* 2012; While *et al.* 2015). Interestingly, *P. cryodroma* is morphologically intermediate to *P. entrecasteauxii* and *P. pagenstecheri*, and more hybrids had *P. cryodroma* ancestry than either of the two other species. If intermediate hybrids are selected against in either parental habitat, then this should have a greater impact on *P. entrecasteauxii* – *P. pagenstecheri* crosses than those involving *P. cryodroma*. Nevertheless, many other pre- and post-zygotic mechanisms are likely to influence the propensity to hybridise. In wall lizards (*Podarcis* spp.), where hybridisation has been documented in a contact zone with no known ecological or temporal barriers (Pinho *et al.* 2009), previous studies suggest that chemosensory cues likely serve as pre-zygotic barriers to hybridisation (Barbosa *et al.* 2006; Gabriot *et al.* 2010b). Since all three *Pseudemoia* spp. have overlapping breeding seasons (Hutchinson & Donnellan 1992) and none exhibit mutually exclusive breeding colouration, the strength of chemosensory cues and other previously unmeasured sexually selected traits may determine hybridisation levels. Moreover, geographic variation in such traits may further explain the observed patterns of hybridisation in this study.

External factors, such as the spatial extent of suitable habitat and degree of anthropogenic habitat loss or modification, may also influence hybridisation frequencies. For

example, rising global temperatures have allowed both flora (Díaz-Varela *et al.* 2010; Grabherr *et al.* 1994) and fauna (Brereton *et al.* 1995; Pounds *et al.* 1999; Sinervo *et al.* 2010) to expand their ranges upward, creating overlaps with higher elevation species that are unable to realize an equivalent shift in elevation (Brereton *et al.* 1995; Parmesan 2006). Suitable sub-alpine habitats for *Pseudemoia* spp. are expected to shrink and/or disappear with rising average temperatures (IPCC 2013). Spatially smaller habitats provide less opportunities for microhabitat segregation, increasing the likelihood of direct contact between species and, therefore, hybridisation (Rhymer & Simberloff 1996). Correspondingly, the locality with the most restricted area of suitable habitat (HT) had the highest percentage of hybrids. All samples from HT were collected within 400 m of commercial buildings, and the site itself was intersected by the region's only major roadway. Increased habitat openness has already been linked with hybridisation between syntopic sister species of anoles (Jezkova *et al.* 2013). In contrast, the localities with more extensive unmodified habitat (BHP and DHP) had fewer or no hybrids. Thus, additional habitat fragmentation may increase hybridisation among populations and should be considered when planning future ski resort development. Multiple, additive threats to these species (such as climate change and habitat loss and fragmentation) compound their conservation status.

Conclusion

This study provides rare insight into the current dynamics of hybridisation between three montane species whose distributions are likely to shift with anthropogenic climate change (Green *et al.* 1992; Hennessy *et al.* 2003; Pickering 2007). Consistent with theoretical expectations (Turelli *et al.* 2001), the frequency of hybrids was greatest between the most recently diverged sister species pair *P. cryodroma* and *P. entrecasteauxii*. However, there was no consistent relationship between the propensity for species to hybridise and their genetic distance based on microsatellites, nor the strength of the potential pre-zygotic barriers examined (morphology and microhabitat). Despite clear evidence of hybridisation at two of the three sites, we found strong genetic structure among species. Future research should therefore explore other potential pre-zygotic reproductive barriers, such as chemosensory cues. Furthermore, subsequent studies should explore the possibility that *P. cryodroma* may serve as a bridging species, facilitating hybridisation between *P. entrecasteauxii* and *P. pagenstecheri* where the three species co-occur. We stress the importance of continuing to monitor admixture among these species to determine whether the frequency of hybridisation shifts in response to future changes in habitat size and quality. Current suitable habitat for the

montane endemic *P. cryodroma* is expected to shrink as average temperatures continue to rise and may disappear as early as 2050 (Green *et al.* 1992; Hennessy *et al.* 2003; Pickering 2007). Thus, such information is critical to understanding the evolutionary implications of hybridisation among *Pseudemoia* spp. as well as the conservation management of the threatened *P. cryodroma*.

Acknowledgements

We thank A. Moussalli for insightful discussions and we thank L. Beilharz, R. Groenewegen, S. Heap, K. Howard, and M. Scroggie for assistance with specimen and tissue collection. We thank the Department of Environment and Primary Industries for in-kind support. C. Moritz, C. Schneider, and anonymous reviewers provided useful comments on earlier drafts of this manuscript. Funding was provided to M.L.H. through the Holsworth Wildlife Research Foundation, Museum Victoria 1854 Student Scholarship, Dame Margaret Blackwood Soroptimist Scholarship, and Wildlife Preservation Society of Australia. D.G.C. was supported by a Monash University Linkage Project Application Support (LPAS) grant.

References

- Abernathy K (1994) The establishment of a hybrid zone between red and sika deer (genus *Cervus*). *Molecular Ecology* **3**, 551-562.
- Alexandrino J, Baird SJE, Lawson L, *et al.* (2005) Strong selection against hybrids at a hybrid zone in the *Ensatina* ring species complex and its evolutionary implications. *Evolution* **59**, 1334-1347.
- Aljanabi SM, Martinez I (1997) Universal and rapid salt-extraction of high quality genomic DNA for PCR-based techniques. *Nucleic Acids Research* **25**, 4692-4693.
- Allendorf FW, Leary RF (1988) Conservation and Distribution of Genetic Variation in a Polytypic Species, the Cutthroat Trout. *Conservation Biology* **2**, 170-184.
- Anderson EC, Thompson EA (2002) A model-based method for identifying species hybrids using multilocus genetic data. *Genetics* **160**, 1217-1229.
- Arnold ML, Hodges SA (1995) Are natural hybrids fit or unfit relative to their parents. *Trends in Ecology and Evolution* **10**, 67-71.

- Awise JC, Bermingham E, Kessler LG, Saunders NC (1984) Characterization of mitochondrial DNA variability in a hybrid swarm between subspecies of bluegill sunfish (*Lepomis macrochirus*). *Evolution* **38**, 931-941.
- Barbosa D, Font E, Desfilis E, Carretero MA (2006) Chemically mediated species recognition in closely related *Podarcis* wall lizards. *Journal of Chemical Ecology* **32**, 1587-1598.
- Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society B* **57**, 289-300.
- Bogdanowicz W, Piksa K, Tereba A (2012) Hybridization hotspots at bat swarming sites. *Plos One* **7**.
- Brereton R, Bennett S, Mansergh I (1995) Enhanced greenhouse climate-change and its potential effect on selected fauna of South-eastern Australia - a trend analysis. *Biological Conservation* **72**, 339-354.
- Brownstein MJ, Carpten JD, Smith JR (1996) Modulation of non-templated nucleotide addition by taq DNA polymerase: Primer modifications that facilitate genotyping. *Biotechniques* **20**, 1004-1006.
- Burgarella C, Lorenzo Z, Jabbour-Zahab R, *et al.* (2009) Detection of hybrids in nature: application to oaks (*Quercus suber* and *Q. ilex*). *Heredity* **102**, 442-452.
- Chapuis M-P, Estoup A (2007) Microsatellite null alleles and estimation of population differentiation. *Molecular Biology and Evolution* **24**, 621-631.
- Chatfield MWH, Kozak KH, Fitzpatrick BM, Tucker PK (2010) Patterns of differential introgression in a salamander hybrid zone: inferences from genetic data and ecological niche modelling. *Molecular Ecology* **19**, 4265-4282.
- Coyne JA, Orr HA (1989) Patterns of speciation in *Drosophila*. *Evolution* **43**, 362-381.
- Coyne JA, Orr HA (2004) *Speciation* Sinauer, Sunderland, Massachusetts.
- de Leon LF, Bermingham E, Podos J, Hendry AP (2010) Divergence with gene flow as facilitated by ecological differences: within-island variation in Darwin's finches. *Philosophical Transactions of the Royal Society B-Biological Sciences* **365**, 1041-1052.
- Díaz-Varela RA, Colombo R, Meroni M, *et al.* (2010) Spatio-temporal analysis of alpine ecotones: A spatial explicit model targeting altitudinal vegetation shifts. *Ecological Modelling* **221**, 621-633.

- 736 Dobzhansky (1937) *Genetics and the Origin of Species* Columbia University Press, New
737 York.
- 738 Dowling TE, Childs MR (1992) Impact of hybridization on a threatened trout of the
739 southwestern United States. *Conservation Biology* **6**, 355-364.
- 740 DSE (2013) Advisory List of Threatened Vertebrate Fauna in Victoria (ed. Environment
741 DoSa). Victorian Government Department of Sustainability and Environment,
742 Melbourne.
- 743 Earl DA, von Holdt BM (2012) STRUCTURE HARVESTER: a website and program for
744 visualizing STRUCTURE output and implementing the Evanno method.
745 *Conservation Genetics Resources* **4**, 359-361.
- 746 Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using
747 the softwards STRUCTURE: a simulation study. *Molecular Ecology* **14**, 2611-2620.
- 748 Excoffier L, Lischer HEL (2010) Arlequin suite ver 3.5: a new series of programs to perform
749 population genetics analyses under Linux and Windows. *Molecular Ecology*
750 *Resources* **10**, 564-567.
- 751 Fisch KM, Mahardja B, Burton RS, May B (2013) Hybridization between delta smelt and
752 two other species within the family Osmeridae in the San Francisco Bay-Delta.
753 *Conservation Genetics*, 1-6.
- 754 Gabirot M, Castilla AM, Lopez P, Martin J (2010b) Differences in chemical signals may
755 explain species recognition between an island lizard, *Podarcis atrata*, and related
756 mainland lizards, *P. hispanica*. *Biochemical systematics and ecology* **38**, 521-528.
- 757 Gardner MG, Fitch AJ, Bertozzi T, Lowe AJ (2011) Rise of the machines - recommendations
758 for ecologists when using next generation sequencing for microsatellite development.
759 *Molecular Ecology Resources* **11**, 1093-1101.
- 760 Garroway CJ, Bowman J, Cascaden TJ, *et al.* (2010) Climate change induced hybridization in
761 flying squirrels. *Global Change Biology* **16**, 113-121.
- 762 Goodman BA, Miles DB, Schwarzkopf L (2008) Life on the rocks: habitat use drives
763 morphological and performance evolution in lizards. *Ecology* **89**, 3462-3471.
- 764 Grabherr G, Gottfried M, Pauli H (1994) Climate effects on mountain plants. *Nature* **369**,
765 448-448.
- 766 Green K, Mansergh IM, Osborne WS (1992) The fauna of the Australian Alps: conservation
767 and management. *Review Geographie Alpine* **2 and 3**, 381-407.
- 768 Guindon S, Dufayard JF, Lefort V, *et al.* (2010) A simple, fast, and accurate algorithm to
769 estimate large phylogenies by maximum likelihood. *Systematic Biology* **59**, 307-321.

- Haines ML, Moussalli A, Stuart-Fox D, Clemann N, Melville J (2014) Phylogenetic evidence of historic mitochondrial introgression and cryptic diversity in the genus *Pseudemoia* (Squamata: Scincidae). *Molecular Phylogenetics and Evolution* **81**, 86-95.
- Harrison RG, Rand DM (1989) Mosaic hybrid zones and the nature of species boundaries. In: *Speciation and its Consequences* (eds. Otte D, Endler JA), pp. 111-133. Sinauer Associates, Inc., Massachusetts, USA.
- Hasselman DJ, Argo EE, McBride MC, *et al.* (2014) Human disturbance causes the formation of a hybrid swarm between two naturally sympatric fish species. *Molecular Ecology* **23**, 1137-1152.
- Hennessy K, Whetton P, Smith I, *et al.* (2003) The impact of climate change on snow conditions in mainland Australia. CSIRO Atmospheric Research, Aspendale, Victoria, Australia.
- Hewitt GM (2011) Quaternary phylogeography: the roots of hybrid zones. *Genetica* **139**, 617-638.
- Hutchinson MN, Donnellan SC (1992) Taxonomy and genetic variation in the Australian lizards of the genus *Pseudemoia* (Scincidae, Lygosominae). *Journal of Natural History* **26**, 215-264.
- IPCC (2013) Climate Change 2013: the Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change eds. Stocker TF, Qin D, Plattner G-K, *et al.*), p. 1535, Cambridge, U. K.
- Jacobsson M, Rosenberg NA (2007) CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics* **23**, 1801-1806.
- James EA, Brown GK, Citroen R, Rossetto M, Porter C (2011) Development of microsatellite loci in *Triglochin procera* (Juncaginaceae), a polyploid wetland plant. *Conservation Genetics Resources* **3**, 103-105.
- Jezkova T, Leal M, Rodriguez-Robles JA (2013) Genetic drift or natural selection? Hybridization and asymmetric mitochondrial introgression in two Caribbean lizards (*Anolis pulchellus* and *Anolis krugi*). *Journal of Evolutionary Biology* **26**, 1458-1471.
- Johnson JR, Johnson BB, Shaffer HB (2010) Genotype and temperature affect locomotor performance in a tiger salamander hybrid swarm. *Functional Ecology* **24**, 1073-1080.
- Kalinowski ST (2005) HP-Rare 1.0: a computer program for performing rarefaction on measures of allelic richness. *Molecular Ecology Notes* **5**, 187-189.

803 Kleindorfer S, O'Connor JA, Dudaniec RY, *et al.* (2014) Species collapse via hybridization in
804 Darwin's tree finches. *American Naturalist* **183**, 325-341.

805 Latch EK, Kierepka EM, Heffelfinger JR, Rhodes OE (2011) Hybrid swarm between
806 divergent lineages of mule deer (*Odocoileus hemionus*). *Molecular Ecology* **20**, 5265-
807 5279.

808 Leary RF, Allendorf FW, Forbes SH (1993) Conservation genetics of bull trout in the
809 Columbia and Klamath river drainages. *Conservation Biology* **7**, 856-865.

810 Liou LW, Price TD (1994) Speciation by reinforcement of premating isolation. *Evolution* **48**,
811 1451-1459.

812 Losos JB, Jackman T, Larson A, de Queiroz K, Rodriguez-Schettino L (1998) Contingency
813 and determinism in replicated adaptive radiations of island lizards. *Science* **279**, 2115-
814 2118.

815 Mallet J (2007) Hybrid speciation. *Nature* **446**, 279-283.

816 Margulies M, Egholm M, Altman WE, *et al.* (2005) Genome sequencing in microfabricated
817 high-density picolitre reactors. *Nature* **437**, 376-380.

818 Marino IAM, Benazzo A, Agostini C, *et al.* (2013) Evidence for past and present
819 hybridization in three Antarctic icefish species provides new perspectives on an
820 evolutionary radiation. *Molecular Ecology* **22**, 5148-5161.

821 Mayr E (1942) *Systematics and the Origin of Species* Columbia University Press, New York.

822 McDevitt AD, Mariani S, Hebblewhite M, *et al.* (2009) Survival in the Rockies of an
823 endangered hybrid swarm from diverged caribou (*Rangifer tarandus*) lineages.
824 *Molecular Ecology* **18**, 665-679.

825 McDonald DB, Parchman TL, Bower MR, Hubert WA, Rahel FJ (2008) An introduced and a
826 native vertebrate hybridize to form a genetic bridge to a second native species.
827 *Proceedings of the National Academy of Sciences* **105**, 10837-10842.

828 Meirmans PG (2006) Using the AMOVA framework to estimate a standardized genetic
829 differentiation measure. *Evolution* **60**, 2399-2402.

830 Melville J, Swain R (2003) Evolutionary correlations between escape behaviour and
831 performance ability in eight species of snow skinks (*Niveoscincus*: Lygosominae)
832 from Tasmania. *Journal of Zoology* **261**, 79-89.

833 Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for
834 inference of large phylogenetic trees. In: *Proceedings of the Gateway Computing*
835 *Environments Workshop (GCE)*, pp. 1-8, New Orleans, Louisiana.

836 Nice CC, Gompert Z, Fordyce JA, *et al.* (2013) Hybrid speciation and independent evolution
837 in lineages of alpine butterflies. *Evolution* **67**, 1055-1068.

838 Nielsen EE, Bach LA, Kotlicki P (2006) hybridlab (version 1.0): a program for generating
839 simulated hybrids from population samples. *Molecular Ecology Notes* **6**, 971-973.

840 Noor MAF (1999) Reinforcement and other consequences of sympatry. *Heredity* **83**, 503-
841 508.

842 Nylander JAA (2004) MrModeltest. Evolutionary Biology Centre, Uppsala University.
843 Available at: <http://www.abc.se/~nylander/>.

844 Olave M, Martinez LE, Avila LJ, Sites JW, Morando M (2011) Evidence of hybridization in
845 the Argentinean lizards *Liolaemus gracilis* and *Liolaemus bibronii* (Iguania:
846 Liolaemini): An integrative approach based on genes and morphology. *Molecular*
847 *Phylogenetics and Evolution* **61**, 381-391.

848 Orr HA (1995) The population genetics of speciation: the evolution of hybrid
849 incompatibilities. *Genetics* **139**, 1805-1813.

850 Parmesan C (2006) Ecological and evolutionary responses to recent climate change. *Annual*
851 *Review of Ecology, Evolution, and Systematics* **37**, 637-669.

852 Peakall R, Smouse PE (2012) GenAlEx 6.5: genetic analysis in Excel. Population genetic
853 software for teaching and research-an update. *Bioinformatics* **28**, 2537-2539.

854 Pickering CM (2007) Climate change and other threats in the Australian Alps. In: *Protected*
855 *Areas: Buffering Nature Against Climate Change*. (eds. Taylor M, Figgis P), pp. 28-
856 34. Proceedings of a WWF and IUCN World Commission on Protected Areas
857 Symposium, 18-19 June 2007, Canberra, Sydney.

858 Pinho C, Kaliontzopoulou A, Carretero MA, Harris DJ, Ferrand N (2009) Genetic admixture
859 between the Iberian endemic lizards *Podarcis bocagei* and *Podarcis carbonelli*:
860 evidence for limited natural hybridization and a bimodal hybrid zone. *Journal of*
861 *Zoological Systematics and Evolutionary Research* **47**, 368-377.

862 Pounds JA, Fogden MPL, Campbell JH (1999) Biological response to climate change on a
863 tropical mountain. *Nature* **398**, 611-615.

864 Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using
865 multilocus genotype data. *Genetics* **155**, 945-959.

866 Pritchard JK, Wen X, Falush D (2009) *Documentation for Structure Software Version*
867 2.3. <http://computing.bio.cam.ac.uk/local/doc/structure.pdf>.

868 Pritchard VL, Edmands S (2013) The genomic trajectory of hybrid swarms: outcomes of
869 repeated crosses between populations of *Tigriopus californicus*. *Evolution* **67**, 774-
870 791.

871 Quirt KC, Blouin-Demers G, Howes BJ, Loughheed SC (2006) Microhabitat selection of Five-
872 Lined Skinks in northern peripheral populations. *Journal of Herpetology* **40**, 335-342.

873 Rhymer JM, Simberloff D (1996) Extinction by hybridization and introgression. *Annual*
874 *Review of Ecology and Systematics* **27**, 83-109.

875 Rice WR (1989) Analyzing tables of statistical analyses. *Evolution* **43**, 223-225.

876 Robbins TR, Walker LE, Gorospe KD, *et al.* (2014) Rise and fall of a hybrid zone:
877 implications for the roles of aggression, mate choice, and secondary succession.
878 *Journal of Heredity* **105**, 226-236.

879 Ronquist F, Huelsenbeck JP (2003) MRBAYES 3: Bayesian phylogenetic inference under
880 mixed models. *Bioinformatics* **19**, 1572-1574.

881 Schuelke M (2000) An economic method for the fluorescent labeling of PCR fragments. *Nat*
882 *Biotech* **18**, 233-234.

883 Schulte U, Veith M, Hochkirch A (2012) Rapid genetic assimilation of native wall lizard
884 populations (*Podarcis muralis*) through extensive hybridization with introduced
885 lineages. *Molecular Ecology* **21**, 4313-4326.

886 Seehausen O, van Alphen JJM, Witte F (1997) Cichlid fish diversity threatened by
887 eutrophication that curbs sexual selection. *Science* **277**, 1808-1811.

888 Servedio MR, Noor MAF (2003) The role of reinforcement in speciation: Theory and data.
889 *Annual Review of Ecology Evolution and Systematics* **34**, 339-364.

890 Sinervo B, Mendez-de-la-Cruz F, Miles DB, *et al.* (2010) Erosion of lizard diversity by
891 climate change and altered thermal niches. *Science* **328**, 894-899.

892 Singhal S, Moritz C (2013) Reproductive isolation between phylogeographic lineages scales
893 with divergence. *Proceedings of the Royal Society B-Biological Sciences* **280**.

894 Stapley J, Hayes CM, Keogh JS (2003) Population genetic differentiation and multiple
895 paternity determined by novel microsatellite markers from the Mountain Log Skink
896 (*Pseudemoia entrecasteauxii*). *Molecular Ecology Notes* **3**, 291-293.

897 Tarroso P, Pereira RJ, Martínez-Freiría F, Godinho R, Brito JC (2014) Hybridization at an
898 ecotone: ecological and genetic barriers between three Iberian vipers. *Molecular*
899 *Ecology* **23**, 1108-1123.

- Taylor EB, Boughman JW, Groenenboom M, *et al.* (2006) Speciation in reverse: morphological and genetic evidence of the collapse of a three-spined stickleback (*Gasterosteus aculeatus*) species pair. *Molecular Ecology* **15**, 343-355.
- Team RDC (2013) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Teasdale LC, Stevens M, Stuart-Fox D (2013) Discrete colour polymorphism in the tawny dragon lizard (*Ctenophorus decresii*) and differences in signal conspicuousness among morphs. *Journal of Evolutionary Biology* **26**, 1035-1046.
- Tsy J, Lumaret R, Flaven-Noguier E, *et al.* (2013) Nuclear microsatellite variation in Malagasy baobabs (*Adansonia*, Bombacoideae, Malvaceae) reveals past hybridization and introgression. *Annals of Botany* **112**, 1759-1773.
- Turelli M, Barton NH, Coyne JA (2001) Theory and speciation. *Trends in Ecology and Evolution* **16**, 330-343.
- Untergrasser A, Cutcutache I, Koressaar T, *et al.* (2012) Primer3 - new capabilities and interfaces. *Nucleic Acids Research* **40**, e115.
- Vähä JP, Primmer CR (2006) Efficiency of model-based Bayesian methods for detecting hybrid individuals under different hybridization scenarios and with different numbers of loci. *Molecular Ecology* **15**, 63-72.
- Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P (2004) micro-checker: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* **4**, 535-538.
- While GM, Michaelides S, Heathcote RJ, *et al.* (2015) Sexual selection drives asymmetric introgression in wall lizards. *Ecology Letters* **18**, 1366-1375.
- Wilson S, Swan G (2013) *A Complete Guide to Reptiles of Australia*, Fourth edition edn. New Holland Publishers, Sydney.

Data Accessibility

- Morphological data, ecological data, and microsatellite genotypes are available at <http://datadryad.org>: doi 10.5061/dryad.kt70b
- Mitochondrial sequences have been deposited in GenBank
- Sampling locations are available in Supporting Information (Table S1).

Author Contributions

M.L.H. contributed to the study design, conducted field sampling, performed laboratory work, analysed the data, and drafted the manuscript. J.M. contributed to study design, data analysis, and manuscript preparation. J.S. provided assistance with laboratory work and data analysis. N.C. contributed to project conception, interpretation, logistic support, fieldwork and provided invaluable insight into the study system. D.G.C. contributed to project conception and molecular marker development. D.S-F. contributed to the study design, data analyses and manuscript preparation. All authors provided editorial input to the final manuscript.

Table 1. Genetic differentiation between population pairs for each site: *BHP* Bogong High Plains, *HT* Mt Higginbotham, *DHP* Dargo High Plains. F_{ST} (above diagonal) and Jost's D (below diagonal). All values are significant at $p < 0.01$.

		<i>P. cryodroma</i>	<i>P. entrecasteauxii</i>	<i>P. pagenstecheri</i>
<u>BHP</u>	<i>P. cryodroma</i>	--	0.060	0.084
	<i>P. entrecasteauxii</i>	0.417	--	0.080
	<i>P. pagenstecheri</i>	0.567	0.552	--
<u>HT</u>	<i>P. cryodroma</i>	--	0.059	0.109
	<i>P. entrecasteauxii</i>	0.412	--	0.057
	<i>P. pagenstecheri</i>	0.653	0.374	--
<u>DHP</u>	<i>P. entrecasteauxii</i>	--	--	0.108
	<i>P. pagenstecheri</i>	--	0.706	--

Table 2. Pooled within class canonical coefficients for canonical variables 1 and 2 for males and females from discriminant analysis of morphological variables. Morphometric variables most strongly correlated with canonical variables are in bold. Refer to text for explanations of morphological variables.

Morphometric	<u>Males</u>		<u>Females</u>	
Variable	Can 1	Can 2	Can 1	Can 2

SVL*	0.020	0.061	0.016	-0.374
HL	0.457	0.014	0.410	0.175
HWE	0.215	-0.192	0.395	0.058
HWJ	0.346	-0.086	0.192	0.134
HD	0.167	-0.155	0.152	-0.159
ILL	-0.118	0.278	-0.137	0.487
PW	-0.252	0.149	-0.193	0.210
UFL	0.401	0.359	0.358	0.454
LFL	0.473	0.404	0.668	0.471
FFL	0.184	0.481	0.453	0.307
UHLL	0.448	0.288	0.417	0.317
LHLL	0.287	0.042	0.436	-0.016
HFL	0.237	-0.118	0.429	0.183
F value	5.39	4.19	2.47	1.13
p value	< 0.001	< 0.001	< 0.001	0.327
Eigenvalue	1.33	0.96	1.38	0.35
Proportion of Variance	0.526	0.378	0.755	0.190

* With the exception of SVL, morphometric variables refer to residuals calculated by regressing measurements against SVL.

Table 3. Pooled within class canonical structure for canonical variables 1 and 2 for BHP, HT, and DHP from discriminant analysis of habitat variables. Habitat variables strongly correlated with canonical variables are in bold. Site abbreviations: *BHP* Bogong High Plains, *HT* Mt Higginbotham, *DHP* Dargo High Plains.

Habitat Variable	<u>BHP</u>		<u>HT</u>		<u>DHP</u>
	Can 1	Can 2	Can 1	Can 2	Can 1
Litter depth	-0.642	0.226	0.785	0.042	0.339
Ground cover	-0.185	0.425	0.437	0.212	0.965
Shrub cover	0.452	-0.294	-0.082	0.059	-0.419
Tree cover	-0.081	0.378	-0.664	0.525	-0.128
Rock cover	-0.057	-0.796	0.083	-0.727	-0.647

F value	2.31	1.80	4.92	3.06	5.41
p value	0.004	0.079	< 0.001	0.004	0.002
Eigenvalue	0.228	0.128	0.933	0.430	0.569
Proportion of variance	0.589	0.331	0.678	0.312	1

Figure Captions

Fig. 1. Elevation map depicting sampling sites within the Victorian Alps in south-eastern Australia. Site names are abbreviated: *BHP* Bogong High Plains, *DHP* Dargo High Plains, *HT* Mt Higginbotham.

Fig. 2. Number of individuals classified as each species or hybrid-cross based on individual proportion of membership (*Q*) estimated in STRUCTURE at A) Bogong High Plains, B) Mt Higginbotham, and C) Dargo High Plains. Classification is based on the following criteria: pure individuals had $Q \geq 0.90$ for one species, hybrid-crosses had $0.90 > Q \geq 0.10$ for two species, and unclassified hybrids had either $0.90 > Q \geq 0.10$ for all three species or $0.90 > Q \geq 0.10$ for one species and $Q > 0.10$ for the other two species.

Fig. 3. Hybrid individuals identified from *BHP* Bogong High Plains, *HT* Mt Higginbotham, and *DHP* Dargo High Plains in: A) STRUCTURE and B) NewHybrids. Individuals are represented by a single vertical line, with the percentage of each colour representing the individual proportion of membership (*Q*) for each of species: *Pseudemoia cryodroma* (yellow), *P. entrecasteauxii* (red), and *P. pagenstecheri* (blue). In B) three additional categories are present: F1 hybrid (white), F2 hybrid (black) and F1 backcross (white and black stripes). Within each site, individuals are grouped by mitochondrial lineage (P.c. = *P. cryodroma*, P.e. = *P. entrecasteauxii*, P.p. = *P. pagenstecheri*). * Individual classified as a hybrid cross for two different sets of species pairs. The data presented was averaged over both analyses.

Fig. 4. Discriminant analyses based on the morphological dataset for A) males and B) females. Individuals are represented by the following symbols: yellow circles, *P. cryodroma*;

red squares, *P. entrecasteauxii*; black asterisks, hybrids; and blue triangles, *P. pagenstecheri*. Ellipses represent 95% confidence levels.

Fig. 5. Discriminant analyses based on the microhabitat dataset for each site. In A) Bogong High Plains and B) Mt Higginbotham, individuals are represented by the following symbols: yellow circles, *P. cryodroma*; red squares, *P. entrecasteauxii*; black asterisks, hybrids; and blue triangles, *P. pagenstecheri*. Ellipses represent 95% confidence levels. In C) Dargo High Plains, individuals are represented by horizontal bars: *P. entrecasteauxii* (red) and *P. pagenstecheri* (blue).

A) BHP

B) HT

C) DHP

P. cryodroma

P. entrecasteauxii

P. pagenstecheri

P. cryodroma x *P. entrecasteauxii*

P. entrecasteauxii x *P. pagenstecheri*

P. cryodroma x *P. pagenstecheri*

Unclassified hybrid

A) Males

B) Females

A) Bogong High Plains

B) Mt Higginbotham

C) Dargo High Plains

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Haines, ML;Melville, J;Sumner, J;Clemann, N;Chapple, DG;Stuart-Fox, D

Title:

Geographic variation in hybridization and ecological differentiation between three syntopic, morphologically similar species of montane lizards

Date:

2016-06-01

Citation:

Haines, M. L., Melville, J., Sumner, J., Clemann, N., Chapple, D. G. & Stuart-Fox, D. (2016). Geographic variation in hybridization and ecological differentiation between three syntopic, morphologically similar species of montane lizards. *MOLECULAR ECOLOGY*, 25 (12), pp.2887-2903. <https://doi.org/10.1111/mec.13652>.

Persistent Link:

<http://hdl.handle.net/11343/291327>