Pharmacological Lactation Suppression with D₂ Receptor Agonists and Risk of Postpartum Psychosis: A Systematic Review.

SNELLEN Martien, POWER Josephine, BLANKLEY Gaynor, GALBALLY Megan.

Dr SNELLEN Martien: Consultant Psychiatrist, Perinatal Mental Health, Mercy Hospital for Women, 163 Studley Road, Heidelberg VIC 3084
msnellen@iprimus.com.au (Phone 03 8458 4834)

Dr POWER Josephine: Consultant Psychiatrist, Perinatal Mental Health, Mercy Hospital for Women, 163 Studley Road, Heidelberg VIC 3084
JPower@mercy.com.au (Phone 03 8458 4834)

Dr BLANKLEY Gaynor: Consultant Psychiatrist and Head of Unit, Perinatal Mental Health, Mercy Hospital for Women, 163 Studley Road, Heidelberg VIC 3084
GBlankley@mercy.com.au (Phone 03 8458 4834)

Prof. GALBALLY Megan: Foundation Chair in Perinatal Psychiatry
School of Psychology and Exercise Science & School of Medicine, University of Notre Dame
Consultant Psychiatrist, Fiona Stanley Hospital
m.galbally@murdoch.edu.au (Phone 08 9360 2844)
Murdoch University, 90 South Street, MURDOCH WA 6150

Short Title: Lactation Suppression and Postpartum Psychosis

Word Count Main Text and Abstract: 2035

The Authors report no conflict of interest. No financial assistance was provided to assist with the article's creation.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/AJO.12479](https://doi.org/10.1111/AJO.12479)

This article is protected by copyright. All rights reserved

Received Date : 11-Mar-2016

Revised Date : 08-Apr-2016

Accepted Date : 19-Apr-2016

Article type : Review

Corresponding author mail id : msnellen@iprimus.com.au

Pharmacological Lactation Suppression with D₂ Receptor Agonists and Risk of Postpartum Psychosis: A Systematic Review.

Short Title: Lactation Suppression and Postpartum Psychosis

Word Count Main Text and Abstract: 244 and 2007

Tables: One

Keywords: lactation suppression, ab lactation, D₂ receptor agonists, postpartum, psychosis.

Abstract

Background: It has been suggested that D₂ receptor agonists commonly used postpartum for the physiological suppression of lactation, such as bromocriptine and cabergoline, may increase the risk of illness onset or relapse in women where there is a predisposition for, or history of schizophrenia, bipolar disorder or postpartum psychosis. This is based on two lines of reasoning: current models of psychosis assume episodes are triggered by dysregulation of brain dopaminergic activity and treated by medications that universally have D₂ receptor antagonist properties; and limited research suggesting these agents may be associated with psychotic episodes in vulnerable individuals outside of the postpartum period.

Aim: To examine whether D₂ agonists trigger psychosis in previously well mothers, or psychotic relapse or exacerbation of symptoms in mothers with known psychotic illnesses, when used to suppress lactation during the early postpartum period.

Materials and Methods: A systematic review of the literature was undertaken of electronic databases including: MEDLINE, EMBASE and PsychINFO from 1950 to 2015 using the keywords.

Results: 8 case reports, 3 case series and a pharmacovigilance survey were identified.

Conclusion: Whilst D₂ receptor agonists appear to increase the risk of triggering psychosis in previously well mothers and those previously diagnosed with schizophrenia, bipolar disorder and postpartum psychosis bromocriptine appears to pose a much greater risk than cabergoline. When considering the use of pharmacological agents to suppress lactation physicians should carefully screen patients for a history of psychosis and consider alternatives in order to moderate this risk.

Introduction

The World Health Organization gives a clear recommendation that breastfeeding is the preferred option for infant nutrition and should be supported exclusively for 6 months and partially up until 2 years of age. However, only 38% of infants are exclusively breastfed globally from 0 to 6 months of age ¹. There are a range of individual and societal reasons and rates vary across nations. However, for women with schizophrenia, bipolar disorder or a history of postpartum psychosis the choice to breastfeed is more complex as it is influenced by both the illness and its treatment. For instance, there is evidence that suggests that the sleep deprivation associated with breastfeeding during the early postpartum may increase the risk of psychotic relapse for women with bipolar disorder or a history of postpartum psychosis, and pharmacological agents such as lithium and clozapine are considered potentially incompatible with breastfeeding. In addition, there are women with serious mental illnesses who require lactation suppression following neonatal loss or child removal by protective services.

In order to prevent breast engorgement, leakage of milk, discomfort and pain, a range of non-pharmacological and pharmacological options exist to aid lactation suppression. Non-pharmacological means include: tight brassieres or binders, minimizing nipple stimulation, ice packs and cabbage leaves. Pharmacological options mainly involve ergot derivative medications that are dopaminergic D₂ receptor agonists such as bromocriptine and cabergoline. These medications have established indications for both pathological and physiological hyperprolactinaemia. Pharmacological agents prescribed

in this context are used not to treat pathology, but to suppress an undesired normal physiological reaction. In this context pharmacological suppression of lactogenesis should generally be considered optional rather than a medical necessity, and a number of concerns have been raised regarding the practice.

The first Cochrane systematic review assessing the effectiveness of all forms of treatments for lactation suppression found the evidence was based on small trials of low methodological quality ². They conclude that there is weak evidence for pharmacological or non-pharmacological lactation suppression in the first week postpartum.

Additionally, debate regarding the safety of bromocriptine in particular has existed since the 1980s when concerns were raised about cardiovascular and neurological risks. The manufacturers of bromocriptine withdrew its use for lactation suppression in the US in 1994 following a series of adverse events including 9 deaths. The FDA has since recommended that medication not be used to suppress lactation, however the practice continues internationally.

Contemporary pathophysiological models of severe psychotic forms of mental illness (such as schizophrenia and bipolar disorder) assume that they are triggered by a dysregulation of dopaminergic activity in the brain arising from the adverse interaction of predisposing risk genes and environmental factors ³. What all successful pharmacological treatments of psychosis have in common is an ability to act as dopamine D₂ receptor antagonists. This naturally raises the question as to whether patients with psychosis, or at risk of psychosis, are susceptible to onset or exacerbation of psychosis when they are prescribed D₂ agonists.

The early postnatal period represents a high-risk period for psychotic illness onset or relapse. Women are approximately 22 times more likely to experience the onset of a manic or psychotic episode in the first postpartum month than at any other time in their lives and the prevalence of postpartum psychosis in the general population is estimated as 1-2 per 1,000 deliveries ⁴. A Danish population-based registry study found a relative risk of 5 for hospital admissions in women with a diagnosis of schizophrenia within the first postpartum month ⁵. Women with a diagnosed bipolar disorder are at extremely

high risk of affective psychosis in the early postnatal period ^{6 7} and experience a nearly sevenfold higher risk of admission for a first episode and a nearly twofold higher risk for a recurrent episode when compared with non-postpartum and non-pregnant women ⁸. The most important predictors for postpartum psychosis are a personal or family history of bipolar disorder or previous episode of postpartum psychosis. However, whilst women with bipolar disorder or schizophrenia are recognized to have an elevated risk of relapse during the early postnatal period, a significant portion of women who experience postpartum psychosis have no previous history of prior episodes ⁸. The underlying pathophysiological mechanisms are poorly understood, however alterations in oestrogen levels and inflammatory processes, and fluctuations in biorhythms and functioning are currently under investigation.

Given that psychosis onset or relapse has been associated with dysregulation of brain dopaminergic activity and this risk appears greater during the early postpartum period it needs to be considered as to whether lactation suppression with D₂ agonists increases this risk.

First we need to examine whether there is any evidence that D₂ agonists precipitate psychosis onset or relapse when used to treat pathological processes. The primary indications for the use of ergot derivatives such as bromocriptine and cabergoline are for pathological hyperprolactinaemia (i.e. prolactinomas). They have also been used in Acromegaly, Parkinson's Disease, Restless Leg Syndrome, and amenorrhea and galactorrhoea secondary to neuroleptic use ¹⁰. The main difference between the two agents is their duration of action and degree of dopamine receptor selectivity. When used to suppress physiological postpartum lactation bromocriptine is traditionally given as 2.5mg twice daily for 14 days and cabergoline (which has greater D₂ selectivity and longer half-life) as a single dose of 1mg. When used to treat pathological hyperprolactinaemia the doses tend to be higher and used for a prolonged period of time and in a comparison study the latter medication has been shown to have a lower incidence of adverse effects overall ¹¹.

In a study of 600 patients treated with the D₂ agonists bromocriptine and lisuride for functioning pituitary tumours at lower dosage eight developed medication-induced psychosis ¹². What is notable is that all of the eight had either a pre-existing history of

psychosis or displayed considerable changes in behaviour or mood prior to initiation of therapy. There have also been numerous case reports of patients treated with ergot derivatives developing a psychotic illness as a direct consequence of exposure to these agents when treated for pathological hyperprolactinaemia and other non-postpartum indications^{13 14 15}. It is unclear if these cases represent cases of ergotism (a medication side effect)¹⁶, or whether the medication uncovered an underlying predisposition to psychosis. Further case reports suggest that these agents may trigger relapse or symptom exacerbation in patients with an established psychotic illness^{17 18} or trigger a first episode of psychosis in predisposed individuals^{19 20 21}. Once again, bromocriptine appears to be implicated more than cabergoline. The main problem with such case reports is that they are anecdotal and do not report on long-term outcome.

There has been some interest in the use of dopamine agonists in the treatment of neuroleptic induced hyperprolactinaemia²². An 8-week randomized, single-blind, placebo-controlled, multicentre study of 60 treated women with schizophrenia found that none experienced an exacerbation of their symptoms when bromocriptine was introduced in order to treat neuroleptic-induced hyperprolactinaemia²³. Similarly, a pilot study of 19 patients with schizophrenia given cabergoline over a 6-week period to treat neuroleptic-induced hyperprolactinaemia found none had a worsening of their psychopathology²⁴. This finding has been supported by other case and case series studies^{25 26 27}. There appears to be only one report in the literature of cabergoline induced psychotic exacerbation in two schizophrenic patients when used to treat neuroleptic induced hyperprolactinaemia²⁸.

Overall, the risk of D₂ agonists precipitating the onset or relapse of psychosis appears to be related to the drug, the indication, the dose and the individual.

Aims

This systematic review aims to examine whether D₂ agonists trigger psychosis in previously well mothers, or psychotic relapse or exacerbation of symptoms in mothers with known psychotic illnesses, when used to suppress lactation during the early postpartum period.

Methods

A systematic review was undertaken of electronic databases including: MEDLINE, EMBASE and PsycINFO which were searched for articles published between 1950 and December 2015 using the terms “bromocriptine”, “cabergoline”, “ab lactation”, “lactation suppression” and “psychosis”. Studies were eligible if they were original case reports, case series or cohort studies. In addition, manual searches of the references of included articles and review of the bibliographies of reference books and review articles was undertaken. Studies in languages other than English were translated using an online program prior to analysis and inclusion in the review. There were no exclusion criteria.

Results

The systematic review of the literature identified 8 case reports, 3 case series, and a pharmacovigilance survey^{29 30 31 32 33 34 35 36 37 38 39 40}. Most reports were of women experiencing a postpartum psychosis following prescription of bromocriptine for lactation suppression during the early postpartum period. These women had either been previously well or experienced a relapse or exacerbation of an underlying mood disorder or schizophrenia. Two studies reported that women with schizophrenia experienced no deterioration in psychotic symptoms following administration of bromocriptine in the early postpartum period. A French pharmacovigilance study for the period 1994 to 2010 reported nine cases of acute psychiatric illness potentially associated with lactation suppression using bromocriptine. Four cases were reported as puerperal psychosis, two as deterioration of schizophrenia, two as confusion and one as mania. There has only been one report of cabergoline triggering an episode of mania in a patient with a pre-existing psychotic illness and none in well women. Results are summarised in Table 1.

Conclusion and Clinical Recommendations

Whilst both bromocriptine and cabergoline have been associated with triggering postpartum psychotic illness onset, relapse or exacerbation it has been proposed that the latter medication confers a lower risk.⁴¹ This view is supported by this systematic review of the literature. The apparent lower incidence of cabergoline induced psychosis, when compared with bromocriptine, may reflect the benefits of its greater receptor specificity, lower overall exposure given the single dose regimen, or the fact that as a newer agent it has been subjected to a briefer time of post-release surveillance.

The literature identified within this systematic review has a number of limitations. In particular, there is a relatively low level of evidence as the bulk of research consists of case reports or case series. The sole randomized control trial was performed outside of the setting of parturition and lactation, and therefore may not be applicable to the perinatal context. There is insufficient evidence available to allow clinicians to quantify individual risk based upon the severity of a woman's pre-existing illness, the antipsychotic agent she is already prescribed or its dose at the time of treatment. It would be important for future studies to examine these issues.

Despite the absence of prospective double blind randomised trials there is enough evidence from these identified studies to suggest that caution be applied when prescribing lactation suppressants to women who have a personal or family history of psychotic illness, and particularly if the illness has previously had a postpartum onset or exacerbation. Cabergoline, with its greater dopaminergic selectivity appears to offer a lower risk of precipitating psychosis than bromocriptine and therefore would be the preferred agent if pharmacological suppression is indicated for an individual woman.

In conclusion, when considering the use of pharmacological agents to suppress lactation physicians should carefully screen patients for a personal as well as family history of schizophrenia, bipolar disorder and postpartum psychosis. Given the potential risk of psychotic relapse these agents may confer it is recommended that pharmacological lactation suppressants be avoided when there is a personal or family history of bipolar disorder or postpartum psychosis and used only with caution in patients with treated schizophrenia. In this context it is also important to note the overall limited evidence for their efficacy to suppress physiological lactation and the availability of viable non-pharmacological treatments. When a risk: benefit analysis is being performed it should also be considered that lactation suppressants used in the context of physiological lactation are not being used to treat pathology, but are used to assist with postpartum breast comfort and thus may be considered to be optional rather than essential. If a decision is made to prescribe lactation suppressants, then close monitoring of the patient's mental state is recommended.

References

- ¹ World Health Organization. World Health Statistics 2013 (www.who.int/gho/publications/world_health_statistics/2013/en/, accessed 10 January 2016).
- ² Oladapo OT & Fawole B. Treatments for suppression of lactation. Cochrane Database Syst Rev. 2009 Jan 21;(1):CD005937
- ³ Tost H, Alam T, Meyer-Lindenberg A. Dopamine and psychosis: theory, pathomechanisms and intermediate phenotypes. *Neurosci Biobehav Rev*. 2010 Apr;34(5):689-700.
- ⁴ Bergink V & Kushner SA. Postpartum Psychosis. In: *Psychopharmacology and Pregnancy: Treatment Efficacy, Risks and Guidelines*. Eds. Galbally M, Snellen M & Lewis A. Springer 2014. p.p.139-150.
- ⁵ Munk-Olsen T, Laursen TM, Pedersen CB, Mors O et al. New parents and mental disorders: a population-based register study. *JAMA*. 2006;296(21):2582-9.
- ⁶ Viguera AC, Tondo L, Koukopoulos AE, Reginaldi D et al. Episodes of mood disorders in 2,252 pregnancies and postpartum periods. *Am J Psychiatry*. 2011 Nov;168(11):1179-85.
- ⁷ Snellen M & Malhi GS. Bipolar Disorder, Psychopharmacology and Pregnancy. In: *Psychopharmacology and Pregnancy: Treatment Efficacy, Risks and Guidelines*. Eds. Galbally M, Snellen M & Lewis A. Springer 2014. p.p.103-118.
- ⁸ Terp IM, Mortensen PB. Post-partum psychoses. Clinical diagnoses and relative risk of admission after parturition. *Br J Psychiatry*. 1998 Jun;172:521-6.
- ⁹ Boyce P, Barriball E. Puerperal psychosis. *Arch Womens Ment Health*. 2010 Feb;13(1):45-7.

- ¹⁰ Rains CP, Bryson HM, Fitton A. Cabergoline. A review of its pharmacological properties and therapeutic potential in the treatment of hyperprolactinaemia and inhibition of lactation. *Drugs*. 1995 Feb;49(2):255-79.
- ¹¹ Webster J, Piscitelli G, Polli A, Ferrari CI et al. A comparison of cabergoline and bromocriptine in the treatment of hyperprolactinemic amenorrhea. Cabergoline Comparative Study Group. *N Engl J Med*. 1994 Oct 6;331(14):904-9.
- ¹² Turner TH, Cookson JC, Wass JA, Drury PL et al. Psychotic reactions during treatment of pituitary tumours with dopamine agonists. *Br Med J (Clin Res Ed)*. 1984 Oct 27;289(6452):1101-3.
- ¹³ Peter SA, Autz A, Jean-Simon ML. Bromocriptine-induced schizophrenia. *J Natl Med Assoc*. 1993 Sep;85(9):700-1.
- ¹⁴ Harris YT, Harris AZ, Deasis JM, Ferrando SJ et al. Cabergoline associated with first episode mania. *Psychosomatics*. 2012 Nov-Dec;53(6):595-600.
- ¹⁵ Jimeno Valdés A, Ruiz Otazo AM, Jimeno Bulnes N. A case of toxic psychosis caused by bromocriptine. *Actas Luso Esp Neurol Psiquiatr Cienc Afines*. 1989 Nov-Dec;17(6):386-90.
- ¹⁶ Iffy L, Lindenthal JJ, McArdle JJ, McNamara RE et al. Ergotism: a possible etiology for puerperal psychosis. *Obstet Gynecol*. 1989 Mar;73(3 Pt 2):475-7.
- ¹⁷ Dorevitch A, Aronzon R, Stark M. Psychotic exacerbation attributed to low-dose bromocriptine treatment of galactorrhea and hyperprolactinemia. *Acta Obstet Gynecol Scand*. 1991;70(4-5):375-6.
- ¹⁸ Johnson JM. Treated mania exacerbated by bromocriptine. *Am J Psychiatry*. 1981 Jul;138(7):980-2.

- ¹⁹ Seeman MV. Transient psychosis in women on clomiphene, bromocriptine, domperidone and related endocrine drugs. *Gynecol Endocrinol*. 2015 Oct;31(10):751-4
- ²⁰ Bilal L, Ching C. Cabergoline-induced psychosis in a patient with undiagnosed depression. *J Neuropsychiatry Clin Neurosci*. 2012 Fall;24(4):E54.
- ²¹ Wang F, Zheng S, Zhang Y. Serious Bromocriptine-induced Psychosis: Suicide in the Treatment of Prolactinoma. *Neurosurgery Quarterly*. November 2013 Nov: 23(4):249–251
- ²² Tollin SR. Use of the dopamine agonists bromocriptine and cabergoline in the management of risperidone-induced hyperprolactinemia in patients with psychotic disorders. *J Endocrinol Invest*. 2000 Dec;23(11):765-70.
- ²³ Lee MS, Song HC, An H, Yang J et al. Effect of bromocriptine on antipsychotic drug-induced hyperprolactinemia: eight-week randomized, single-blind, placebo-controlled, multicenter study. *Psychiatry Clin Neurosci*. 2010 Feb;64(1):19-27.
- ²⁴ Cavallaro R, Cocchi F, Angelone SM, Lattuada E et al. Cabergoline treatment of risperidone-induced hyperprolactinemia: a pilot study. *J Clin Psychiatry*. 2004 Feb;65(2):187-90.
- ²⁵ Pollice R, Di Giovambattista E, Tomassini A, Di Pucchio A et al. Risperidone-induced symptomatic hyperprolactinemia in youth with schizophrenia: efficacy and tolerability of cabergoline treatment. *Clin Ter*. 2007 Mar-Apr;158(2):121-6.
- ²⁶ Cohen LG, Biederman J. Treatment of risperidone-induced hyperprolactinemia with a dopamine agonist in children. *J Child Adolesc Psychopharmacol*. 2001 Winter;11(4):435-40.
- ²⁷ Coronas R, Cobo J, Giménez-Palop O, Ortega E et al. Safety of cabergoline in the management of pituitary prolactin-induced symptoms with patients treated with atypical neuroleptics. *Curr Drug Saf*. 2012 Apr;7(2):92-8.

- ²⁸ Chang SC, Chen CH, Lu ML. Cabergoline-induced psychotic exacerbation in schizophrenic patients. *Gen Hosp Psychiatry*. 2008 Jul-Aug;30(4):378-80.
- ²⁹ Kemperman CJ, Zwanikken GJ. Psychiatric side effects of bromocriptine therapy for postpartum galactorrhoea. *J R Soc Med*. 1987 Jun; 80(6): 387–388.
- ³⁰ Lake CR, Reid A, Martin C, Chernow B. Cyclothymic disorder and bromocriptine: predisposing factors for postpartum mania? *Can J Psychiatry*. 1987 Nov;32(8):693-4.
- ³¹ Durst R, Dorevitch A, Ghinea C, Ginath Y. Bromocriptine-associated postpartum psychotic exacerbation. *Harefuah*. 1990 Feb 15;118(4):203-4.
- ³² Olbrich HM, Martin P. Drug treatment of schizophrenic psychoses in puerperium. *Nervenarzt*. 1994 Jul;65(7):482-5.
- ³³ Reeves RR, Pinkofsky HB. Postpartum psychosis induced by bromocriptine and pseudoephedrine. *J Fam Pract*. 1997 Aug;45(2):164-6.
- ³⁴ Canterbury RJ, Haskins B, Kahn N, Saathoff G et al. Postpartum psychosis induced by bromocriptine. *Actas Luso Esp Neurol Psiquiatr Cienc Afines*. 1989 Nov-Dec;17(6):386-90.
- ³⁵ Iffy L, Lindenthal J, Szodi Z, Griffin W. Puerperal psychosis following ablation with bromocriptine. *Med Law*. 1989;8(2):171-4.
- ³⁶ Pinardo Zabala A, Alberca Muñoz ML, Giménez García JM. Postpartum psychosis associated with bromocriptine. *An Med Interna*. 2003 Jan;20(1):50-1.
- ³⁷ Misdrahi D, Chalard R, Verdoux H. Postpartum mania induced by Bromocriptine: a case report. *J Gynecol Obstet Biol Reprod (Paris)*. 2006 Feb;35(1):79-81.
- ³⁸ Gahr M, Kolle A, Freudenmann RW. Disorganized schizophrenia does not deteriorate with dopamine agonists cabergoline and bromocriptine used for ablactation. *Pharmacopsychiatry* 2011; 44:74.

³⁹ Salinas Botrán A, Urgelés Puértolas D, Peregrin Abad E. Postpartum psychosis associated with cabergoline. *Med Clin (Barc)*. 2013 Jan 5;140(1):45-6

⁴⁰ Bernard N, Janzen H, Becker M, Pecriaux C et al. Severe adverse effects of bromocriptine in lactation inhibition: a pharmacovigilance survey. *BJOG*. 2015 Aug;122(9):1244-51.

⁴¹ Baeza Pertegaz I, Goikolea Alberdi JM, Parellada Rodón E. Is cabergoline a better drug to inhibit lactation in patients with psychotic symptoms? *J Psychiatry Neurosci*. 2002 Jan;27(1):54.

Study	Design	Medication and number of subjects	Findings
Kemperman 1987 ²⁹	Case report	Bromocriptine n = 1	Postpartum mania occurring in a woman with no pre-existing psychiatric illness
Lake 1987 ³⁰	Case report	Bromocriptine n = 1	Postpartum mania in a woman with a premorbid diagnosis of cyclothymia
Durst 1990 ³¹	Case series	Bromocriptine n = 2	Deterioration in mental state in 2 women with pre-existing psychosis following bromocriptine administration for ab lactation
Olbrich 1994 ³²	Retrospective case series	Bromocriptine n = 4	No postpartum worsening of psychosis in 4 women with a diagnosis of schizophrenia
Reeves 1997 ³³	Case report	Bromocriptine n = 1	Postpartum psychosis in a woman with no pre-existing psychiatric history
Canterbury 1989 ³⁴	Retrospective case series	Bromocriptine n = 2	Postpartum psychosis in a woman with no prior psychiatric history
Iffy 1989 ³⁵	Case report	Bromocriptine n = 1	Postpartum psychosis in a woman with no premorbid psychiatric history
Pinardo Zabala 2003 ³⁶	Case report	Bromocriptine n = 1	Postpartum psychosis in a woman with no known psychiatric history
Misdrahi 2006 ³⁷	Case report	Bromocriptine n = 1	Postpartum psychosis in a woman with a diagnosis of mood disorder
Gahr 2011 ³⁸	Case report	Cabergoline n= 1 Bromocriptine n=1	A woman with schizophrenia treated serially with cabergoline then bromocriptine with no worsening of psychotic symptoms

Salinas Botran 2013 ³⁹	Case report	Cabergoline n= 1	Precipitation of postpartum mania in a woman with a history of psychosis
Bernard 2015 ⁴⁰	Pharmacovigilance Survey	105 Severe adverse events reported with Bromocriptine	Among women prescribed bromocriptine for ablactation there were reports of 9 psychiatric adverse events: 2 women with acute deterioration of schizophrenia, 4 with postpartum psychosis, 2 with confusion and 1 with mania. There were also 5 reports of “other nervous system disorders” without clarification

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Snellen, M;Power, J;Blankley, G;Galbally, M

Title:

Pharmacological lactation suppression with D-2 receptor agonists and risk of postpartum psychosis: A systematic review

Date:

2016-08-01

Citation:

Snellen, M., Power, J., Blankley, G. & Galbally, M. (2016). Pharmacological lactation suppression with D-2 receptor agonists and risk of postpartum psychosis: A systematic review. AUSTRALIAN & NEW ZEALAND JOURNAL OF OBSTETRICS & GYNAECOLOGY, 56 (4), pp.336-340. <https://doi.org/10.1111/ajo.12479>.

Persistent Link:

<http://hdl.handle.net/11343/291412>