Received Date : 01-Jul-2016

Accepted Date : 01-Jul-2016

Article type : Invited Concise Review

Title: Refugee child oral health

Authors

Dr Elisha Riggs

Research Fellow, Healthy Mothers Healthy Families

Murdoch Childrens Research Institute

The Royal Children's Hospital

Flemington Rd Parkville, Victoria 3052 AUS

Email: elisha.riggs@mcri.edu.au

Dr Sadna Rajan

Senior Lecturer

Melbourne Dental School

The University of Melbourne

Email: rsadna@gmail.com

Ms Sue Casey

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/odi.12530](https://doi.org/10.1111/odi.12530)

This article is protected by copyright. All rights reserved

Manager, Sector Development & Partnerships

Foundation House, The Victorian Foundation for Survivors of Torture Inc

Email: caseys@foundationhouse.org.au

Associate Professor Nicky Kilpatrick.

Director, Cleft Services Royal Children's Hospital

Senior Research Fellow, Murdoch Childrens Research Institute

Department of Paediatrics, University of Melbourne,

Email: nicky@bassdata.com.au

Running title: Refugee child oral health

Keywords: refugee, oral health, children

Author Manuscript

Introduction

Globally, the movement of people between countries is increasing. Not only does this include economic or skilled migrants moving for employment and education reasons but also a rapidly growing refugee and asylum seeking population. As of 2015, a staggering 60 million people were forcibly displaced from their homes due to persecution, conflict, violence and human rights abuse. Of these, children and adolescents below the age of 18 years comprise over half of the world's refugees. Whilst many people remain displaced and languish in appalling circumstances, each year approximately 60,000 children are resettled in high-income countries (United Nations High Commissioner for Refugees, 2015).

Children from refugee backgrounds are likely to have poor physical, mental and social health outcomes, and their parents commonly experience barriers accessing appropriate and meaningful health information and services (Woodland et al., 2010, Davidson et al., 2004). In particular, these children are at increased risk of poor oral health due to the adversity they have experienced early in life, which may impact on their long-term health, development and wellbeing (Shonkoff et al., 2012). Health services, including providers of dental health services, need to recognise, understand and be responsive to the issues emerging with this changing and growing demographic in order to minimise poor oral health outcomes.

This review summarises the evidence surrounding the oral health (dental status, oral hygiene practices, and access to care) of children of refugee backgrounds living in developed countries. Based upon this evidence we will discuss potential strategies available to oral health service providers in high-income refugee-receiving countries to optimise provision of responsive dental care to this growing population.

Background

People from refugee backgrounds may arrive in their new country in a variety of ways, through formal migration programs, including humanitarian resettlement programs or seek asylum on or soon after arrival. Many experience protracted periods in refugee camps or living precariously in urban settings, time spent in detention centres, and multiple displacements seeking safety. Whatever the pathway, the refugee experience is characterised by displacement, conflict, human rights violations, persecution, family separation, prolonged times in transit with limited or no access to services or basic necessities (shelter, food, water and safety) and often includes torture as well as physical and sexual violence (Victorian Foundation for the Survivors of Torture, 1998). Refugee populations have an increased risk of health problems related to their traumatic and stressful

experiences which can be exacerbated during settlement by separation, loss of family members and poverty (Victorian Foundation for the Survivors of Torture, 1998).

Dental caries is one of the most common chronic diseases of childhood (Selwitz et al., 2007) affecting between 60-90% of children globally (World Health Organization, 2012). Poor oral health in the form of untreated dental caries is associated with chronic pain, infection and sepsis which impacts on both function e.g. speech, eating and sleep as well as school performance e.g. absenteeism, concentration and achievement (Gussy et al., 2006). In addition, having poor oral health can have a psychological impact on children contributing to lower self-esteem and confidence which can cause social isolation and exclusion.

On arrival in their new country, children from refugee backgrounds may never have had access or been exposed to oral health care or common preventive oral health measures, such as a toothbrush or fluoridated toothpaste (Riggs et al., 2015a). Further, families who arrive and subsequently have

children in their new country, may be unaware of preventative oral health practices for their children (Riggs et al., 2012). Another risk factor for poor oral health is the context of family social deprivation associated with socio economic status that may be compounded by low levels of parental education and health literacy (Riggs et al., 2016a). In addition, for some, previous experiences of torture and trauma may create a fear of seeking dental care (Davidson et al., 2006b).

Consistently, studies demonstrate the disparities experienced by people from refugee background in relation to untreated decayed teeth and major oral health issues (Davidson et al., 2006a). Diversity within refugee populations also suggests that acculturation (degree to which practices within the host country are adopted) may also play a role in the deterioration of oral health over time (Cote et al., 2004). Understanding these disparities is important in order to address modifiable oral health practices, however, there is little evidence available to inform interventions to improve oral health, particularly for children from refugee backgrounds.

Methods

The specific aims of this review are:

1. To report the oral health status of children of refugee background resettled in high-income refugee-receiving countries.
2. To describe the oral health beliefs, knowledge and practices of resettled refugee children and their families.
3. To describe the interventions that aim to improve refugee child oral health conducted in high-income refugee-receiving countries?

To inform this review, a systematic search strategy was developed and conducted to explore the international literature over the past ten years (2005-2016, Medline, Embase and Pubmed databases). The search strategy was developed in Medline and modified for each subsequent database (see appendix 1 in supplemental materials).

There is no agreed definition of 'who is a refugee?', however, the refugee experience is often characterised by persecution, displacement, conflict, human rights violations, family separation, prolonged uncertainty and limited or no access to services or basic necessities (shelter, food, water and safety). Experiences may include torture, physical and sexual violence (Ager and

Strang, 2008). For the purpose of this review, the term 'refugee background' is used as it recognises the varying experiences of forcibly displaced people.

A total of 281 papers were identified. Exclusion criteria (See Table 1) were applied, resulting in 13 papers (12 studies) that included children of likely refugee background or their parents.

Table 1 Inclusion and exclusion criteria

	Inclusion Criteria	Exclusion Criteria
Study design	intervention, cross sectional, epidemiological, qualitative, reviews	Case reports, commentaries. Papers without a clear description of the methods
Type of paper	Peer reviewed published papers	Non-peer reviewed. Grey literature.
Participants	Refugee background children aged 0-18.	Children or their parents not of refugee background.

	Or refugee background parents of children aged 0-18 included*	Studies in which the child data could not be separated from the adults Focus on adults only
Outcome of interest	Oral health status, behaviours, knowledge or attitudes	
Setting	High-income refugee receiving countries	Refugee camps, low-income countries, developing countries

Most of the studies were excluded because they focussed on refugee adult populations and did not describe their population sufficiently to identify whether the participants were of refugee background. In total, there were 1 longitudinal, 2 intervention, 4 cross-sectional, and 5 qualitative studies conducted in Canada, Australia and Israel (See Tables 2 and 3).

Author Manuscript

Table 2: Characteristics of longitudinal and cross sectional studies

Studies	Study design	Country of Study	Population	Sample size	Data collected	Outcomes
Vered et al (2008)(Vered et al., 2008)	Longitudinal observational – five year follow-up.	Israel	Ethiopian ‘immigrants’	BL 6 yrs and FU 11 yrs (n=48) BL 12 yrs and FU 17 yrs (n=81)	DMFT index Calibrated examiners	DMFT: Age 6-11 BL=0.10 ± 0.59; FU 0.31 ± 0.97 Age 12-17 BL=0.61 ± 1.52; FU 0.81 ± 1.75
Melvin et al (2006)(Melvin, 2006)	Intervention: school based dental screening for high risk children followed by targeted referral and supported by classroom education. 3-year prospective service utilisation evaluation.	United States	Mix of refugee and otherwise ‘impoverished’ children not further defined	n=635, age 6-11	Clinical screening by dental hygienists and referred to dentist for treatment Record review for treatment provided	% of children with a dental home; YR1 increased 59% to 78% YR2 increased 51% to 87% Proportion of children receiving restorative care YR1 = 22%, YR2 = 11% YR3 = 46%
Gibbs et al (2015)(Gibbs et al., 2015)	Intervention: bilingual peer educator led health promotion programme with service re-	Australia	39% Iraqi, 17% Lebanese - refugee background 44% Pakistani - non	n=341, age 1-4 yrs	Clinical examination with calibrated examiners.	Outcomes for refugee/non-refugee not separated. Intervention group showed sig

	orientation. Prospective before/after intervention with 18 month FU and control non-intervention group as comparison.		refugee background.		Questionnaires of parents.	improvement in both Debris Index (OR=0.44 (0.22 to 0.88)) and Modified Gingival Index (OR=0.34 (0.19 to 0.61)) 47% of intervention families and 49% of control families failed to attend FU
--	--	--	---------------------	--	----------------------------	--

Johnston et al (2012)(Johnston et al., 2012)	Cross-sectional: health assessment on arrival.	Australia	Refugees (59% Asia, 16% West and central Africa, 25% East Africa – no breakdown of children)	n=67, aged < 15 years	Retrospective review of medical records. Information collected by GMPs	19% (CI 9.6-29) dental disease (including dental pain and gum disease)
Mutch et al (2012)(Mutch et al., 2012)	Cross-sectional: health assessment on arrival	Australia	Refugees (35 self-reported ethnicities from 10 geographic regions)	n=1026 children (median age 7.8 years, range 2months-17.3years)	Retrospective review of medical records. Information collected by non-dental healthcare professionals	27.4% dental disease 9.4% reported toothache
Shah et al (2014)(Shah et al., 2014)	Cross-sectional: health assessment on arrival	United States	Refugees (80% Asia, 15% Africa, 4% Middle East, 1% Americas)	Age 0-18 Total=555	Retrospective review of medical records. Visual examination done by nurses	44.8% dental caries with significant differences between ethnic groups. African, Bhutanese & Burmese.
Nicol et al (2015)(Nicol et al., 2015)	Intervention: bespoke referral from initial settlement assessment to dental services	Australia	Refugees (Burma, Iran, Sri Lanka, Afghanistan, Iraq, Africa, Nth Africa, Middle East)	n=105 (mean age 3.2 years, range 2.9-3.4)	dmft Attendance at dental services.	62% had dmft > 0 46% attended dental services.

BL = Baseline assessment; FU = Follow Up assessment; GMP = General Medical Practitioner

Author Manuscript

Table 3 Characteristics of the qualitative studies

Studies	Study type and method	Country of Study	Population	Sample size
Amin et al (2012)(Amin and Perez, 2012)	Qualitative Focus groups	Canada	Refugee - Ethiopian, Eritrean and Somali	n=48 mothers (mean=33.6±8.1) with children age 3-5 yrs
Nicol et al (2014)(Nicol et al., 2014)	Qualitative Focus groups and interviews	Australia	Refugee - 41% Burma, 23% Middle East (Iraq, Kuwait), 13% Sudan, 8% Afghanistan, 5% Burundi, 5% Congo, 5% Rwanda & Nigeria	n=39 women with children age 0-5 yrs
Prowse et al (2014)(Prowse et al., 2014)	Qualitative Focus groups	Canada	Aboriginal, Hutterite, immigrant (Congo, Eritrea, Nigeria, Sudan), refugee (Chad, Congo, Ethiopia, Iraq, Morocco, Nigeria and Somalia)	n=40 parents/caregivers of children age 0-6 yrs n=9, Aboriginal; n=14, Hutterite; n=11, Refugee; n=6, Immigrant
Riggs et al (2016)(Riggs et al., 2016b)	Qualitative Focus groups	Australia	Refugee background from Afghanistan and Sri Lanka	n=25 mothers and fathers with children age 0-3 yrs 14 Afghan; 11 Sri Lankan
Riggs et al (2014, 2015) (Riggs et al.,	Qualitative Focus groups	Australia	Refugee and migrant background from Iraq, Pakistan and Lebanon	n=122 women with children age 0-6 yrs (n=22 Assyrian Chaldean (from Iraq)

2015a, Riggs et al., 2014a)	and interviews			n=24 Pakistani n=14 Iraqi n=29 Lebanese n=33 Lebanese and Iraqi - breakdown not reported)
--------------------------------	-------------------	--	--	---

Results

1. Oral health status of refugee children

Three studies included a clinical oral health examination undertaken by an oral healthcare professional (Vered et al., 2008, Gibbs et al., 2015, Nicol et al., 2015). The remaining four quantitative studies reported only retrospective clinical data recorded as part of a refugee general health assessment (Melvin, 2006, Mutch et al., 2012, Johnston et al., 2012, Shah et al., 2014). The origin of these audit data varied but in only one study was an oral healthcare professional involved in the dental screening (Melvin, 2006). Consequently there is very limited evidence surrounding the current oral health status of resettled refugee children.

Only one longitudinal study was identified (Vered et al., 2008) and whilst reported within the timeframes set out in the search strategy (i.e. between 2005 and 2016), the data were collected in Israel between 1999 and 2005 and as such may not be representative of current refugee child populations. The study consisted of 672 Ethiopian immigrants (refugee-like background) of whom 129 were children (under 18 years of age). Each child was examined at baseline (aged 6 and 12 years) and again after five years (aged 12 and 17 years). While DMFT was reported for these two sub-groups at both baseline and follow-up, the relatively small numbers and wide age range (with the associated difference in number of erupted permanent teeth) render these data uninformative. However, for the study sample as a whole there was a statistically significant increase in caries experience with prevalence increasing from 30% to 43% over the period of the study.

In all, no studies compared refugee oral health to any population of children born in the host country.

2. Oral health behaviours, knowledge and beliefs

Five qualitative studies (6 papers) were identified that describe the oral health knowledge, practices and beliefs of parents of children from refugee backgrounds. All are exploratory in nature and have qualitative designs (Amin and Perez, 2012, Nicol et al., 2014, Prowse et al., 2014, Riggs et al., 2016b, Riggs et al., 2015a, Riggs et al., 2014a). Several studies provide accounts of the traditional oral health practices that parents used in their home countries and how they have changed (or not) in their new country (Nicol et al., 2014, Riggs et al., 2015a, Prowse et al., 2014). Generally, parents acknowledged the importance of maintaining good oral practices, however barriers in accessing traditional forms of oral hygiene were noted. Interestingly, the Israeli longitudinal study reported that at baseline 74% of the study sample (both children and adults included at 6-18 months post arrival) used chewing and cleaning sticks and while five years later, 97% used toothbrushes exclusively (Vered et al., 2008).

There was no intervention conducted, this change in behaviour is likely to be a result of acculturation i.e. adopting practices of the host country and relinquishing those of their home country.

In most of the qualitative studies, parents suggested that the change in their children's diet was a factor that caused dental decay. Parents described a shift from a diet low in sugar in their home country, to a diet high in sugar and processed foods in their new country. Furthermore parents often felt they had little control over their child's changing diet (Amin and Perez, 2012, Nicol et al., 2014, Prowse et al., 2014, Riggs et al., 2015a).

The qualitative studies also highlighted the challenges that families face when accessing dental services in their new country. Barriers to service access were related to parental knowledge of services and entitlements, attitudes towards dentists and oral health services, English language skills, interpreter provision, dental insurance and cost of services, social support and transportation (Riggs et al., 2016b, Riggs et al., 2014a).

None of the included studies asked children themselves about their oral health.

3. Learnings from intervention studies

All the studies suggest there is a high need for dental care when children first arrive in their new country. Three studies describe interventions to facilitate access to dental care for this population (Melvin, 2006, Nicol et al., 2015, Gibbs et al., 2015). However, the outcomes and strength of evidence is mixed not least because only one study clearly describes the participants' refugee background (Nicol et al., 2015).

Set in the United States, The Tooth Tutor program involved a dental hygienist visiting schools to conduct dental screenings to identify children without a 'dental home' (defined as a child who had not seen a dentist within the previous 12 months) (Melvin, 2006). Once identified these children, who were not exclusively from refugee backgrounds, were exposed to targeted oral health education and referred for restorative care when required. Over the course of two academic years, the proportion of high risk children reported to have a 'dental home' rose from between 51-59% to 78-87%. There was also an initial decrease in the proportion of children requiring restorative care over the first two years. However by the third year, the need for restorative care increased to 46% which was attributed (without any supporting evidence) to an influx of newly arrived refugees to the schools with high dental needs.

Nicol and co-workers in Western Australia, report on a similarly targeted oral health screening programme integrated in to a general health assessment as part of a Refugee Health Service (Nicol et al., 2015). A research paediatric dentist completed an oral health screening and referred high risk children to local community dental services using bespoke referral pathways. Disappointingly less than half (46%) of those children referred actually engaged with the community dental services despite the provision of interpreting services and support from refugee health nurses. However, of those that did attend nearly two-third had caries. Financial cost and other competing community priorities were deemed barriers associated with non-attendance.

The Teeth Tales is a community based intervention involving culturally and linguistically matched peer education program with Iraqi and Lebanese from refugee background families and a third group of Pakistani non-refugee (economic) migrants in Melbourne, Australia (Gibbs et al., 2015, Gibbs et al., 2014). Simultaneously, a health service intervention was implemented that targeted aspects of cultural competence in dental service delivery. 521 families (692 children) were recruited at baseline and allocated to either intervention or comparison arms. 275 families (365 children) were followed-up 18 months later with a similarly low response rate between the intervention arm (54%) and the

comparison arm (50%). While the impact of the intervention on rates of dental caries was not reported, the intervention did increase parent oral health knowledge with a decrease in the presence of plaque on the children's teeth being interpreted as an indicator of improved tooth brushing practices.

Discussion

It is acknowledged that children from refugee backgrounds have high oral health needs and that they face specific and complex barriers to accessing dental services. Furthermore it appears that caries experience may increase with acculturation. Parental report suggests the low sugar diet and oral hygiene practices in their home country for children was positive, however, the acculturative nature of settlement is associated with an increased availability of high sugar food. This, coupled with the inability of refugee parents to access their traditional oral hygiene practices for example miswak, and a perceived loss of control over their children's dietary intake, contributes to the worsening levels of

disease over time. This has also been reported in adult Somali refugees in the United States (Geltman et al., 2013, Geltman et al., 2014). This young and vulnerable population requires responsive and empathetic care that includes working with their parents, families and communities. However there is a dearth of evidence to support effective interventions that address these issues, with this review suggesting that refugee child oral health continues to be a neglected area of research and intervention. Consequently, we are left to extrapolate from the limited evidence available, intervention strategies that appear promising and to advocate for appropriate resourcing and robust evaluation.

Multi-disciplinary collaborative workforce

It has been argued that prompt oral health care plays a significant part in supporting recovery and settlement in a new country (Davidson et al., 2007). To adequately engage families with young children, a multi-disciplinary approach is required to provide on-arrival health assessment and care that includes timely provision of preventative healthcare and information for families that includes their oral health needs. Such an approach should involve all practitioners that interact with families with young children, i.e. maternal and child health nurses, community health services, teachers, case workers, general practitioners, and counsellors along with the dental workforce.

For instance, on arrival to their new country, an oral health assessment should be available to all children with appropriate referral and follow-up in place. This should be offered regardless of reported dental pain. Dental problems should be treated and if no toothache or dental problem is identified oral health education should be provided given their new context and high risk of developing poor oral health. However, as the findings from Nicol and colleagues demonstrate, once referred, there are still barriers for families to attend dental care. A better understanding of these barriers at a service level, through community engagement and potential service changes is required to address ongoing issues related to access (Nicol et al., 2015). Barriers identified by other studies include limited access to transport, long waiting times, reminder letters sent in English, as well as staff understanding of policies and entitlements (Riggs et al., 2014a, Amin and Perez, 2012, Riggs et al., 2016b). Establishing a dental home for the newly arrived family would help assist to overcome pressing settlement priorities (psychosocial needs, housing, schooling etc) (Mutch et al., 2012, Johnston et al., 2012). The dental home, however, should be 'refugee friendly' by way of offering a flexible and approachable high quality service. Woodland and colleagues outline 10 elements of good practice to promote improved access, equity and quality of care in service delivery for newly arrived refugee children (Woodland et al., 2010). These elements of good practice delivery need to be considered for delivery in traditional and non-traditional oral health service settings.

As such, identifying opportunities in existing appropriate and acceptable non-dental settings to promote oral health and dental service access to refugee families may be a more sustainable strategy. For example, given that maternal characteristics undoubtedly play a role in their child's oral health (George et al., 2010), greater collaboration between dental and maternity health services provides a platform for oral health care. Although maternity services vary world-wide, in many countries, including Australia maternity care is government funded and provided universally to all women and families. Combining this with the high birth rate associated with many refugee background women, (Paxton et al., 2011) antenatal care provides an opportunity for sustained contact with healthcare services such that pregnant refugee women could be offered oral health care and education during this time. However, research has found that refugee women may not be aware of the significance of their own oral health; the importance of establishing a dental home; the safety of dental care during pregnancy and the recommended dental hygiene practices for their new born

(Riggs et al., 2016b). This further highlights the need for local agency collaborations and community engagement.

Caring for children typically requires communicating and working with their parents. Enabling people to communicate freely in their own language supports the development of trust, respect, rapport, cultural safety and relationship-centred care (Riggs et al., 2016a). Providing the best possible standards of professional interpreting requires service systems and management to ensure timely access to language services, accompanied by professional development, and staff opportunities to reflect and learn from their practice. The broader refugee literature demonstrates the important role of interpreters in healthcare to aid communication between clinicians and their clients (Yelland et al., 2016, Karliner et al., 2007).

The refugee population may well have had disrupted education owing to protracted time in precarious living situations, so it cannot be assumed that everyone can read in their own language let alone in that of the 'host' country. Further, due to displacement and changes in family composition, traditional ways of sharing health information may be fragmented. A range of communication strategies may need to be considered for information provision. Bicultural staff may assist families to navigate health services, get to appointments and negotiate expectations they may have for services and treatment. They also act as an aide for clinicians to understand differing health beliefs and social circumstances that may affect access to services and decision making. Internationally, the benefits of a community health worker workforce are recognised, (Glenton et al., 2013) and with support, capacity building and training this workforce could be extended and integrated into the oral health sector.

Community engagement and participation

The benefits of a considered approach to working in partnership with communities from refugee backgrounds to address health issues are not only increasingly self-evident but necessary to improve outcomes and eliminate disparities (Riggs et al., 2014b). A commitment by the oral health sector to embed participatory strategies in the engagement of communities is worthwhile. Community education is necessary to promote the importance of oral health prevention to communities where this may be an unfamiliar concept. To do this, community-based participatory strategies which are effective in engaging communities are essential, not only to conduct research but also to develop valid, effective and sustainable interventions and improve service delivery. For example, Community Advisory Groups have proved to be a fruitful and effective strategy for the engagement of refugee background communities in the context of maternal and child health and service engagement (Riggs et al., 2015b). This involves bringing together community advisors for a particular community, to

discuss community priorities and appropriate responses. Advisors enable community consultation that is inclusive and rigorous allowing community voices to be heard and dissemination of knowledge to be shared between services and the community.

Lack of attention to adequate community engagement may limit community acceptance of both the outcomes of the research and/or subsequent intervention efforts. This is inefficient and unethical and also potentially counterproductive to reducing health inequalities. A major consequence of the underrepresentation of diverse communities in research means that the available evidence is skewed, and recommendations for services, policy and programs may exacerbate inequalities in subpopulations already adversely affected (Renzaho et al., 2012).

Identifying refugee populations

There are many limitations of the research conducted with refugee populations, not least the need to define study participants accurately. Several manuscripts failed to adequately define the 'refugee'

population and it is therefore possible that studies may have been excluded that actually did include a refugee population but were inadequately described. In the course of this review, we made some assumptions based upon the participant's country of origin and our knowledge and understanding of the patterns of global humanitarian movement. For example, Vared's longitudinal study of Ethiopian migrants in Israel (Vered et al., 2008) are of 'likely-refugee background' as the statistics are evidence of the movement of this population for humanitarian reasons in previous decades (United Nations High Commissioner for Refugees, 2016).

Many papers were excluded from this review because they described their populations as 'migrant, immigrant, foreign-born, Asian, Hispanic etc'. Without further details it is impossible to infer specific refugee populations. Adequate identification in research and clinical settings is critical to establishing effective responsive interventions/care that is mindful of the health issues and trauma that may impact a child's oral health. Identifying populations of refugee backgrounds is not straightforward as they are often 'invisible' in administrative health datasets (Paxton et al., 2012) with similar challenges relating to identification also likely in clinical encounters. Professional development for staff, including both clinicians and administrators, would support better identification and triage into appropriate models of care. Identification of a 'refugee' study population must go beyond adequate documentation of country of origin, year of arrival and language spoken. An understanding of the migration history and settlement priorities would enable researchers/practitioners to be mindful of the intersecting factors (for example culture, language, financial hardship and other life circumstances) that contribute to social disparities experienced by refugee background populations that may impact on their oral health status and inform approaches to addressing these disparities. Again, adopting strategies that support bicultural community workers and community advisors are an appropriate way of gaining this knowledge. This is a complex and challenging area of research and further work is needed to assist in adequate ascertainment of refugee background people both in research, dental and other healthcare services.

Reliable measures of dental caries

Disappointingly, but consistent with much of the wider dental literature, there was a lack of consistency and quality of both the screening protocols and the instruments used to diagnose dental caries. Most of the studies use data from either retrospective review of medical records (Johnston et al., 2012, Mutch et al., 2012, Shah et al., 2014) or 'dental screenings' that are not comprehensive hence caries experience may be under reported (Melvin, 2006, Nicol et al., 2015). International standards for epidemiological oral health studies encourage consistency through the use of the International Caries Detection and Assessment System (ICDAS II) (Ismail et al., 2007). This was used

in only one study and while Gibbs and co-workers describe the implementation of ICDAS II in detail (including training and calibration of examiners) they do not report any caries outcome data in their paper (Gibbs et al., 2015). A validated and reliable method of caries assessment (that can be conducted in non-clinical settings) is required so that interventions can be evaluated and population level outcomes can be monitored overtime.

Conclusion

Overall there are limited longitudinal and intervention studies with refugee background children to support evidence-based strategies for this population. A comprehensive health assessment can assist in identifying children at risk of poor health and may help provide them with timely and effective care, advocacy and appropriate referral. The ultimate goal for this population would be to establish the burden of disease and to subsequently provide families with oral health care emphasising preventative and life-long self-management (self-efficacy). To achieve this, populations, and their

needs, have to be adequately identified, engaged with and heard, by developing partnerships and building models of community engagement. Once this is in place, disease prevalence can be assessed and culturally appropriate oral health care and education can be provided to parents and their children. Whilst it is not at all clear what might work to prevent the oral health issues experienced by refugee background children, community engagement is undoubtedly important in identifying the issues, building capacity within these communities and enabling ongoing access to a suitable dental home.

Conflicts of Interest.

The authors have no conflicts of interest to declare.

References

- Ager, A. & Strang, A. (2008). Understanding integration: A conceptual framework. *J Refug Stud*, **21**: 166-191.
- Amin, M. & Perez, A. (2012). Is the wait-for-patient-to-come approach suitable for African newcomers to Alberta, Canada? *Community Dent Oral Epidemiol*, **40**: 523-31.
- Cote, S., Geltman, P., Nunn, M., Lituri, K., Henshaw, M. & Garcia, R. I. (2004). Dental caries of refugee children compared with US children. *Pediatrics*, **114**: 733-40.
- Davidson, N., Skull, S., Burgner, D., Kelly, P., Raman, S., Silove, D., Steel, Z., Vora, R. & Smith, M. (2004). An issue of access: Delivering equitable health care for newly arrived refugee children in Australia. *J Paediatr Child H*, **40**: 569-575.
- Davidson, N., Skull, S., Calache, H., Chesters, D. & Chalmers, J. (2007). Equitable access to dental care for an at-risk group: a review of services for Australian refugees. *Aust N Z J Public Health*, **31**: 73-80.
- Davidson, N., Skull, S., Calache, H., Murray, S. & Chalmers, J. (2006a). Holes a plenty: oral health status a major issue for newly arrived refugees in Australia. *Aust Dent J*, **51**: 306-311.

Davidson, N., Skull, S., Calache, H., Murray, S. S. & Chalmers, J. (2006b). Holes a plenty: oral health status a major issue for newly arrived refugees in Australia. *Aust Dent J*, **51**: 306-11.

Geltman, P. L., Adams, J. H., Cochran, J., Doros, G., Rybin, D., Henshaw, M., Barnes, L. L. & Paasche-Orlow, M. (2013). The impact of functional health literacy and acculturation on the oral health status of Somali refugees living in Massachusetts. *Am J Public Health*, **103**: 1516-23.

Geltman, P. L., Hunter Adams, J., Penrose, K. L., Cochran, J., Rybin, D., Doros, G., Henshaw, M. & Paasche-Orlow, M. (2014). Health literacy, acculturation, and the use of preventive oral health care by Somali refugees living in massachusetts. *J Immigr Minor Health*, **16**: 622-30.

George, A., Johnson, M., Blinkhorn, A., Ellis, S., Bhole, S. & Ajwani, S. (2010). Promoting oral health during pregnancy: current evidence and implications for Australian midwives. *J Clin Nurs*, **19**: 3324-33.

Gibbs, L., Waters, E., Christian, B., Gold, L., Young, D., De Silva, A., Calache, H., Gussy, M., Watt, R., Riggs, E., Tadic, M., Hall, M., Gondal, I., Pradel, V. & Moore, L. (2015). Teeth Tales: a community-based child oral health promotion trial with migrant families in Australia. *BMJ Open*, **5**: e007321.

Gibbs, L., Waters, E., De Silva, A., Riggs, E., Moore, L., Armit, C., Johnson, B., Morris, M., Calache, H., Gussy, M., Young, D., Tadic, M., Christian, B., Gondal, I., Watt, R., Pradel, V., Truong, M. & Gold, L. (2014). An exploratory trial implementing a community-based child oral health promotion intervention for Australian families from refugee and migrant backgrounds: a protocol paper for Teeth Tales. *BMJ Open*, **4**: e004260.

- Glenton, C., Colvin, C. J., Carlsen, B., Swartz, A., Lewin, S., Noyes, J. & Rashidian, A. (2013). Barriers and facilitators to the implementation of lay health worker programmes to improve access to maternal and child health: qualitative evidence synthesis. *Cochrane Database of Systematic Reviews*.
- Gussy, M. G., Waters, E. G., Walsh, O. & Kilpatrick, N. M. (2006). Early childhood caries: current evidence for aetiology and prevention. *J Paediatr Child Health*, **42**: 37-43.
- Ismail, A., Sohn, W., Tellez, M., Amaya, A., Sen, A., Hasson, H. & Pitts, N. (2007). The International Caries Detection and Assessment System (ICDAS): an integrated system for measuring dental caries. *Community Dent Oral Epidemiol*, **35**: 170-178.
- Johnston, V., Smith, L. & Roydhouse, H. (2012). The health of newly arrived refugees to the Top End of Australia: results of a clinical audit at the Darwin Refugee Health Service. *Aust J Prim Health*, **18**: 242-7.
- Karliner, L. S., Jacobs, E. A., Chen, A. H. & Mutha, S. (2007). Do professional interpreters improve clinical care for patients with limited English proficiency? A systematic review of the literature. *Health Serv Res*, **42**: 727-54.
- Melvin, C. S. (2006). A collaborative community-based oral care program for school-age children. *Clinical Nurse Specialist*, **20**: 18-22.
- Mutch, R. C., Cherian, S., Nemba, K., Geddes, J. S., Rutherford, D. M., Chaney, G. M. & Burgner, D. P. (2012). Tertiary paediatric refugee health clinic in Western Australia: analysis of the first 1026 children. *J Paediatr Child Health*, **48**: 582-7.
- Nicol, P., Al-Hanbali, A., King, N., Slack-Smith, L. & Cherian, S. (2014). Informing a culturally appropriate approach to oral health and dental care for pre-school refugee children: a community participatory study. *BMC Oral Health*, **14**: 69.
- Nicol, P., Anthonappa, R., King, N., Slack-Smith, L., Cirillo, G. & Cherian, S. (2015). Caries burden and efficacy of a referral pathway in a cohort of preschool refugee children. *Aust Dent J*, **60**: 73-9.
- Paxton, G., Smith, N., Win, A., Mulholland, N., & Hood, S. (2011). Refugee Status Report. A report on how refugee children and young people in Victoria are faring. Melbourne, Victorian Department of Education and Early Childhood Development.

- Paxton, G. A., Kay, M. P. & Correa-Velez, I. (2012). Lost and found: improving ascertainment of refugee-background Australians in population datasets. *Med J of Aust*, **197**: 552-3.
- Prowse, S., Schroth, R. J., Wilson, A., Edwards, J. M., Sarson, J., Levi, J. A. & Moffatt, M. E. (2014). Diversity considerations for promoting early childhood oral health: a pilot study. *Int J Dent*, doi: 10.1155/2014/175084.
- Renzaho, A., Renzaho, C. & Polonsky, M. (2012). Left out, left off, left over: Why migrants from non-English speaking backgrounds are not adequately recognised in health promotion policy and programs. *Health Promot J Austr*, **23**: 84-85.
- Riggs, E., Gibbs, L., Kilpatrick, N., Gussy, M., Van Gemert, C., Ali, S. & Waters, E. (2015a). Breaking down the barriers: a qualitative study to understand child oral health in refugee and migrant communities in Australia. *Ethn Health*, **20**: 241-57.
- Riggs, E., Gussy, M., Gibbs, L., Van Gemert, C., Waters, E. & Kilpatrick, N. (2014a). Hard to reach communities or hard to access services? Migrant mothers' experiences of dental services. *Aust Dent J*, **59**: 201-207.
- Riggs, E., Gussy, M., Gibbs, L., Van Gemert, C., Waters, E., Priest, N., Watt, R., Renzaho, A. M. & Kilpatrick, N. (2014b). Assessing the cultural competence of oral health research conducted with migrant children. *Community Dent Oral Epidemiol*, **42**: 43-52.
- Riggs, E., Van Gemert, C., Gussy, M., Waters, E. & Kilpatrick, N. (2012). Reflections on cultural diversity in oral health promotion and prevention. *Glob Health Promot*, **19**: 60-63.
- Riggs, E., Yelland, J., Duell-Piening, P. & Brown, S. J. (2016a). Improving health literacy in refugee populations. *Med J of Aust*, **204**: 9-10.

- Riggs, E., Yelland, J., Shankumar, R. & Kilpatrick, N. (2016b). 'We are all scared for the baby': promoting access to dental services for refugee background women during pregnancy. *BMC Pregnancy Childbirth*, **16**: 12.
- Riggs, E., Yelland, J., Szwarc, J., Casey, S., Chesters, D., Duell-Piening, P., Wahidi, S., Fouladi, F. & Brown, S. (2015b). Promoting the inclusion of Afghan women and men in research: reflections from research and community partners involved in implementing a 'proof of concept.'. *Int J Equity Health*, **14**: 13.
- Selwitz, R. H., Ismail, A. I. & Pitts, N. B. (2007). Dental caries. *Lancet*, **369**: 51-59.
- Shah, A. Y., Suchdev, P. S., Mitchell, T., Shetty, S., Warner, C., Oladele, A. & Reines, S. (2014). Nutritional status of refugee children entering DeKalb County, Georgia. *J Immigr Minor Health*, **16**: 959-67.
- Shonkoff, J. P., Siegel, B. S., Garner, A. S., Dobbins, M. I., Earls, M. F., McGuinn, L., Pascoe, J. & Wood, D. L. (2012). The lifelong effects of early childhood adversity and toxic stress. *Pediatrics*, **129**: (1):e232-46
- United Nations High Commissioner for Refugees (2015). World at War. Global Trends. Forced Displacement in 2014. Geneva, Switzerland: UNHCR.
- United Nations High Commissioner for Refugees. (2016). *Ethiopia* [Online]. Available: <http://www.unhcr.org/ethiopia.html> [Accessed 2016].
- Vered, Y., Zini, A., Livny, A., Mann, J. & Sgan-Cohen, H. D. (2008). Changing dental caries and periodontal disease patterns among a cohort of Ethiopian immigrants to Israel: 1999-2005. *BMC Public Health*, **8**: 345.
- Victorian Foundation for the Survivors of Torture (1998). Rebuilding Shattered Lives. Melbourne.
- Woodland, L., Burgner, D., Paxton, G. & Zwi, K. (2010). Health service delivery for newly arrived refugee children: a framework for good practice. *J Paediatr Child Health*, **46**: 560-7.
- World Health Organization. (2012). *Oral Health* [Online]. Available: <http://www.who.int/mediacentre/factsheets/fs318/en/> [Accessed 2016].

Yelland, J., Riggs, E., Szwarc, J., Casey, S., Duell-Piening, P., Chesters, D., Wahidi, S., Fouladi, F. & Brown, S. (2016). Compromised communication: a qualitative study exploring Afghan families and health professionals' experience of interpreting support in Australian maternity care. *BMJ Qual Saf*, **25**: (4):e1

Author Manuscript

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Riggs, E;Rajan, S;Casey, S;Kilpatrick, N

Title:

Refugee child oral health

Date:

2017-04-01

Citation:

Riggs, E., Rajan, S., Casey, S. & Kilpatrick, N. (2017). Refugee child oral health. ORAL DISEASES, 23 (3), pp.292-299. <https://doi.org/10.1111/odi.12530>.

Persistent Link:

<http://hdl.handle.net/11343/291573>