LETTER TO THE EDITOR

The Role of Health Literacy in the Treatment of Osteoporosis

SM Hosking,¹ R Buchbinder,^{2,3} JA Pasco,^{1,2,4} LJ Williams,¹ and SL Brennan-Olsen^{1,5,6}

¹School of Medicine, Deakin University, Geelong, Australia

²Department of Epidemiology and Preventive Medicine, Monash University, Melbourne, Australia

³Monash Department of Clinical Epidemiology, Cabrini Institute, Malvern, Australia

⁴Melbourne Medical School–Western Campus, University of Melbourne, St Albans, Australia

⁵Australian Institute for Musculoskeletal Science, University of Melbourne, St Albans, Australia

⁶Institute of Health and Ageing, Australian Catholic University, Melbourne, Australia

Address correspondence to: <!--<query>AQ: Please supply name of corresponding author and highest academic degree(s). Please provide complete mailing address and E-mail address. Phone and fax numbers are not needed.</query>-->

Journal of Bone and Mineral Research, Vol. XX, No. X, Month 2016, pp XXXX–XXXX<!--<query>PROD: Please fill in volume number, issue number, month, publication year, and page range.</query>-->

DOI: 10.1002/jbmr. 2919

© 2016 American Society for Bone and Mineral Research

To the Editor

We noted with interest the recent Editorial by Khosla and Shane⁽¹⁾ that drew attention to the poor uptake of, and adherence to, therapeutic interventions in patients at risk of fragility fracture. We wish to emphasize the authors' statement that highlighted the importance of

ensuring patients have access to accurate information to inform their healthcare decisions, and we agree that the consequences of osteoporosis and the importance of medication adherence are not necessarily clear to patients. We add that health literacy is likely to play a major role in the ability of patients to access, understand, and apply information in order to make informed decisions about osteoporosis treatment.

To date, very little research has been undertaken regarding health literacy, osteoporosis, and medication adherence. One population-based study found that individuals with osteoporosis had poorer functional health literacy (health-related literacy and numeracy skills) compared with the general population.⁽²⁾ Although another study found no association between health literacy and medication adherence in women with osteoporosis,⁽³⁾ studies of people with other chronic diseases have shown an association between lower levels of functional health literacy and poorer medication adherence.⁽⁴⁾

Functional health literacy tools are limited in their ability to capture the full breadth of the health literacy concept and focus principally on literacy, with some including numeracy skills.⁽⁵⁻⁷⁾ Additional aspects of health literacy that may be important in the decision to adhere to osteoporosis medication include the doctor–patient relationship as well as the doctor’s ability to impart information and an individual’s ability to understand and appraise additional information on the risks and benefits of treatment well enough to make an informed decision. It is likely that poor medication adherence is multifaceted and complex.

Understanding health literacy needs of individual patients might provide greater insight into the barriers of medication adherence, as discussed by Khosla and Shane. This may be achieved by the application of more recently developed health literacy tools. As an exemplar, the Health Literacy Questionnaire (HLQ)⁽⁸⁾ is a multidimensional measure of health literacy that was developed using a grounded approach. The HLQ includes nine

independent domains of health literacy, and can be used to identify an individual's health literacy needs. This information would inform a more person-centered approach to addressing the problem and may be one method for improving medication adherence and patient outcomes.

References

1. Khosla S, Shane E. A crisis in the treatment of osteoporosis. *J Bone Miner Res.* 2016;1–7<!--<query>AQ: Please supply Vol.(issue) numbers.</query>-->.
2. Hill CL, Appleton SL, Black J, et al. Role of health literacy in self-reported musculoskeletal disorders. *Arthritis.* 2015;2015:607472. Epub 2015 Aug 18. DOI: 10.1155/2015/607472<!--<query>AQ: Any pub update on this?</query>-->.
3. Shehadeh-Sheeny A, Eilat-Tsanani S, Bishara E, Baron-Epel O. Knowledge and health literacy are not associated with osteoporotic medication adherence, however income is, in Arab postmenopausal women. *Patient Educ Couns.* 2013;93(2):282–8.
4. Berkman ND, Sheridan SL, Donahue KE, Halpern DJ, Crotty K. Low health literacy and health outcomes: an updated systematic review. *Ann Intern Med.* 2011;155(2):97–107.
5. Weiss BD, Mays MZ, Martz W, et al. Quick assessment of literacy in primary care: the newest vital sign. *Ann Fam Med.* 2005;3:514–22.
6. Parker RM, Baker DW, Williams M V, Nurss JR. The test of functional health literacy in adults: a new instrument for measuring patients' literacy skills. *J Gen Intern Med.* 1995;10:537–41.
7. Murphy PW, Davis TC, Long SW, Jackson RH. Rapid estimate of adult literacy in medicine (REALM): a quick reading test for patients. *J Read.* 1993;37(2):124–30.
8. Osborne RH, Batterham RW, Elsworth GR, Hawkins M, Buchbinder R. The grounded

psychometric development and initial validation of the health literacy questionnaire (HLQ).

BMC Pub Health. 2013;13(1):1–17.

Author Manuscript

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Hosking, SM;Buchbinder, R;Pasco, JA;Williams, LJ;Brennan-Olsen, SL

Title:

The Role of Health Literacy in the Treatment of Osteoporosis

Date:

2016-10-01

Citation:

Hosking, S. M., Buchbinder, R., Pasco, J. A., Williams, L. J. & Brennan-Olsen, S. L. (2016). The Role of Health Literacy in the Treatment of Osteoporosis. JOURNAL OF BONE AND MINERAL RESEARCH, 31 (10), pp.1909-1909. <https://doi.org/10.1002/jbmr.2919>.

Persistent Link:

<http://hdl.handle.net/11343/291591>