Received Date : 09-Mar-2015

Revised Date : 31-Oct-2015

Accepted Date : 27-Nov-2015

Article type : Original Article

Can heat-killed *Gordonia bronchialis* enhance growth and immunity in rainbow trout (*Oncorhynchus mykiss*)

Najmeh Sheikhzadeh^{1,*}, Keyvan Tahapour¹, Katayoon Nofouzi^{2,*}, Amir Tukmechi³, Sadigheh Shabanzadeh⁴, Ali Khani Oushani⁵, John Stanford⁶, Graham McIntyre⁶, Karim Mardani⁷

¹ Department of Food Hygiene and Aquatic Animals, Faculty of Veterinary Medicine, University of Tabriz, Tabriz, Iran; ² Department of Pathobiology, Faculty of Veterinary Medicine, University of Tabriz, Tabriz, Iran; ³ Department of Pathobiology and Quality Control, Artemia and Aquatic Animals Research Institute, Urmia University, Iran; ⁴ Department of Fisheries, Babol Branch, Islamic Azad University, Babol, Iran; ⁵ Department of Fisheries, Science and Research Branch, Islamic Azad University, Tehran, Iran; ⁶ BioEos Ltd, Kent, UK. Ex: (Department of Medical Microbiology, Royal Free and University College Medical School, London, UK); ⁷ Department of Food Hygiene and Quality Control, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Correspondences: N. Sheikhzadeh, Department of Food Hygiene and Aquatic Animals, Faculty of Veterinary Medicine, University of Tabriz, Tabriz, Iran. E-mail: nsheikh@tabrizu.ac.ir

K. Nofouzi, Department of Pathobiology, Faculty of Veterinary Medicine, University of Tabriz, Tabriz, Iran. E-mail: nofouzi@tabrizu.ac.ir

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/anu.12445](https://doi.org/10.1111/anu.12445)

This article is protected by copyright. All rights reserved

Abstract

In the present study, the efficacy of heat-killed *Gordonia bronchialis* on growth performance, immune system and gastrointestinal structure in rainbow trout (*Oncorhynchus mykiss*) were evaluated. Fish (mean weight 30 g) were fed basal diet (control), or treatment diets containing 2.48×10^8 (low dose) or 1.24×10^9 (high dose) cells kg^{-1} feed of heat-killed *Gordonia bronchialis* on a pulse basis (5 days on treatment diet; 10 days on basal diet) for 95 days. On days 95 and 105, some of the fish were sampled for analysis of mentioned parameters. On days 110 and 120, the remaining fish were injected intraperitoneally with a 20 ml L^{-1} suspension of chicken red blood cells. Results showed that growth performance was significantly enhanced in both treatment groups compared to the control group. Serum complement and lysozyme activities and hemagglutination antibody titer were higher in both treatment groups compared with controls. The length of the intestinal and pyloric caeca folds were increased in the high dose group. Meanwhile, the number of goblet cells was increased in both treatment groups. This study suggests that heat-killed *Gordonia bronchialis* has the potential to enhance growth, immunological parameters and the gastrointestinal structure in rainbow trout.

KEY WORDS: *Gordonia bronchialis*, rainbow trout, immune system, growth, gastrointestinal histology.

Introduction

It is widely demonstrated that farmed fish are more susceptible to disease agents than their wild counterparts due to the artificial conditions posed by intensive rearing (Price 1999; Huntigford 2004; Salinas *et al.* 2006). When infectious outbreaks appear they may be fought by means of chemotherapeutics, vaccines or immunostimulants (Salinas *et al.* 2006). The administration of immunostimulants to fish through the diet has appeared as a very promising control measure in fish farms. In fact, these additives augment the innate and adaptive immune responses and remain the key strategies for prevention of diseases in fish aquaculture.

Among the aerobic, near-mycobacterial genera, within the Actinomycetales, are some species with adjuvant activities and antigens very similar to those of *Mycobacterium vaccae*, but with subtle differences (Stanford & Stanford 2012). More recently, a number of these species, including *Gordonia bronchialis* (*G. bronchialis*), *Rhodococcus coprophilus*, and *Tsukamurella inchonensis*, capable of exerting subtly different adjuvant or immunostimulatory activities in human, veterinary and agricultural medicine have been identified (Tarrés *et al.* 2012). For example, published studies with heat-killed *G. bronchialis* showed that it could limit parasitaemias in rats challenged with live *Trypanosoma cruzi* (Fontanella *et al.* 2007). Pregnant rats treated with *G. bronchialis* also gave birth to offspring protected from subsequent challenge with *Trypanosoma cruzi* (Davila *et al.* 2011). In dogs, a preparation of *G. bronchialis* was particularly effective against an allergy induced by flea-bites (Marro *et al.* 2011). Mice treated with heat-killed *G. bronchialis* controlled an IgE-mediated food allergy in a mouse model (Smaldini *et al.* 2013). These findings can be attributed to immune modulation resulting from the effects of powerful cell wall-associated adjuvants in *G. bronchialis* (Smaldini *et al.* 2013).

Some studies have so far addressed the effects of heat-killed *G. bronchialis* in aquaculture. Oral administration of *G. bronchialis* improved survival and weight gain in shrimp and young fish and enhanced growth rate and color and reduced skeletal abnormalities in koi carp (*Cyprinus carpio*) (Stanford & Stanford 2012). In this study, the effects of dietary heat-killed *G. bronchialis* on some immune parameters, growth performance and gastrointestinal structure in juvenile rainbow trout (*Oncorhynchus mykiss*) were evaluated.

Materials and methods

Fish

The experiment was carried out at a fish farm in Firoozkooh, Iran. Rainbow trout, of 30g average wet weight, were acclimatized for 10 days on the basal diet produced in a commercial mill (Faradaneh, Isfahan, Iran). The open formula of this commercial diet is given in Table 1. Fish were kept in indoor cement tanks ($1.8 \times 0.22 \times 0.35$ m) in flow-through system supplied with aerated well water with following characteristics: flow rate 0.5 l s^{-1} , water temperature $12 \pm 1^\circ\text{C}$, dissolved oxygen >8 ppm, $\text{NH}_3 < 0.01 \text{ mg l}^{-1}$, $\text{NO}_2 < 0.1 \text{ mg l}^{-1}$, hardness 275 mg l^{-1} and pH 7.8. A total of 162 fish were randomly distributed into three experimental groups following a completely randomized design.

Experimental diet

Cultured and heat-killed *G. bronchialis* by autoclaving was obtained from BioEos Ltd (Kent, UK). Experimental diets were prepared by adding 2.48×10^8 and 1.24×10^9 cells kg^{-1} feed of heat-killed *G. Bronchialis* into commercial basal diet which was equivalent to 2.074 g and 10.370 g powder in potato starch, according to the manufacturer. The heat-killed organism was added to the diet by spraying 20 ml fish oil kg^{-1} feed. A control diet was prepared by spraying 20 ml fish oil kg^{-1} only to the basal diet. The modified feed was stored in tightly sealed plastic bags at 8 - 10°C until they were used in the feeding experiments. Three groups in triplicate with 18 fish in each tank were fed the experimental diets formulated with 2.48×10^8 or 1.24×10^9 cells of dead *G. bronchialis*, or the basal diet for 95 days with pulse administration of treatment diet for 5 days followed by 10 days on the basal diet as following: day 1 till 5: treatment diet; day 6 till 15: basal diet; day 16 till 20: treatment diet; day 21 till 30: basal diet. Similar pattern was followed till day 95. From day 95 till day 105, fish were reverted back to the basal diet. Fish were sampled on days 95 and 105 to evaluate whether the effects of using this immunostimulant would last till that time. In this study, fish were initially fed at a feeding rate of 25 g kg^{-1} of body weight daily; this rate was designed to assure apparent satiation. Till the end of the trial, fish were weighed once every 2 weeks to adjust the feeding during the experiment.

Growth performance

All fish in each tank were taken and weighed at the beginning and the termination of the feeding trial on day 95. Feed conversion ratio (FCR) and thermal growth coefficient (TGC) were determined for each tank using the following formula:

FCR = total feed supplied / weight gain.

TGC = $[(W_2^{1/3} - W_1^{1/3}) / (T_2 - T_1) \times ^\circ\text{C}] \times 1000$, where W_1 and W_2 are the initial and final weight (g), respectively, $T_2 - T_1$ is the number of days in the feeding period and $^\circ\text{C}$ is the mean daily temperature (Sanden *et al.* 2006).

Immunological parameters

On days 95 and 105 of the trial, four fish from each tank were anaesthetized in a clove oil bath (50 $\mu\text{l L}^{-1}$) and blood was collected from the caudal vein. Fish bloods were allowed to clot at room temperature for 1 h and stored in a refrigerator overnight. The clot was then centrifuged at 1500 g for 5 min. Then the serum was collected and stored in sterile eppendorf tubes at -20°C until used for assays. Noteworthy, all sampling fish for immunological parameters were removed from study after blood collection.

Serum lysozyme activity was measured according to methods described by Demers & Bayne (1997). Briefly, 25 μl of individual serum samples was mixed with a 175 μl *Micrococcus lysodieticus* (Sigma) suspension at 75 $\mu\text{g ml}^{-1}$ in 0.1 M phosphate citrate buffer, pH 5.8. After rapid mixing, turbidity changes were measured every 30 s for 5 min at 450 nm at approximately 20°C using a microplate reader. The dilutions of hen eggwhite lysozyme (Sigma) ranging from 0 to 20 $\mu\text{g ml}^{-1}$ in the same buffer served as the standard. The equivalent unit of activity of the sample as compared to the standard was determined and expressed as $\mu\text{g ml}^{-1}$ serum.

Alternative complement serum activity was determined using previously reported method (Andani *et al.* 2012). Rabbit red blood cells (RaRBC) were washed three times in ethylene glycol tetra acetic acid–magnesium–gelatin veronal buffer (0.01 M EGTA–Mg–GVB, pH 7) and resuspended in this buffer to a concentration of 2×10^8 cells ml^{-1} . A 100 μl suspension of RaRBC was added to 250 μl of serially diluted serum in buffer. Samples were incubated at 20°C for 90 min with regular shaking. Approximately 3.15 ml of 0.85% NaCl solution was added and the tubes centrifuged at 1600 g for 10 min at 4°C . The extent of hemolysis was estimated by measuring the optical density of the supernatant at 414 nm using a spectrophotometer (Awariness, Palm, FA). ACH50 units were defined as the concentration of serum giving 50% hemolysis of RaRBC.

Total antibody level in the serum was determined following the method of Siwicki *et al.* (1994). The difference in total protein content prior to and after precipitation of the antibody component with 12% polyethylene glycol (PEG; Sigma) was determined by the Bradford method (Kruger 1996).

The total peroxidase content present in serum was measured according to Cuesta *et al.* (2005). Serum samples (15 μl) were placed in each well of a 96 well plate. 135 μl of HBSS without

Ca²⁺ or Mg²⁺ was then added to each well. Finally 50 µl of 20 mM 3,3',5,5'-tetramethylbenzidine hydrochloride (TMB) (Sigma) and 5 mM H₂O₂ was added. The color-change reaction was stopped after 2 min by adding 50 µl of 2 M sulfuric acid and the optical density was read at 450 nm by ELISA reader.

Gastrointestinal histological parameters

Tissues were collected from two fish in each tank on days 95 and 105 of the trial. **Noteworthy, sampling fish for histology were in addition to the fish used for immunological parameters. After necropsy, macroscopic observations were performed.** Then, liver, kidney, spleen, gills, skeletal muscle, intestine, pyloric caeca samples were taken, separately. Samples were fixed in buffered formalin for 48 h, dehydrated in alcohols and xylene and then embedded in paraffin. A five micron subsample was then rehydrated in alcohol and stained with haematoxylin-eosin. Length and thickness of proximal intestinal and pyloric caeca folds were measured using a graded ocular lens. The percentage of goblet cells among all epithelial cells in fish proximal intestine and pyloric caeca were also defined. Goblet cell percentage was determined by counting 20 folds in each fish and about five sections per sample. The magnification used was ×200.

Challenge test with chicken red blood cells (C-RBC) suspension

Red blood cells were collected from 28 days old, Ross strain broiler chickens raised under recommended rearing requirements (Ross Broiler Management handbook, Aviagen, 2014). Two milliliters of citrated (3.8% NaCit solution at 1:10 dilution) chicken blood was suspended in 10 ml Tris buffered saline (TBS, 50 mM TrisHCl, pH 8.0, 150 mM NaCl) and centrifuged at 2500 g for 10 min. The supernatant was aspirated without disturbing the blood cells. 10 ml TBS was added and mixed by inverting. Following the centrifugation and washing two more times, the supernatant was aspirated and 2.5% chicken red blood cell (C-RBC) suspension in TBS was prepared (Sheikhzadeh *et al.* 2012). On day 110 of trial, all remaining fish in each tank (**six fish per tank**) were anesthetized in a clove oil bath (50 µl L⁻¹) and injected intraperitoneally with 0.5 mL of the C-RBC suspension in PBS. 10 days later, these fish received boosting injection. On day 135 of trial, 15 days after boosting injection, blood was collected from these injected fish

and serum was obtained with the same procedure described above and kept for further use at - 20°C.

Hemagglutination assay

The antigen-specific antibody response was determined by hemagglutination assay (Rao & Chakrabarti 2005). Blood from chicken was collected in citrate solution and the cells were washed in PBS and resuspended to 20 ml L⁻¹ in PBS. Serum of control or test fish (50 µL) was serially diluted in PBS in 96 well round-bottomed microtiter plates. An equal volume of 20 ml L⁻¹ C-RBC was added to all wells and kept for 1 h at room temperature and then overnight at 4°C. Hemagglutinating antibody titer was expressed as the reciprocal of the highest dilution giving visible agglutination.

Statistical analysis

The results were expressed as a mean ± standard error (SEM). The statistical significance of data was evaluated by two-way and one-way analysis of variance (ANOVA), using the statistical package for social sciences (SPSS) software version 19.0. The data were analyzed by LSD analysis with the least square difference to compare the means. Differences were considered significant at $P < 0.05$.

Results

Growth performance

Final weight, final length, FCR, and TGC improved significantly in both treatment groups compared to the control group (Table 2).

Immunological parameters

No significant differences in all immunological responses between day 95 and 105 were shown during this trial. Dietary intake of heat-killed *G. bronchialis* in both treatment groups increased the serum complement activity compared to the control group on days 95 of the feeding trial. Similar increase was also shown on day 105 of the trial. No significant impact on serum total antibody and peroxidase contents in any treatment group was noted on days 95 and 105 of the study. In the assay of lysozyme activity in serum of fish treated with both low and high dose of *G. bronchialis*, significant elevation was observed compared with the control group on days 95 and 105 of the study (Table 3). After immunization, the antigen-specific response was significantly higher in the high dose groups than the control group on day 130 of trial whereas no significant differences were noticed between treatment groups (Figure 1).

Histological findings

Upon post-mortem examination, no gross lesions or microscopic changes were detected in any group. No pathological changes were also noted in the intestine and pyloric caeca of fish from all groups that shows the normal architectures of these tissues. In fish that received the high dose of heat-killed *G. bronchialis*, the intestinal fold length increased compared to low dose group and control group on days 95 and 105 of feeding. No significant change in fold thickness in any group was shown on day 95 of trial where as on day 105 fold thickness increased in both treatment groups in comparison with control group. Significant increase in mucous cell numbers for fish in both treatment groups were shown compared to the control group on day 95 of the

trial. Highest elevation was just noted in fish receiving the high dose of heat-killed *G. bronchialis*. Meanwhile, on day 105 of the study, mucous cell numbers was only increased in fish received the high dose *G. bronchialis* compared with other groups. Comparing the intestinal morphology on days 95 and 105, significant differences in goblet cells number were only noted between control fish (Table 4).

On days 95 and 105 of the trial, the fold length of the pyloric caeca increased in fish fed the high dose of heat-killed *G. bronchialis* compared with control and low dose group. On day 95, higher fold thickness of the pyloric caeca was observed in both treatment groups compared to the control group. Whereas on day 105 fold thicknesses did not change in any treatment group compared with the control group. Higher numbers of mucous cells in pyloric caeca were exhibited in both treatment groups compared with the control group on days 95 and 105 of trial. Additionally, highest mucous cell number was just noted in fish fed the high dose of heat-killed *G. bronchialis*. Comparing the pyloric caeca morphology on days 95 and 105, similar pattern to intestine were shown in a way that significant differences in goblet cells number between control fish were noted (Table 5).

Discussion

The results show that heat-killed *G. bronchialis* in both treatment groups enhanced the growth performance in rainbow trout. Similar results were observed in koi carp and shrimp (Stanford & Stanford 2012). In fish species, higher growth performance can happen by different mechanisms (Heidarieh *et al.* 2012). Influencing nutrient especially protein digestibility by maintaining the function and structure of the small intestine leads to increased digestive capacity of the gut. Improved digestive enzymes, including lipase, amylase and protease could also result in better growth performance. In the present study, histological examination showed some differences in small intestine and pyloric caeca structures. In the high dose group, increased fold length in the intestine and pyloric caeca were noted. Thicker folds in the pyloric caeca and intestine in both

treatment groups were also shown on days 95 and 105 of the feeding trial, respectively.

Numerous anatomical features determine the total absorptive surface area of the gastrointestinal tract. For example, taller, narrower and regularly shaped fold and higher number of fold per unit area are indicators that the function of the intestinal folds is activated (Heidarieh *et al.* 2013). In this study, taller shaped fold in the high dose group were a good indicator but the thicker folds should be evaluated further to elucidate the effects, which will happen to fish administrated with heat-killed *G. bronchialis*. Comparing the effects of intestinal morphology with growth performance, it is clear that most intestinal changes were exhibited with a higher dose of administration. However, similar growth performance in both treatment groups was observed compared to the control group. Therefore, other factors may also have affected the growth performance in this study. Noteworthy, this trial was carried out on commercial fish farm; different additives inside the extruded diet might also interact with heat-killed *G. bronchialis*. Therefore, to elucidate the exact mechanisms through which the heat-killed *G. bronchialis* can affect the growth in fish, administration of this heat-killed bacterium in a diet without any additives is needed.

In this study, large amount of goblet cells in fish gastro-intestine after feeding dead *G. bronchialis* were noted. Even though higher goblet cell density in fish skin and intestinal tract after challenge with infectious diseases have been widely studied (Buchmann & Bresciani 1998; Ringø *et al.* 2003) increase in mucus production in response to the feed additives were also shown (Torresillas *et al.* 2011; Covello *et al.* 2012; Heidarieh *et al.* 2012; Xueqin *et al.* 2012; Shin *et al.* 2014). Goblet cells in the gastrointestinal tract provide protection against damage by forming a viscous, hydrated blanket on the surface of the mucosa which acts as a sensitive first line of immune defense parameter in fish (Shin *et al.* 2014). Meanwhile, it was suggested that goblet cells positively affect the absorption of digestible substances (Shin *et al.* 2014) that can partly explain the improvement in growth performance in fish administrated with heat-killed *G. bronchialis*.

Heat-killed *G. bronchialis* in both treatment groups were effective in enhancing serum lysozyme and complement activities. On other hand, total antibody and peroxidase levels were not affected by administration of dead *G. bronchialis*. The lysozyme and alternative pathway of complement are powerful non-specific defense mechanisms for protecting fish against a wide range of

potentially invasive organisms, such as bacteria, fungi, viruses, and parasites (Choi *et al.* 2008; Son *et al.* 2009). In fact, the innate immune system of fish comprises several humoral and cellular factors that may differentially respond to the offered immunostimulants (Shin *et al.* 2014). After immunization, higher antigen-specific response was shown in high dose group, showing the positive effect of heat-killed *G. bronchialis* on specific immune responses in rainbow trout. Immunostimulatory effects of dead *G. bronchialis* have been earlier reported in laboratory animal and veterinary medicine (Fontanella *et al.* 2007; Davila *et al.* 2011; Marro *et al.* 2011; Stanford & Stanford 2012; Smaldini *et al.* 2013). The effectiveness of preparations of heat-killed Actinomycetales is attributed to highly active immunostimulatory adjuvants in their cell walls, with subtle differences between species (Marro *et al.* 2011).

Considering the long period of fish culture, continual feeding of immunostimulants might have harmful effects on the immune system (Bricknell & Dalmo 2005). Therefore, a pulse feeding strategy during 95 days, with treatment diet for 5 days followed by basal diet for 10 days, was evaluated. Results showed up-regulation of the immune system after administration of heat-killed *G. bronchialis* on day 95 of trial. Meanwhile, fish were also sampled on day 105 to analyze the effects of 10 days withdrawal time on different immunological and histological parameters. In fact, on day 105 an enhanced response of the immune system and histological parameters fairly similar to day 95 were noted even though goblet cell percentage in low dose group returned to baseline which shows the minor decrease in immune stimulation. Noteworthy, comparing the immunological and histological parameters on days 95 and 105, significant differences in goblet cells number in intestine and pyloric caecum between control fish were noted whereas no significant differences in treatment groups between day 95 and 105 were shown during this trial. The exact reason for differences in goblet cells number between control fish is unknown and needs more considerations to clarify these observations. Converse to this substance, immune stimulation by feeding a probiotic is more transient and a few days after cessation of probiotic administration, it decreases dramatically (Andani *et al.* 2012). Reduction in immune responses following the administration of probiotics can be attributed to a decrease in bacterial count after feeding on the basal diet (Andani *et al.* 2012). In general, immune stimulation by feeding different substances even in withdrawal time can provide an immense flexibility in fish farming especially during periods of increased disease risk.

In conclusion, this preliminary study showed that administration of heat-killed *G. bronchialis* has beneficial effects in rainbow trout, affecting parameters like growth performance, immunity and gastrointestinal structure. Further investigations are needed to fully understand the interaction between this organism and different fish species even on the molecular level.

Acknowledgements

The authors wish to express their gratitude to BioEos Ltd. for supplying heat-killed *G. bronchialis*. The authors are also grateful to Research affairs of Tabriz University for the financial support.

References

- [1] Andani, H.R.R., Tukmechi, A., Meshkini, S. & Sheikhzadeh, N. (2012) Antagonistic activity of two potential probiotic bacteria from fish intestines and investigation of their effects on growth performance and immune response in rainbow trout (*Oncorhynchus mykiss*). *J. Appl. Ichthyol.*, **28**, 728-734.
- [2] Bricknell, I. & Dalmo, R.A. (2005) The use of immunostimulants in fish larval aquaculture. *Fish Shellfish Immunol.*, **19**, 457-472.

- [3] Buchmann, K. & Bresciani, J. (1998) Microenvironment of *Gyrodactylus derjavini* on rainbow trout *Oncorhynchus mykiss*: association between mucous cell density in skin and site selection. *Parasitol. Res.*, **84**, 17-24.
- [4] Choi, S.H., Park, K.H., Yoon, T.J., Kim, J.B., Jang, Y.S. & Choe, C.H. (2008) Dietary Korean mistletoe enhances cellular non-specific immune responses and survival of Japanese eel (*Anguilla japonica*). *Fish Shellfish Immunol.*, **24**, 67–73.
- [5] Covello, J.M., Friend, S.E., Purcell, S.L., Burka, J.F., Markham, R.J.F., Donkin, A.W., Groman, D.B. & Fast, M.D. (2012) Effects of orally administered immunostimulants on inflammatory gene expression and sea lice (*Lepeophtheirus salmonis*) burdens on Atlantic salmon (*Salmo salar*). *Fish Shellfish Immunol.*, **366-367**, 9-16.
- [6] Cuesta, A., Rodriguez, A., Esteban, M.A. & Meseguer, J. (2005) *In vivo* effects of propolis, a honeybee product, on gilthead seabream innate immune responses. *Fish Shellfish Immunol.*, **18**, 71-80.
- [7] Davila, H., Didoli, G., Bottasso, O. & Stanford, J. (2011) Maternal immunization with Actinomycetales immunomodulators reduces parasitaemias in offspring challenged with *Trypanosoma cruzi*. *Immunotherapy*, **3**, 577-583.
- [8] Demers, N.E. & Bayne, C.J. (1997) The immediate effects of stress on hormones and plasma lysozyme in rainbow trout. *Develop. Comp. Immunol.*, **21**, 363–373.
- [9] Fontanella, G.H., Pascutti, M.F., Daurelio, L., Perez, A.R., Nocito, A.L., Wojdyla, D., Bottasso, O., Revelli, S.S. & Stanford J.L. (2007) Improved outcome of *Trypanosoma cruzi* infection in rats following treatment in early life with suspensions of heat-killed environmental Actinomycetales. *Vaccine*, **25**, 3492–3500.
- [10] Heidarieh, M., Mirvaghefi, A.R., Akbari, M., Farahmand, H., Sheikhzadeh, N., Shahbazfar, A.A. & Behgar, M. (2012) Effect of dietary Ergosan on growth performance, digestive enzymes, intestinal histology, hematological parameters and body composition of rainbow trout (*Oncorhynchus mykiss*). *Fish Physiol. Biochem.*, **38**, 1169-1174.

- [11] Heidarieh, M., Mirvaghefi, A.R., Sepahi, A., Sheikhzadeh, N., Shahbazfar, A.A. & Akbari, M. (2013) Effects of dietary Aloe vera on growth performance, skin and gastrointestinal morphology in rainbow trout (*Oncorhynchus mykiss*). *Turk. J. Fish Aquat. Sci.*, **13**, 367-373.
- [12] Huntigford, F.A. (2004) Implications of domestication and rearing conditions for the behaviour of cultivated fishes. *J. Fish. Biol.*, **65**, 122-142.
- [13] Kruger, N.J. (1996) The Bradford method for protein quantitation. In: The protein protocols handbook. (Walker, J.M. ed), pp. 11–15. Humana Press, Totowa, NJ, USA.
- [14] Marro, A., Pirles, M., Schiaffino, L., Bin, L., Davila, H., Bottasso, O.A., McIntyre, G., Ripley, P.R., Stanford, C.A. & Stanford, J.L. (2011) Successful immunotherapy of canine flea allergy with injected Actinomycetales preparations. *Immunotherapy*, **3**, 971-978.
- [15] Nootash, S., Sheikhzadeh, N., Baradaran, B., Khani Oushani, A., Maleki Moghadam, M.R., Nofouzi, K., Monfaredan, A., Aghebati, L., Zare, F. & Shabanzadeh, S. (2013) Green tea (*Camellia sinensis*) administration induces expression of immune relevant genes and biochemical parameters in rainbow trout (*Oncorhynchus mykiss*). *Fish Shellfish Immunol.*, **35**, 1916-1923.
- [16] Price, E.O. (1999) Behavioral development in animals undergoing domestication. *Appl. Anim. Behav. Sci.*, **65**, 245–271.
- [17] Rao, V. & Chakrabarti, R. (2005) Stimulation of immunity in Indian major carp *Catla catla* with herbal feed ingredients. *Fish Shellfish Immunol.*, **18**, 327-334.
- [18] Ringø, E., Olsen, R.E., Mayhew, T.M. & Myklebust, R. (2003) Electron microscopy of the intestinal microflora of fish. *Aquaculture*, **227**, 395–415.
- [19] Ross broiler management handbook. Aviagen, UK 2014. Page 1-130. http://en.aviagen.com/assets/Tech_Center/Ross_Broiler/Ross-Broiler-Handbook-2014i-EN.pdf

- [20] Salinas, I., Rosales, P.D., Cuesta, A., Meseguer, J., Chabrillo'n, M., Morin'igo, M.A. & Esteban, M.A. (2006) Effect of heat-inactivated fish and non-fish derived probiotics on the innate immune parameters of a teleost fish (*Sparus aurata* L.). *Vet. Immunol. Immunopathol.*, **111**, 279-286.
- [21] Sanden, M., Krogdahl, A., Bakke-Mckellep, A.M., Buddington, R.K. & Hemre G.I. (2006) Growth performance and organ development in Atlantic salmon, *Salmo salar* L. parr fed genetically modified (GM) soybean and maize. *Aquacult. Nutr.*, **12**, 1-14.
- [22] Sheikhzadeh, N., Heidarieh, M., Karimi Pashaki, A., Nofouzi, K., Ahrab Farshbafi, M. & Akbari, M. (2012) HilysesTM, fermented *Saccharomyces cerevisiae*, enhances the growth performance and skin non-specific immune parameters in rainbow trout (*Oncorhynchus mykiss*). *Fish Shellfish Immunol.*, **32**, 1083-1087.
- [23] Shin, C.H., Cha, J.H., Rahimnejad, S., Jeong, J.B., Yoo, B.W., Lee, B.K., Ahn, H.J., Choi, S.I., Choi, Y.J., Park, Y.H., Kim, J.D. & Lee, K.J. (2014) Effects of dietary supplementation of Barodon, an anionic alkali mineral complex, on growth performance, feed utilization, innate immunity, goblet cell and digestibility in olive flounder (*Paralichthys olivaceus*). *Asian Australas J. Anim. Sci.*, **27**, 383-390.
- [24] Siwicki, A.K., Anderson, D.P. & Rumsey, G.L. (1994) Dietary intake of immunostimulants by rainbow trout affects non-specific immunity and protection against furunculosis. *Vet. Immunol. Immunopathol.*, **41**, 125–139.
- [25] Smaldini, P., Stanford, J., Fossati, C.A. & Docena, G.H. (2013) Modulatory effect of Actinomycetales genera on activation of gut epithelial cells and inhibition of Nuclear Factor-kB pathway. *Innate Immun.*, **20**, 626-638.
- [26] Son, V.M., Chang, C.C., Wu, M.C., Guu, Y.K. & Chiu, C.H. (2009) Dietary administration of the probiotic, *Lactobacillus plantarum*, enhanced the growth, innate immune responses, and disease resistance of the grouper *Epinephelus coioides*. *Fish Shellfish Immunol.*, **26**, 691–698.

- [27] Stanford, J. & Stanford, C. (2012) Mycobacteria and their world. *Int. J. Mycobiol.*, **1**, 3-12.
- [28] Tarrés, M.C., Gayol, M.D.C., Picena, J.C., Alert, N., Bottasso, O., McIntyre, G., Stanford, C. & Stanford, J. (2012) Beneficial effects of immunotherapy with extracts derived from Actinomycetales on rats with spontaneous obesity and diabetes. *Immunotherapy*, **4**, 1-11.
- [29] Torrecillas, S., Makol, A., Caballero M.J., Montero, D., Gines, R., Sweetman, J. & Izquierdo, M. (2011) Improved feed utilization, intestinal mucus production and immune parameters in sea bass (*Dicentrarchus labrax*) fed mannan oligosaccharides (MOS). *Aquat. Nutr.*, **17**, 223-233.
- [30] Xueqin, J., Kania, P.W. & Buchmann, K. (2012) Comparative effects of four feed types on white spot disease susceptibility and skin immune parameters in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *J. Fish Dis.*, **35**, 127-135.

Table 1 Ingredients composition of **the commercial basal diet** used in this study

Ingredients (g kg⁻¹, DM)	Control diet
Fish meal ^a	195
Soybean oil ^b	41
Corn flour	86.5
Soybean meal	200
Wheat	173
Poultry byproduct meal	270
Vitamin mixture ^c	5
Mineral mixture ^d	5
Zeolit	10
L-Carnithine ^e	0.3
Vitamin C ^f	0.2
Molasses	10
Lysine	0.6
Methionine	0.6
Active-mos ^g	1.5
Natuzyme-p ^h	0.3

NaCl	1
Proximate chemical analysis (g kg⁻¹)	
Dry matter	910
Crude protein	430
Crude lipid	178
Ash	90
Crude fiber	42
Phosphorus	14

^a North Sefid Mahi Corporation, Iran with 540 g kg⁻¹ crude protein

^b North Sefid Mahi Corporation, Iran

^c Unit kg⁻¹ of mixture: Vitamin: retinol acetate (A), 1 600 000 IU; Cholecalciferol (D3), 400 000 IU; DL- α -tocopheryl acetate (E), 40 IU; Menadione sodium bisulfate (K3), 2 000 mg; Biotin (H2), 240 mg; Thiamin mononitrate (B1), 6 000 mg; Riboflavin (B2), 8 000 mg; Calcium d-pantothenate (B3), 12 000 mg; Niacinamide (B5), 40 000 mg; Pyridoxine hydrochloride (B6), 4 000 mg; Folic acid (B9), 2 000 mg; Cyanocobalamin (B12), 8 000 mg; Vitamin C, 60 000 mg; Inositol, 20 000 mg; B.H.T., 20 000 mg; Carrier up to 1 kg

^d Unit kg⁻¹ of mixture: Mineral: Fe, 26 000 mg; Zn, 12 500 mg; Se, 2 000 mg; Co, 480 mg; Cu, 4 200 mg; Mn, 15 800 mg; I, 1 000 mg; Choline chloride, 12 000 mg; Carrier up to 1 kg

^e Science Laboratories, Iran. Contains: 50 g L-Carnithin, Carrier up to 100 g

^f Canavit Company, Canada

^g Biorigin Company, Brazil. Contains: 100% yeast cell wall

^h T.F.A. Company, Iran. Conatins Unit kg⁻¹ of mixture: Cellulase, 6 000 000 U; Xylanase, 10 000 000 U; β-glucanase, 700 000 U; α-amylase, 700 000 U; .Pectinase 70 000 U; Phytase, 1 500 000 U; Lipase, 30 000 U; Proteases (Acid & Neutral), 3 000 000 U

Table 2 Growth performance in rainbow trout after feeding different doses of heat-killed *Gordonia bronchialis*

<i>Gordonia bronchialis</i> in diet (cells kg ¹	Initial Weight (g)	Initial Length (cm)	Final Weight (g)	Final Length (cm)	Feed Conversion Ratio	Thermal Growth Coefficient
---	--------------------------	---------------------------	------------------------	-------------------------	-----------------------------	----------------------------------

feed)					(FCR)	(TGC)
Control	31.05 ± 0.29	13.99 ± 0.03	135.82 ± 3.40 ^a	22.90 ± 0.27 ^a	0.91 ± 0.04 ^a	1.75 ± 0.05 ^a
2.48 × 10⁸	30.98 ± 0.23	13.99 ± 0.03	149.76 ± 1.62 ^b	24.08 ± 0.23 ^b	0.80 ± 0.01 ^b	1.91 ± 0.01 ^b
1.24 × 10⁹	30.89 ± 0.20	13.97 ± 0.03	149.83 ± 2.13 ^b	24.15 ± 0.16 ^b	0.80 ± 0.01 ^b	1.90 ± 0.01 ^b

Data are mean ± SEM. Those within a column superscripted by * are significantly different (P<0.05); for FCR and TGC n= 3 in each treatment group, for remaining parameters n = 54 in each treatment group.

Table 3 Immunological parameters in rainbow trout serum after feeding different doses of heat-killed *Gordonia bronchialis*

Gordonia bronchialis in diet (cells kg ⁻¹ feed)	Complement titer (units mL ⁻¹)		Total antibody (mg mL ⁻¹)		Peroxidase (450 nm)		Lysozyme activity (µg mL ⁻¹)	
	Day 95	Day 105	Day 95	Day 105	Day 95	Day 105	Day 95	Day 105
Control	0.075 ± 0.004 ^a	0.081 ± 0.001 ^a	3.17 ± 0.20	3.10 ± 0.14	0.256 ± 0.002	0.249 ± 0.001	33.01 ± 6.77 ^a	34.89 ± 1.50 ^a
2.48 × 10⁸	0.149 ± 0.009 ^b	0.138 ± 0.007 ^b	3.12 ± 0.19	2.99 ± 0.23	0.262 ± 0.003	0.260 ± 0.002	112.30 ± 19.55 ^b	115.60 ± 14.51 ^b
1.24 × 10⁹	0.141 ± 0.012 ^b	0.136 ± 0.007 ^b	3.08 ± 0.20	3.09 ± 0.19	0.257 ± 0.002	0.254 ± 0.002	102.00 ± 8.92 ^b	107.09 ± 8.93 ^b

Data are mean ± SEM. Means within a column superscripted by different small letters are significantly different (P<0.05); Means within a row superscripted by different capital letters are significantly different (P<0.05); n = 12 for each treatment group.

Table 4 Intestinal morphology in rainbow trout after feeding different doses of heat-killed *Gordonia bronchialis*

Gordonia bronchialis in diet (cells kg⁻¹ feed)	Fold length (µm)		Fold thickness (µm)		Goblet cell percentage	
	Day 95	Day 105	Day 95	Day 105	Day 95	Day 105
Control	432.50 ± 17.10 ^a	480.10 ± 19.88 ^a	116.66 ± 4.97	112.08 ± 4.32 ^a	33.50 ± 0.98 ^{aA}	41.58 ± 2.00 ^{aB}
2.48 × 10⁸	465.20 ± 19.48 ^a	470.08 ± 13.70 ^a	122.50 ± 6.41	124.16 ± 4.99 ^b	39.50 ± 0.88 ^b	42.25 ± 0.87 ^a
1.24 × 10⁹	570.10 ± 13.25 ^b	570.01 ± 12.49 ^b	127.50 ± 6.97	126.66 ± 5.55 ^b	46.33 ± 1.22 ^c	46.50 ± 0.96 ^b

Data are mean ± SEM. Means within a column superscripted by different small letters are significantly different (P<0.05); Means within a row superscripted by different capital letters are significantly different (P<0.05); n = 6 for each treatment group.

Table 5 Pyloric caeca morphology rainbow trout after feeding different doses of heat-killed *Gordonia bronchialis*

<i>Gordonia bronchialis</i> in diet (cells⁻¹ kg feed)	Fold length (µm)		Fold thickness (µm)		Goblet cell percentage	
	Day 95	Day 105	Day 95	Day 105	Day 95	Day 105
Control	727.50 ± 21.07 ^a	780.83 ± 25.41 ^a	115.10 ± 3.37 ^a	120.83 ± 3.12	28.50 ± 1.24 ^{aA}	34.67 ± 1.63 ^{aB}
2.48 × 10⁸	761.66 ± 17.44 ^a	728.50 ± 27.17 ^a	133.33 ± 5.41 ^b	121.66 ± 6.49	33.42 ± 0.82 ^b	37.92 ± 1.05 ^b

1.24×10^9	833.33 $\pm$ 24.72 ^b	918.33 $\pm$ 28.11 ^b	141.66 $\pm$ 4.40 ^b	125.83 $\pm$ 4.51	40.50 $\pm$ 1.33 ^c	41.02 $\pm$ 1.41 ^c
--------------------------------------	------------------------------------	------------------------------------	-----------------------------------	----------------------	-------------------------------	-------------------------------

Data are mean $\pm$ SEM. Means within a column superscripted by different small letters are significantly different ($P < 0.05$); Means within a row superscripted by different capital letters are significantly different ($P < 0.05$); n = 6 for each treatment group.

Figure 1 Hemagglutination antibody titer in rainbow trout after feeding different doses of heat-killed *Gordonia bronchialis*. Data are mean ± SEM. Those superscripted by different letters are significantly different ($P < 0.05$); $n = 18$ for each treatment group.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Sheikhzadeh, N;Tahapour, K;Nofouzi, K;Tukmechi, A;Shabanzadeh, S;Oushani, AK;Stanford, J;McIntyre, G;Mardani, K

Title:

Can heat-killed *Gordonia bronchialis* enhance growth and immunity in rainbow trout (*Oncorhynchus mykiss*)?

Date:

2017-08-01

Citation:

Sheikhzadeh, N., Tahapour, K., Nofouzi, K., Tukmechi, A., Shabanzadeh, S., Oushani, A. K., Stanford, J., McIntyre, G. & Mardani, K. (2017). Can heat-killed *Gordonia bronchialis* enhance growth and immunity in rainbow trout (*Oncorhynchus mykiss*)?. *AQUACULTURE NUTRITION*, 23 (4), pp.788-795. <https://doi.org/10.1111/anu.12445>.

Persistent Link:

<http://hdl.handle.net/11343/291630>