Received Date : 26-Mar-2016

Revised Date : 28-Jun-2016

Accepted Date : 20-Aug-2016

Article type : General Article

EVJ-GA-16-075.R1

Risk factors in equine transport-related health problems: a survey of the Australian equine industry.

B. Padalino^{1,2,3*}, S. L. Raidal³, E. Hall¹, P. Knight⁴, P. Celi^{5,6}, L. Jeffcott¹ and G. Muscatello¹

¹School of Life and Environmental Sciences, The Faculty of Veterinary Science, The University of Sydney, Camden, NSW, Australia;

²Department of Veterinary Medicine, The University of Bari, Italy, EU;

³School of Animal and Veterinary Sciences, Charles Stuart University, Wagga Wagga, NSW, Australia;

⁴Discipline of Biomedical Science, School of Medical Sciences, Sydney Medical School, University of Sydney, Australia;

⁵DSM Nutritional Products, Animal Nutrition and Health, Columbia, Maryland, USA;

⁶Faculty of Veterinary and Agricultural Sciences, The University of Melbourne, Parkville, VIC, Australia.

***Corresponding author email:** barbara.padalino@sydney.edu.au

Keywords: horse; transportation; illness; risk; journey duration; survey.

Summary

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/evj.12631](https://doi.org/10.1111/evj.12631)

This article is protected by copyright. All rights reserved

Background: Transportation can affect equine health and is a potential source of economic loss to the industry.

Objectives: To identify journey (duration, vehicle, commercial/non-commercial) and horse (sex, age, breed, use, amateur or professional status) characteristics associated with development of transport-related health problems.

Study design: Cross-sectional online survey.

Methods: An online survey was conducted targeting amateur and professional participants in the Australian equine industry; eligible respondents organised horse movements at least monthly. They provided details of the last case of a transport-related health problem that had affected their horse(s). Associations between type of health problem, journey and horse characteristics were examined with multivariable multinomial regression analysis.

Results: Based on 214 responses, health problems were classified as injuries, muscular problems, heat stroke, gastrointestinal and respiratory problems, or death/euthanasia. Respiratory problems were reported most frequently (33.7%) followed by gastrointestinal problems (23.8%) and traumatic injuries (16.3%). The type of health problem was associated with journey-duration ($P < 0.001$) and horse breed ($P = 0.001$). Injuries were more likely to occur on short journeys, while more severe illnesses (gastrointestinal and respiratory problems or death/euthanasia) were more likely to occur on long journeys. Using Standardbreds as the reference group, Thoroughbreds, Arabians and Warmbloods were more likely to experience a severe illness than injury.

Main limitation: Self-selected participation in the study and self-reported nature of transport-related problems.

Conclusion: Horses undertaking journeys longer than 24 hours were at greater risk of developing severe disease or dying. Further studies are needed on long-haul transportation effects to safeguard the welfare of horses moved over long distances.

Introduction

Transportation is an integral part of many horse related activities, with horses being moved frequently [1] and for a wide range of reasons [2]. Horses transported frequently may become habituated to travel [3]. However, for other horses, the challenges associated with transport, such as confinement, noise and vibration [4] may trigger fear [5]. The physical and mental stressors

associated with transportation can result in adverse effects on the horses' health [6]. Fear may also trigger behaviours that put horses at risk of injury [7], ranging from small abrasions to catastrophic fractures [8]. The stress associated with transport, and the way in which horses are managed during transport [9-11], can contribute also to the development of potentially fatal infections of the respiratory [12] or gastrointestinal systems [13].

Transport's effects on physiological [14], behavioural [15], endocrine [16], reproductive [17], muscular [18], gastric [19], inflammatory [20] and respiratory parameters [9] have been described in many observational studies. In contrast, few epidemiological studies have been conducted. The incidence of and risk factors for health problems has been reported in horses being transported for slaughter [21-23]. The only epidemiological study on risk factors for the development of transport-related health problems during long haul transportation of horses for other purposes identified journey duration (more than 20 hours) and season (spring) as risk factors. However, the study reported only 47 cases [24]. Consequently, the relationship between journey and horse characteristics, and the risk of developing disease remains poorly defined. We collected data with an online questionnaire and examined associations between transport related health problems and journey and horse characteristics across a number of Australian equine industry sectors.

Material and methods

Study design and data collection

The study was a cross-sectional online survey conducted in Australia from June to September 2015. Detail of the design and distribution of the cross sectional survey and the description of the demographic characteristics of the study population have been reported previously [25]. Briefly, the survey was digitised using SurveyMonkey^a. The target population was people involved in any equine industry sector who had organised or participated in the movement of horses (commercially or non-commercially) at least monthly over the past two years. The respondents classified themselves as either professionals (people who were involved for financial reward, such as trainers, stud/farm managers) or amateurs (people involved as a hobby, such as riders, owners). An invitation letter and the link to the survey (<https://www.surveymonkey.com/r/SM9F9SJ>) were provided to several Australian horse sport

associations and were published on their websites. The link was also promoted through several horse magazines, relevant Facebook pages and online horse forums.

The questionnaire was divided into four parts: respondent details; management strategies pre-, during and post-transport; transport-related behavioural and health problems identified during and after transportation in the past two years; details of the most recent case including horse sex, age, breed, the use of the horse, the vehicle in which the horse was transported, the journey duration, and whether the horse was moved by a commercial or non-commercial carrier. The results for the first three parts are presented elsewhere [25]. The data collected within the first and fourth part of the questionnaire (Supplementary Item 1) are presented in the current report.

Risk factors and outcome

Horse-level predictive variables were sex (mare/filly, gelding, stallion/colt), age (8-24 months, 2-5 years, 6-10 years, >10 years), breed (Arabian, Quarter horse, Standardbred, Thoroughbred, Warmblood, use (breeding, recreational non-competitive activities, Standardbred racing, Thoroughbred racing, endurance, equestrian competitive sport), and amateur or professional status.

Journey-level predictive variables were categorised according to the type of vehicle used (truck, horse trailer) and operator (commercial versus non-commercial transporter). The journey duration was categorised as: short (less than 8 hours), intermediate (8-24 hours) and long (more than 24 hours). These cut-offs were chosen on the basis of the European and Australian Code of Animal Transportation, in which the maximum journey durations without watering are of 8 and 24 hours respectively [26; 27]. In Australia a rest stop of 12 hours is recommended after 12 and compulsory after 24 hours of journey [26]. Thus all reported multiday trips in this dataset included mandatory rest stops.

To categorise outcome, there was an open question in which the respondents reported a description of the signs and their veterinarians' diagnosis of any transport-related health problem that affected the horse. Based on the respondent's description, health problems were classified into six categories (injuries, muscular problems, heat stroke, gastrointestinal problems, respiratory problems, death/euthanasia) by an experienced equine veterinarian (B.P.; Table 1).

Data analysis

Initial descriptive analysis was undertaken using statulator^{beta} (<http://statulator.com/descriptive.html>). Associations between the predictive variables were explored using Contingency tables and χ^2 tests using GenStat[®] Version 14^b. A multivariable multinomial regression analysis was constructed using SPSS Version 22^c with health problem category as outcome with injuries as the reference level for comparisons. Vehicle and operator were excluded as they were found to be collinear with journey duration. Horse age, breed, sex, use, amateur or professional status and journey-duration were considered for inclusion in the final multinomial model. A stepwise backward elimination was then conducted to remove the least significant variable one at a time until all variables within the model had P value < 0.05. The data met with all assumptions for multinomial regression including that of IIA. The findings are presented as odds ratio (OR), confidence interval (95% CI) and P value for each predictive variable value.

Results

Population

Of the 797 responses to the survey, 214 included details of a transport-related health problem and these 214 records make up the database. The distribution of the data (including missing values) is reported in Supplementary Item 2.

The frequency of the health outcomes according to the predictive variables is shown in Supplementary Item 3. Ten horses died during transit: 8 were humanely destroyed due to fractures (5 limb, 1 pelvis, and 2 neck), and 2 were found dead. A further 15 horses were humanely destroyed within one week after the journey due to colic (5 cases), colitis (5 cases), and pneumonia (5 cases). Additionally, 7 deaths occurred within 24 hours after arrival, of which 5 underwent post mortem examination, one was diagnosed with water intoxication; no cause of death was identified in the other four cases.

Journey variables

Journey duration was associated with vehicle (Pearson chi-square: 71.51, df = 2; P<0.001) and transport by a commercial company (Pearson chi-square: 78.74, df = 2; P<0.001). Companies transported fewer horses on short journeys and more horses on long journeys. The number of horses moved by horse trailer was larger for short journeys, and smaller for long journeys (Supplementary Item 4).

Factors associated with Health Outcomes

The final multivariate multinomial model for risk factors associated with health problems included journey-duration (χ^2 : 88.153, df:10, $P < 0.001$) and breed (χ^2 : 46.087, df:20, $P = 0.001$). None of the other predictive variables considered for inclusion reached significance. Figure 1 shows the distribution of the different transport-related illness according to the journey duration category. Using injuries as reference, death/euthanasia (odds ratio, OR: 101.6, 95% confidence interval (CI):10.2-1010.5, $P < 0.001$), gastrointestinal (OR:14.2, CI:1.5-133.8, $P = 0.02$) and respiratory (OR:113.9; CI:12.2-1060.7, $P < 0.001$) problems were more likely to occur on long journeys than on short journeys. Respiratory problems were also more likely (OR: 15.7, CI:4.3-56.7, $P < 0.001$) to occur on intermediate journeys than on short journeys. Using the injury group as the reference, muscular problems were more likely to occur on an intermediate journey than on a short one (OR:5.8, CI:1.1-29.5, $P = 0.03$). There was no significant difference among the journey-duration categories comparing injuries versus heat stroke (Supplementary Item 5). Using injuries as reference group, gastrointestinal problems were more likely to occur in Arabians (OR: 95.8, CI: 4.6-1990.3, $P = 0.003$) and Warmbloods (OR: 43.0, CI: 3.8-485.9, $P = 0.002$) compared with Standardbreds. Respiratory problems were more likely to occur in Arabians (OR: 20.8, CI: 1.2-345.2, $P = 0.034$), Warmbloods (OR: 18.5, CI: 2.5-136.89, $P = 0.004$), and Thoroughbreds (OR: 7.4, CI: 1.2-45.7, $P = 0.031$) compared with Standardbreds. Death/euthanasia was more likely to occur in Thoroughbreds than in Standardbreds (OR; 7.5, CI: 1.0-56.0, $P = 0.048$) (Supplementary Item 5).

Discussion

This is the first study to investigate whether journey and horse characteristics were associated with transport-related health problems across a diverse range of Australian horses used for various activities undertaking different journeys. Journey duration and breed were identified as risk factors for the development of transport-related health problems, while horse sex, age, use and amateur or professional status were not predictors. The main finding of this study was the association between journey-duration and the nature of transport-related health problems, confirming that journeys longer than 24 hours pose the greatest risk of horses having serious health outcomes [24; 28]. The association between health problem category and breed should be considered preliminary, and warrants future research using a larger dataset.

The observation that shorter trips are associated with a higher risk of injury is in agreement with previous reports. In an epidemiological study conducted in Australia, injuries occurred more often at the beginning of a 3.5 day journey and that they were often related to behavioural problems [24]. Previous studies have also identified that behavioural problems and movement of the horse within the vehicle are greatest during the first hour of transportation, and that horses become habituated after 5 hours of transport [14; 17; 29; 30]. The higher risk of injuries during short trips is likely to be associated with behavioural problems and lack of habituation.

In this study muscular problems were reported to occur more often during non-commercial transport of non-racing horses, and the risk of muscular problem was greater for intermediate journeys compared with short journeys. The reasons for this cannot be ascertained from the available data, although it could be speculated that limitations in driver ability and in horse fitness could have contributed. In an electromyographic study on the effects of transportation on muscle, horses transported by less experienced drivers required more muscular effort to maintain balance compared with horses transported by expert drivers [18]. In another study it was found that the effort required to maintain balance during a 300 km journey had the same impact on muscles as a 1,500 metres canter, and caused a comparable increase in serum muscle enzyme activities [31]. The effects of journey-duration on muscular problems warrant future research.

Respiratory diseases were the most commonly identified problem in this study, a finding that agrees with previous studies [32; 33]. In our study the likelihood of respiratory problems was approximately 15 times greater on intermediate journeys and approximately 100 times greater on long journeys than on short journeys. The relationship between duration of transport and incidence of respiratory disease has been previously reported [6; 12; 34; 35] and our data confirm it. This relationship may relate to the head position of the transported horses, vehicle ventilation or air quality. If horses are restrained in a way that prevents them lowering their head, mucociliary clearance will be adversely affected [9]. Ventilation may be inadequate in many types of vehicles [36] resulting in an accumulation of dust, bacteria and noxious gases in the vehicle as journey length increases [11]. The ongoing high incidence of respiratory diseases suggest that more research is needed to identify how ventilation systems can be improved and how any periods of enforced head elevation can be shortened.

In agreement with our previous study [24], the risk of gastrointestinal and respiratory disorders and death/euthanasia was greater than the risk of injury for journeys longer than 24 hours. In

livestock the association between adverse outcomes and journey duration is influenced by the physiological and clinical state of the animal before and during journey, the management of feeding and watering, the opportunities animals have to rest and the thermal environment rather than journey length per se [28]. These factors may also be important in determining whether horses experience adverse outcomes as a result of transport. However, the reasons why some horses develop fatal diseases during and after a multi-day journey, while others remain healthy under the same conditions are unknown. Protracted transport stress may compromise the immune system and lead to psychological and physical exhaustion and death [35]. The use of immunostimulants before shipping has been found to be useful in reducing the incidence of transport-related pneumonia in horses transported for more than 24 hours [37]. Thus, the higher risk of severe diseases in horses transported for longer than 24 hours might be related to immunosuppression, and the relationship between long journeys and the immune system requires further investigation.

Compared with Standardbreds, Thoroughbred, Arabian and Warmblood horses were more likely to develop gastrointestinal and respiratory diseases than to be injured during transportation. Thoroughbreds were found at higher risk of transport pleuropneumonia in a previous study [12]. Arabians have been found at higher risk of colic compared with other breeds [38]. There might therefore be a breed-predisposition for developing a particular type of transport-related diseases. However, our data should be considered preliminary and a larger data set would be required to determine the actual effects of breed on different types of transport related illnesses.

This study has a number of limitations that must be considered in interpreting the results. The problems of bias associated with self-selected participation in the study could not be addressed, nor could the possibility of response bias in the answers provided. It was not possible to check the diagnoses reported, nor was there any standardisation of the processes by which the diagnoses were made. The target population was not estimated and it was not possible to calculate a response rate, a common problem with online surveys [39]. Notwithstanding these limitations, this is the first study which analysed 214 cases with a novel approach and our findings may be important in helping reduce the negative impact of transportation on horse health.

Conclusions

There is an association between transport-related health problems and journey-duration and the likelihood of developing a more severe illness (i.e. respiratory and gastrointestinal problem or death/euthanasia) was higher on journeys over 24 hours than on journeys of less than 8 hours, suggesting the need to decrease the maximum journey time in Australia. This study also highlights the need for further research into the effects of long haul transport on the respiratory, gastrointestinal and immune systems to assist in proposing improved management practices for safeguarding horse welfare during travel, particularly over long distances.

Authors' declaration of interests

No competing interests have been declared.

Ethical animal research

The ethical aspects of this study have been approved by the Human Research Ethics Committee of the University of Sydney [2015/308].

Source of funding

This study was funded by the University of Sydney International Postgraduate Research Scholarship (IPRS).

Acknowledgements

The authors would like to thank Goldners Horse Transport, Hunter Valley Thoroughbred Breeders Association, Equestrian Australia association, and the International Society of Equitation Science for emailing the link of the survey to their members. This project would not have been possible without the voluntary contribution of the respondents who gave their time to respond in the online survey. Thanks also to Jayne Ion for her technical assistance with SurveyMonkey program.

Authorship

B. Padalino, E. Hall, S. Raidal, P. Knight, P. Celi, L. Jeffcott, and G. Muscatello conceived and designed the survey; B. Padalino and E. Hall analysed the data, B. Padalino wrote the paper; E. Hall, S. Raidal, P. Knight, L. Jeffcott, and G. Muscatello edited the paper.

Manufacturers' addresses

^aSurveyMonkey Inc., California, USA. (www.surveymonkey.com)

^bVSNi International, Hemel Hempstead, Hertfordshire, UK.

^cIBM SPSS

Supplementary Information

Supplementary Item 1: Questionnaire.

Supplementary Item 2: Data distribution.

Supplementary Item 3: Health outcomes.

Supplementary Item 4: Associations between journey variables.

Supplementary Item 5: Associations between health outcomes and horse and journey variables.

References

[1] Rosanowski, S.M., Cogger, N., Rogers, C.W., Bolwell, C.F., Benschop, J. and Stevenson, M.A. (2013) Analysis of horse movements from non-commercial horse properties in New Zealand. *New Zealand Vet. J.* **61**, 245-253.

[2] Leadon, D.P. (1994) Studies of the effects of transporting horses: better to arrive than to travel. *Equine Vet. J.* **26**, 346-347.

- [3] Houpt, K.A. and Wickens, C.L. (2014) Handling and transport of horses. In: *Livestock Handling and Transport: Fourth Edition*. pp 315-341.
- [4] Padalino, B. (2015) Effects of the different transport phases on equine health status, behavior, and welfare: A review. *J. Vet. Behav.: Clinical Applications and Research* **10**, 272-282.
- [5] Grandin, T. and Shivley, C. (2015) How Farm Animals React and Perceive Stressful Situations Such As Handling, Restraint, and Transport. *Animals* **5**, 04-09.
- [6] Oikawa, M.A., Takagi, S. and Yashiki, K. (2004) Some aspects of the stress responses to road transport in thoroughbred horses with special reference to shipping fever. *J. Equine Sci.* **15**, 99-102.
- [7] Ferguson, D.L. and Rosales-Ruiz, J. (2001) Loading the problem loader: the effect of target training and shaping on trailer-loading behavior of horses. *J. Appl. Behav. Analysis* **34**, 409-423.
- [8] Mansmann, R.A. and Woodie, B. (1995) Equine transportation problems and some preventives: A review. *J. Equine Vet. Sci.* **15**, 141-144.
- [9] Raidal, S.L., Bailey, G.D. and Love, D.N. (1997) Effect of transportation on lower respiratory tract contamination and peripheral blood neutrophil function. *Aust. Vet. J.* **75**, 433-438.
- [10] Stull, C.L. and Rodiek, A.V. (2002) Effects of cross-tying horses during 24 h of road transport. *Equine Vet. J.* **34**, 550-555.
- [11] Oikawa, M., Hobo, S., Oyamada, T. and Yoshikawa, H. (2005) Effects of orientation, intermittent rest and vehicle cleaning during transport on development of transport-related respiratory disease in horses. *J. Comp. Path.* **132**, 153-168.

- [12] Austin, S.M., Foreman, J.H. and Hungerford, L.L. (1995) Case-control study of risk factors for development of pleuropneumonia in horses. *J. Am. Vet. Med. Assoc.* **207**, 325-328.
- [13] McClintock, S.A. and Begg, A.P. (1990) Suspected salmonellosis in seven broodmares after transportation. *Aust. Vet. J.* **67**, 265-267.
- [14] Tateo, A., Padalino, B., Boccaccio, M., Maggiolino, A. and Centoducati, P. (2012) Transport stress in horses: Effects of two different distances. *J. Vet. Behav.: Clinical Applications and Research* **7**, 33-42.
- [15] Padalino, B., Maggiolino, A., Boccaccio, M. and Tateo, A. (2012) Effects of different positions during transport on physiological and behavioral changes of horses. *J. Vet. Behav.: Clinical Applications and Research* **7**, 135-141.
- [16] Fazio, E., Medica, P., Cravana, C., Aveni, F. and Ferlazzo, A. (2013) Comparative endocrinological responses to short transportation of Equidae (*Equus asinus* and *Equus caballus*). *Anim. Sci. J.* **84**, 258-263.
- [17] Baucus, K.L., Squires, E.L., Ralston, S.L., McKinnon, A.O. and Nett, T.M. (1990) Effect of transportation on the estrous cycle and concentrations of hormones in mares. *J. Anim. Sci.* **68**, 419-426.
- [18] Giovagnoli, G., Trabalza Marinucci, M., Bolla, A. and Borghese, A. (2002) Transport stress in horses: An electromyographic study on balance preservation. *Livestock Production Science* **73**, 247-254.
- [19] McClure, S.R., Carithers, D.S., Gross, S.J. and Murray, M.J. (2005) Gastric ulcer development in horses in a simulated show or training environment. *J. Am. Vet. Med. Assoc.* **227**, 775-777.

- [20] Wessely-Szponder, J., Bełkot, Z., Bobowiec, R., Kosior-Korzecka, U. and Wójcik, M. (2015) Transport induced inflammatory responses in horses. *Polish J. Vet. Sci.* **18**, 407-413.
- [21] Stefancic, I. and Martin, D. (2005) Influence of transport conditions on animal welfare. In: *ISAH*, Warsaw, Poland. pp 148-152.
- [22] Marlin, D., Kettlewell, P., Parkin, T., Kennedy, M., Broom, D. and Wood, J. (2011) Welfare and health of horses transported for slaughter within the European Union Part 1: Methodology and descriptive data. *Equine Vet. J.* **43**, 78-87.
- [23] Roy, R.C., Cockram, M.S. and Dohoo, I.R. (2015) Welfare of horses transported to slaughter in Canada: Assessment of welfare and journey risk factors affecting welfare. *Can. J. Anim. Sci.* **95**, 509-522.
- [24] Padalino, B., Hall, E., Raidal, S., Celi, P., Knight, P., Jeffcott, L. and Muscatello, G. (2015) Health problems and risk factors associated with long haul transport of horses in Australia. *Animals* **5**, 1296-1310.
- [25] Padalino, B., Raidal, S., Hall, E., Knight, P., Celi, P., Jeffcott, L. and Muscatello, G. (in press) Survey of horse transportation in Australia: issues and practices. *Aust. Vet. J.* Doi: 10.1111/avj.12486
- [26] <http://www.animalwelfarestandards.net.au/land-transport/>.
- [27] <http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=uriserv:f83007>.
- [28] Nielsen, B.L., Dybkjær, L. and Herskin, M.S. (2011) Road transport of farm animals: effects of journey duration on animal welfare. *Animal* **5**, 415-427.

- [29] Fazio, E., Medica, P., Aronica, V., Grasso, L. and Ferlazzo, A. (2008) Circulating β -endorphin, adrenocorticotrophic hormone and cortisol levels of stallions before and after short road transport: Stress effect of different distances. *Acta Vet. Scand.* **50**, 1-7.
- [30] Schmidt, A., Möstl, E., Wehnert, C., Aurich, J., Müller, J. and Aurich, C. (2010) Cortisol release and heart rate variability in horses during road transport. *Hormones and Behavior* **57**, 209-215.
- [31] Codazza, D., Maffeo, G. and Redaelli, G. (1974) Serum enzyme changes and hematochemical levels in Thoroughbreds after transport and exercise. *J. S. African Vet. Assoc.* **45**, 331-335.
- [32] Leadon, D., Frank, C. and Backhouse, W. (1989) A preliminary report on studies on equine transit stress. *J. Equine Vet. Sci.* **9**, 200-202.
- [33] Leadon, D.P. and Hodgson, D.R. (2014) Transport of horses. In: *The Athletic Horse: Principles and Practice of Equine Sports Medicine: Second Edition*, Elsevier-Saunders, St. Louis, USA. pp 155-161.
- [34] Kohn, C.W. (2000) *Guidelines for horse transport by road and air*, AHSA/MSPCA.
- [35] Marlin, D.J. (2004) Transport of horses. In: *Equine Sports Medicine and Surgery*, Elsevier-Saunders, St. Louis, USA. pp 1239-1250.
- [36] Purswell, J.L., Gates, R.S., Lawrence, L.M., Jacob, J.D., Stombaugh, T.S. and Coleman, R.J. (2006) Air exchange rate in a horse trailer during road transport. *Transactions of the ASABE* **49**, 193-201.
- [37] Nestved, A. (1996) Evaluation of an immunostimulant in preventing shipping stress related respiratory disease. *J. Equine Vet. Sci.* **16**, 78-82.

[38] Cohen, N.D. and Peloso, J.G. (1996) Risk factors for history of previous colic and for chronic, intermittent colic in a population of horses. *J. Am. Vet. Med. Assoc.* **208**, 697-703.

[39] Christley, R. (2016) Questionnaire survey response rates in equine research. *Equine Vet. J.* **48**, 138-139.

Table 1: Categorisation of health problems observed in transported horses.

Health problem category	Definition
Injuries	Laceration, abrasion, contusion, swelling.
Muscular problems	Typing up, sore muscle, stiffness.
Heat stroke	Rectal temperature >38.5°C, sweating, lethargy.
Gastrointestinal problems	Oesophagal obstruction, gastric ulceration, diarrhoea, colic, enterocolitis.
Respiratory problems	Nasal discharge, coughing, inflammation/infection of the upper or lower respiratory tract, and pneumonia.
Death	Horses found dead or humanely destroyed.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Padalino, B;Raidal, SL;Hall, E;Knight, P;Celi, P;Jeffcott, L;Muscatello, G

Title:

Risk factors in equine transport-related health problems: A survey of the Australian equine industry

Date:

2017-07-01

Citation:

Padalino, B., Raidal, S. L., Hall, E., Knight, P., Celi, P., Jeffcott, L. & Muscatello, G. (2017). Risk factors in equine transport-related health problems: A survey of the Australian equine industry. EQUINE VETERINARY JOURNAL, 49 (4), pp.507-511. <https://doi.org/10.1111/evj.12631>.

Persistent Link:

<http://hdl.handle.net/11343/291844>