Introduction

The advent of immune checkpoint blockade is considered a revolution by some in cancer therapy progress. Nivolumab and pembrolizumab are anti PD-1 antibodies evaluated in randomised trials demonstrating improvements in survival in patients with metastatic melanoma, renal cell carcinoma and non-small cell lung cancer (1-3). Therapy is generally well tolerated and responses can be durable.

In the era of rapid dissemination of information regarding advances in cancer care, desperate patients are increasingly seeking to access drugs on the basis of early data. They are ill equipped to critique or contextualise the evidence of benefit of these therapies. The desire to access such high cost therapy is exacerbated by the understandable enthusiasm of the medical, pharmaceutical and general community for these therapies in the media.

Encouraging efficacy signals with immune checkpoint inhibitors have been observed for a number of tumour types in early phase trials (4-8) leading to patients with solid tumours pursuing access to self-funded pembrolizumab. The outcome for patients receiving this treatment for non-approved indications is unknown.

The aim of this retrospective study is to describe outcomes and associated toxicity of pembrolizumab in patients with non-melanoma solid organ malignancies treated at Chris O'Brien Lifehouse. These patients were not enrolled in clinical trials and consequently did not receive the drug under trial conditions.

Methods

Patients with non-melanoma solid organ malignancies who received or were planned to receive self-funded pembrolizumab were included. Pembrolizumab was administered intravenously at a dose of 2mg per kg of body weight every three weeks. Immune related adverse events (irAE) were graded according to the National Cancer Institute CTCAE, version 4.0 (9). Time to progression (TTP) was calculated from the date of pembrolizumab commencement until date of progression or death for patients that received at least one dose of pembrolizumab. Time to response (TTR) was calculated from the date of pembrolizumab commencement to the date of radiological response. Response was assessed as per RECIST 1.1 (10). Ethics committee approval was obtained for publication of this retrospective data (X15-0422; LH16.004).

Results

Between April and December 2015, 21 patients received or were planned to receive self-funded pembrolizumab. Patient and disease characteristics are described in Table 1. Malignancy type included 6 patients with a diagnosis of mesothelioma and 9 patients with sarcoma. The remaining 6

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/imj.13232](https://doi.org/10.1111/imj.13232)

patients had diagnoses of an undifferentiated carcinoma of the parotid, urothelial carcinoma, mucoepidermoid carcinoma, nasopharynx squamous cell cancer, non-small cell lung adenocarcinoma and an anal squamous cell cancer. The median age was 50, 57% had an Eastern Cooperative Oncology Group (ECOG) performance status of 0-1, 28% an ECOG of 2 and 10% had an ECOG of 3-4. Twenty-eight percent and 62% of patients had received 0-1 and at least 2-4 lines of prior treatment respectively. In 9% of patients the number of prior treatment lines was unknown but described as heavily pre-treated in the medical record. The median duration of follow up was 3.0 months (range, 0.4-9.6). At the time of last follow up, 11 patients had died.

The median number of pembrolizumab doses received was 3 (range 0-9), with 2 patients still receiving treatment at the time of last follow up and 1 patient undergoing a treatment break after an excellent metabolic response on Positron Emission Tomography (PET) scan.

Of the 21 patients, 14 (67%) requested pembrolizumab therapy and for 7 (33%) patients, this was offered by the clinician. Patients who requested pembrolizumab therapy had worse outcomes with a shorter TTP of 39 days (7-57) compared to 75 days (17-90) (Table 2).

Three patients died before receiving pembrolizumab. No patients achieved a complete response to pembrolizumab. Of the 18 patients that received at least 1 dose of pembrolizumab, a partial response (PR) was confirmed in 3 (17%). Fifteen of the 18 patients that received at least 1 dose of pembrolizumab (83%) had progressive disease (PD) (Table 3).

Patients achieving Partial Response

Three (17%) of 18 patients treated with pembrolizumab had a PR.

Patient 1:

A 65 year old male with undifferentiated carcinoma of the parotid with lung metastases. This patient had an ECOG performance status of 2 at the time of commencing pembrolizumab and had received 3 prior lines of therapy. He had a reduction in tumour burden of -84% determined by RECIST 1.1 (Figure 1). Time to response (TTR) was 1.7 months and is ongoing after 5.4 months of follow up. This patient developed a dermatological irAE of grade 1 severity.

Patient 2:

A 75 year old male with a urothelial malignancy with lung, retroperitoneal and mesenteric metastases. This patient had not received any prior lines of therapy as chemotherapy was contraindicated due to pre-existing myelodysplasia. He had an ECOG of 0 when commenced on pembrolizumab. He had an excellent metabolic response on PET with TTR of 2 months and is maintained after 6.5 months of follow up. This patient developed grade 1 dermatological and thyroid toxicity.

Treatment with pembrolizumab was ongoing for the above 2 patients at date of last follow up.

Patient 3:

A 25 year old male with a diagnosis of Ewing Sarcoma with lung and bone metastases. This patient had received multiple prior lines of therapy although the precise number was not documented. He had an ECOG performance status of 0 at commencement of pembrolizumab and achieved a sustained metabolic response on PET (11). TTR was 2 months and continues after 8.4 months of follow up. He developed grade 2 hypothyroidism requiring thyroxine replacement as an irAE. Due to disease response and significant financial costs of self-funding therapy, the patient received 9 cycles of pembrolizumab followed by observation with regular imaging to assess for disease progression.

Non-responders to pembrolizumab

Fifteen (83%) of the 18 patients that had received at least 1 dose of anti PD-1 therapy experienced PD. Median TTP was 39 days (range 7-90). Of these 15 patients, 5 received subsequent lines of anti-cancer therapy. These 5 patients had a median age of 31 (range 16-62) and an ECOG performance status ranging from 0-4. All had received prior lines of treatment (range 1-3). One had a temporary improvement in performance status, 3 remained clinically stable and 1 had symptomatic decline while receiving pembrolizumab.

The 9 patients that did not receive further lines of treatment had a median age of 62 (range 24-76) and ECOG performance status ranging from 0-2. They had received 0-4 prior lines of therapy. Eight patients had symptomatic decline with 4 experiencing rapid decline. In 1 patient clinical symptoms were not documented due to transfer of care overseas however this patient died less than 2 months after starting pembrolizumab.

In 1 patient, documentation of any subsequent treatment was not recorded.

Four patients received only 1 cycle of pembrolizumab (Table 4) and all experienced PD. These patients had received 2-4 prior lines of treatment. They had a baseline ECOG of 1-2 followed by rapid clinical decline. These 4 patients died a median of 27 days (range 13-43) after receiving a single dose of therapy.

Death before receiving pembrolizumab

Three (14%) patients died prior to receiving pembrolizumab (Table 2).

Patient 1:

A 61 year male with epithelioid pleural mesothelioma with diffuse peritoneal disease. He had received 3 prior lines of therapy. Pembrolizumab was requested by the patient. His ECOG performance status was 1 as per medical record documentation but disease symptoms were rapidly increasing and cachexia was present at the time of seeking to self-fund pembrolizumab. Best supportive care had been offered as well as palliative chemotherapy with gemcitabine as alternatives to self-funded pembrolizumab.

Patient 2:

A 47 year old male with Ewing Sarcoma with an ECOG of 2 at the time of discussing self-funded pembrolizumab. He had received 3 prior lines of therapy. Pembrolizumab was offered by the clinician. Within 4 weeks before his death, he was admitted with a symptomatic haemorrhagic brain metastasis causing focal neurology right sided weakness and seizures. He proceeded to craniotomy and resection followed by eventual decline in performance status and death.

Patient 3:

A 50 year old man with a mucoepidermoid carcinoma and a past history of glomerulonephritis. The patient had an ECOG of 1 documented within 3 weeks before his death and had received 2 prior lines of therapy. Pembrolizumab was requested by the patient. He experienced rapid progression of hepatic and brain disease. Liver derangement occurred with GGT elevated at 19 times the upper limit of normal (ULN) and AST 17 times the ULN equating to grade 3 severity. Possibly inefficacy of pembrolizumab had been discussed as well as not administering any further systemic therapy due to decline in clinical status.

Adverse Events

IrAE of any grade occurred in 6 (33%) of patients (Table 5). Toxicities reported were mild including 4 patients with skin toxicity, 2 patients with thyroid irAE and 1 patient experiencing joint and muscle toxicity. One patient with thyroid toxicity demonstrated a hyperthyroid phase followed by hypothyroid phase and the other required thyroxine replacement for grade 2 hypothyroidism. There were no grade 3-4 irAE.

Discussion

This retrospective review identified 21 patients with non-melanoma cancers that received or were planned to receive self-funded pembrolizumab at Chris O'Brien Lifehouse. Over two thirds of patients in our cohort were heavily pre-treated, having received at least 2-4 prior lines of therapy, and 38% had an ECOG of 2-4. There is little data regarding the use of pembrolizumab in patients with these poor prognostic characteristics. Initial data from KEYNOTE-001 assessed pembrolizumab in multiple solid cancers (including melanoma) with 50% of patients having received 3 or more prior therapies (5). However, no patients had an ECOG of 2-4 with 33% and 67% of patients having an ECOG of 0 and 1 respectively. In these initial 30 patients, 2 (7%) patients achieved a CR, 3 (10%) achieved a PR and 15 (50%) experienced SD.

In our cohort, 3 (17%) of patients who received at least one dose of pembrolizumab achieved meaningful responses. The responses were both radiologically and clinically significant and further demonstrate immunotherapy efficacy in multiple solid cancer types. Interestingly, an excellent metabolic response was seen in a patient with Ewing sarcoma. In early data from SARC028, PR rates to pembrolizumab were 19% for soft tissue sarcomas and 5% for bone sarcomas with no PR seen for

Ewing Sarcoma (8). No responses were seen in any patient with non-small cell lung cancer or mesothelioma in this cohort. No patient experienced grade 3-4 immune related toxicity and may reflect the short duration that pembrolizumab was prescribed.

Out of the 21 patients that sought to self-fund pembrolizumab, 3 (14%) patients died before the approved drug could be supplied and 4 (19%) died within 13-43 days of receiving a single dose of pembrolizumab. This outcome highlights several issues regarding the careful selection of patients, futility of anti-cancer therapy at the end of life and patient's perceived benefit of receiving this therapy.

Indicators of overly aggressive end-of-life cancer care have been identified with respect to chemotherapy use but are lacking for immunotherapy agents. This includes but is not limited to, receiving chemotherapy in the last 2 weeks of life, commencing a new line of chemotherapy in the last 30 days of life, the timing and delivery of hospice care (12). Cancer treatment in patients nearing death has been reported to be increasingly aggressive in an analysis of more than 28,000 patients monitored by the SEER registry (13). The use of chemotherapy within the last 4 weeks of life has been documented at rates of 12-18% (14, 15) and 8% within the last 2 weeks of life (15). The reasons for this are multifactorial and may include duration of palliative care involvement, gender (14) and individual clinician (15). With data for immunotherapy showing promise in solid organ malignancies (1-3), being better tolerated than cytotoxic agents with the possibility of durable responses, patients may feel compelled to pursue such therapy. In turn, this may delay appropriate transition to palliative care.

Despite documentation of best supportive care as an option to some of the patients in this cohort, patients persisted with seeking self-funded immunotherapy. We were not able to explore the level of understanding that patients had regarding their prognosis, perceived benefit of seeking self-funded therapy or possible futility. Nor were we able to document the level to which withdrawal of anti-cancer therapy was discussed. Medical futility and the complex circumstances' surrounding communicating this to patients has been described (16). It involves undertaking a clinical action when it is clear that this action will not achieve the proposed goal and is doomed to fail.

The discrepancy between patients and physicians estimates of survival has been documented (17). In a prospective study of 917 patients with advanced non-small cell lung or colon cancer, it was observed that patients overestimated their chance of surviving 6 months. Additionally, patients who believed they would survive for at least 6 months were more inclined to opt for life prolonging therapies rather than best supportive care. It is unknown if this influenced a proportion of the patients in our cohort but would seem plausible. It may also be simpler for physicians to offer another line of therapy than it is to have a difficult discussion around withdrawal of treatment (15, 18).

Economic harm from massive out of pocket costs is a distinct issue for patients desperate to fund non-approved therapies (19). Kantarjian et al describes the crossing of a moral line between

profiteering and the need for a fair price being applied to cancer drugs whereby it is affordable to those in need while reasonable profits are attained by the supplier (20). During the period that this cohort of patients self-funded their therapy through a cost-share program, the cost per 50mg vial of pembrolizumab was \$2,230. For the average 70kg patient, this equates to \$6,700 per cycle or \$20,000 for 3 cycles. Patients were required to finance 3 of the first 6 cycles of treatment and all cycles of treatment thereafter were charged. The current cost-share program charges \$1472 per 50mg vial with no free cycles. The degrees of financial burden by grades 1-5 (mild to catastrophic) has been described as a tool to identify patients at financial risk (19). Financial burden was not assessed in this cohort and although a difficult issue to raise when discussing a terminal illness, it has influenced treatment decisions with 1 patient achieving a partial response temporarily ceasing therapy as a consequence.

There are of course limitations of this data with our review being retrospective and descriptive with small patient numbers. Nevertheless, the information gleaned from this cohort demonstrates occasional activity of anti PD-1 therapy in non-approved solid cancers with a low toxicity profile and sheds light on the shortcomings and difficulties in the management of cancer patients at the end of life.

Acknowledgements

A J Lomax: Position supported in 2015 and 2016 by the Simon Kerr Memorial Fellowship

References

1. Robert C, Schachter J, Long GV, Arance A, Grob JJ, Mortier L, et al. Pembrolizumab versus Ipilimumab in Advanced Melanoma. *N Engl J Med*. 2015;372(26):2521-32.
2. Borghaei H, Paz-Ares L, Horn L, Spigel DR, Steins M, Ready NE, et al. Nivolumab versus Docetaxel in Advanced Nonsquamous Non-Small-Cell Lung Cancer. *N Engl J Med*. 2015 DOI: 10.1056/NEJMoa1507643.
3. Motzer RJ, Escudier B, McDermott DF, George S, Hammers HJ, Srinivas S, et al. Nivolumab versus Everolimus in Advanced Renal-Cell Carcinoma. *N Engl J Med*. 2015;DOI: 10.1056/NEJMoa1510665.
4. Alley EW, Molife LR, Santoro A, Beckey K, Yuan S, Cheng JD, et al. Abstract CT103: Clinical safety and efficacy of pembrolizumab (MK-3475) in patients with malignant pleural mesothelioma: Preliminary results from KEYNOTE-028. *Cancer Research*. 2015;75(15 Supplement):CT103.
5. Patnaik A, Kang SP, Rasco D, Papadopoulos KP, Elassaiss-Schaap J, Beeram M, et al. Phase I Study of Pembrolizumab (MK-3475; Anti-PD-1 Monoclonal Antibody) in Patients with Advanced Solid Tumors. *Clinical cancer research : an official journal of the American Association for Cancer Research*. 2015;21(19):4286-93. Epub 2015/05/16.
6. Topalian SL, Hodi FS, Brahmer JR, Gettinger SN, Smith DC, McDermott DF, et al. Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. *N Engl J Med*. 2012;366(26):2443-54.
7. Plimack ER, Bellmunt J, Gupta S, Berger R, Montgomery RB, Heath K, et al. Pembrolizumab (MK-3475) for advanced urothelial cancer: Updated results and biomarker analysis from KEYNOTE-012. *J Clin Oncol* 2015;33 (suppl; abstr 4502).
8. Tawbi H. Safety and efficacy of PD-1 blockade using pembrolizumab in patients with advanced soft tissue (STS) and bone sarcomas (BS): Results of SARC028—A multicenter phase II study. *ASCO Annual Meeting 2016;Abstract Number: 11006*
9. Common Terminology Criteria for Adverse Events (CTCAE) Version 4.0. Published: May 28, 2009 (v4.03: June 14, 2010).
10. Eisenhauer EA, Therasse P, Bogaerts J, Schwartz LH, Sargent D, Ford R, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer* 2009. 2009;45(2):228-47.
11. McCaughan GJ, Fulham MJ, Mahar A, Soper J, Hong AM, Stalley PD, et al. Programmed cell death-1 blockade in recurrent disseminated Ewing sarcoma. *J Hematol Oncol*. 2016;9(48).
12. Earle CC, Neville BA, Landrum MB, Souza JM, Weeks JC, Block SD, et al. Evaluating claims-based indicators of the intensity of end-of-life cancer care. *Int J Qual Health Care*. 2005;17(6):505-9.
13. Earle CC, Neville BA, Landrum MB, Ayanian JZ, Block SD, Weeks JC. Trends in the aggressiveness of cancer care near the end of life. *J Clin Oncol*. 2004;22(2):315-21.
14. Zdenkowski N, Cavenagh J, Ku YC, Bisquera A, Bonaventura A. Administration of chemotherapy with palliative intent in the last 30 days of life: the balance between palliation and chemotherapy. *Intern Med J*. 2013;43(11):1191-8.
15. Kao S, Shafiq J, Vardy J, Adams D. Use of chemotherapy at end of life in oncology patients. *Ann Oncol*. 2009;20(9):1555-9.

16. Kasman DL. When is medical treatment futile? A guide for students, residents, and physicians. *Journal of general internal medicine*. 2004;19(10):1053-6. Epub 2004/10/16.
17. Weeks JC, Cook E, O'Day SJ, et al. Relationship between cancer patients' predictions of prognosis and their treatment preferences. *JAMA*. 1998;279(21):1709-14.
18. Earle C, Landrum MB, Souza JM, Neville BA, Weeks JC, Ayanian JZ. Aggressiveness of cancer care near the end of life: is it a quality-of-care issue? *J Clin Oncol*. 2008;26(23):3860-6.
19. Carrera PM, Olver I. The financial hazard of personalized medicine and supportive care. *Support Care Cancer*. 2015;23(12):3399-401.
20. Kantarjian H, Steensma D, Rius SJ, Elshaug A, Light D. High cancer drug prices in the United States: reasons and proposed solutions. *J Oncol Pract*. 2014;10(4):e208-11.

Figure 1. a) Coronal view of exophytic tumour at the left angle of the mandible measuring 37mm (red arrow) b) poorly visualised after 4 months of pembrolizumab (blue arrow) c) axial view of tumour, 27x25mm (red arrow) d) difficult to appreciate after 4 months of pembrolizumab (blue arrow)

Table 1. Patient characteristics

Patient characteristics	Self-funded pembrolizumab n = 21	Patient requested n=14/21 (67%)	Clinician offered n=7/21 (33%)
Median age (range) - yr	50 (16-76)	61 (25-76)	43 (16-75)
Male sex – no. (%)	17 (81%)	11 (52%)	6 (29%)
ECOG performance status – no. (%)			
0-1	12 (57%)	9 (43%)	3 (14%)
2	6 (28%)	3 (14%)	3 (14%)
3-4	2 (10%)	1 (5%)	1 (5%)
Unknown	1 (5%)	1 (5%)	0
Pre-existing autoimmune condition	1 (5%)	1 (5%)	0
Malignancy type			
Mesothelioma*	6 (28%)	6 (28%)	0
Sarcoma [†]	9 (43%)	5 (24%)	4 (19%)
Other ^{††}	6 (28%)	3 (14%)	3 (14%)
Prior lines of treatment – no. (%)			
0-1	6 (28%)	5 (24%)	1 (5%)
2-4	13 (62%)	8 (38%)	5 (24%)
Unknown 'heavily pre-treated'	2 (9%)	1 (5%)	1 (5%)

*Epithelioid n=5; Sarcomatoid n=1

[†]Osteosarcoma n=3; Synovial sarcoma n=1; Ewing sarcoma n=3; Leiomyosarcoma n=1, Angiosarcoma n=1

^{††} Undifferentiated parotid n=1; Urothelial n=1; Mucoepidermoid n=1; Nasopharynx squamous cell n=1; NSCLC adenocarcinoma ca n=1; Anal squamous cell n=1

Table 2. Pembrolizumab therapy: Clinician offered vs Patient requested

Self-funded pembrolizumab n=21 (%)			
	Requested by patient 14/21 (67%)	Offered by clinician 7/21 (33%)	Total
Died before receiving pembrolizumab	2 (9%)	1 (5%)	3 (14%)
Partial Response (PR)	1 (5%)	2 (9%)	3 (14%)
Progressive Disease (PD)	11 (52%)	4 (19%)	15 (71%)
Median TTP/death days (range)	39 (7-57)	75 (17-90)	39 (7-90)

Table 3. Response to Treatment

Tumour response	Received ≥ 1 dose pembrolizumab n = 18
Best overall response at any assessment	
CR	0 (0%)
PR [*]	3 (17%)
SD	0 (0%)
PD ^{**}	15 (83%)

*RECIST 1.1 n=1, PET metabolic response n=2

**RECIST 1.1 n=7, PET n=4, clinical n=4

Table 4. Pembrolizumab cycles received

Planned to receive pembrolizumab n = 21	Outcome
Median number pembrolizumab cycles (range)	3 (0-9)*
Received 1 dose pembrolizumab only	4 (19%)
Median time from receiving 1 st dose to death, days (range)	27 (13-43)
Died before receiving 1 st dose pembrolizumab	3 (14%)

*n=2 treatment is ongoing, n=1 treatment break due to excellent response and financial cost

Table 5. Immune-related adverse events

irAE (received ≥ 1 dose pembrolizumab n=18)	Any grade (%)	Grade 3-4
Any	6 (33%)	0
Skin	4 (22%)	0
Thyroid*	2 (11%)	0
Joint and muscle	1 (6%)	0

*n=1 Thyroiditis (hyperthyroid followed by hypothyroid phase); n=1 Hypothyroid

Outcomes of patients with non-melanoma solid tumours receiving self-funded pembrolizumab at Chris O'Brien Lifehouse

A J Lomax¹, J Beith^{1,2,3}, V Bhadri^{1,2,3}, M Boyer^{1,2,3}, P Grimison^{1,2,3}, L G Horvath^{1,2,3}, S Kao^{1,2,3}, M Tattersall^{1,2,3}, D Thomas^{1,4} and C McNeil^{1,2,3}

¹Chris O'Brien Lifehouse, Camperdown, New South Wales, Australia

²Royal Prince Alfred Hospital, Camperdown, New South Wales, Australia

³University of Sydney, Camperdown, New South Wales, Australia

⁴Kinghorn Cancer Centre, Darlinghurst, New South Wales, Australia

Corresponding author:

Anna J Lomax

anna.lomax@lh.org.au

Chris O'Brien Lifehouse

119-143 Missenden Road

Camperdown NSW 2050

PO BOX M33 Missenden Road NSW 2050

Phone: +61 2 8514 0641

Abstract

Background: Immunotherapy agents show anti-cancer activity in several solid cancers. Efficacy in non-melanoma solid tumours for non-approved indications is unknown.

Method: Retrospective review describing outcomes and toxicity of self-funded pembrolizumab in patients with non-melanoma solid cancers at Chris O'Brien Lifehouse.

Results: From April 2015 to December 2015, 21 patients received/planned to receive self-funded pembrolizumab. Median age was 50 (16-76), 28% and 10% had an ECOG of 2, and 3-4 respectively. 62% received at least 2-4 lines of prior drug treatment. Median follow-up was 3.0 months (range, 0.4-9.6). Fourteen (67%) patients requested pembrolizumab. Pembrolizumab was clinician offered for 7 (33%) patients. Patients who requested pembrolizumab had worse outcomes. 3 patients died before receiving pembrolizumab. Of the 18 patients that received at least 1 dose, a partial response was observed in 3 (17%). Progressive disease occurred in 83%. 4 patients received only 1 cycle of pembrolizumab and died a median of 27 days (range 13-43) after. Immune related adverse events of any grade occurred in 33%. No grade 3-4 events were observed.

Conclusion: Pembrolizumab was well tolerated. Meaningful responses were observed in 17% of treated patients. Response continues after 5-6.5 months follow-up in 11% and >8 months of follow-up for the other responding patient. Financial impact to the patient can be substantial. Outcomes for 33% were poor with 3 patients dying prior to receiving therapy and 4 dying within weeks of receiving one dose. This highlights issues regarding the careful selection of patients, futility of anti-cancer therapy at the end of life and patient's perceived benefit of receiving this therapy.

Keywords: pembrolizumab; immunotherapy; end of life; advanced cancer; futility

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Lomax, AJ;Beith, J;Bhadri, V;Boyer, M;Grimison, P;Horvath, LG;Kao, S;Tattersall, M;Thomas, D;McNeil, C

Title:

Outcomes of patients with non-melanoma solid tumours receiving self-funded pembrolizumab at Chris O'Brien Lifehouse.

Date:

2016-12

Citation:

Lomax, A. J., Beith, J., Bhadri, V., Boyer, M., Grimison, P., Horvath, L. G., Kao, S., Tattersall, M., Thomas, D. & McNeil, C. (2016). Outcomes of patients with non-melanoma solid tumours receiving self-funded pembrolizumab at Chris O'Brien Lifehouse.. Intern Med J, 46 (12), pp.1392-1398. <https://doi.org/10.1111/imj.13232>.

Persistent Link:

<http://hdl.handle.net/11343/292191>