Rectal adenocarcinoma perforation following palliative colorectal stenting

Tamara M Vu, MBBS

Jonathan Alastair Simpson, BMBS, PhD, FRCS

Saleh Alzarhani, MBBS, MS

A Craig Lynch, MBChB, MMedSci, FRACS

Satish Warriar, MBBS, MS, FRACS

Alexander Heriot, MD, MBA, FRACS, FACS, FRCS (Gen.)

Department of Surgical Oncology, Peter MacCallum Cancer Centre, East Melbourne, Victoria, Australia

Figures: 2

Word count: 674

Corresponding author:

Tamara M Vu

tamara.vu@petermac.org

03 9565 1111

Department of Surgical Oncology

Peter MacCallum Cancer Centre

St Andrew's Place

East Melbourne Victoria 3002

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/ans.13480

Stenting of rectal cancers is a recognised management strategy in malignant bowel obstruction. Placement of a self-expanding metal stent can be used as a bridging measure prior to definitive surgery, or as a palliative solution to an obstructing tumour in patients not suitable for resection. However, colonic perforation is a small but significant risk associated with stent insertion, with a reported rate of 7.4% across stent recipients.ⁱ We present a case where a rectal stent inserted for palliation of a near obstructing rectosigmoid tumour caused perforation of the tumour and faeculent peritonitis three months following its insertion.

A 73-year-old man with a past history of prostate cancer treated with brachytherapy presented with bleeding per rectum and microcytic anaemia. Colonoscopy demonstrated a bulky tumour 25cm from the anal verge that could not be passed with the colonoscope. Histology was suspicious for invasive carcinoma. Staging CT revealed an abnormally thick-walled sigmoid with infiltration of the mesorectal fat, as well as multiple low density liver lesions consistent with metastases on MRI liver. The decision was made to perform rectosigmoid stenting prior to further investigation. A 25x120mm WallFlex Colonic (Boston Scientific) uncovered titanium alloy stent was inserted without complication.

PET scan demonstrated intense activity in the sigmoid primary, the liver metastases and a single left upper lobe metastasis. After discussion in the Lower GI Malignancy MDM, the patient proceeded to palliative chemotherapy with 5-FU and leukovorin.

Three months after stent insertion, the patient presented to his local hospital with acute abdominal pain and fever. CT scan demonstrated perisigmoid stranding associated with small locules of extraluminal gas, although gross free gas was not seen. He was transferred urgently to the quaternary centre where the stent had been inserted. On arrival he was found to be peritonitic, febrile and tachycardic, and the decision was made for laparotomy. At

operation he was found to have a perforated sigmoid, with the proximal end of the stent extruding through the defect and frank stool in the peritoneal cavity. A Hartmann's procedure was performed, with the tumour and stent removed en bloc and an end colostomy raised, followed by copious washout. His recovery was complicated by postoperative ileus and an intraabdominal collection requiring radiologically guided drainage.

Perforation is an infrequent but serious complication of colorectal stenting. An early randomised controlled trial was terminated prematurely due to stent-related morbidity,ⁱⁱ although subsequent evidence has shown comparable outcomes to defunctioning surgery.ⁱⁱⁱ A recent meta-analysis demonstrated an overall perforation rate of 7.4% (including both bridge to surgery and palliative patients).^{iv} Perforation can result from the ends of the stent penetrating the bowel wall, as here; or from tumour ingrowth causing stenosis and catastrophic proximal re-obstruction. This risk is increased with concomitant use of bevacizumab (which our patient did not receive), but not other chemotherapeutic agents.^v Insertion technique and surgeon experience can also contribute to the likelihood of perforation.^{vi}

There is considerable literature to suggest that stenting in the bridge to surgery setting is safe and viable, with one recent prospective observational multicentre study showing a perforation rate as low as 1.6%.^{vii} In the palliative setting, the current retrospective data suggests that stents can be used safely with palliative chemotherapy, with a perforation rate of 8% seen in one recent review of 38 patients receiving 5-FU based chemotherapy.^{viii} A recent Australian study also suggests that stenting in the palliative setting results in significantly less impact on quality of life than defunctioning surgery.^{ix} Stenting does has an

established role in the management of obstructing colonic cancers, however perforation remains a potential risk of which clinicians need to be aware.

ⁱ Van Halsema EE, van Hooft JE, Small AJ et al. Perforation in colorectal stenting: a meta-analysis and search for risk factors. *Gastrointest Endosc.* 2014; **79**(6): 970–82.

ⁱⁱ Van Hooft JE, Fockens P, Marinelli AW et al. Dutch Colorectal Stent Group: Early closure of a multicentre randomised clinical trial of endoscopic stenting versus surgery for stage IV leftsided colorectal cancer. *Endoscopy.* 2008; **40**(3): 184–91.

ⁱⁱⁱ Atukorale YN, Church JL, Hoggan BL. Self-expanding metallic stents for the management of emergency malignant large bowel obstruction: A systematic review. *J Gastrointest Surg.* 2015; **26** (ahead of print).

^{iv} Van Halsema EE, van Hooft JE, Small AJ et al. Perforation in colorectal stenting: a meta-analysis and search for risk factors. *Gastrointest Endosc.* 2014; **79**(6): 970–82.

^v Imbulgoda A, MacLean A, Heine J et al. Colonic perforation with intraluminal stents and bevacizumab in advanced colorectal cancer: retrospective case series and literature review. *Can J Surg.* 2015; **58**(3): 167–71. .

^{vi} Van Hooft JE, Fockens P, Marinelli AW et al. Dutch Colorectal Stent Group: Early closure of a multicentre randomised clinical trial of endoscopic stenting versus surgery for stage IV leftsided colorectal cancer. *Endoscopy.* 2008; **40**(3): 184–91.

^{vii} Saito S, Yoshida S, Isayama H. A prospective multicentre study on self-expanding metallic stents as a bridge to surgery for malignant colorectal obstruction in Japan: efficacy and safety in 312 patients. *Surg Endosc.* 2015; **8** (ahead of print).

^{viii} Ceze N, Charachon A, Locher C. Safety and efficacy of palliative systemic chemotherapy combined with colorectal self-expandable metallic stents in advanced colorectal cancer: A multicentre study. *Clin Res Hepatol Gastroenterol.* 2015; **20** (ahead of print).

^{ix} Young CJ, De-Lyde KJ, Young JM et al. Improving quality of life for people with incurable large bowel obstruction: Randomised control trial of colonic stent insertion. *Dis Col Rect.* 2015; **58**: 838–49.

stent perf photo.tif

stent perforated annotated.tif

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Vu, TM;Simpson, JA;Alzarhani, S;Lynch, AC;Warrier, S;Heriot, A

Title:

Rectal adenocarcinoma perforation following palliative colorectal stenting

Date:

2018-06-01

Citation:

Vu, T. M., Simpson, J. A., Alzarhani, S., Lynch, A. C., Warrier, S. & Heriot, A. (2018). Rectal adenocarcinoma perforation following palliative colorectal stenting. ANZ JOURNAL OF SURGERY, 88 (6), pp.E558-E559. <https://doi.org/10.1111/ans.13480>.

Persistent Link:

<http://hdl.handle.net/11343/292235>