Objectives: To assess the impact of four rabbit diets (hay only; extruded diet with hay; muesli with hay; muesli only) on faecal pellet size, faecal output and caecotrophy.

Methods: Thirty-two Dutch rabbits were studied over 17 months. Faecal pellet size and weight were measured in weeks 3, 9, 21 and 43 and faecal output in weeks 10, 22 and 45. Number of uneaten caecotrophs was recorded weekly.

Results: Faecal pellets were consistently smaller and lighter in rabbits fed muesli only, and the size of pellets produced by those fed muesli with hay decreased over the course of the study. Faecal output was greatest in rabbits with the highest hay intake. Uneaten caecotrophs were found in greatest frequency in rabbits fed muesli.

Clinical Significance: Muesli diets have a negative effect on faecal output and caecotroph ingestion and may therefore predispose to digestive disorders. Higher hay intake is associated with greater faecal output and fewer uneaten caecotrophs and may assist in preventing gastrointestinal stasis.

Keywords: Rabbit, Digestive, Gastrointestinal Stasis, Caecotrophs, Fly strike

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1111/jsap.12620](https://doi.org/10.1111/jsap.12620)

Introduction

Whilst rabbits are popular pets (PFMA 2015, PDSA 2015), studies to determine their specific nutritional requirements have not been conducted. Published nutritional guidelines (NRC 1977, FEDIAF 2013) are based on research conducted on laboratory and commercially-farmed rabbits. Pet rabbits were traditionally fed muesli-type diets but the number fed in this way has decreased, with more being fed hay, pelleted foods and vegetables (PDSA 2015). Muesli-type diets, especially if fed without hay, have been suggested to play a role in a range of diseases including gastrointestinal stasis (Mullan and Main 2006, Meredith 2012, RSPCA 2013), and previous studies have found that they are also associated with obesity (Prebble et al. 2015a), dental disease (Meredith et al. 2015), and behavioural changes (Prebble et al. 2015b).

Dietary fibre particles are separated in the proximal colon of the rabbit according to their size (Jilge 1982), with small particles (<0.3mm) retained in the caecum for microbial fermentation (Gidenne 1993) and larger particles passed rapidly through the colon to be expelled in faecal pellets. Larger fibre particles are necessary to maintain normal caecal-colic motility (Cheeke 1994). Increasing dietary fibre levels increases the rate of passage of ingesta (Hoover and Heitmann 1972, Fraga et al. 1991, Gidenne 1992, Bellier and Gidenne 1996) and, conversely, fine grinding of fibre results in prolonged retention times and reduced food intake (Laplace et al. 1977; Fraga et al. 1991; Gidenne 1992). The caecal contents are expelled once or twice in a 24-hour period as mucus coated caecotrophs that are consumed intact directly from the anus (Madsen 1939, Taylor 1939). Caecotrophy provides a source of B-complex vitamins, vitamin K and microbial protein (Kulwich et al. 1953, Hirakawa 2001). The amount of caecotrophs consumed is influenced by diet; all are eaten when food is scarce but, when food is provided *ad*

libitum, protein and fibre content affect consumption. Increased levels of dietary fibre increase consumption and high protein levels reduce consumption. Increased dietary fibre content also reduces the protein and volatile fatty acid content of caecotrophs (Carabaño et al. 1988).

Gastrointestinal (GI) disorders are common in pet rabbits (Varga 2014, Harcourt-Brown 2014) and often considered to be related to diet rather than enteric pathogens. GI stasis is characterised by progressive hypomotility and eventual cessation of intestinal movement with clinical signs including anorexia, reduced or absent faecal production and accumulation of gas within the digestive tract (Harcourt-Brown 2002, Oglesbee and Jenkins 2012). Faecal pellets gradually reduce in size and frequency of production and may also be misshapen (Saunders and Rees Davies 2005). Pet rabbits also commonly present with uneaten caecotroph accumulation around the anus (Saunders and Rees Davies 2005, Eatwell 2006, Harcourt-Brown 2014, Varga 2014) which predisposes to flystrike (Harcourt-Brown 2002, Meredith 2014). Low fibre diets are frequently implicated in both GI stasis (Rees Davies and Rees Davies 2003, Saunders & Rees Davies 2005, Varga 2014) and reduced caecotroph consumption (Fekete and Bokori 1985).

A low fibre level alone may not cause GI stasis but is believed to act as a predisposing factor; other factors such as stress, pain, infectious disease (e.g. coccidiosis) or medication may be required to cause clinical disease (Harcourt-Brown 2002). Provision of high fibre diets to commercially-farmed rabbits reduces the prevalence of diarrhoea and associated mortality (de Blas et al. 1986, Bennegadi et al. 2001, Blas et al. 2010) and may therefore act protectively and promote health. Reduced faecal output and faecal pellet size are often seen in the early stages of gastrointestinal stasis before faecal

output ceases completely (Oglesbee and Jenkins 2012). Wild rabbits (*Oryctolagus cuniculus*) generally produce between 300 and 500 pellets/24 hours (Lockley 1962, Simonetti 1989, Gonzalez-Redondo 2009) although up to 820 have been reported (Taylor and Williams 1956), and mean faecal pellet diameter of wild rabbits was 7.90 ± 0.14 mm in one study (Simonetti 1989). Diet may be responsible for these differences in numbers as, although not studied in European rabbits (*Oryctolagus cuniculus*), higher fibre diets significantly increase the number of faecal pellets produced by wild cottontails (*Sylvilagus floridanus*) (Cochran and Stains 1961). Feeding muesli and, or, a lack of hay may contribute to gastrointestinal stasis and lack of caecotrophy in pet rabbits. This study aims to assess the effect of feeding two concentrate types, extruded nugget (EH) and muesli (MH) alongside *ad lib* hay with *ad lib* hay (HO) fed alone and *ad lib* muesli (MO), on faecal output, faecal pellet size and the quantity of caecotrophs left uneaten.

Materials and methods

This study was conducted as a part of a long-term study to assess the effect of diet on the health and welfare of pet rabbits as previously described by Prebble & Meredith (2014), Prebble et al. (2015a.b) and Meredith et al (2015). Study design and methodology was approved by the Ethical Review Committees of the Royal (Dick) School of Veterinary Studies and the Food and Environment Research Agency (FERA). The rabbits were housed in a facility licensed by the Home Office, although, a project licence under the Animals (Scientific Procedures) Act 1986 (ASPA) was not required for this study that examined diet changes alone. The study was continually monitored by the FERA Ethics Committee and Home Office inspector throughout its duration.

Diets

On arrival, 32 eight week old rabbits were housed in pairs and acclimatised over 40 days (days –54 to –14) by maintaining their weaning diet of 50 g per rabbit of an extruded diet (Burgess® Excel-Junior and Dwarf Rabbit; Burgess Pet Care, Thornton Le Dale, North Yorkshire, UK) once a day plus *ad lib* Timothy Hay provided in wall-mounted hayracks. Water was provided *ad lib* in 700 ml bottles. At Day –14, the rabbits were randomly allocated to one of four diet groups each consisting of eight rabbits:

1. Hay only (HO) – *ad lib* supply of Timothy hay (n=8);
2. Extruded diet and hay (EH) – 50 g per rabbit Burgess Excel Adult Rabbit (Burgess Pet Care, Thornton Le Dale, North Yorkshire, UK) with *ad lib* supply of hay (n=8);
3. Muesli and hay (MH) – 60 g per rabbit Russell Rabbit Complete Muesli (Supreme Petfoods Limited, Ipswich, Suffolk, UK) with *ad lib* hay (n=8);
4. Muesli only (MO) – *ad lib* supply (125 g per rabbit) of Russell Rabbit Complete Muesli (Supreme Petfoods Limited, Ipswich, Suffolk, UK) (n=8).

The rabbits were gradually transitioned on to the new diet over a 2-week period (Day –14 to 0) and remained on that diet for 17 months. All concentrates were weighed out daily to ensure accurate and consistent weights of food were offered. Food remaining after 24 hours was weighed to determine food consumed. The nutritional composition of the four diets is detailed in Table 1. Hay intake and selective feeding of muesli were recorded in months 3, 6 and 12 as reported by Prebble & Meredith (2014).

A rabbit in the EH group died suddenly on day 209, but the cause of death was not evident on post-mortem examination. A rabbit in the MO group was removed at month 7 following the development of clinical dental disease. Data from these two rabbits were not included in any subsequent analyses.

Data collection

Faecal samples were collected from individual rabbits in weeks 3, 9, 21 and 43. The rabbits were separated for up to three hours in the light period using a Perspex divider within their home pen. Fifteen (or the total number produced if less than 15) of these faecal pellets were then randomly selected and weighed using digital scales (Fisherbrand DP 300) and a mean faecal pellet weight calculated. The size of the faecal pellet was measured using digital callipers; as the faecal pellets are not spherical or symmetrical the longest diameter was measured.

Following the observation that longer periods of separation (>3 hrs) were required in the MO group in order to get sufficient quantities of faecal pellets, total faecal output in all groups was measured in weeks 10, 22 and 45. Bedding was removed from the pens for a 24-hour period and all faeces passed during this period collected and weighed. A sample of 20 pellets was weighed and the mean faecal pellet weight calculated. The total weight of faecal pellets produced was then divided by the mean faecal pellet weight to give an indication of the number of pellets produced.

A count of caecotrophs found in the pens during weekly cleaning was done by the same researcher on 25 of the weeks during the period between week 4 and week 31.

Caecotrophs were not measured in weeks 7 and 12 because the researcher was unable to be present during cleaning.

Statistical Analysis

Data were analysed using Minitab (v16.1.1 © 2010 Minitab Inc.) and R software (v2.15.1 © 2012 The R Foundation for Statistical Computing). For most of the measurements, data from each time point were analysed using standard analysis of variance (ANOVA) also taking the sex of the rabbit into account, using standard Tukey's *post hoc* tests to assess pair-wise differences between the groups where overall differences were obtained. Residuals were examined for adequate normality before analysis. A Pearson product moment correlation was carried out to assess the relationship between rabbit weight, faecal pellet weight and faecal pellet diameter and also between faecal weight and number of faecal pellets. $P < 0.05$ was taken to indicate statistical significance. The number of caecotrophs produced from particular time points were analysed with general linear models with Poisson errors used to incorporate the integer nature of the caecotroph data with pairwise post-hoc Tukey analysis carried out if there were overall differences.

Results

Faecal pellet weight and diameter

Faecal pellets were consistently lighter and smaller in rabbits fed muesli only, and greater hay intake was associated with higher faecal pellet weight and size. There was a statistically significant positive linear relationship between hay intake and both faecal pellet weight ($t_{1,30}=4.96$, $P<0.001$, $R^2=45.0$) and diameter ($t_{1,30}=4.25$, $P<0.001$, $R^2=37.6$, figure 1). Faecal pellet weight was positively correlated with faecal pellet diameter at all timepoints ($r>0.635$, $P<0.001$). Rabbits fed muesli had the smallest faecal pellet size over the course of the study. Significant changes in mean faecal pellet weight and diameter occurred over the duration of the study ($F_{3,25}=3.24$, $P=0.039$, Table 2), but this was most noticeable in the MH group (62% reduction in faecal pellet weight and 32% reduction in diameter) and MO groups (47% reduction in faecal pellet weight and 14% reduction in diameter). In contrast, smaller changes were observed in the HO (38% reduction in faecal pellet weight and 4% increase in diameter) and EH (26% reduction in faecal pellet weight and 3% reduction in diameter) (Figures 1 and 2) groups.

Significant differences were present in both faecal pellet weight and diameter at each time point ($F_{3,25}=29.92$, $P<0.017$, Figure 2). In week 3, faecal pellets produced by the HO group were 26% heavier than those in the EH group and 23% heavier than the MH group, while the MO group produced pellets that were 21% smaller than the MH group ($P<0.028$). The EH and MH groups were not different from each other ($P>0.127$).

In week 9, faecal pellets produced by the HO group ($0.34g\pm0.09$, $10.53mm\pm1.53$), continued to be significantly heavier and larger than all other groups ($P<0.009$) while

those produced by the MO group (0.14 ± 0.05 , $8.52 \text{ mm} \pm 0.85$) were both smaller ($P < 0.002$) and lighter ($P < 0.035$). There remained no difference between the EH ($0.2 \text{ g} \pm 0.05$, $9.72 \text{ mm} \pm 0.87$) and MH (0.23 ± 0.03 , $9.82 \text{ mm} \pm 0.71$) groups ($P = 0.920$). The faecal pellets in the MH group reduced in size more than those produced by other groups and by week 21 they were 29% smaller than those produced by the EH group ($P = 0.011$) and by week 43 pellets produced by the MH group were 47% smaller than the EH group and were similar in size to those produced by MO group ($P = 0.90$, Figure 3).

Total faecal output

Increased hay intake was associated with production of more faecal pellets, with a positive linear relationship between hay intake and both the weight of faeces produced ($t_{1,13} = 3.45$, $P < 0.004$, $R^2 = 47.7$) and the number of faecal pellets ($t_{1,13} = 3.11$, $P < 0.008$, $R^2 = 42.6$). Significant differences in the number of pellets produced over a 24-hour period were present between groups at all time points ($F_{3,11} = 18.89$, $P < 0.003$, Table 3). The HO group produced 46 - 62% more faecal pellets over a 24-hour period than those in the MO group ($P < 0.001$), 24-29% more than the EH group ($P < 0.07$) and 32-43% more than the MH group ($P < 0.02$). The EH produced (12- 25%) more faecal pellets than the MH group although this difference was not significant ($P > 0.06$). Likewise there was no difference in the number of faecal pellets produced by the MH and MO groups ($P > 0.13$, Figure 4).

In week 52 a single rabbit in the MO group had signs of gastrointestinal stasis including reduced food intake, hunched posture, tooth grinding, reduced faecal pellet production and altered behaviour and was removed from the study for treatment.

Caecotrophs

Uneaten caecotrophs were found with greatest frequency in rabbits fed muesli. No caecotrophs were found in pens in the HO group for 11 of the 25 weeks in which counts were performed, and the caecotrophs found in the HO group in the remaining 14 weeks were only from one pen. Analysis was therefore only performed on the three remaining groups. Overall, significant differences between groups were seen throughout the trial ($\chi^2_{3,15} = 5.35$, $P < 0.001$, Figure 5). The MO group left more caecotrophs (18 ± 2) than the EH (4 ± 3) and MH (7 ± 1) in week four ($P < 0.001$), whilst there was no difference between the EH and MH groups ($P = 0.273$). However, by the end of the trial the reduction in the number of caecotrophs left by the EH group (1 ± 1) was significantly less than those left by the MH group (5 ± 2 , $P < 0.02$). While the number of caecotrophs left by the MO group had fallen (14 ± 3) this was still significantly more than all other groups ($P < 0.001$).

Discussion

This study demonstrates that diet has a significant impact on faecal pellet weight and size, total faecal output and caecotrophy. Significantly heavier and larger faecal pellets were produced by those rabbits consuming hay, in contrast to the smaller, lighter pellets produced by the MO group, which had no access to hay. Faecal pellets produced by the MO group were similar in appearance to the small irregular pellets observed in the early stages of GI stasis (Lord 2012, Harcourt-Brown 2014). Although the mean faecal diameter in the MO group was comparable with that reported in wild rabbits of 7.90mm, the range included pellets as small as 5.80mm (Figure 3). As low fibre levels are frequently implicated in GI stasis in pet rabbits (Varga 2014, Rees Davies and Rees

Davies 2003, Lord 2012) the finding that low hay diets are associated with smaller faecal pellet diameter and faecal output is of clinical relevance.

The smaller faecal pellets produced by the MO group throughout the study and the MH group towards the end of the study are similar in size and appearance to those observed by the authors and described by Harcourt-Brown (2014) in early clinical cases of gastrointestinal stasis. However, the low occurrence of GI stasis (one rabbit) in this study supports the assertion that low fibre diets are only one contributing factor in the development of GI stasis (Harcourt- Brown 2002) and that low fibre diets are not directly causative. Dietary fibre has a protective effect on the morbidity and mortality from digestive diseases observed in farmed rabbits and at weaning (Bennegadi et al. 2001, Gidenne et al. 2001) and this may also be true for GI stasis. This protective effect may be a result of the effect of dietary fibre on volatile fatty acid (VFA) production. Rabbits fed high fibre, low starch diets produce less VFAs than those on low fibre, high starch diets (Carabaño et al, 1988, Gidenne et al, 2004). This would suggest that rabbits fed the low fibre, high starch muesli diet would have higher VFA levels which have been associated with reduced colon motility in *in vitro* studies (Squires et al. 1992, Dass et al. 2007). This reduction in motility may make rabbits fed on a muesli-only diet more susceptible to the development of GI stasis if other factors associated with reduced motility such as pain or stress (Harcourt-Brown, 2014) are present.

Faecal output over a 24-hour period was greater in rabbits consuming more hay (hay intake reported in Prebble and Meredith (2014)). As large fibre particles pass rapidly through the colon and are excreted in faecal pellets (Jilge 1982, Carabaño et al. 1988), the differences in hay (and therefore fibre) intake between the groups can explain these

difference in faecal output. The number of faecal pellets produced by the EH and MH groups fall within the lower end of the range (300-500 daily) generally reported in wild rabbits (Lockley 1962, Simonetti 1989, Gonzalez-Redondo 2009). The HO groups produced numbers closer to, or exceeding, the top end of this range, but did not reach levels in wild rabbits reported by Taylor and Williams (1956). The MO group had considerably lower faecal output, similar to the findings of Arnold and Reynolds (1943) in Arizona and antelope jack rabbits fed a highly digestible mash, which produced half as much as those fed grass. In addition, differences in faecal pellet colour were observed between groups; the HO group produced faecal pellets that were lighter in colour, whilst those produced in the MO group were darker. The faecal pellets produced by the EH and HO groups were similar in colour. Objective quantification of these differences, for example using colorimetry, was outside the scope of this project.

Uneaten caecotrophs were regularly found in all groups fed concentrates but infrequently in the HO group. Highest numbers were found in the MO group (up to 30), whilst the MH (up to 10) and EH (up to 5) groups that consumed hay left less caecotrophs, supporting findings that caecotroph consumption is associated with forage-based (Clauss et al. 2012) and high fibre, low protein diets (Fekete and Bokori 1985). Caecotrophy contributes to protein, energy and vitamin requirements (Kulwich et al. 1953, Fraga et al. 1991, Kerti et al. 2005), and the higher protein and energy content of concentrates allows nutritional requirements to be met solely from the diet, resulting in increased numbers of uneaten caecotrophs. Other factors implicated in reduced caecotroph consumption include obesity, perineal dermatitis, dental and musculoskeletal diseases (Harcourt-Brown 2014). Obesity was observed in the MO group (Prebble et al. 2015), but was not sufficient to prevent grooming (which was observed frequently by the

researcher) and should not have impaired caecotrophy. No other conditions were apparent in the rabbits before or during data collection. In the absence of other contributory factors, the differences in uneaten caecotrophs within the pens could be attributed to dietary factors alone. The cause of the reduction in uneaten caecotrophs in the MO group by the end of the trial is not known, but possibilities include an age-related effect, or changes in dry matter (DM) or water intake and would require further investigation..

Smaller faecal pellet size and weight, lower total faecal output and greater number of non-consumed caecotrophs may indicate that muesli-based diets contribute to the development of digestive disorders. In particular, feeding muesli in the absence of hay cannot be recommended. Further study is required to assess the significance of diet in the development of gastrointestinal stasis and the interaction of other factors. A higher fibre diet promotes a larger faecal pellet size and weight and higher total faecal output, therefore feeding hay alone or alongside concentrate diets is recommended for pet rabbits and may be beneficial in preventing protective against GI stasis.

Reference List

Arnold, J.F. and Reynolds, H.G., (1943) Droppings of Arizona and Antelope Jack Rabbits and the "Pellet Census." *The Journal of Wildlife Management*, 7(3), 322

Bellier, R. and Gidenne, T., (1996) Consequences of reduced fibre intake on digestion, rate of passage and caecal microbial activity in the young rabbit. *British Journal of Nutrition*, 75(3), 353–363

Bennegadi, N., Gidenne, T. and Licois, D., (2001) Impact of fibre deficiency and sanitary status on non-specific enteropathy of the growing rabbit. *Animal Research*, 50(5), 401–413

Blas, E., Cervera, C. and Fernández-Carmona, J., (2010) Effect of two diets with varied starch and fibre levels on the performances of 4-7 weeks old rabbits. *World Rabbit Science*, 2(4), 117–121

de Blas, J.C., Santomá, G., Carabaño, R. and Fraga, J., (1986) Fiber and starch levels in fattening rabbit diets. *Journal of Animal Science*, 63(6), 1897–1904

Carabaño, R., Fraga, M. J., Santomá, G. and de Blas, J.C., (1988) Effect of diet on composition of cecal contents and on excretion and composition of soft and hard feces of rabbits. *Journal of Animal Science*, 66(4), 901–910

Cheeke, P.R., (1994) Nutrition and Nutritional Diseases. In: The Biology of the Laboratory rabbit. Eds D. H. Ringer and P.J. Manning. Academic Press, New York pp 321–333

Clauss, M., Burger, B., Liesegang, A., et al., (2012) Influence of diet on calcium metabolism, tissue calcification and urinary sludge in rabbits (*Oryctolagus cuniculus*). Journal of Animal Physiology and Animal Nutrition, 96(5), 798–807

Cochran, G.A. and Stains, H.J., (1961) Deposition and Decomposition of Fecal Pellets by Cottontails. The Journal of Wildlife Management, 25(4), 432

Dass, N. B., John, A. K., Bassil, A. K., Crumbley, C. W., Shehee, W. R., Maurio, F. P., Moor, G. B. T., Taylor, C. M. and Sanger, G. J. (2007) The relationship between the effects of short-chain fatty acids on intestinal motility in vitro and GPR43 receptor activation. Neurogastroenterology and Motility, 19 (1) 66-74

Eatwell, K., (2006) Perineal Soiling in the Rabbit. Proceedings of the British Small Animal Veterinary Congress. Birmingham, UK.

FEDIAF, (2013) Nutritional guidelines for feeding pet rabbits, European Pet Food Industry Federation, Brussels, Belgium

Fekete, S. and Bokori, J., (1985) The effect of the fiber and protein level of the ration upon cecotrophy of the rabbit. Journal of Applied Rabbit Research, 8, 68–71

Fraga, M.J., Pérez de Ayala, P., Carabaño, R., and de Blas J. C., (1991) Effect of type of fiber on the rate of passage and on the contribution of soft feces to nutrient intake of finishing rabbits. *Journal of Animal Science*, 69(4), 1566–74

Gidenne, T., (1992) Effect of fibre level, particle size and adaptation period on digestibility and rate of passage as measured at the ileum and in the faeces in the adult rabbit. *British Journal of Nutrition*, 67(1), 133– 288 146

Gidenne, T., (1993) Measurement of the rate of passage in restricted fed rabbits: effect of dietary cell wall level on the transit of fibre particles of different sizes. *Animal Feed Science and Technology*, 42(1), 151–163

Gidenne, T., Arveux, P. and Madec, O., (2001) The effect of the quality of dietary lignocellulose on digestion, zootechnical performance and health of the growing rabbit. *Animal Science*, 73, 97-104

Gidenne, T., Jehl, N., Lapanouse, A. and Segura, M. (2004) Inter-relationship of microbial activity, digestion and gut health in the rabbit: effect of substituting fibre by starch in diets having a high proportion of rapidly fermentable polysaccharides. *British Journal of Nutrition*. 92. 95-104

Gonzalez-Redondo, P., (2009) Number of faecal pellets dropped daily by the wild rabbit (*Oryctolagus cuniculus*). *Journal of Animal and Veterinary Advances*, 8(12), 2635–2637

Harcourt-Brown, F., (2014) Digestive system disease. In : *BSAVA Manual of Rabbit Medicine*. Eds A. Meredith and B. Lord. BSAVA, Gloucester. pp 168-190

Hirakawa, H., (2001) Coprophagy in leporids and other mammalian herbivores. *Mammal Review*, 31(1), .61–80

Hoover, W.H. and Heitmann, R.N., (1972) Effects of dietary fiber levels on weight gain, cecal volume and volatile fatty acid production in rabbits. *The Journal of nutrition*, 102(3), 375–379

Jilge, B., (1982) Rate of movement of marker particles in the digestive tract of the rabbit. *Laboratory Animals*, 16(1), 7–11

Kerti, A., Bárdos, L., Deli, J., and Oláh, P., (2005) Investigation of the changes in vitamin metabolism without caecotrophy in rabbits. *Proceedings of the ISAH International Congress*. Warsaw, Poland pp. 475–478.

Kulwich, R., Struglia, L. and Pearson, P.B., (1953) The effect of coprophagy on the excretion of B vitamins by the rabbit. *The Journal of Nutrition*, 49(4), 639–645

Laplace, J.P., Lebas, F. and Germain, C., (1977) Le transit digestif chez le Lapin. VII.– Influence de la finesse du broyage des constituants d'un aliment granulé. *Annales de zootechnie*. 26(3) 413–420

Lockley, R., (1962) Production of faecal pellets in the wild rabbit. *Nature*, 194, 988–989

Lord, B., (2012) Gastrointestinal disease in rabbits 1. Gastric diseases. In Practice, 34(2), 90–96

Madsen, H., (1939) Does the rabbit chew the cud. Nature, 143, 981–982.

Meredith, A., (2014) Dermatoses. In: BSAVA Manual of Rabbit Medicine gery. Eds A. Meredith and B. Lord. BSAVA, Gloucester. pp 255-263

Meredith, A., (2012) Is obesity a problem in pet rabbits? Veterinary Record, 171(8), 192–193.

Meredith A. L., Prebble J. L., and Shaw D. J., (2015) Impact of diet on incisor growth and attrition and the development of dental disease in pet rabbits. Journal of Small Animal Practice 56(6):37-82

Mullan, S.M. and Main, D.C.J.,(2006) Survey of the husbandry, health and welfare of 102 pet rabbits. Veterinary Record, 159(4), 103–109.

NRC, (1977) Nutrient of Requirements of Rabbits 2nd ed. National Research Council, Washington

Oglesbee, B.L. and Jenkins, J.R., (2012) Gastrointestinal Diseases. In Queensbury, K E and Carpenter, J W, ed. Ferrets, rabbits and rodents clinical medicine and surgery. St. Louis, Missouri: Elsevier., pp. 193–204.

PDSA (2015) PDSA Pet Animal Wellbeing Report: The state of our pet nation, Peoples Dispensary for Sick Animals.

PFMA (2015) <http://www.pfma.org.uk/historical-pet-population> [Accessed June 29, 2015].

Prebble, J. and Meredith, A., (2014) Food and water intake and selective feeding in rabbits on four feeding regimes. *Journal of Animal Physiology and Animal Nutrition*, 98(5), 991–1000.

Prebble, J., Shaw, D. and Meredith, A., (2015a) Bodyweight and body condition score in rabbits on four different feeding regimes. *Journal of Small Animal Practice*. 56(3), 207-12

Prebble, J., Langford, F., Shaw, D., and Meredith, A. (2015b) The effect of four different feeding regimes on rabbit behaviour. *Applied Animal Behaviour Science*. 169, 86–92

Rees Davies, R. and Rees Davies, J.A., (2003) Rabbit gastrointestinal physiology. *Veterinary Clinics of North America: Exotic Animal Practice*, 6(1), 139–153.

RSPCA (2013) <http://www.rspca.org.uk/adviceandwelfare/pets/rabbits> [Accessed June 28, 2015].

Saunders, R. and Rees Davies, R., (2005) Notes on rabbit internal medicine, Blackwell Publishing, Oxford

Simonetti, J.A., (1989) Tasas de defecación y descomposición de fecas de *Oryctolagus cuniculus* en Chile central. *Medio Ambiente (Chile)*, 10, 92–95

Squires, P. E., Ramsey, R. D., Edwards, C. A. and Read, N. W. (1992) Effect of short-term fatty acids on contractile activity and fluid flow in rat colon in vitro. *American Journal of Physiology*. 262 (5 pt 1). G813-7

Taylor, E.L., (1939) Do rabbits chew the cud? [Appendix to Madsen's note]. *Nature*, 143, 983

Taylor, R. and Williams, R., (1956) The use of pellet counts for estimating the density of populations of the wild rabbit, *Oryctolagus cuniculus* (L.). *New Zealand Journal of Science and Technology*, 38, 236–256.

Varga, M., (2014) Rabbit Basic Science. In *Textbook of Rabbit Medicine* 2nd ed. Elsevier., pp 3-110

Figure legends

Figure 1 Scatter plot of hay intake (g) by pen and faecal pellet diameter (mm) of Dutch rabbits (n=30) fed four diets (HO, EH, MH and MO) in week 43 of the trial period with linear regression line fitted to data.

Figure 2 a) Mean faecal pellet weight (g) of Dutch rabbits (n=32) fed four diets (HO, EH, MH and MO) measured at four points over a 40 week period. b) Mean faecal pellet diameter (mm) of Dutch rabbits (n=32) fed four diets (HO, EH, MH and MO) measured at four points over a 40 week period.

Figure 3 Boxplot of faecal pellet diameter (mm) of Dutch rabbits (n=30) fed four diets (HO, EH, MH and MO) in week 43 of the trial period. The horizontal bar represents the median, the box is the interquartile range and whiskers represent the range of values.

*** significant at $P < 0.001$, * significant at $P < 0.05$

Figure 4 Boxplot of weight of faeces produced over a 24 hour period (g) by Dutch rabbits (n=30) fed four diets (HO, EH, MH and MO) in week 45 of the trial period.

Boxplots as for Figure 3. *** significant at $P < 0.001$, * significant at $P < 0.05$

Figure 5 Mean number of caecotrophs found on a weekly basis in the pens of Dutch rabbits (n=32) fed four diets (HO, EH, MH and MO) measured at over a 40 week period.

Author Manuscript

figure 1.jpg

figure 2a.jpg

Faecal pellet diameter (mm) (Week 43)

Author Manuscript

figure 3.jpg

Faeces produced in 24 hours (g) (week 45)

Author Manuscript

figure 4.jpg

figure 5.jpg

Figure 2b.jpg

Impact of diet on faecal output and caecotroph consumption in rabbits.

Anna L Meredith MA VetMB PhD CertLAS DZooMed DipECZM MRCVS

Royal (Dick) School of Veterinary Studies and The Roslin Institute, University of Edinburgh,
Easter Bush Campus, Midlothian, EH25 9RG

Corresponding author; Email: Anna.Meredith@ed.ac.uk Tel:0131 651 7457

Jennifer L Prebble¹ BVSc MRCVS

Askham Bryan College, Askham Fields Lane, York, YO23 3FR

¹ JP was employed on a KTP partnership between the Royal (Dick) School of Veterinary Studies and Burgess Pet Care , Victory Mill, Priestmans Lane, Thornton-Le-Dale, Pickering, North Yorkshire, YO18 7RU

Keywords

Rabbit, Digestive, Gastrointestinal Stasis, Caecotrophs, Fly strike

Acknowledgements

The authors acknowledge financial support from the Technology Strategy board provided through a Knowledge Transfer Partnership between the Royal (Dick) School of Veterinary Studies and Burgess Pet Care and also thank Burgess Pet Care for additional financial support. We also thank FERA for allowing us the use of their facilities, ASIST for providing routine rabbit care and Station House Vets for neutering and providing treatment to the

rabbits. We also thank Dr Darren Shaw for his advice and assistance in performing the statistical analysis.

Word Count=3113

Author Manuscript

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Meredith, AL;Prebble, JL

Title:

Impact of diet on faecal output and caecotroph consumption in rabbits

Date:

2017-01-01

Citation:

Meredith, A. L. & Prebble, J. L. (2017). Impact of diet on faecal output and caecotroph consumption in rabbits. JOURNAL OF SMALL ANIMAL PRACTICE, 58 (3), pp.139-145.
<https://doi.org/10.1111/jsap.12620>.

Persistent Link:

<http://hdl.handle.net/11343/292292>