COVERSHEET

(1) Title**Tract-specific atrophy in focal epilepsy: disease, genetics or seizures?**Running head**Tract-specific atrophy in focal epilepsy**(2) Author names & affiliations

David N Vaughan MBBS^{1,2,3}, David Raffelt PhD¹, Evan Curwood PhD¹, Meng-Han Tsai MD, PhD^{4,5}, Jacques-Donald Tournier PhD^{6,7}, Alan Connelly PhD^{1,2}, Graeme D Jackson BSc MD^{1,2,3}

1. Florey Institute of Neuroscience and Mental Health, Melbourne, Australia
2. Department of Florey, University of Melbourne, Victoria, Australia
3. Department of Neurology, Austin Health, Heidelberg, Victoria, Australia
4. Department of Neurology, Kaohsiung Chang Gung Memorial Hospital, Kaohsiung, Taiwan
5. Department of Nursing, Meiho University, Pingtung, Taiwan
6. Department of Biomedical Engineering, Division of Imaging Sciences and Biomedical Engineering, King's College London, UK
7. Centre for the Developing Brain, Division of Imaging Sciences and Biomedical Engineering, King's College London, King's College London, UK

(3) Corresponding Author**David Vaughan****Melbourne Brain Centre, 245 Burgundy St, Heidelberg, Victoria, Australia 3084****david.vaughan@florey.edu.au**

- (4) Characters in title: 72
Characters in running head: 40

- (5) Word count: Abstract 249, Introduction 433, Discussion 1463, Body 4419.
Figures/tables: 1 table & 5 color figures

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version record](#). Please cite this article as [doi:10.1002/ana.24848](https://doi.org/10.1002/ana.24848).

Tract-specific atrophy in focal epilepsy: disease, genetics or seizures?

ABSTRACT

Objective: To investigate whether genetics, underlying pathology or repeated seizures contribute to atrophy in specific white matter tracts.

Methods: Medically-refractory unilateral temporal lobe epilepsy with hippocampal sclerosis (HS-TLE, n=26) was studied as an archetype of focal epilepsy, using voxel-based analysis of diffusion-weighted imaging. A genetic effect was assessed in first-degree relatives of HS-TLE who did not have epilepsy themselves (HS-1°Rel; n=26). The role of disease process was uncovered by comparing HS-TLE to unilateral TLE with normal clinical MRI (MRI-neg TLE; n=26, matched for seizure severity). The effect of focal seizures was inferred from lateralized atrophy common to both HS-TLE and MRI-neg TLE, in comparison to healthy controls (n=76).

Results: HS-1°Rel had bilaterally small hippocampi, but no focal white matter atrophy was detected, indicating a limited effect of genetics. HS-TLE had lateralized atrophy of most temporal lobe tracts, and hippocampal volumes in HS-TLE correlated with parahippocampal cingulum and anterior commissure atrophy, indicating an effect of the underlying pathology. Ipsilateral atrophy of the tapetum, uncinate and inferior fronto-occipital fasciculus was found in both HS-TLE and MRI-neg TLE, suggesting a common lateralized effect of focal seizures. Both epilepsy groups had bilateral atrophy of the dorsal cingulum and corpus callosum fibers, which we interpret as a consequence of bilateral insults (potentially generalized seizures and/or medications).

Interpretation: Underlying pathology, repeated focal seizures and global insults each contribute to atrophy in specific tracts. Genetic factors make less of a contribution in this cohort. A multi-factorial model of white matter atrophy in focal epilepsy is proposed.

1 INTRODUCTION

2 Abnormal white matter structure is an important aspect of focal epilepsy, which
3 relates to both the causes and the consequences of the disease. Such changes are often
4 widespread, may occur with a seizure focus in different brain regions¹⁻³, and have
5 been found in the presence of diverse pathologies including hippocampal sclerosis⁴,
6 focal dysplasia⁵, malformations of cortical development⁶ and even when no visible
7 lesion is present⁷⁻⁹.

8 Temporal lobe epilepsy with hippocampal sclerosis (HS-TLE) is often considered the
9 archetype of focal epilepsy¹⁰ as it has been extensively studied, and accounts for
10 around one third of patients with chronic medically-refractory TLE¹¹. The
11 hippocampus is the focus for seizure onset¹², and has the diagnostic histopathology of
12 pyramidal neuronal loss, granule cell dispersion and reactive gliosis¹³. Patients with
13 TLE who have no evidence of hippocampal sclerosis on imaging (MRI-neg TLE)
14 provide an informative comparison group. They have clinical seizures that are
15 individually indistinguishable from HS-TLE, but several group-level differences
16 indicate that there is different underlying pathological process. Seizures begin at a
17 later age in MRI-neg TLE, there is a lower proportion of antecedent childhood febrile
18 convulsions^{14,15} and histological examination does not show the classical pattern of
19 hippocampal neuronal loss^{16,17}. Genetic factors may contribute to the structural brain
20 abnormalities in HS-TLE, and this can be isolated by investigating first degree
21 relatives of people of HS-TLE who do not have seizures themselves (HS-1°Rel)¹⁸.

22 White matter abnormalities in TLE have previously been investigated with either
23 volumetry of T1-weighted imaging, or microstructural analysis of diffusion weighted
24 imaging. Voxel-based morphometry in HS-TLE has shown widespread atrophy
25 affecting bilateral temporal and extratemporal regions¹⁹⁻²² or a more restricted pattern
26 involving the ipsilateral temporal lobe^{1,4}. Relatives of HS-TLE show the genetic traits
27 of bilaterally small hippocampi¹⁸ and reduced total white matter volume²³. In MRI-
28 neg TLE no consistent alteration of white matter morphometry has been detected²⁴.
29 Previous diffusion weighted imaging studies of TLE have all employed the diffusion
30 tensor model (DTI) within various analysis frameworks, although DTI examines only

31 microstructural changes in the white matter and does not deal well with the crossing
32 fibre problem²⁵.

33 Therefore, to assess atrophy of specific white matter tracts in this study we use the
34 recently developed metric of Fibre Density and Cross-section (FDC)²⁶. This measure
35 is calculated from diffusion-weighted MRI, for each *fibre* population within a *voxel* (a
36 *fixel*)²⁷, and combines the morphometric information of Fibre Cross-section (FC) with
37 the microstructural information of Fibre Density (FD)²⁸.

38 Thus we investigate three well-defined cohorts (HS-TLE, MRI-neg TLE and HS-
39 1°Rel) asking the question of whether genetics, the underlying disease process and the
40 effects of ongoing seizures produce injury to specific tracts.

41 **METHODS**

42 ***Participants***

43 Patients with medically-refractory unilateral TLE, between the ages of 14 and 65
44 years, were studied prospectively during work-up for epilepsy surgery between 2007
45 and 2015, at the Austin Hospital, Melbourne, Australia. This included a subset of
46 patients (17 HS-TLE and 16 MRI-neg TLE) for whom functional MRI data has
47 already been reported²⁹. The diagnosis of TLE was based on clinical history, video-
48 EEG recording of typical seizures, and a congruent abnormality on nuclear medicine
49 studies (temporal lobe hypometabolism on FDG-PET, or ictal hyperperfusion on ictal-
50 interictal SPECT). Patients with independent seizures from both left and right
51 temporal lobes were excluded.

52 Routine clinical MRIs acquired with our epilepsy protocol³⁰ were reviewed by both a
53 Neuroradiologist and an Epileptologist. Patients were only included if there were
54 characteristic features of unilateral hippocampal sclerosis (HS-TLE) or if the mesial
55 temporal regions appeared normal and no other potentially epileptogenic abnormality
56 was seen (MRI-neg TLE). Patients were excluded if there was previous neurosurgery
57 or any other potentially epileptogenic intracranial abnormality.

58 Twenty-nine HS-TLE and 28 MRI-neg TLE patients met inclusion criteria and
59 consented to participate. Two MRI-neg TLE participants were excluded; one had
60 inadequate image acquisition, and the other had subtle but pathological enlargement

of the amygdala on subsequent imaging. To balance epilepsy groups for gender and epileptic side, three female patients with right HS-TLE were randomly selected and removed. The final analysis included 26 HS-TLE, and 26 MRI-neg TLE (Table 1).

First-degree relatives of HS-TLE, who had no personal history of seizures themselves, were drawn from our previous study¹⁸. Of 32 participants, 2 were excluded to meet age criteria, and a further 4 males were randomly selected and removed to balance the cohort for gender. The final analysis included 26 HS-1°Rel participants drawn from 15 families. In four of these families (27%) the proband had a known first-degree family history of seizures, compared to 8 families (31%) in the HS-TLE cohort.

Healthy adult volunteers were selected from historical controls that had diffusion weighted imaging performed at our institution. An equal number of males and females between ages 14 and 65 years of age were selected, resulting a group of 76 controls.

Consent was obtained from all participants, or their legal guardian in the case of minors. The study was approved by the Austin Health Human Research Ethics Committee.

MRI acquisition

Diffusion-weighted images were acquired on a 3T Siemens Tim Trio MRI scanner (60 directions, b-value 3000s/mm², 2.5mm isotropic voxels, echo time 110ms, repetition time 8300ms, parallel acceleration factor 2). An anatomical T1-weighted MPRAGE image was also acquired (voxel size 0.9mm isotropic, repetition time 1900ms, inversion time 900ms, echo time 2.6ms, flip angle 9°).

T1-weighted image processing

Hippocampal volumes were evaluated by manual tracing on the coronal T1-weighted image, using ImageJ software^{18,31}, with operators blinded to subject group and side of the brain. Estimated total intracranial volume was calculated using FreeSurfer 5.3³². Volumes were compared between groups using a linear mixed model, with intracranial volume as a covariate and family membership as a random effect.

Diffusion image pre-processing

Diffusion images were analysed using MRtrix3³³ (www.mrtrix.org). Pre-processing included correction for head motion, bias fields, and spatial up-sampling by a factor of two²⁸. Global intensity normalization was performed by scaling all volumes by the median intensity of white-matter voxels on $b=0$ s/mm² images. The fibre orientation distribution (FOD) was calculated at each voxel using robust constrained spherical deconvolution³⁴ with a group average response function²⁸.

Patients with right-sided TLE had their FOD images flipped right-to-left, with re-orientation of FOD lobes, so that the epileptic side was aligned across all patients. An equal proportion (50%) of participants within the Control and HS-1°Rel cohorts were randomly selected, stratified by gender and age-by-decade, and their FOD images flipped right-to-left also.

Study-specific FOD template and tractogram

A left-right symmetrical study-specific FOD template was created from a subset of participants (12 HS-TLE, 12 MRI-neg TLE, 12 controls), randomly selected with stratification for gender and epileptic side. FOD images ($l_{\max}=4$) and their left-to-right flipped mirror counterparts, were repeatedly non-linearly registered to the previous group mean³⁵. The registration step included reorientation of the FOD³⁶. This process was iterated until sufficient convergence of the group mean FOD image, obtained after 10 iterations. FOD images for all participants were then registered to this unbiased symmetrical template.

Whole-volume probabilistic tractography was performed on the final group mean FOD image (at $l_{\max} = 8$) to generate 20 million streamlines, and filtered to 2 million streamlines using the SIFT algorithm³⁷.

Fixel-based calculation of Fibre Density and Cross-section (FDC)

The amplitude of an FOD lobe, at the relatively high b-value used here, is proportional to the volume of restricted water within axons of the given orientation, and can be interpreted as the apparent Fibre Density (FD) in that direction²⁸. The FD is calculated for each fixel by integrating over the respective FOD lobe³⁷.

Additional valuable information about tract morphology is available from the registration of each image to the template. By considering the local expansion or

contraction of the warp field perpendicular to the fixel orientation, a relative measure of the fibre cross-section (FC) is obtained²⁶.

Multiplying FD and FC at each fixel gives a combined measure of fibre density and cross-section (FDC)²⁶. FDC is a more comprehensive measure of intra-axonal volume within a pathway, as it accounts for both microscopic axonal changes (detected as differences in diffusion signal), as well as macroscopic changes in a fibre-bundle (detected as relative differences in the registration warps).

Fixel-based statistical comparison between groups

Statistical comparisons of FDC between groups were performed at each fixel, by one-way ANCOVA with the contrasts of (i) HS-TLE versus controls, (ii) MRI-neg TLE versus controls, and (iii) 1st-Rel-TLE versus controls. Covariates of no interest were age, the estimated total intracranial volume and whether the image had been flipped right-to-left. Connectivity-based Fixel Enhancement (CFE) was used for statistical inference, using streamlines from the template tractogram and default parameters (smoothing=10mm FWHM, C=0.5, E=2, H=3)²⁷. A corrected p-value was assigned to each fixel using permutation testing, with 5000 permutations and preserving family grouping within each permutation.

Definition of tracts-of-interest

Seventeen named white matter tracts were selected for further characterization: the fornix, dorsal cingulum, parahippocampal cingulum, uncinate, arcuate, inferior fronto-occipital and inferior longitudinal fasciculi on each side, and the anterior commissure, bi-temporal fibers of the tapetum (traversing the splenium of the corpus callosum), and bi-frontal fibers traversing the anterior corpus callosum. For each tract, appropriate streamlines were selected from the template tractogram using a semi-automated rule-based approach, and used to create a fixel-mask (see images in Fig. 4).

Comparison of mean FDC within tracts-of-interest

The weighted mean FDC was calculated for each tract-of-interest, by summing along each streamline, dividing by streamline length, and averaging over all streamlines. Each group was compared to controls using a linear mixed model ('lme4', 'lsmeans' and 'pbrktest' packages in R) with covariates for intracranial volume, age and

whether right-to-left flipped, and with family membership included as a random effect. Multiple comparisons were corrected using the Dunnett procedure between groups ('multcomp' package) then Bonferroni correction for 24 tracts. Results are displayed as the percentage difference in mean FDC from controls, with 95% confidence intervals.

An individual asymmetry index (AI) was calculated for each bilateral tract-of-interest, with results displayed as difference from controls $\left(\overline{AI}_{\text{HS-TLE}} - \overline{AI}_{\text{Controls}}\right)$.

$$AI = \frac{\text{tractFDC}_{\text{Left}} - \text{tractFDC}_{\text{Right}}}{\text{tractFDC}_{\text{Left}} + \text{tractFDC}_{\text{Right}}}$$

White matter atrophy versus hippocampal volumes, seizure frequency and epilepsy duration

An exploratory post-hoc analysis was performed to examine the relationship between hippocampal volumes and tract-specific white matter atrophy (in 'R' notation):

```
tractFDC ~ Group + RLFlip + Age + IntracranialVolume + HippVol:Group
```

Ipsilateral hippocampal volumes (HippVol) were mean centered within-group to remove collinearity. For each tract-of-interest, a partial F-test was first performed for any difference between groups, and significant tracts were further tested for pairwise differences in the HippVol coefficient (unpaired t-tests with Dunnett correction, uncorrected for the number of tracts examined).

A second post-hoc linear model was used to explore the relationship between FDC and patient-estimated seizure frequency and epilepsy duration, again with these predictors mean-centered within group:

```
tractFDC ~ Group + SeizureFrequency:Group + EpilepsyDuration:Group
+ RLFlip + IntracranialVolume
```

RESULTS

TABLE 1 AROUND HERE

Clinical findings

Patients with HS-TLE had a longer duration of epilepsy than those with MRI-neg TLE (mean 22.2 years versus 14.0 years, 2-tailed t-test $p=0.004$, Table 1), reflecting

generally earlier onset of epilepsy in the HS-TLE group. Frequency of focal and generalized seizures was not significantly different between the TLE groups. Most patients were on 2 or 3 anti-epileptic medications. This was not significantly different between groups and included similar medications and dose ranges. Of the HS-TLE patients, 22 (12 right, 10 left) subsequently proceeded to anterior temporal lobe resection, with hippocampal sclerosis demonstrated in all, and a good outcome with respect to seizures in 86% (19 of 22). In MRI-neg TLE, 8 patients (6 right, 2 left) proceeded to surgery with 7 having anterior temporal lobectomy and one a focal posterior temporal resection. Seven had no evidence of hippocampal sclerosis or dysplasia, however in one patient neuronal loss consistent with mild hippocampal sclerosis was found. A good outcome with respect to seizures was achieved in 63% (5 of 8).

FIGURE 1 AROUND HERE

Hippocampal Volumetry

In HS-TLE, the volume of the affected hippocampus was significantly smaller than in controls (Fig. 1a; Left hippocampal volume [mean±sd] in Left HS-TLE 1981±273mm³ versus Controls 3364±318mm³ adjusted for intracranial volume, 95% CI on difference [-1102, -1667mm³]; Right hippocampal volume in Right HS-TLE 2194±486mm³ versus Controls 3450±345mm³, 95% CI on difference [-956, -1557mm³]). There was no significant difference between HS-TLE and controls at the contralateral hippocampus.

In MRI-neg TLE, hippocampal volumes were not different to controls (Fig. 1b; Left hippocampal volume in Left MRI-neg TLE 3312±434mm³; Right hippocampal volume in Right MRI-neg TLE 3622±350mm³).

In HS-1°Rel, left and right hippocampal volumes were both significantly smaller than controls (Fig. 1c; Left hippocampal volume in HS-1°Rel 2938±244mm³, 95% CI on difference [-188, -710mm³]; Right hippocampal volume in HS-1°Rel 2856±259mm³, 95% CI on difference [-322, -853mm³]). Volumes were also smaller than the contralateral hippocampus in HS-TLE (95% CI on difference [-155, -813mm³]).

FIGURE 2 AROUND HERE

Fixel Based Analysis

209 Reduced white matter fiber density and cross-section (FDC) was seen in both HS-
210 TLE and MRI-neg TLE compared to controls. In the HS-1°Rel group no fixels were
211 identified as significantly different from controls.

212 Fixel based analysis: HS-TLE versus Controls

213 In HS-TLE, decreased FDC was seen ipsilateral to the epileptic focus in fixels of the
214 fornix, uncinate fasciculus, ILF, IFOF, parahippocampal cingulum and arcuate
215 fasciculus (Fig. 2a). Bilateral reduction was seen at fixels of the anterior commissure,
216 tapetum, anterior corpus callosum and dorsal cingulum (more ipsilateral than
217 contralateral). The fixels showing greatest magnitude in reduction of FDC, typically
218 30-40% smaller than controls, were at the temporal pole, inferior temporal white
219 matter, and at the anterior commissure. The reduction in FDC in most tracts was
220 produced by both reduced fiber density (FD) and reduced cross-sectional area (FC).
221 Exceptions were the fornix and anterior commissure, which are thin tracts only a few
222 voxels across and show reduction primarily in FD.

223 Fixel based analysis: MRI-neg TLE versus Controls

224 In MRI-neg TLE, lateralized reduction in FDC ipsilateral to the epileptic focus was
225 seen in a small number of fixels, at the ILF and tapetum fibers of the affected
226 temporal lobe. Bilateral reduction in FDC was more extensive, seen symmetrically in
227 the tapetum, anterior corpus callosum, and in the dorsal cingulum bilaterally (Fig. 2).
228 The greatest FDC reduction was at the ipsilateral tapetum and bilateral dorsal
229 cingulum (~25% decrease). Reduced FDC in the corpus callosum and the tapetum
230 was primarily a consequence of reduced fiber density (FD). Reduced FDC in the
231 bilateral dorsal cingulum bilaterally had similar contributions from FD and FDC.

232 FIGURE 3 AROUND HERE

233 ***Tract-of-Interest Analysis***

234 Within the tracts-of-interest, HS-TLE showed greatest atrophy at the ipsilateral
235 uncinate, parahippocampal cingulum, fornix, arcuate and at the anterior commissure
236 (Fig. 3a). The MRI-neg TLE group had mildly reduced FDC at the uncinate, but the
237 remainder of these tracts were not significantly different from controls.

Both HS-TLE and MRI-neg TLE showed similar atrophy at the bilateral dorsal cingulum, frontal corpus callosum, tapetum and ipsilateral IFOF. The magnitude of FDC reduction here was similar in both forms of TLE (Fig. 3b).

FIGURE 4 AROUND HERE

Left-right asymmetry in tracts-of-interest showed that atrophy was strongly lateralized to the epileptic side in HS-TLE (Fig. 5). This effect was strongest in the uncinate fasciculus, parahippocampal cingulum and hippocampal fornix. No significant asymmetry was seen in MRI-neg TLE, although a non-significant trend toward ipsilateral atrophy may be present at the uncinate fasciculus and ILF.

White matter atrophy within tracts-of-interest in HS-1°Rel

The HS-1°Rel group did not show any significant tract-of-interest atrophy compared to controls (Fig. 3) nor any significant asymmetry (Fig. 4). However, in all tracts tested the group-mean FDC was around 4-8% smaller than in controls, suggesting a global trend in this direction.

Relationship between hippocampal volume and atrophy within tracts-of-interest

The post-hoc linear model relating ipsilateral hippocampal volume to white matter atrophy showed a difference between groups at the ipsilateral parahippocampal cingulum ($F_{139,3}=2.93$, $p=0.036$) and at the anterior commissure ($F_{139,3}=2.82$, $p=0.041$). This was driven by the HS-TLE group, where a smaller affected hippocampus corresponded to the extent of white matter atrophy, a relationship that was not seen in controls (Ipsilateral parahippocampal cingulum: -19.3% FDC per mL of hippocampal atrophy, -2.0% per mL in controls, $p=0.028$; Anterior commissure: -9.6% FDC per mL of hippocampal atrophy, +4.4% per mL in controls, $p=0.049$). Surprisingly there was no detected difference between the groups for this relationship at the hippocampal fornix ($F_{139,3}=0.80$, $p=0.49$).

Relationships between seizure frequency, epilepsy duration and white matter atrophy

The linear relationships between seizure frequency and FDC, and between epilepsy duration and FDC, did not reach statistical significance at any of the tracts-of-interest. However the estimated effect size for patient-reported seizure frequency was maximal at the bilateral dorsal cingulum in both epilepsy groups (around -1.4% per seizure/month in HS-TLE and -0.8% per seizure/month in MRI-neg TLE). Epilepsy

duration in HS-TLE had greatest effect size at the hippocampal fornix, ipsilateral parahippocampal cingulum and anterior commissure (range -2.0% to -2.5% per decade of epilepsy). In MRI-neg TLE the maximal effect of epilepsy duration was at the ipsilateral uncinate fasciculus, anterior commissure and tapetum fibers (range -2.7% to -4.6% per decade of epilepsy).

DISCUSSION

In this study we considered HS-TLE as an archetype of focal epilepsy. Atrophy of specific white matter tracts was identified using the metric of Fiber Density & Cross-section in temporal lobe tracts ipsilateral to the seizure focus (parahippocampal cingulum, uncinate, fornix, inferior longitudinal and inferior fronto-occipital fasciculi, tapetum and anterior commissure), as well as in bilateral tracts that are less directly associated with the seizure focus (bilateral dorsal cingulum, frontal and parieto-occipital fibers of the corpus callosum).

To interpret these findings we applied the following premises: (1) abnormalities that are common to HS-1°Rel and HS-TLE are likely to have a genetic basis and predate the onset of epilepsy; (2) abnormalities that are unique to HS-TLE compared to MRI-neg TLE are likely to be related to the underlying pathological process of HS; (3) correlation between severity of white matter abnormalities and amount of hippocampal atrophy is further evidence of a relationship to the underlying pathological process; (4) abnormalities that are common to both HS-TLE and MRI-neg TLE are likely to be the consequence of seizures and/or anti-epilepsy medications.

Common abnormalities in HS-1°Rel and HS-TLE: weak evidence for a trend toward diffuse mild white matter atrophy as a familial trait?

Bilaterally reduced hippocampal volumes in the relatives of people with HS-TLE (by ~15%) indicates a familial trait with a likely genetic basis¹⁸. In HS-TLE hippocampal atrophy was greatest at the affected hippocampus (by 38%), but no atrophy was found on the contralateral side. Although reduced contralateral volume might be expected in HS-TLE given our findings in the relatives, this was largely precluded by our recruitment criteria of strictly unilateral hippocampal atrophy in the HS-TLE group. An alternative speculative explanation may be that contralateral hippocampal volume increases in HS-TLE following the development of epilepsy.

No significant white-matter difference was found between HS-1°Rel and controls, either in the fixel-based or the tracts-of-interest analysis. However there is a trend toward reduced FDC in HS-1°Rel in all tracts studied (by ~4-8%), indicating that a mild but diffuse reduction in white matter as identified by Scanlon²³ could also be present in our cohort. Overall our interpretation is that the white matter atrophy found in HS-TLE is related to the disease and/or treatment, rather than primarily being a familial trait.

Unique abnormalities in HS-TLE: specific associations of the underlying disease

White matter atrophy in HS-TLE, but not in MRI-neg TLE, is found at the antero-mesial temporal lobe tracts on the epileptic side: the ipsilateral parahippocampal cingulum, fornix, uncinate, ILF, arcuate and the anterior commissure (Figs. 2a & 3a). Possible causative factors that are unique to HS-TLE include the occurrence of childhood febrile convulsions (more frequent in HS-TLE than MRI-neg TLE), the age of onset of epilepsy (earlier in HS-TLE), hippocampal pathology (neuronal loss and gliosis in HS, but minimal changes in MRI-neg TLE), and subtle differences in functional network abnormalities (with a mesial temporal rather than neocortical emphasis²⁹).

Correlated hippocampal atrophy and white matter atrophy in HS-TLE: an indicator of common causation in a subset of tracts?

In HS-TLE, the parahippocampal cingulum and the anterior commissure both show an association between the hippocampal volume and the extent of white matter atrophy. This proportional relationship between the extent of injury at these tracts and in the hippocampus suggests that a common cause is at work, with the underlying process driving hippocampal sclerosis also being a contributor to white matter atrophy in these tracts.

We did not find a correlation between hippocampal volume and the extent of atrophy at the corpus callosum or cingulum, unlike Scanlon et al.⁸. This may relate to our different statistical approach which employed a separate linear regression slope for each group. Otherwise our results are in agreement with previous studies that have not shown any association between hippocampal volume and the ILF, uncinate or arcuate fasciculus³⁸ or temporal lobe white matter³⁹. It was surprising that we found no linear

relationship at the ipsilateral fornix given the well-established histological abnormalities of the fornix in HS-TLE⁴⁰. This may represent a “floor” effect due to the severe epilepsy in our cohort and the extent of fornix atrophy.

Common bilateral abnormalities in HS-TLE and MRI-neg TLE: a non-specific effect of global insults

Regions of common white matter atrophy were found in both forms of TLE at the bilateral dorsal cingulum, and at the inter-hemispheric fibres of the anterior corpus callosum (projecting symmetrically toward bilateral dorsolateral frontal cortex) and posterior corpus callosum (projecting symmetrically toward bilateral parietal cortex). Similar diffusion changes affecting the cingulum and/or corpus callosum have also been detected in other forms of epilepsy, including frontal lobe epilepsy with cortical dysplasia^{1,3}, and in a mixed cohort of both focal and generalized epilepsy⁴¹, suggesting that the causation is independent of both the underlying disease pathology and the location of the epileptic focus. The location and connections of these tracts makes it difficult to explain this atrophy as a direct consequence of unilateral focal seizures in TLE, particularly as the expected connecting pathway from the temporal lobe of the parahippocampal cingulum appears to be relatively spared in MRI-neg TLE. Neither patient-reported seizure frequency nor disease duration showed a significant correlation at these tracts in our analysis, although a correlation with epilepsy duration has been found at the corpus callosum by some diffusion studies^{39,42}. Furthermore, greater change at the corpus callosum has been found with refractory versus benign epilepsy⁴³, in association with bilateral temporal interictal epileptiform discharges⁴⁴, as well as in an experimental rat model of focal epilepsy where no anti-epileptic medications were used⁴⁵. Thus our interpretation is that atrophy in these tracts is likely to be a consequence of factors that have a bilateral and symmetrical impact, with generalized seizures, anti-epileptic medications, and other global factors such as psychological and socioeconomic status all being possible causes.

Common ipsilateral abnormalities in HS-TLE and MRI-neg TLE: focal seizures as a cause of lateralized white matter atrophy

A common unilateral finding in both TLE groups was reduced FDC of the ipsilateral tapetum, where this tract leaves the posterior temporal lobe (Fig. 2). The spatial

association of this tract with the expected seizure focus points to propagation of focal seizures as the possible cause. As the only finding that was clearly lateralized in MRI-neg TLE, targeted assessment of this region may be useful in future studies that aim to lateralize TLE on the basis of white matter atrophy. In the tract-of-interest analyses, the ipsilateral uncinate and IFOF were also abnormal in MRI-neg TLE, but as they did not show abnormal asymmetry are less likely to be useful markers for determining the epileptic side.

The idea that recurrent focal seizures can cause localized chronic white matter injury has been widely discussed but remains controversial, with some longitudinal studies detecting no change^{46,47}. We did not find a significant association between FDC and the retrospective patient-report of seizure frequency, although this measure may not be an individually accurate estimate of the true number of events⁴⁸. If seizures are assessed prospectively on EEG, white matter measurements at the ipsilateral uncinate, arcuate and ILF in HS-TLE can be associated with time since last seizure³⁸, indicating that seizure-related vasogenic oedema can occur in propagation tracts. Other studies have compared lateralized diffusion abnormalities with duration of epilepsy and found ipsilateral correlations at the temporal lobe, parahippocampal gyrus and internal capsule³⁹, the uncinate and arcuate in left TLE⁴⁹, or in the ipsilateral cingulum, fornix and external capsule in right TLE⁵⁰. Taken together, these findings support recurrent focal seizures as a cause of progressive ipsilateral white atrophy, but that demonstrating this conclusively is susceptible to variability between cohorts and methodologies.

FIGURE 5 AROUND HERE

Factors contributing to white matter atrophy: a model for focal epilepsy

Our findings in HS-TLE suggest a multi-factorial model for the development of tract-specific white matter atrophy in focal epilepsy. Genetic factors predisposing to HS-TLE (Fig. 5A) are associated with bilaterally small hippocampi but produce minimal white matter change. An initial triggering insult (Fig. 5B), such as a febrile convulsion, may cause both grey and white matter injury at the hippocampus and closely related tracts (especially the parahippocampal cingulum and anterior commissure), with the development of epilepsy. Repeated focal seizures (Fig. 5C) may cause progressive white matter injury in tracts radiating away from the seizure

397 focus (eg. the tapetum, uncinate fasciculus and IFOF). Meanwhile, repeated global
398 insults such as generalized seizures and/or the effects of medications (Fig. 5D)
399 contribute to bilateral changes in tracts distant from the focus, particularly at the
400 bilateral dorsal cingulum and the corpus callosum.

401 Several aspects of this model require further validation, as the main limitation of this
402 study is its cross-sectional design. Longitudinal studies in both human focal epilepsy
403 and in experimental animal models will be needed to further isolate and confirm the
404 consequences of repeated focal seizures, generalized seizures and anti-epileptic
405 medications in specific white-matter tracts.

ACKNOWLEDGEMENTS

We thank the patients and clinicians of the Austin Hospital Comprehensive Epilepsy Programme; Farnoosh Sadeghian, Mira Semmelroch and Susan Palmer for assistance with coordination of participants and data processing; and Dr Heath Pardoe for assistance with initial steps of the T1-weighted imaging data analysis. This study was supported by the National Health and Medical Research Council (NHMRC) of Australia (program grant 628952 and project grant 1081151), a computation grant from the Victorian Life Sciences Computation Initiative, and the Victorian Government Operational Infrastructure Support Program. DNV was supported by an NHMRC Postgraduate Scholarship (1055877) and a Windermere Foundation Doctoral Scholarship. JDT acknowledges financial support from the Department of Health via the National Institute for Health Research (NIHR) comprehensive Biomedical Research Centre award to Guy's & St Thomas' NHS Foundation Trust in partnership with King's College London and King's College Hospital NHS Foundation Trust. GDJ is supported by an NHMRC practitioner fellowship (1060312).

AUTHOR CONTRIBUTIONS

Concept and study design: D.N.V. and G.D.J. Data acquisition and analysis: D.N.V., D.R., E.C., M-H.T. and J-D.T. Drafting the manuscript and figures: D.N.V., D.R., J-D.T., A.C., G.D.J.

POTENTIAL CONFLICTS OF INTEREST

Nothing to report

REFERENCES

1. Campos BM, Coan AC, Beltramini GC, et al. White matter abnormalities associate with type and localization of focal epileptogenic lesions. *Epilepsia* 2015;56(1):125–132.
2. Diehl B, Tkach J, Piao Z, et al. Diffusion tensor imaging in patients with focal epilepsy due to cortical dysplasia in the temporo-occipital region: Electro-clinico-pathological correlations. *Epilepsy Res.* 2010;90(3):178–187.
3. Fonseca V de C, Yasuda CL, Tedeschi GG, et al. White Matter Abnormalities in Patients with Focal Cortical Dysplasia Revealed by Diffusion Tensor Imaging Analysis in a Voxelwise Approach. *Front. Neurol.* 2012;3
4. Bernasconi N, Duchesne S, Janke A, et al. Whole-brain voxel-based statistical analysis of gray matter and white matter in temporal lobe epilepsy. *NeuroImage* 2004;23(2):717–723.
5. Kim H, Harrison A, Kankirawatana P, et al. Major white matter fiber changes in medically intractable neocortical epilepsy in children: A diffusion tensor imaging study. *Epilepsy Res.* 2013;103(2–3):211–220.
6. Eriksson SH, Rugg-Gunn FJ, Symms MR, et al. Diffusion tensor imaging in patients with epilepsy and malformations of cortical development. *Brain* 2001;124(3):617–626.
7. Keller SS, Ahrens T, Mohammadi S, et al. Voxel-Based Statistical Analysis of Fractional Anisotropy and Mean Diffusivity in Patients with Unilateral Temporal Lobe Epilepsy of Unknown Cause. *J. Neuroimaging* 2013;23(3):352–9.
8. Scanlon C, Mueller SG, Cheong I, et al. Grey and white matter abnormalities in temporal lobe epilepsy with and without mesial temporal sclerosis. *J. Neurol.* 2013;260(9):2320–2329.
9. Whelan CD, Alhusaini S, O'Hanlon E, et al. White matter alterations in patients with MRI-negative temporal lobe epilepsy and their asymptomatic siblings. *Epilepsia* 2015;56(10):1551–1561.
10. Richardson MP. Large scale brain models of epilepsy: dynamics meets connectomics. *J. Neurol. Neurosurg. Psychiatry* 2012;83(12):1238–1248.
11. Li LM, Fish DR, Sisodiya SM, et al. High resolution magnetic resonance imaging in adults with partial or secondary generalised epilepsy attending a tertiary referral unit. *J. Neurol. Neurosurg. Psychiatry* 1995;59(4):384–387.
12. Wieser H-G. Mesial Temporal Lobe Epilepsy with Hippocampal Sclerosis. *Epilepsia* 2004;45(6):695–714.
13. Blümcke I, Thom M, Wiestler OD. Ammon's Horn Sclerosis: A Maldevelopmental Disorder Associated with Temporal Lobe Epilepsy. *Brain Pathol.* 2002;12(2):199–211.
14. Kim SE, Andermann F, Olivier A. The clinical and electrophysiological characteristics of temporal lobe epilepsy with normal MRI. *J. Clin. Neurol.* 2006;2(1):42–50.

- 469 15. Cohen-Gadol AA, Bradley CC, Williamson A, et al. Normal magnetic resonance
470 imaging and medial temporal lobe epilepsy: the clinical syndrome of paradoxical
471 temporal lobe epilepsy. *J. Neurosurg.* 2005;102(5):902–909.
- 472 16. Immonen A, Jutila L, Muraja-Murro A, et al. Long-term epilepsy surgery outcomes
473 in patients with MRI-negative temporal lobe epilepsy. *Epilepsia*
474 2010;51(11):2260–2269.
- 475 17. Carne RP, O'Brien TJ, Kilpatrick CJ, et al. MRI-negative PET-positive temporal lobe
476 epilepsy: a distinct surgically remediable syndrome. *Brain J. Neurol.* 2004;127(Pt
477 10):2276–2285.
- 478 18. Tsai M-H, Pardoe HR, Perchyonok Y, et al. Etiology of hippocampal sclerosis:
479 Evidence for a predisposing familial morphologic anomaly. *Neurology*
480 2013;81(2):144–149.
- 481 19. Diniz PRB, Velasco TR, Salmon CEG, et al. Extratemporal damage in temporal lobe
482 epilepsy: magnetization transfer adds information to volumetric MR imaging. *AJNR*
483 *Am. J. Neuroradiol.* 2011;32(10):1857–1861.
- 484 20. Yasuda CL, Valise C, Saúde AV, et al. Dynamic changes in white and gray matter
485 volume are associated with outcome of surgical treatment in temporal lobe
486 epilepsy. *NeuroImage* 2010;49(1):71–79.
- 487 21. Yu A, Li K, Li L, et al. Whole-brain voxel-based morphometry of white matter in
488 medial temporal lobe epilepsy. *Eur. J. Radiol.* 2008;65(1):86–90.
- 489 22. Braga B, Yasuda CL, Cendes F. White Matter Atrophy in Patients with Mesial
490 Temporal Lobe Epilepsy: Voxel-Based Morphometry Analysis of T1- and T2-
491 Weighted MR Images. *Radiol. Res. Pract.* 2012;e481378.
- 492 23. Scanlon C, Ronan L, Doherty CP, et al. MRI-Based Brain Structure Volumes in
493 Temporal Lobe Epilepsy Patients and their Unaffected Siblings: A Preliminary
494 Study. *J. Neuroimaging* 2013;23(1):64–70.
- 495 24. Mueller SG, Laxer KD, Cashdollar N, et al. Voxel-based Optimized Morphometry
496 (VBM) of Gray and White Matter in Temporal Lobe Epilepsy (TLE) with and
497 without Mesial Temporal Sclerosis. *Epilepsia* 2006;47(5):900–907.
- 498 25. Farquharson S, Tournier J-D, Calamante F, et al. White matter fiber tractography:
499 why we need to move beyond DTI. *J. Neurosurg.* 2013;118(6):1367–1377.
- 500 26. Raffelt DA, Tournier J-D, Smith RE, et al. Investigating white matter fibre density
501 and morphology using fixel-based analysis. *NeuroImage*
502 2016;DOI:10.1016/j.neuroimage.2016.09.029
- 503 27. Raffelt DA, Smith RE, Ridgway GR, et al. Connectivity-based fixel enhancement:
504 Whole-brain statistical analysis of diffusion MRI measures in the presence of
505 crossing fibres. *NeuroImage* 2015;117:40–55.
- 506 28. Raffelt D, Tournier J-D, Rose S, et al. Apparent Fibre Density: A novel measure for
507 the analysis of diffusion-weighted magnetic resonance images. *NeuroImage*
508 2012;59(4):3976–3994.

- 509 29. Vaughan DN, Rayner G, Tailby C, Jackson GD. MRI-negative temporal lobe epilepsy:
510 A network disorder of neocortical connectivity. *Neurology*
511 2016;DOI:10.1212/WNL.0000000000003289
- 512 30. Jackson GD, Badawy RAB. Selecting patients for epilepsy surgery: Identifying a
513 structural lesion. *Epilepsy Behav.* 2011;20(2):182–189.
- 514 31. Pardoe HR, Pell GS, Abbott DF, Jackson GD. Hippocampal volume assessment in
515 temporal lobe epilepsy: How good is automated segmentation? *Epilepsia*
516 2009;50(12):2586–2592.
- 517 32. Fischl B, Dale AM. Measuring the thickness of the human cerebral cortex from
518 magnetic resonance images. *Proc. Natl. Acad. Sci. U. S. A.* 2000;97(20):11050–
519 11055.
- 520 33. Tournier J-D, Calamante F, Connelly A. MRtrix: Diffusion tractography in crossing
521 fiber regions. *Int. J. Imaging Syst. Technol.* 2012;22(1):53–66.
- 522 34. Tournier J-D, Calamante F, Connelly A. A robust spherical deconvolution method
523 for the analysis of low SNR or low angular resolution diffusion data. In:
524 Proceedings of the International Society for Magnetic Resonance in Medicine. Salt
525 Lake City, USA: 2013
- 526 35. Raffelt D, Tournier J-D, Fripp J, et al. Symmetric diffeomorphic registration of fibre
527 orientation distributions. *NeuroImage* 2011;56(3):1171–1180.
- 528 36. Raffelt D, Tournier J-D, Crozier S, et al. Reorientation of fiber orientation
529 distributions using apodized point spread functions. *Magn. Reson. Med.*
530 2012;67(3):844–855.
- 531 37. Smith RE, Tournier J-D, Calamante F, Connelly A. SIFT: Spherical-deconvolution
532 informed filtering of tractograms. *NeuroImage* 2013;67:298–312.
- 533 38. Concha L, Kim H, Bernasconi A, et al. Spatial patterns of water diffusion along white
534 matter tracts in temporal lobe epilepsy. *Neurology* 2012;79(5):455–462.
- 535 39. Keller SS, Schoene-Bake J-C, Gerdes JS, et al. Concomitant fractional anisotropy and
536 volumetric abnormalities in temporal lobe epilepsy: cross-sectional evidence for
537 progressive neurologic injury. *PloS One* 2012;7(10):e46791.
- 538 40. Concha L, Livy DJ, Beaulieu C, et al. In vivo diffusion tensor imaging and
539 histopathology of the fimbria-fornix in temporal lobe epilepsy. *J. Neurosci.*
540 2010;30(3):996–1002.
- 541 41. Hutchinson E, Pulsipher D, Dabbs K, et al. Children with new-onset epilepsy exhibit
542 diffusion abnormalities in cerebral white matter in the absence of volumetric
543 differences. *Epilepsy Res.* 2010;88(2–3):208–214.
- 544 42. Yin X, Qiu S, Liu Z, et al. Extratemporal abnormalities of brain parenchyma in young
545 adults with temporal lobe epilepsy: A diffusion tensor imaging study. *Clin. Radiol.*
546 2014;69(6):589–596.
- 547 43. Caligiuri ME, Labate A, Cherubini A, et al. Integrity of the corpus callosum in
548 patients with benign temporal lobe epilepsy. *Epilepsia* 2016;57(4):590–596.

549 44. Pustina D, Doucet G, Skidmore C, et al. Contralateral interictal spikes are related to
550 tapetum damage in left temporal lobe epilepsy. *Epilepsia* 2014;55(9):1406–1414.

551 45. Otte WM, Dijkhuizen RM, Meer MPA van, et al. Characterization of Functional and
552 Structural Integrity in Experimental Focal Epilepsy: Reduced Network Efficiency
553 Coincides with White Matter Changes. *PLOS ONE* 2012;7(7):e39078.

554 46. Holtkamp M, Schuchmann S, Gottschalk S, Meierkord H. Recurrent seizures do not
555 cause hippocampal damage. *J. Neurol.* 2004;251(4):458–463.

556 47. Liu RSN, Lemieux L, Bell GS, et al. Cerebral damage in epilepsy: a population-based
557 longitudinal quantitative MRI study. *Epilepsia* 2005;46(9):1482–1494.

558 48. Hoppe C, Poepel A, Elger CE. Epilepsy: accuracy of patient seizure counts. *Arch.*
559 *Neurol.* 2007;64(11):1595–1599.

560 49. Govindan RM, Makki MI, Sundaram SK, et al. Diffusion tensor analysis of temporal
561 and extra-temporal lobe tracts in temporal lobe epilepsy. *Epilepsy Res.*
562 2008;80(1):30–41.

563 50. Chiang S. White matter structural connectivity changes correlate with epilepsy
564 duration in temporal lobe epilepsy. *Epilepsy Res.* 2016;120:37–46.

565

566 **FIGURE LEGENDS**

567 **Figure 1 - Hippocampal volumes in Patients and Controls**

568 Hippocampal volumes for each participant, measured by manual tracing on T1-
569 weighted images, and corrected for intracranial volume by linear regression. (a-c)
570 Crosshairs show the mean for each group, surrounded by the 95% confidence ellipse.
571 Crosshairs that fall outside another ellipse indicate a significant difference in group
572 means ($p < 0.05$). (d) Boxplots show the median and interquartile range for each group,
573 with whiskers to 1.5x interquartile range.

574 **Figure 2 - Fixels with significant white matter atrophy in TLE**

575 Fixels where Fiber Density & Cross-section (FDC) is less than in Controls ($p < 0.05$
576 FWE corrected), after accounting for intracranial volume, age and brain side (right-to-
577 left flip). The background image is the symmetrical study template, showing the
578 scalar magnitude of the fiber orientation distribution at each voxel. Tracts affected in
579 both HS-TLE and MRI-neg TLE are labelled in bold.

Accepted Article

Figure 3 - White matter atrophy in Tracts-of-interest

Fiber Density & Cross-section (FDC) averaged within tracts-of-interest, displayed as percentage difference from the Control mean, adjusted for age, intracranial volume and brain side. Horizontal bars show the 95% confidence interval. Non-significant results cross the control mean and are shown in lighter grey. (Left column) Tracts-of-interest with significant atrophy in HS-TLE compared to Controls. (Right column) Tracts-of-interest which show atrophy in both HS-TLE and MRI-neg TLE, of similar magnitude in both groups. Tracts-of-interest that did not reach statistical significance are omitted. 'HS'=TLE with Hippocampal Sclerosis. 'MRI-n'=MRI-negative TLE. '1°Rel': First degree relatives of people with HS-TLE. ILF: inferior longitudinal fasciculus. IFOF: inferior fronto-occipital fasciculus.

591 **Figure 4 - Asymmetry of white matter atrophy in tracts-of-interest**

592 Relative asymmetry index of Fiber Density & Cross section (FDC) for each tract-of-
593 interest. Relative asymmetry index = $(FDC_{left} - FDC_{right}) / (FDC_{left} + FDC_{right}) -$
594 $mean(AI_{controls})$.

Accepted Article

595 **Figure 5 - Factors contributing to white matter atrophy in Focal Epilepsy**

596

Accepted Article

597 **TABLES**

598 **Table 1:** Study participants

	N (M:F)	Epilepsy Side L:R	Age mean (range)	Onset age mean (range)	Focal Sz per month med (range)	Gen Sz per year med (range)	# AEDs mean (range)
HS-TLE	26 (13:13)	13:13	39.2 (14-65)	14.0 (1-31)	7.6 (0.1-35)	0.3 (0-39)	2.5 (1-5)
HS-1°Rel	26 (13:13)	-	37.6 (14-65)	-	-	-	-
MRI-neg TLE	26 (12:14)	13:13	34.8 (16-52)	22.2 (4-52)	8.7 (0.3-46)	0.7 (0-12)	2.2 (1-4)
Controls	72 (36:36)	-	36.5 (17-65)	-	-	-	-
			NS*	p=0.004†	NS†	NS†	NS

599 (*one-way ANOVA, †Mann-Whitney U test. Sz=seizures. Gen=Generalized)

600

601

Figure 1 - Hippocampal volumes in Patients and Controls

Hippocampal volumes for each participant, measured by manual tracing on T1-weighted images, and corrected for intracranial volume by linear regression. (a-c) Crosshairs show the mean for each group, surrounded by the 95% confidence ellipse. Crosshairs that fall outside another ellipse indicate a significant difference in group means ($p < 0.05$). (d) Boxplots show the median and interquartile range for each group, with whiskers to 1.5x interquartile range.

Fig. 1

Acc

Figure 1 - Hippocampal volumes in Patients and Controls
Hippocampal volumes for each participant, measured by manual tracing on T1-weighted images, and corrected for intracranial volume by linear regression. (a-c) Crosshairs show the mean for each group, surrounded by the 95% confidence ellipse. Crosshairs that fall outside another ellipse indicate a significant difference in group means ($p < 0.05$). (d) Boxplots show the median and interquartile range for each group, with whiskers to 1.5x interquartile range.

Acc

Figure 2 - Fixels with significant white matter atrophy in TLE

Fixels where Fiber Density & Cross-section (FDC) is less than in Controls ($p < 0.05$ FWE corrected), after accounting for intracranial volume, age and brain side (right-to-left flip). The background image is the symmetrical study template, showing the scalar magnitude of the fiber orientation distribution at each voxel. Tracts affected in both HS-TLE and MRI-neg TLE are labelled in bold.

Fig. 2

Accepted

Figure 2 - Fixels with significant white matter atrophy in TLE
Fixels where Fiber Density & Cross-section (FDC) is less than in Controls ($p < 0.05$ FWE corrected), after accounting for intracranial volume, age and brain side (right-to-left flip). The background image is the symmetrical study template, showing the scalar magnitude of the fiber orientation distribution at each voxel. Tracts affected in both HS-TLE and MRI-neg TLE are labelled in bold.

Accept

Figure 3 - White matter atrophy in Tracts-of-interest

Fiber Density & Cross-section (FDC) averaged within tracts-of-interest, displayed as percentage difference from the Control mean, adjusted for age, intracranial volume and brain side. Horizontal bars show the 95% confidence interval. Non-significant results cross the control mean and are shown in lighter grey. (Left column) Tracts-of-interest with significant atrophy in HS-TLE compared to Controls. (Right column) Tracts-of-interest which show atrophy in both HS-TLE and MRI-neg TLE, of similar magnitude in both groups.

Tracts-of-interest that did not reach statistical significance are omitted. 'HS'=TLE with Hippocampal Sclerosis. 'MRI-n'=MRI-negative TLE. '1°Rel': First degree relatives of people with HS-TLE. ILF: inferior longitudinal fasciculus. IFOF: inferior fronto-occipital fasciculus.

Fig. 3

Acc

Figure 3 - White matter atrophy in Tracts-of-interest
Fiber Density & Cross-section (FDC) averaged within tracts-of-interest, displayed as percentage difference from the Control mean, adjusted for age, intracranial volume and brain side. Horizontal bars show the 95% confidence interval. Non-significant results cross the control mean and are shown in lighter grey. (Left column) Tracts-of-interest with significant atrophy in HS-TLE compared to Controls. (Right column) Tracts-of-interest which show atrophy in both HS-TLE and MRI-neg TLE, of similar magnitude in both groups. Tracts-of-interest that did not reach statistical significance are omitted. 'HS'=TLE with Hippocampal Sclerosis. 'MRI-n'=MRI-negative TLE. '1°Rel': First degree relatives of people with HS-TLE. ILF: inferior longitudinal fasciculus. IFOF: inferior fronto-occipital fasciculus.

ACC

Figure 4 - Asymmetry of white matter atrophy in tracts-of-interest
 Relative asymmetry index of Fiber Density & Cross section (FDC) for each tract-of-interest. Relative asymmetry index = $(FDC_{left} - FDC_{right}) / (FDC_{left} + FDC_{right}) - \text{mean}(AI_{controls})$.

Fig. 4

Figure 4 - Asymmetry of white matter atrophy in tracts-of-interest
Relative asymmetry index of Fiber Density & Cross section (FDC) for each tract-of-interest. Relative asymmetry index = (FDCleft-FDCright)/(FDCleft+FDCright) -mean(AIcontrols).

Figure 5 - Factors contributing to white matter atrophy in Focal Epilepsy
Fig. 5

Accepted A

Figure 5 - Factors contributing to white matter atrophy in Focal Epilepsy

Accepted A

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Vaughan, DN;Raffelt, D;Curwood, E;Tsai, M-H;Tournier, J-D;Connelly, A;Jackson, GD

Title:

Tract-specific atrophy in focal epilepsy: Disease, genetics, or seizures?

Date:

2017-02-01

Citation:

Vaughan, D. N., Raffelt, D., Curwood, E., Tsai, M. -H., Tournier, J. -D., Connelly, A. & Jackson, G. D. (2017). Tract-specific atrophy in focal epilepsy: Disease, genetics, or seizures?. ANNALS OF NEUROLOGY, 81 (2), pp.240-250. <https://doi.org/10.1002/ana.24848>.

Persistent Link:

<http://hdl.handle.net/11343/292384>