Received Date : 18-Oct-2016

Revised Date : 15-Feb-2017

Accepted Date : 20-Feb-2017

Article type : Original Manuscript

Corresponding author mail id : louise.quesnel@usherbrooke.ca

SIZE, SEASON AND OFFSPRING SEX AFFECT MILK COMPOSITION AND JUVENILE SURVIVAL IN WILD KANGAROOS

Louise Quesnel^a, Allison MacKay^a, David M. Forsyth^{b,c}, Kevin R. Nicholas^d, and Marco
Festa-Bianchet^{a,c}

^aDépartement de biologie, Université de Sherbrooke, Sherbrooke, Canada

^bVertebrate Pest Research Unit, New South Wales Department of Primary Industries,
Orange, Australia

^cSchool of BioSciences, University of Melbourne, Melbourne, Australia

^dDepartment of Anatomy and Developmental Biology, Monash University, Melbourne,
Australia

Page heading: Milk composition in a seasonal environment

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/jzo.12453](https://doi.org/10.1111/jzo.12453)

This article is protected by copyright. All rights reserved

23

24 Key words: lactation, *Macropus giganteus*, maternal effects, multimodel inference,
25 timing of reproduction, Trivers-Willard hypothesis, sex allocation, milk composition

26

27 ABSTRACT

28 The main component of mammalian maternal care is milk production. The composition
29 of milk has been hypothesised to determine offspring survival, and to vary with offspring
30 sex. Few studies, however, have examined variation in milk composition of wild
31 mammals in seasonal environments, where environmental conditions can impact the
32 ability of mothers to provide care. We investigated individual differences in milk
33 composition and offspring survival in wild eastern grey kangaroos (*Macropus giganteus*).
34 We analyzed total protein and lipid concentrations in 103 milk samples from 91 females
35 over two years. In a year of low forage production, the few females that lactated were in
36 poorer condition and produced milk of lower energy content compared to females
37 lactating in a year of high forage production. Females nursing in late winter produced
38 milk with more lipids compared to females at the same stage of lactation in late spring,
39 whose milk had a higher proportion of protein. The milk of larger females contained
40 more protein than lipid, and females in better condition allocated higher proportions of
41 protein to sons than to daughters. Increased protein concentration in milk was correlated
42 with offspring longevity, and this effect was stronger for sons than daughters. A
43 seasonally-modulated lactation strategy enables the adjustment of milk composition to
44 available resources. Condition-specific protein allocation in favour of sons for mothers of
45 higher caring ability suggests adaptive sex-biased maternal care in this highly sexually
46 dimorphic mammal.

47 INTRODUCTION

48 Lactation is the most costly component of maternal care in mammals (Oftedal, 1985), and
49 the prolonged dependence of offspring on milk results in ubiquitous maternal effects
50 (Maestripieri & Mateo, 2009) with consequences for evolutionary and population
51 ecology. Maternal traits (Georges & Guinet, 2000; Bowen *et al.*, 2001) and differential

care for sons and daughters (Landete-Castillejos *et al.*, 2005; Robert & Braun, 2012) may contribute to variability in milk provisioning. Milk composition is correlated with maternal condition in pinnipeds (Iverson *et al.*, 1993; Crocker *et al.*, 2001), primates (Nommsen *et al.*, 1991; Tardif *et al.*, 2001), and ungulates (Landete-Castillejos *et al.*, 2005; Bovolenta *et al.*, 2013), but not in tammar wallabies (*Macropus eugenii*) or Columbian ground squirrels (*Urocitellus columbianus*) (Robert & Braun, 2012; Skibieli & Hood, 2015). Moreover, differences in milk provisioning affect offspring size (Mellish, Iverson & Bowen, 1999; Landete-Castillejos, Garcia & Gallego, 2001; Robert & Braun, 2012) and survival (Skibieli & Hood, 2015).

Sex-biased allocation (Trivers & Willard, 1973) is one potential source of variation in milk composition. In polygynous mammals, males have higher reproductive variance than females, therefore sons should benefit more than daughters from greater milk provisioning (Trivers & Willard, 1973). Hence, mothers with higher caring ability should favour sons, whereas mothers with lower ability should favour daughters. Some studies of milk composition have provided support for sex-biased allocation by reporting that sons received more (Landete-Castillejos *et al.*, 2005) or higher-energy milk (Powe, Knott & Conklin-Brittain, 2010) with greater lipid (Fujita *et al.*, 2012) or protein concentrations (Robert & Braun, 2012) than daughters. Other studies found no sex-related differences (Hinde, 2009; Hinde *et al.*, 2014).

Experimentally restricted diets have revealed the importance of nutrition on lactation performance (Roberts, Cole & Coward, 1985; Landete-Castillejos *et al.*, 2003; Bovolenta *et al.*, 2013). If the timing of reproduction affects milk provisioning (Landete-Castillejos *et al.*, 2001; Skibieli & Hood, 2015) then there could be large seasonal variation in maternal caring ability in the wild. Most studies investigating maternal care through milk, however, were performed in captive (Hinde, 2009; Robert & Braun, 2012), or semi-captive (Landete-Castillejos *et al.* 2001) environments. Hence, little is known about sex-biased milk allocation in wild mammals.

We studied milk composition in wild eastern grey kangaroos (*Macropus giganteus*) in a temperate environment. Similar to ungulates, kangaroos are large herbivores and males have higher reproductive variance than females (Fisher, Blomberg & Owens, 2002). The

short gestation and prolonged lactation of marsupials, however, make kangaroos ideal for studying postnatal care (Sharp *et al.*, 2016). Stage-specific changes in milk composition during marsupial lactation have been extensively studied in the congeneric tammar wallaby (Green, Newgrain & Merchant, 1980; Green, Griffiths & Leckie, 1983), with milk components changing substantially throughout lactation (Tyndale-Biscoe & Janssens, 1988). We identified determinants of milk lipid and protein, two macronutrients with high caloric density and sensitivity to maternal nutritional state (Parker, Barboza & Gillingham, 2009). We expected that females at the same stage of lactation would vary milk composition according to forage production. Male body size in kangaroos affects mating success (Rioux-Paquette *et al.*, 2015); we predicted that mothers with a higher ability to care would provide milk with higher protein content to sons than to daughters. Finally, we tested whether protein concentration at peak lactation affected offspring survival (Skibieli & Hood, 2015).

METHODS

Study area and data collection

The study area in Wilsons Promontory National Park, Victoria, Australia (38°57'S, 146°17'E) is a 1.1-km² grassland surrounding an emergency airstrip (Davis, Coulson & Forsyth, 2008), where grasses, sedges and herbs are the main forage (Davis, Forsyth & Coulson, 2010). The kangaroo population has been monitored since 2008, with annual recaptures of marked individuals (Gélin *et al.*, 2013). The climate is temperate, with mean maximum monthly temperatures ranging from 12.5 °C in July to 26.8 °C in February, and annual precipitation of 453–966 mm between 2012 and 2015 (Corner Inlet Station, 38°81'S, 146°19'E; www.bom.gov.au/climate/data/).

We used a pole syringe (King *et al.* 2011) to capture females carrying pouch young aged six to 10 months from mid-winter (July) to early summer (December) in 2014 and 2015. Young were transferred to a cloth bag, sexed and their head, foot and leg lengths (mm) measured. They were marked with coloured plastic ear tags (Leader Products Pty Ltd, Craigieburn, Australia). Mothers were weighed to the closest 0.25 kg and their hind leg length measured (mm). To stimulate milk letdown, oxytocin (Oxyto-Sure, Vetoquinol, concentration = 20 USP·mL⁻¹, dose = 12 µl·kg) was injected intramuscularly. After 1–4

minutes, milk was expressed into a screw-top vial by gently stripping the teat. Samples were refrigerated for 1–4 hours, then stored at -20°C until analysis. The time of milk collection (morning, 0600–0900 hours; evening, 1600–2300 hours) was recorded and young were returned to the pouch. We collected 103 milk samples from 91 females. Of these, 12 and 11 were newly marked in 2014 and 2015, respectively, with colored Allflex ear tags and a flexible collar (Gélin *et al.*, 2013).

Milk analysis

Milk samples were assayed in triplicate for total protein (Thermo Scientific-Pierce™ BCA Protein Assay Kit #23227) and lipid (Atwood & Hartmann, 1992) concentrations. The partial energy (carbohydrate content not included) calculated per unit volume was derived from lipid and protein concentrations, assuming energy values of $24.6\text{ kJ}\cdot\text{g}^{-1}$ for protein and $38.1\text{ kJ}\cdot\text{g}^{-1}$ for lipid (Ofstedal, 1984). At this stage of marsupial lactation, carbohydrates decrease and remain at low concentrations for the remainder of lactation (Tyndale-Biscoe & Janssens, 1988), therefore should not greatly influence milk energy. Milk yield was not quantified, and time since last suckling could not be measured because nursing occurred inside the pouch.

Explanatory variables

Environmental, maternal and offspring characteristics could affect milk composition. The maternal characteristics we considered were leg length and relative body condition, measured as the ratio of observed body mass to mass predicted by an OLS regression of log female mass on log leg length (Le Cren, 1951). Age was unknown for 73% of mothers and hence was not included in analyses. The offspring characteristics we considered were sex and age. Age was the mean of estimates from sex-specific growth models for head, foot and leg lengths (Poole *et al.* 1982) and was included in analyses to account for increases in both lipid and total protein concentrations in milk (Green *et al.*, 1980, 1983). Environmental variables included year of sampling, sampling date and birthdate of young, the two latter characteristics representing seasonal variables. Birthdate was expressed as Julian day, with 1 August as day 1, because the peak of births occurs during spring and early summer (60% were born in November-January, 9% were

born in May-October). Year of sampling indirectly considered forage production, that was not seasonally predictable. Forage production was measured in January, April, July and October by harvesting palatable vegetation from 50 systematically-distributed, 0.79-m² circular wire-mesh exclusion cages. We calculated forage production between estimated birthdate and milk sampling date for each mother-young pair assuming constant daily vegetation growth between measures.

We examined how milk composition explained offspring survival using age of juveniles in days at their last sighting. We searched for marked young on most days from late July to early December, and again in early March. Maximum longevity was 550 days (18 months), the minimal age of weaning (Poole, 1975).

Model development and analysis

We first evaluated the sources of variation in milk composition by considering multiple hypotheses. We tested sex-biased allocation by including interactions between maternal characteristics and offspring sex. We considered an alternative hypothesis, environmental modulation of milk provisioning at annual and seasonal levels, by including year and offspring birthdate in analyses, and interactions between year and both maternal condition and offspring sex were tested. Preliminary analyses indicated a curvilinear relationship between milk macronutrients and age of young, therefore age was included as a quadratic term. A total of 19 hypotheses were modelled using linear mixed regressions (Table 4; Appendix S1), and three response variables describing milk composition, total protein and lipid concentrations, and the lipid-to-protein ratio (LPR), were separately investigated. High LPR indicates a greater proportion of lipids than proteins, and low LPR indicates greater protein content regardless of milk solids concentrations. Using LPR accounts for the confounding effect of milk volume diluting macronutrients (Hinde, 2009).

We formulated 11 hypotheses to explain the effect of milk composition on offspring longevity (Table 4; Appendix S1), including offspring sex and interactions between sex and macronutrients. Age of young at capture was included in models because it was correlated with longevity ($r_p=0.20$, $P<0.001$). Maternal condition was included because it

is linked to juvenile survival in agile wallabies (*M. agilis*) (Bolton, Newsome & Merchant, 1982). We also considered birthdate and an interaction between birthdate and year.

To establish the relative strength of evidence for each candidate model, we performed model selection using Akaike's Information Criterion corrected for sample size (AIC_c) (Burnham & Anderson, 2002) for the four candidate sets. Models were ordered according to Δi values relative to the model with the lowest AIC_c . Akaike weights (w_i), indicating the probability that a model was the best in the candidate set (Burnham & Anderson, 2002), were computed for each model. A model had significant support if $w_i \geq 0.05$. We then performed multimodel inference to estimate averaged effect sizes; parameter estimates for variables that best described variance in milk composition and offspring longevity had 95% confidence intervals that did not overlap zero (Burnham, Anderson & Huyvaert, 2011).

All models included maternal identity as a random variable to account for the use of non-averaged triplicates. We used the *lmer* function in the *lme4* package (Bates *et al.*, 2014) with R version 3.2.2 (R Core Team 2015) to fit models to the data, and the *aictab* function in the *AICcmodavg* package (Burnham *et al.*, 2011) to obtain AIC_c values, perform model averaging and estimate model parameters.

RESULTS

Proximate milk composition

We analyzed 103 samples, 78 from 2014 and 25 from 2015 (Table 1). Among females marked in previous years, 49% (70 of 144) carried pouch young in 2014 and 11% in 2015 (15 of 131) ($\chi^2=44.48$, $df=1$, $P<0.0001$). Young in 2014 were estimated to be seven days older at capture, on average, than in 2015 (range: 196–284 days; Table 2). The sampling period was 37 days longer in 2014 (31 July to 5 December) than in 2015 (29 July to 29 October). Estimated birthdates spanned five months in 2014 (23 November 2013 to 29 April 2014) and four months in 2015 (5 December 2014 to 29 March 2015).

Forage production was 48% lower in 2015 than in 2014 (Table 2), and maternal condition (Table 2) and partial energy (Fig. 1) were also higher in 2014 than in 2015. Maternal

body condition increased in spring (Table 3). Further examinations of relationships between explanatory variables (Tables 2,3) provided additional evidence of the lack of bias in sampling. Average age of young, sampling date and birthdate were not different according to offspring sex (Table 2). Time of day did not affect protein concentrations (Welch two-sample t-test: $t = -0.42$, $df = 261.02$, $P = 0.68$), but lipid concentrations were higher in the morning than in the evening (Welch two-sample t-test: $t = 2.84$, $df = 207.46$, $P = 0.005$). On average, evening samples were collected later in the season (Welch two-sample t-test: $t = -2.26$, $df = 258.58$, $P = 0.02$).

Determinants of milk composition

Lipid and protein

The concentrations of protein and lipids varied substantially (24.04 – $151.50 \text{ mg}\cdot\text{mL}^{-1}$ and 46.76 – $316.10 \text{ mg}\cdot\text{mL}^{-1}$, respectively), with significant year effects (Table 2). Detailed distributions, model selection and multimodel inference summaries for these macronutrients are provided in Appendices S2&S5 in the Supporting Information.

Lipid:protein ratio

Because sampling date and birthdate of young were highly correlated (Table 3), models fitted with each variable yielded similar results. We report models using birthdate because they had greater strength of evidence than those including sampling date (Likelihood-ratio test: $X^2 = 7.11$, $df = 1$, $P < 0.001$). There was support ($w_i \geq 0.05$) for three models explaining LPR (Table 5). Model averaging (Table 7) suggested that LPR was best explained by negative interactions between maternal condition and both year and offspring sex (Fig. 2a, c), and by negative effects of birthdate (Fig. 2b) and maternal size (Fig. 2d). Detailed multimodel inference summaries are provided in Appendix S3.

Juvenile survival

Longevity estimates were accurate and precise for young that died before mid-December because we located every marked individual almost daily. We excluded 12 late-born pouch young from the analyses to avoid underestimating longevity because they were last seen alive when we left the study area. Most young disappeared before permanent pouch

emergence and only 11 (10%) survived to 550 days (2014: six females, two males; 2015: two females, one male). Mean longevity was 321 ± 97 (range: 201–550) days. Two models had significant support (i.e. $w_i \geq 0.05$; Table 6). Model-averaged estimates (Table 7) confirmed that offspring longevity increased with maternal condition and lipid concentration (Fig. 3a), but increased with protein concentration only for sons (Fig. 3b). Finally, longevity decreased with birthdate and was lower in 2015 than in 2014 (Table 7). Detailed multimodel inference results are presented in Appendix 4.

DISCUSSION

Forage productivity and milk composition of eastern grey kangaroos differed markedly between years. The substantial annual and seasonal differences in forage production during lactation in the two years of the study were associated with differences in maternal condition and milk composition, that we attribute to much lower forage production in 2015. Milk composition was further explained by offspring sex through an interaction with maternal condition, as mothers in better condition allocated more protein to sons than to daughters. Interestingly, larger mothers in our study area tend to have more sons (Le Gall-Payne, Coulson & Festa-Bianchet, 2015), and during our study they produced milk with more protein than smaller mothers. Moreover, milk composition had a direct and sex-dependent effect on offspring survival.

Forage production during lactation in 2015 was half of that in 2014, and mothers in 2015 were in poorer condition and produced lower-energy milk than mothers in 2014. Similarly, food-restricted red deer hinds (*Cervus elaphus*) produced less milk, containing less lipids, protein and glucose, than control hinds (Landete-Castillejos *et al.*, 2003), and female tammar wallabies increased food intake to avoid metabolizing fat reserves as lactation progressed (Cork, 1991). The concomitant differences in milk composition and forage production suggests that eastern grey kangaroos relied at least partly on current nutrition to meet lactational requirements. Lactating kangaroos increase their bite rate and midday foraging time compared to non-lactating females (Gélin *et al.*, 2013). Increasing foraging effort may not, however, compensate for the costs of lactation when forage is scarce (Parker *et al.*, 2009). Low forage production likely lowered protein intake, but

256 females that maintained relatively high condition in the poor year (2015) produced high-
257 protein milk, suggesting that they were able to buffer against forage scarcity.

258 Milk composition was also affected by birthdate. At mid lactation, early-born kangaroos
259 received more milk lipids than proteins, independently of maternal condition. For
260 temperate herbivores, dietary protein might not be enough for milk production before the
261 onset of spring vegetation growth (Taillon, Barboza & Côté, 2013). Also, mothers of
262 early-born pouch young face both low food abundance and high thermoregulatory costs
263 because of cooler temperatures in late winter and there might be higher costs to deplete
264 protein stores than fat stores (Barboza & Parker, 2008). Our results provide a rare
265 example of how forage production, and potentially temperature, affect maternal condition
266 and milk composition in a wild terrestrial mammal. Eastern grey kangaroos exhibit a
267 conservative reproductive strategy and are partly income breeders (Gélin, Coulson &
268 Festa-Bianchet, 2016); lower milk partial energy in a low-forage year suggests that
269 females attempt to maintain condition by transferring environmental costs to their young.
270 Females, however, acted as capital breeders early in the season by using body stores to
271 meet the high lactation costs, suggesting a resource-modulated lactation strategy.

272 Kangaroos are polygynous (Fisher *et al.*, 2002), and successful sons could provide higher
273 fitness returns compared to daughters (Trivers & Willard, 1973). Since milk protein is of
274 paramount importance to pre-weaning growth (Landete-Castillejos *et al.*, 2001; Parker *et al.*, 2009), greater protein transfer by mothers could improve sons' reproductive potential
275 by increasing their adult body size (Miller *et al.*, 2010). We found that mothers of sons
276 produced milk with higher proportions of protein than mothers of daughters (Robert &
277 Braun, 2012), but this effect was moderated by maternal body condition. Mothers in
278 better relative condition favoured protein allocation to sons while mothers in poorer
279 condition favoured protein allocation to daughters, revealing a condition-specific and
280 sex-biased care strategy (Trivers & Willard, 1973). This result supports previous findings
281 that heavier macropod mothers tend to have more sons (Sunnucks & Taylor, 1997; Le
282 Gall-Payne *et al.*, 2015), probably because they can provide high-quality care (Robert,
283 Schwanz & Mills, 2010). In marsupials, milk composition is autonomously controlled by
284 the mammary gland and is unaffected by suckling patterns (Trott *et al.*, 2003). It is
285

therefore unlikely that higher protein allowance for sons was a non-adaptive maternal response to higher suckling and nutrient demand (Landete-Castillejos *et al.*, 2005), because despite different compositions, partial energy density per unit volume of milk was similar for sons and daughters. We do not know if females produced different amounts of milk. Data on milk volume would allow a better understanding of sex-biased care, with potential sex differences in production and macronutrient dilution (Nommsen *et al.*, 1991; Hinde, 2009; Hinde *et al.*, 2014).

Surprisingly, structural size affected milk composition independently of maternal condition. Larger females produced high-protein milk compared to smaller females. Kangaroos have indeterminate skeletal growth, so larger individuals tend to be older (Jarman, 1989). Age could therefore partly explain the effect of size, as experienced females may select better food patches. Alternatively, larger females may be able to allocate more resources to lactation (Pomeroy, Fedak & Anderson, 1999) because smaller females face a steeper trade-off between allocating protein stores to structural growth or to milk production (Stearns, 1989).

In 2015, when forage production and maternal condition were poor, offspring survival and population reproductive rate were lower compared to 2014. Model selection revealed that offspring longevity was affected by maternal condition, birthdate and milk composition. Higher lipid concentrations increased longevity for both sexes, but low protein concentrations had a greater negative effect on the longevity of sons. Higher milk protein during mid lactation was also associated with higher offspring mass at 14 months in captive tammar wallabies (Robert & Braun, 2012). In kangaroos, sons require more protein than daughters, as they grow faster during the pouch stage (Poole *et al.*, 1982), and selection favors rapid structural growth in male juveniles of sexually-dimorphic species (Arnould *et al.*, 2001). Marsupial milk composition follows a temporal protein delivery sequence, partly acting as signal for tissue development (Sharp *et al.*, 2016). Apart from retarding structural growth, deficiency in specific compounds could also cause abnormal development (Sharp *et al.*, 2016) and weaken resistance to pathogens (Daly *et al.*, 2007; Joss *et al.*, 2009), jeopardizing survival. Our results are conservative, because it is possible that we overestimated the age of high-provisioned young. Our

analyses could not compare milk attributes to early survival, which was likely affected by poor environmental conditions.

We found that early-born young survived longer. A similar result in mountain goats (*Oreamnos americanus*) was attributed to early-born kids having a longer period of access to high forage quality in a temperate environment (Côté & Festa-Bianchet, 2001). We remain cautious about the negative effect of birthdate on longevity, as seasonal conditions likely fluctuate between years in our study area.

In conclusion, we found strong support for sex-biased milk macronutrient allocation in wild kangaroos. Asynchronous reproduction revealed effects of environmental conditions on a direct component of maternal care. Our study is a rare example of concomitant deterioration in milk quality and forage production in a large wild terrestrial mammal.

ACKNOWLEDGEMENTS

We thank Julie Sharp who provided space and equipment at Deakin University for analyses, and Vengaimu Modepalli and the other graduate students who helped. We acknowledge W. J. King, and two anonymous reviewers for helpful comments on earlier drafts. This study was funded by the National Geographic Society, the Natural Sciences and Engineering Research Council of Canada (NSERC), the Fonds de Recherche Québécois en Nature et Technologies (FRQNT) and the Ministère de l'Éducation, du Loisir et du Sport of Québec (MELS). Handling of animals was authorized by the University of Melbourne Animal Ethics Committee (1312902) and by the Université de Sherbrooke Animal Care Committee (MFB2012-02), affiliated with the Canadian Council on Animal Care.

REFERENCES

- Arnould, J.P.Y., Boyd, I.L., Rawlins, D.R. & Hindell, M.A. (2001). Variation in maternal provisioning by lactating Antarctic fur seals (*Arctocephalus gazella*): Response to experimental manipulation in pup demand. *Behav. Ecol. Sociobiol.* **50**, 461–466.
- Atwood, C.S. & Hartmann, P.E. (1992). Collection of fore and hind milk from the sow and the changes in milk composition during suckling. *J. Dairy Res.* **59**, 287–298.

- Barboza, P.S. & Parker, K.L. (2008). Allocating protein to reproduction in arctic reindeer and caribou. *Physiol. Biochem. Zool.* **81**, 835–855.
- Bates, D., Mächler, M., Bolker, B.M. & Walker, S.C. (2014). Fitting linear mixed-effects models using lme4. *arXiv:1406.5823v1[stat.CO]*.
- Bolton, B.L., Newsome, A.E. & Merchant, J.C. (1982). Reproduction in the agile wallaby *Macropus agilis* (Gould) in the tropical lowlands of the Northern Territory: opportunism in a seasonal environment. *Aust. J. Ecol.* **7**, 261–277.
- Bovolenta, S., Corazzin, M., Messina, M., Focardi, S. & Piasentier, E. (2013). Supplementary feeding of farmed fallow deer : effect on milk composition and fawn performance. *Ital. J. Anim. Sci.* **12**, e98.
- Bowen, W.D., Ellis, S.L., Iverson, S.J. & Boness, D.J. (2001). Maternal effects on offspring growth rate and weaning mass in harbour seals. *Can. J. Zool.* **79**, 1088–1101.
- Burnham, K.P. & Anderson, D.R. (2002). *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*. 2nd edn. New York: Springer.
- Burnham, K.P., Anderson, D.R. & Huyvaert, K.P. (2011). AIC model selection and multimodel inference in behavioral ecology : some background, observations, and comparisons. *Behav. Ecol. Sociobiol.* **65**, 23–35.
- Cork, S.J. (1991). Meeting the energy requirements for lactation in a macropodid marsupial : current nutrition versus stored body reserves. *J. Zool.* **225**, 567–576.
- Côté, S.D. & Festa-Bianchet, M. (2001). Birthdate, mass and survival in mountain goat kids: effects of maternal characteristics and forage quality. *Oecologia* **127**, 230–8.
- Le Cren, E.D. (1951). The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *J. Anim. Ecol.* **20**, 201–219.
- Crocker, D.E., Williams, J.D., Costa, D.P. & Boeuf, B.J. Le. (2001). Maternal traits and reproductive effort in northern elephant seals. *Ecology* **82**, 3541–3555.
- Daly, K.A., Digby, M., Lefe, C., Thomson, P., Nicholas, K. & Williamson, P. (2007).

- Analysis of the expression of immunoglobulins throughout lactation suggests two periods of immune transfer in the tammar wallaby (*Macropus eugenii*). *Vet. Immunol. Immunopathol.* **120**, 187–200.
- Davis, N.E., Coulson, G. & Forsyth, D.M. (2008). Diets of native and introduced mammalian herbivores in shrub-encroached grassy woodland, south-eastern Australia. *Wildl. Res.* **35**, 684–694.
- Davis, N.E., Forsyth, D.M. & Coulson, G. (2010). Facilitative interactions between an exotic mammal and native and exotic plants: hog deer (*Axis porcinus*) as seed dispersers in south-eastern Australia. *Biol. Invasions* **12**, 1079–1092.
- Fisher, D.O., Blomberg, S.P. & Owens, I.P.F. (2002). Convergent maternal strategies and ungulates and macropods. *Evolution.* **56**, 167–176.
- Fujita, M., Roth, E., Lo, Y., Hurst, C., Vollner, J. & Kendell, A. (2012). In poor families, mothers' milk is richer for daughters than sons : a test of Trivers–Willard hypothesis in agropastoral settlements in northern Kenya. *Am. J. Phys. Anthropol.* **59**, 52–59.
- Le Gall-Payne, C., Coulson, G. & Festa-Bianchet, M. (2015). Supersize me: heavy eastern grey kangaroo mothers have more sons. *Behav. Ecol. Sociobiol.* **69**, 795–804.
- Gélin, U., Coulson, G. & Festa-Bianchet, M. (2016). Heterogeneity in reproductive success explained by individual differences in bite rate and mass change. *Behav. Ecol.* **27**, 777–783.
- Gélin, U., Wilson, M.E., Coulson, G.M. & Festa-Bianchet, M. (2013). Offspring sex, current and previous reproduction affect feeding behaviour in wild eastern grey kangaroos. *Anim. Behav.* **86**, 885–891.
- Georges, J.-Y. & Guinet, C. (2000). Maternal care in the subantarctic fur seals on Amsterdam Island. *Ecology* **81**, 295–308.
- Green, B., Griffiths, M. & Leckie, R.M.C. (1983). Qualitative and quantitative changes in milk fat during lactation in the tammar wallaby (*Macropus eugenii*). *Aust. J. Biol. Sci.* **36**, 455–461.

- Green, B., Newgrain, K. & Merchant, J. (1980). Changes in milk composition during lactation in the tammar wallaby (*Macropus eugenii*). *Aust. J. Biol. Sci.* **33**, 35–42.
- Hinde, K. (2009). Richer milk for sons but more milk for daughters : sex-biased investment during lactation varies with maternal life history in rhesus macaques. *Am. J. Hum. Biol.* **519**, 512–519.
- Hinde, K., Carpenter, A.J., Clay, J.S. & Bradford, B.J. (2014). Holsteins favor heifers, not bulls: biased milk production programmed during pregnancy as a function of fetal sex. *PLoS One* **9**, e86169.
- Iverson, S.J., Bowen, W.D., Boness, D.J. & Oftedal, O.T. (1993). The effect of maternal size and milk energy output on pup growth in grey seals (*Halichoerus grypus*). *Physiol. Zool.* **66**, 61–88.
- Jarman, P. (1989). Sexual dimorphism in Macropodidae. In *Kangaroos, wallabies and rat-kangaroos*: 433–447. Griggs, G.C., Jarman, P. & Hume, I.D. (Eds). . Chipping Norton, New South Wales: Surrey Beattie & Sons.
- Joss, J.L., Molloy, M.P., Hinds, L. & Deane, E. (2009). A longitudinal study of the protein components of marsupial milk from birth to weaning in the tammar wallaby (*Macropus eugenii*). *Dev. Comp. Immunol.* **33**, 152–161.
- King, W.J., Wilson, M.E., Allen, T., Festa-Bianchet, M. & Coulson, G. (2011). A capture technique for free-ranging eastern grey kangaroos (*Macropus giganteus*) habituated to humans. *Aust. Mammal.* **33**, 47–51.
- Landete-Castillejos, T., Garcia, A. & Gallego, L. (2001). Calf growth in captive Iberian red deer (*Cervus elaphus hispanicus*): effects of birth date and hind milk production and composition. *J. Anim. Sci.* **79**, 1085–1092.
- Landete-Castillejos, T., García, A., Gómez, J.Á. & Gallego, L. (2003). Lactation under food constraints in Iberian red deer *Cervus elaphus hispanicus*. *Wildlife Biol.* **9**, 131–139.
- Landete-Castillejos, T., García, A., Lopez-Serrano, F.R. & Gallego, L. (2005). Maternal quality and differences in milk production and composition for male and female

- Iberian red deer calves (*Cervus elaphus hispanicus*). *Behav. Ecol. Sociobiol.* **57**, 267–274.
- Maestripieri, D. & Mateo, J.M. (Eds.). (2009). *Maternal effects in mammals*. Chicago: University of Chicago Press.
- Mellish, J.A., Iverson, S.J. & Bowen, W.D. (1999). Variation in milk production and lactation performance in grey seals and consequences for pup growth and weaning characteristics. *Physiol. Biochem. Zool.* **72**, 677–690.
- Miller, E.J., Eldridge, M.D.B., Cooper, D.W. & Herbert, C.A. (2010). Dominance, body size and internal relatedness influence male reproductive success in eastern grey kangaroos (*Macropus giganteus*). *Reprod. Fertil. Dev.* **22**, 539–49.
- Nommsen, L.A., Lovelady, C.A., Heining, M.J., Lönnerdal, B. & Dewey, K.G. (1991). Determinants of energy, protein, lipid, and lactose concentrations in human milk during the first 12 mo of lactation : the DARLING study. *Am. J. Clin. Nutr.* **53**, 457–465.
- Oftedal, O.T. (1984). Milk composition, milk yield and energy output at peak lactation: a comparative review. In *Physiological strategies in lactation: the proceedings of a symposium held at the Zoological Society of London on 11 and 12 November 1982*. Peaker, M., Vernon, R. & Knight, C. (Eds). London: Academic Press.
- Oftedal, O.T. (1985). Pregnancy and lactation. In *Bioenergetics of wild herbivores*: 215–238. White, R.J. & Hudson, R.G. (Eds). Boca Raton: CRC Press.
- Parker, K.L., Barboza, P.S. & Gillingham, M.P. (2009). Nutrition integrates environmental responses. *Funct. Ecol.* **23**, 57–69.
- Pomeroy, P.P., Fedak, M.A. & Anderson, S. (1999). Consequences of maternal size for reproductive expenditure and pupping success of grey seals at North Rona, Scotland. *J. Anim. Ecol.* **68**, 235–253.
- Poole, W.E. (1975). Reproduction in the two species of grey kangaroos , *Macropus giganteus* Shaw and *M . fuliginosus* (Desmarest) II. Gestation , parturition and pouch life. *Aust. J. Ecol.* **23**, 333–353.

- Poole, W.E., Carpenter, S.M. & Wood, J.T. (1982). Growth of grey kangaroos and the reliability of age determination from body measurements I. The eastern grey kangaroo, *Macropus giganteus*. *Aust. Wildl. Res.* **9**, 9–20.
- Powe, C.E., Knott, C.D. & Conklin-Brittain, N. (2010). Infant sex predicts breast milk energy content. *Am. J. Hum. Biol.* **22**, 50–54.
- Rioux-Paquette, E., Garant, D., Martin, A.M., Coulson, G. & Festa-Bianchet, M. (2015). Paternity in eastern grey kangaroos: moderate skew despite strong sexual dimorphism. *Behav. Ecol.* **26**, 1147–1155.
- Robert, K.A. & Braun, S. (2012). Milk composition during lactation suggests a mechanism for male biased allocation of maternal resources in the tammar wallaby (*Macropus eugenii*). *PLoS One* **7**, e51099.
- Robert, K.A., Schwanz, L.E. & Mills, H.R. (2010). Offspring sex varies with maternal investment ability: empirical demonstration based on cross-fostering. *Biol. Lett.* **6**, 242–245.
- Roberts, S.B., Cole, T.J. & Coward, W.A. (1985). Lactational performance in relation to energy intake in the baboon. *Am. J. Anim. Nutr.* **41**, 1270–1276.
- Sharp, J.A., Wanyonyi, S., Modepalli, V., Watt, A., Kuruppath, S., Hinds, L.A., Kumar, A., Abud, H.E., Lefevre, C. & Nicholas, K.R. (2016). The tammar wallaby: A marsupial model to examine the timed delivery and role of bioactives in milk. *Gen. Comp. Endocrinol.* <http://dx.doi.org/10.1016/j.ygcen.2016.08.007>.
- Skibieli, A.L. & Hood, W.R. (2015). Milk matters : offspring survival in Columbian ground squirrels is affected by nutrient composition of mother's milk. *Front. Ecol. Evol.* **3**, 111.
- Stearns, S.C. (1989). Trade-offs in life-history evolution. *Funct. Ecol.* **3**, 259–268.
- Sunnucks, P. & Taylor, A.C. (1997). Sex of pouch young related to maternal weight in *Macropus eugenii* and *M. parma* (Marsupialia: Macropodidae). *Aust. J. Zool.* **45**, 573–578.

- Taillon, J., Barboza, P.S. & Côté, S.D. (2013). Nitrogen allocation to offspring and milk production in a capital breeder. *Ecology* **94**, 1815–1827.
- Tardif, S.D., Power, M., Oftedal, O.T., Power, R.A. & Donna, G. (2001). Lactation, maternal behavior and infant growth in common marmoset monkeys (*Callithrix jacchus*); effects of maternal size and litter size. *Behav. Ecol. Sociobiol.* **51**, 17–25.
- Trivers, R.L. & Willard, D.E. (1973). Natural selection of parental ability to vary the sex ratio of offspring. *Science*. **179**, 90–92.
- Trott, J.F., Simpson, K.J., Moyle, R.L.C., Hearn, C.M., Shaw, G., Nicholas, K.R. & Renfree, M.B. (2003). Maternal regulation of milk composition, milk production, and pouch young development during lactation in the tammar wallaby (*Macropus eugenii*). *Biol. Reprod.* **68**, 929–936.
- Tyndale-Biscoe, C.H. & Janssens, P.A. (Eds.). (1988). *The developing marsupial: models for biomedical research*. Berlin, Heidelberg: Springer Verlag.
- TABLE CAPTIONS
- Table 1. Number of milk samples assayed during a two-year study of eastern grey kangaroos at Wilsons Promontory National Park, Australia.
- Table 2. Mean $\pm$ SD values and two-way analysis of variance by year of sampling and sex of offspring for variables recorded for each milk sample collected at Wilsons Promontory National Park, Australia, in 2014 and 2015.
- Table 3. Pearson's product-moment correlation coefficients between maternal, pouch young and environmental variables characterizing eastern grey kangaroo milk samples (n=103) collected at Wilsons Promontory National Park, Australia, in 2014 and 2015.
- Table 4. Parameters used in linear mixed models fitted to assess determinants of milk composition (m) and effects of milk composition on offspring longevity (Y) in Wilsons Promontory National Park, Australia, in 2014 and 2015.
- Table 5. Model selection summary for the effects of maternal characteristics, age and sex of pouch young, birthdate and year on lipid-to-protein ratio in eastern grey kangaroo

510 milk. LL, log-likelihood; K, number of estimated parameters; AIC_c = Akaike's
511 Information Criterion corrected for small sample size; Δi , difference between model
512 AIC_c and the lowest AIC_c in the model set; w_i , Akaike model weight. Models are ranked
513 in descending order of support, and models with significant support ($w_i \geq 0.05$) are in
514 bold.

515 Table 6. Model selection summary for the effects of milk composition, sex of pouch
516 young, birthdate and year on eastern grey kangaroo survival to weaning. LL, log-
517 likelihood; K, number of estimated parameters; AIC_c = Akaike's Information Criterion
518 corrected for small sample size; Δi , difference between model AIC_c and the lowest AIC_c
519 in the model set; w_i , Akaike model weight. Models are ranked in descending order of
520 support, and models with significant support ($w_i \geq 0.05$) are in bold.

521 Table 7. Model-averaged parameter estimates explaining variation in lipid-to-protein
522 ratio in eastern grey kangaroo milk and offspring longevity, using standardized variables
523 (reduced and centered to the mean).

524

Table 1:

525

	Year		Total
	2014	2015	
Daughters	35	16	51
Sons	43	9	62
TOTAL	78	25	103

Table 2:

	Year					Offspring					Year × Offspring		
	Mean values					Mean values							
	2014	2015	F	P	df	Female	Male	F	P	df	F	P	df
Lipid concentration (mg·mL ⁻¹)	174 ± 56	119 ± 41	58.66	< 0.001	1	161 ± 68	160 ± 45	0.018	0.893	1	2.349	0.126	1
Protein concentration (mg·mL ⁻¹)	90.4 ± 22.4	48.9 ± 8.5	235	< 0.001	1	78.5 ± 26.4	82.4 ± 26.9	1.64	0.201	1	0.949	0.331	1
Date of sampling (1–29 July)	47 ± 35	41 ± 27	2.25	0.135	1	47 ± 35	44 ± 32	0.47	0.492	1	1.849	0.175	1
Pouch young birthdate (1–1 August)	175 ± 41	175 ± 33	0.005	0.944	1	175 ± 41	175 ± 37	0.00	0.988	1	0.880	0.349	1
Maternal relative condition	1.007 ± 0.018	0.998 ± 0.011	10.02	0.002	1	1.004 ± 0.017	1.007 ± 0.017	2.87	0.092	1	1.704	0.192	1
Maternal leg length (mm)	529 ± 14	525 ± 15	3.48	0.063	1	528 ± 15	528 ± 13	0.12	0.733	1	0.746	0.388	1
Age of pouch young at capture (days)	233 ± 19	226 ± 16	7.55	0.006	1	232 ± 21	230 ± 17	0.84	0.362	1	0.327	0.568	1
Total forage production during lactation (g·m ⁻²)	154 ± 15	80 ± 18	1191	< 0.001	1	132 ± 40	142 ± 30	6	< 0.001	1	1	0.229	1

Table 3:

Pearson's product-moment correlations (r_p)										
	Age of pouch young	P	Sampling date	P	Birthdate	P	Maternal relative condition	P	Maternal leg length	P
Sampling date (1=29 July)	-0.06	0.306	-	-	-	-	-	-	-	-
Birthdate (1=1 August)	-0.53	< 0.001	0.87	< 0.001	-	-	-	-	-	-
Maternal relative condition	0.07	0.209	0.28	< 0.001	0.19	< 0.001	-	-	-	-
Maternal leg length (mm)	0.16	0.006	-0.17	0.004	-0.22	< 0.001	-0.13	0.027	-	-
Total forage production during lactation ($\text{g}\cdot\text{m}^{-2}$)	0.48	< 0.001	0.15	0.010	-0.11	0.059	0.19	< 0.001	0.12	0.033

Model	Description
Milk composition	
$m \{.\}$	Milk composition varies randomly (i.e. null model)
$m \{a + a\}$	Milk composition varies curvilinearly with age of young (a)
$m \{bd\}$	Milk composition varies linearly with birthdate of young (bd)
$m \{C\}$	Milk composition varies linearly with maternal relative condition (C)
$m \{L\}$	Milk composition varies linearly with maternal leg length (L)
$m \{S\}$	Milk composition varies with offspring sex (S)
$m \{yr\}$	Milk composition varies with years of sampling (yr)
$m \{all\}$	Milk composition varies additively with all predictor variables
$m \{S \times C\}$	Milk composition varies with an interaction between offspring sex and maternal relative condition
$m \{S \times L\}$	Milk composition varies with an interaction between offspring sex and maternal leg length
$m \{yr + bd\}$	Milk composition varies additively with year of sampling and birthdate of young
$m \{all + S \times C\}$	Milk composition varies additively with all predictors variables and an interaction between offspring sex and maternal relative condition
$m \{all + S \times C + S \times yr\}$	Milk composition varies additively with all predictors variables and an interaction between offspring sex and both maternal relative condition and year of sampling
$m \{all + S \times L\}$	Milk composition varies additively with all predictor variables and an interaction between offspring sex and maternal size
$m \{all + S \times L + S \times yr\}$	Milk composition varies additively with all predictor variables and an interaction between offspring sex and both maternal size and year of sampling
$m \{all + yr \times C + S \times yr\}$	Milk composition varies additively with all predictor variables and an interaction between year of sampling and both maternal relative condition and offspring sex
$m \{all + S \times C + C \times yr\}$	Milk composition varies additively with all predictor variables and an interaction between maternal relative condition and both year and offspring sex
$m \{all + S \times C + C \times yr + S \times yr\}$	Milk composition varies additively with all predictor variables and two-way interactions between offspring sex, year and maternal relative condition
$m \{all + S \times C \times yr\}$	Milk composition varies additively with all predictor variables and a three-way interaction between offspring sex, year and maternal relative condition
Offspring survival	
$Y \{.\}$	Offspring longevity varies randomly (i.e. null model)

$Y\{a\}$	Offspring longevity varies linearly with age of young (a)
$Y\{lip\}$	Offspring longevity varies linearly with milk lipid concentration (lip)
$Y\{prot\}$	Offspring longevity varies linearly with milk protein concentration ($prot$)
$Y\{C\}$	Offspring longevity varies linearly with maternal relative condition (C)
$Y\{S\}$	Offspring longevity varies with offspring sex (S)
$Y\{yr \times bd\}$	Offspring longevity varies with an interaction between year of sampling and birthdate of young
$Y\{all\}$	Offspring longevity varies additively with all predictor variables
$Y\{S \times lip\}$	Offspring longevity varies additively with all predictor variables and an interaction between offspring sex and milk lipid concentration
$Y\{S \times prot\}$	Offspring longevity varies additively with all predictor variables and an interaction between offspring sex and milk protein concentration
$Y\{S \times lip + S \times prot\}$	Offspring longevity varies additively with all predictor variables and an interaction between offspring sex and both milk lipid and protein concentrations

Table 4:

Table 5:

Model ID	LL	K	AIC _c	Δi	w_i
$m\{all + S \times C + C \times yr\}$	-251.08	12	527.24	0.00	0.52
$m\{all + S \times C + C \times yr + S \times yr\}$	-250.43	13	528.11	0.87	0.34
$m\{all + S \times C \times yr\}$	-250.17	14	529.79	2.55	0.15
$m\{all + S \times C + S \times yr\}$	-276.57	12	578.22	50.98	0.00
$m\{all + yr \times C + S \times yr\}$	-276.95	12	578.97	51.73	0.00
$m\{all + S \times C\}$	-281.18	11	585.27	58.03	0.00
$m\{all + S \times L + S \times yr\}$	-301.37	11	625.64	98.40	0.00
$m\{all + S \times L\}$	-303.96	11	630.82	103.58	0.00
$m\{all\}$	-310.36	10	641.47	114.24	0.00
$m\{yr + bd\}$	-320.27	7	654.93	127.69	0.00
$m\{S \times C\}$	-325.20	8	666.89	139.65	0.00
$m\{bd\}$	-339.28	4	686.69	159.45	0.00
$m\{S \times L\}$	-340.14	8	696.76	169.52	0.00
$m\{a + a^2\}$	-346.53	5	703.27	176.03	0.00
$m\{S\}$	-360.84	4	729.82	202.58	0.00
$m\{yr\}$	-362.49	4	733.11	205.88	0.00
$m\{L\}$	-362.97	4	734.07	206.83	0.00
$m\{.\}$	-364.33	3	734.73	207.50	0.00
$m\{C\}$	-364.31	4	736.76	209.53	0.00

550

551

Table 6:

Model ID	LL	K	AIC _c	Δi	w_i
$Y\{S \times lip + S \times prot\}$	-1221.91	12	2468.97	0.00	0.64
$Y\{S \times prot\}$	-1223.58	11	2470.14	1.17	0.36
$Y\{all\}$	-1231.52	10	2483.84	14.87	0.00
$Y\{S \times lip\}$	-1230.59	11	2484.14	15.17	0.00
$Y\{yr \times bd\}$	-1344.58	6	2701.47	232.50	0.00

$Y\{a\}$	-1383.68	4	2775.51	306.54	0.00
$Y\{prot\}$	-1386.23	4	2780.61	311.64	0.00
$Y\{C\}$	-1402.46	4	2813.07	344.09	0.00
$Y\{lip\}$	-1419.23	4	2846.60	377.63	0.00
$Y\{S\}$	-1429.38	4	2866.90	397.93	0.00
$Y\{.\}$	-1438.05	3	2882.18	413.20	0.00

Table 7:

Terms	Parameter value	95% confidence interval
Lipid:protein		
Age of young	-0.23	-0.34, -0.13
Year (reference=2014)	-0.06	-0.26, 0.13
Sex of offspring (reference=female)	0.20	-0.07, 0.46
Maternal relative condition	0.16	-0.03, 0.35
Maternal leg length	-0.31	-0.52, -0.10
Offspring birthdate	-0.38	-0.61, -0.16
Year×Sex	0.41	-0.24, 1.07
Maternal condition×Sex	-1.46	-1.78, -1.14
Maternal condition×Year	-1.23	-1.53, -0.92
Offspring longevity		
Age of young	-5.64	-13.19, 1.91
Year (reference=2014)	-15.38	-28.43, -2.32
Maternal relative condition	52.98	44.65, 61.32
Offspring birthdate	-23.15	-29.24, -17.06
Offspring sex	-35.51	-43.42, -27.59
Lipid concentration	18.99	12.13, 25.85
Protein concentration	45.37	30.72, 60.02
Lipid×Sex	15.66	-0.87, 32.20
Protein×Sex	36.97	19.95, 53.99

555

556 FIGURE CAPTIONS

557 Figure 1. Partial gross energy of eastern grey kangaroo milk by year and offspring sex.
558 Energy derived from lipids and proteins was significantly lower ($F= 8.93$, $df= 1$, $P <$
559 0.001) in 2015 than in 2014 but was independent of offspring sex ($F= 0.049$, $df= 1$, $P=$
560 0.83).

561 Figure 2. Determinants of milk lipid-to-protein ratio in eastern grey kangaroos. (a)
562 Model-averaged effect of scaled maternal relative condition moderated by year (a proxy
563 of forage production). 2014: closed circles, solid line; 2015: open triangles, dotted line.
564 (b) Model-averaged effect of scaled maternal relative condition as a function of offspring
565 sex. The relationships between scaled maternal relative condition x offspring sex for lipid
566 and protein concentrations mirror the effect shown here. Sons: open triangles, dotted line.
567 Daughters: closed circles, solid line. (c) Model-averaged effect of scaled offspring
568 birthdate. (d) Model-averaged effect of scaled maternal leg length (range: 488-562 mm).
569 Details on multimodel inference are provided in Appendix 3 in the Supporting
570 Information.

571 Figure 3. Model-averaged effects of (a) scaled lipid concentration and (b) scaled protein
572 concentration in milk of eastern grey kangaroos on offspring survival to weaning (i.e. 550
573 days). Sons: open triangles, dotted line. Daughters: closed circles, solid line. Details on
574 multimodel inference are provided in Appendix 4 in the Supporting Information.

575

576

Figure 1:

578

579

580

581

582

583

584

Figure 2:

586

587

588

589

590

591

592

Figure 3:

jzo_12453_f1.tiff

jzo_12453_f2.tiff

jzo_12453_f3.tiff

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Quesnel, L;MacKay, A;Forsyth, DM;Nicholas, KR;Festa-Bianchet, M

Title:

Size, season and offspring sex affect milk composition and juvenile survival in wild kangaroos

Date:

2017-08-01

Citation:

Quesnel, L., MacKay, A., Forsyth, D. M., Nicholas, K. R. & Festa-Bianchet, M. (2017). Size, season and offspring sex affect milk composition and juvenile survival in wild kangaroos. JOURNAL OF ZOOLOGY, 302 (4), pp.252-262. <https://doi.org/10.1111/jzo.12453>.

Persistent Link:

<http://hdl.handle.net/11343/292693>