Cervical Cancer: A Global Health Crisis

William Small, Jr., MD, FACRO, FACR, FASTRO^{*}, Monica Bacon, RN[†], Amishi Bajaj, BA^{*}, Linus T. Chuang, MD^{*}, Brandon J. Fisher, DO[‡], Matthew M. Harkenrider, MD^{*}, Anuja Jhingran, MD, FACR[¶], Henry C. Kitchener, MD FRCOG, FMedSci[¶], Linda R. Mileschkin, MD, FRACP[»], Akila N. Viswanathan, MD, MPH^{**}, David K. Gaffney, MD, PhD, FACR, FASTRO[§]

^{}Department of Radiation Oncology, Stritch School of Medicine Loyola University Chicago, IL*

[†]GCIG Operations Manager and Executive Director, Canada

^{}Department of Obstetrics, Gynecology, and Reproductive Science, Icahn School of Medicine at Mount Sinai, New York, NY*

[‡]Gamma West Cancer Services, Ogden, UT

[¶]Department of Radiation Oncology, The University of Texas M.D. Anderson Cancer Center, Houston, TX

[¶]Department of Obstetrics and Gynecology, University of Manchester, Manchester, UK

[»]Division of Hematology and Medical Oncology, Peter MacCallum Cancer Centre, Melbourne, Australia

^{**}Department of Radiation Oncology and Molecular Radiation Sciences, Johns Hopkins University School of Medicine, Baltimore, Maryland

[§]Department of Radiation Oncology, University of Utah, Huntsman Cancer Institute, Salt Lake City, UT

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version record](#). Please cite this article as [doi:10.1002/cncr.30667](https://doi.org/10.1002/cncr.30667).

Corresponding author: William Small, Jr., MD,

Professor and Chairman, Department of Radiation Oncology

Stritch School of Medicine Loyola University Chicago 2160 S 1st Ave

Maguire Center, Rm 2932, Maywood, IL 60153

Phone: 708-216-2559 Fax: 708-216-6076

E-mail: wmsmall@lumc.edu

Running title: Cervical Cancer: A Global Health Crisis

Total Number of Text Pages: 23

Total Number of Tables: 1

Total Number of Figures: 2

Funding Sources: None

Conflicts of Interest Notification: None

Authors Contributions: All authors were involved in the concept, writing and approval of the final manuscript.

Precis for use in Table Contents: Cervical cancer is a leading cause of death worldwide. In this paper, we discuss biology, prevention, and treatment of this disease, in addition to activist efforts aimed at improving treatment and access to care.

Abstract:

Cervical cancer is the fourth most common malignancy in women worldwide. Almost all cases of cervical cancer result from infection with human papilloma virus (HPV), and prevention of cervical cancer includes screening and vaccination. Primary treatment options for cervical cancer may include surgery or a concurrent chemoradiotherapy regimen consisting of cisplatin-based chemotherapy with external beam radiation therapy and brachytherapy.

Cervical cancer causes more than a quarter of a million deaths per year, as a result of grossly deficient treatments in many developing countries. This warrants a concerted global effort to counter shocking loss of life and suffering that largely goes unreported.

This article provides a review of cervical cancer: its biology, prevention, treatment, and discusses the global cervical cancer crisis and efforts to improve the prevention and treatment of cervical cancer in underdeveloped countries.

Keywords: cervical cancer, developing world, HPV, vaccination, GCIG, CCRN, chemotherapy, brachytherapy, activism

Introduction

Cervical cancer is one of the leading causes of cancer death among women [1]. Worldwide, cervical cancer is the fourth most frequently occurring malignancy in women and results in an estimated 530,000 new cases annually with 270,000 deaths. About 85% of the worldwide deaths from cervical cancer occur in underdeveloped or developing countries, and the death rate is 18 times higher in low and middle-income countries (LMIC) compared to wealthier countries. [2]. The highest incidence rates occur in Central and South America, the Caribbean, Sub-Saharan Africa, and Southern Asia [3]. In the United States in 2016, there were an estimated 12,990 cases and 4,120 deaths from cervical cancer [4], and the median age of diagnosis is 47 years.

Standard management of early stage (IA – IB1) cervical cancer is radical hysterectomy and lymph node dissection and/or radiation +/- chemotherapy. [5-7]. Standard management of locally advanced cervical cancer includes external beam radiotherapy with concurrent cisplatin-based chemotherapy with brachytherapy [8-16]. Brachytherapy is critical for curative intent treatment of cervical cancer and when replaced by external beam radiotherapy the results are clearly inferior[17-20]. With state-of-the-art staging and treatment, 3-year local control for early stage and advanced stage cervical cancer is 87-95% and 74-85%, respectively [15,16]. For all stages combined, 3-5-year survival from cervical cancer for many underdeveloped countries is <50% [21]. Death from cervical cancer often involves local progression resulting in significant suffering including ureteral obstruction, pain, and fistulas. The purpose of this article is to thoroughly review the biology, prevention strategies, treatment, and activism regarding cervical cancer with an emphasis on the global impact of these complex issues.

Biology of Cervical Cancer

The cervix is lined by stratified squamous epithelium that covers the exocervix and mucus-secreting columnar epithelium characteristic of the endocervical canal. The transition between these two populations of cells is called the squamocolumnar junction, and it is this area that is believed to be at greatest risk of viral neoplastic transformation. Tumors arising in the ectocervix are most commonly squamous cell carcinomas, which account for 75% of invasive cervical carcinoma. In contrast, tumors arising from the endocervix are more likely to be adenocarcinomas. Adenosquamous, small cell or neuroendocrine, serous papillary, and clear cell carcinomas of the cervix are less common histological subtypes.

The majority of cases of cervical cancer result from infection with HPV, with HPV DNA identified in approximately 95% of malignant cervical lesions [22]. The majority of HPV infections are transient and will be spontaneously cleared. However, in some cases, persistent infection will result in the development of the premalignant conditions of cervical intraepithelial neoplasia or adenocarcinoma in situ. Without treatment, the transition from dysplasia to invasive carcinoma may take years to decades to develop in most women. However, in around 10% of patients this transition can occur in under a year [23]. In addition, adenocarcinoma in situ appears to be more difficult to detect on Pap smear and this is thought to be one of the reasons for the increasing incidence of this subtype of cervical cancer [24].

Various factors have been suggested to increase the likelihood of the development of persistent infection and subsequent malignant transformation including cigarette

smoking, long-term oral contraceptive use, high parity, and co-infection with type 2 herpes simplex virus or human immunodeficiency virus (HIV). HPV serotypes 16 and 18 account for approximately 70% of cases, with the most common serotypes of HPV in women with cervical cancer in descending order of frequency being 16, 18, 45, 31, 33, 52, 58, and 35.

Perhaps due to the relative rarity of locally advanced or metastatic cervical cancer in the developed world, there have been only a few published reports of profiling of cervix tumors to look for actionable driver mutations. The most common finding has been of abnormalities in the PI3Kinase pathway, as reported by Wright et al. [25] who used the Oncomap platform to interrogate 80 cervical tumors for 1250 mutations in 139 genes. They identified PIK3CA mutations in 31% of cases, with shorter survival times seen in those with a mutation [25]. However, targeting this pathway therapeutically has proven difficult. They also identified KRAS mutations in 17.5% of the adenocarcinomas but none of the squamous cell carcinomas, suggesting that these tumor subtypes will need different types of targeted therapies. Ojesina et al. [26] have recently published the findings of deep sequencing of 115 cervix cancers to look for somatic mutations. They identified several novel somatic mutations in the squamous cell carcinomas profiled including E322K substitutions in the MAPK1 gene (8%); inactivating mutations in the HLA-B gene (9%); and mutations in EP300 (16%), FBXW7 (15%), TP53 (5%), and ERBB2 (6%). Somatic mutations in ELF3 (13%) and CBFB (8%) mutations were found in 24 adenocarcinomas [26].

Prevention of Cervical Cancer

Recognition that cervical neoplasia begins as intraepithelial change, which usually takes many years to develop into invasive disease, led to the use of cervical exfoliative cytology to detect cervical intraepithelial neoplasia that can be treated to prevent the development of cervical cancer. With the discovery that cervical cancer is caused by high risk HPV infection, the development of prophylactic vaccination in the 1990s there is now the means to achieve a more global approach to prevention through prophylactic vaccination. Vaccination can be viewed as primary prevention, with screening as secondary prevention.

The pivotal role of HPV in cervical carcinogenesis means that screening with HPV testing can achieve a more accurate risk-based approach. Randomized trials [27-30] have demonstrated that HPV testing is more sensitive than cytology, and for HPV negative women, screening intervals can safely be extended [31]. HPV testing lacks specificity, which means that cytology is required to triage women for referral to colposcopy. Based on limited data, triage of high-risk (hr) HPV-positive women using a combination of genotyping for HPV 16 and 18 and reflex cytology for women positive for the 12 other hrHPV genotypes appears to be a reasonable approach to managing hrHPV-positive women [32]. A challenge for primary HPV screening is the management of women with negative cytology, but various risk-based strategies are being developed based on HPV type and persistence. Screening programs around the world are in the process of switching from primary cytology aided by visual inspection with acetic acid to primary HPV testing.

Prophylactic Vaccination Against HPV

HPV infection of the cervix, thought to occur in most women at some time in their life, is most prevalent following the onset of sexual activity. In the majority of cases, the infection is cleared by the immune system. However, in a significant minority, infection is persistent and the viral genome becomes integrated into host DNA resulting in genomic dysregulation caused largely by the HPV oncogenes E6 and E7. The concept behind prophylactic vaccination is to achieve a high level of type-specific neutralizing antibodies directed against HPV capable of preventing cervical infection. The critical discovery that led to the vaccines we have today is that the major capsid protein of HPV, L1, could self-assemble into so-called virus like particles (VLPs) [33], which were shown to be highly immunogenic. Two vaccines, both based on VLPs made from HPV types 16 and 18, were produced with each using a different adjuvant. One was bivalent (types 16 and 18), and the other was quadrivalent to include the types responsible for genital warts (types 6 and 11). Both of these vaccines have been rigorously tested, initially in phase I and II trials and then in pivotal phase III trials [34,35]. These were performed on cohorts aged 15-26, and they demonstrated very high levels of type-specific antibody, which achieved very high efficacy (greater than 95%) in preventing HPV infection and similar efficacy in preventing type-specific CIN as well as vaginal and vulvar lesions. The data from these trials showed however that the vaccines were ineffective in females who already had an established HPV infection. Additionally, vaccination of boys prior to sexual activity at age 11-12 is recommended by the American Academy of Pediatrics to prevent HPV-induced cancers of the oropharynx, anus, and penis [36]. Vaccination of both sexes will have a large impact on herd immunity.

Most developed countries have introduced vaccination programs for pre-pubescent girls, and there has been early evidence of public health benefit with reduction in the incidence of high-risk infection, reduced incidence of cervical abnormalities, and even a reduction in genital warts in males who have not been vaccinated. This provides clear evidence of herd protection achieved by vaccination.

Recent Developments in Prophylactic Vaccination

The original vaccination regimens were based on three doses, given at time 0, 2, and 6 months. Recently, two doses have been shown to be as effective as three [37], provided the second dose is given 6-12 months after the initial dose. In the United Kingdom, for example, a two-dose regimen has now replaced the three dose regimen in the publicly funded schools-based program, which is achieving 85-90% coverage. Another development has been the production of a nonavalent vaccine that adds types 31, 33, 45, 52 and 58 to the 6, 11, 16 and 18 vaccine. This vaccine has been demonstrated in phase III trials to achieve similar efficacy against types 6, 11, 16 and 18 in addition to achieving high efficacy against the new types [38,39]. The nonavalent vaccine has been licensed in some countries, including the U.S. and the U.K. [40, 41], and may well replace the bi- and quadrivalent vaccines over the next few years [42, 43]. Prophylactic vaccination has the means to save hundreds of thousands of lives, but this will require the political will to ensure that vaccination is implemented in resource poor countries.

There is also increasing interest and research into the possibility of treating established cervix cancer using immunotherapy approaches. The 2 main oncogenes associated with HPV driven cancers, E6 and E7 are considered excellent targets for immunotherapy. Promising results have been seen with clinical trials involving

therapeutic HPV vaccines, adoptive T cell therapy and checkpoint inhibitors, with ongoing trials examining various combination immunotherapy approaches with standard treatments such as radiotherapy [44].

Gynecological Cancer Intergroup (GCIG) and the Cervix Cancer Research Network (CCRN)

The Gynecologic Cancer InterGroup (GCIG), formalized in 1997, aims to promote and facilitate high quality clinical trials in order to improve outcomes for women with gynecological cancer. Currently, there are 29 member groups including representation from North America, Europe, Asia and Australia. GCIG has several standing committees including Ovarian Cancer, Endometrial Cancer, Cervix Cancer, Translational Research, Harmonization (operations & statistics), Rare Tumors, Symptom Benefit, Phase II, and Membership.

The GCIG has also developed a Cervix Cancer Research Network (CCRN) whose aim is to promote high quality clinical research for cancer of the cervix in developing countries. The purpose of CCRN is to bring research in cervical cancer to the countries where the burden is the highest and there is a lack of GCIG cooperative groups (Figure 1). Interested sites complete a pre-qualifying set of Capability questions followed by a Radiologic/Physics check (questionnaire courtesy of IROC, Houston, TX). Site visits are then performed by a review team from GCIG assessing clinical activity, site resources, clinical trial operations, radiation therapy facilities/quality assurance and treatment record, and clinical trials management information. Participation in a beam measurement

program (TLD/OSLD) every 2 years is a requirement. Ongoing QA/QC is performed according to the lead group trial protocols. CCRN currently has 4 active cervical cancer trials described below.

The success of the GCIG relates to pooled intellectual resources and collaboration, rapid and large accrual, evidence-based medicine and application of the results, and opportunity for substantial translational research. GCIG is unique in the world of cancer research and been intensively productive in collaborative trials, intellectual exchanges and learning, brainstorming and consensus conferences.

Figure 1:

[FIGURE 1]

Modern Treatment of Cervical Cancer

The treatment of cervical cancer is dictated by FIGO stage: a clinical staging system [45]. For early cervical cancers, surgery is recommended. A cone biopsy is adequate treatment for stage IA1 whereas, for patients with IA1 with lymphovascular space invasion or IA2 cancer, a cone biopsy with negative margins and pelvic lymph node dissection is recommended. Fertility sparing surgery is an option for patients with early cervical cancers. For high risk IA1 thru IB1 a radical trachelectomy and pelvic lymph node dissection can be considered. An additional option for some would be pelvic radiotherapy and brachytherapy. There are ongoing trials evaluating reduced intensity surgery for patients with early lesions. The SHAPE trial is evaluating simple versus radical hysterectomy for patients with cervical tumors less than 2 cm in size. SHAPE is a CCRN trial which has immediate application to under-resourced countries. A randomized

trial of surgery versus radiotherapy for IB1-IIA cervix cancer showed no difference in survival [5]. Notably, patients in this trial did not receive chemotherapy, and 84% of patients in the surgical arm with tumors greater than 4 cm required postoperative radiotherapy. Morbidity was greater in patients that received both modalities; hence, current recommendations are to try to use a single modality.

Advanced imaging such as CT, MRI, and PET are not permissible in FIGO staging; however, imaging (if available) should be used to appropriately guide treatment. PET scans are helpful at delineating the extent of disease. MRI is superior at showing soft tissue resolution for the extent of cervical cancer within the pelvis. This can be critical for brachytherapy treatment planning or conformal radiotherapy techniques.

An NCI Alert in 1999 demonstrated the superiority of cisplatin-containing concurrent chemoradiotherapy for women with advanced cervical cancer. The hazard rate for reduction and death was approximately 0.52 [11]. Consequently, this was rapidly adopted worldwide, and weekly cisplatin became the worldwide standard [46]. The optimization of chemotherapy is unclear, and the CCRN has 3 trials testing the optimal combination of chemotherapy and radiotherapy [47]. Extended adjuvant chemotherapy in locally advanced disease is being tested currently in the OUTBACK trial. RTOG 0724 is also evaluating extended adjuvant chemotherapy for patients treated with a radical hysterectomy who have positive nodes or positive parametria, although this currently is not a CCRN trial. Dose-intense neoadjuvant chemotherapy is being tested in the phase III INTERLACE trial. Additionally, a phase II trial showed promising results of a higher dose of cisplatin given every three weeks [48]. This is now being compared to weekly

cisplatin in the TACO trial. The TACO trial has been the most successful CCRN trial with significant accrual from Vietnam and Thailand.

In patients with advanced disease receiving curative radiation, an important quality metric is to keep the total treatment course within 8 weeks. Prolonged treatment after 8 weeks in multiple studies have documented an approximate 1% loss in local control for every day beyond 8 weeks. Adherence to a few quality metrics such as receipt of concurrent chemoradiotherapy, brachytherapy and completion of treatment within 8 weeks will markedly improve survival worldwide.

Brachytherapy

Brachytherapy is an integral component of the treatment of advanced cervical cancer and is the standard of care in combination with external beam radiation therapy in all-national guidelines. [49-51] The advantage of brachytherapy comes from its dosimetric benefits including the ability to deliver a locally high and conformal dose to the site of disease with a rapid dose fall-off, sparing adjacent structures such as the bladder, rectum, sigmoid, and small bowel. [52] Brachytherapy remains unavailable in many countries. Even in countries where brachytherapy is easily accessible, utilization is declining [53-63]. An analysis of the Surveillance, Epidemiology, and End Results (SEER) database found a decline in the use of brachytherapy from 83% in 1988 to 58% in 2009 ($p < 0.001$), though brachytherapy was independently associated with better cause-specific survival (hazard ratio [HR] 0.64; 95% confidence interval [CI], 0.57-0.71) and overall survival (OS) (HR 0.66; 95% CI, 0.60-0.74) [52]. A similar study of the National Cancer Database (NCDB) found brachytherapy utilization decreased from 97%

in 2004 to 86% in 2011 [63,64]. In one of these studies, the impact of the use of brachytherapy was greater than that seen for the use of concurrent chemotherapy. [63] Brachytherapy is a complex procedure that necessitates significant resources and infrastructure that is particularly challenging in resource limited countries. Only 20 of the 52 African countries had brachytherapy in 2010 [52,64]. Of 12 centers in Latin America, three do not perform gynecological brachytherapy [52].

One high-dose rate brachytherapy machine can treat approximately 10-12 cases a day. In Ethiopia, a country of 94.1 million people with 60,000 new cancer cases per year, there is one after-loader for the whole country. In Thailand, in one hospital, 1,000 brachytherapy procedures are done in one year by one after-loader. In Honduras, where 1,000 new cases of cervical cancer are diagnosed annually, there is no brachytherapy in the entire country. Increasing the worldwide availability of brachytherapy should be a global health priority.

Treatment of Cervical Cancer in the Developing World

The majority of cervical cancer patients in the developing world present with advanced stage disease with limited access to adequate treatment. As a result, the mortality rates are high for women in the developing world. (Fig. 2) Because of the unpredictable availability of resources, the guidelines that are used to treat cervical cancer in high-income countries are not applicable to many of the developing countries.

Figure 2:

[FIGURE 2]

Figure 2. Estimated age-standardized rates (world) of deaths, females, cervical cancer, worldwide in 2012

Data source: GLOBOCAN 2012

Graph production: IARC (<http://gco.iarc.fr/today>)

World Health Organization

Two resource-stratified guidelines were recently published by National Comprehensive Cancer Network (NCCN) and American Society of Clinical Oncology (ASCO). [45] The NCCN Guidelines provided evidence-based recommendations by the representatives from NCCN Member Institutions. ASCO established a process including mixed methods of guideline development, adaptation of the clinical practice guidelines of other organizations, and formal consensus by the international expert panels. Recommendations were made on management of cervical cancer based on four different resource stratifications (Table 1). Included in Table 1 is the four resource settings that included a basic setting where bare essential services are available to provide gynecologic cancer care. The other three settings, limited, enhanced to maximal, offer additional capacities that are essential in providing care and improving survival for cervical cancers. Both guidelines stressed that the highest level of care be provided to women whenever available.

Table 1:

[Table 1]

A Global Call to Action

It is no exaggeration to say that cancer represents an imminent and severe crisis for developing countries, with cervical cancer leading as the one of the most prevalent. A resounding call to action for this crisis is building; advocates are needed to save lives. For example, a recent report from the *Lancet Oncology* presented a body of evidence that quantifies the worldwide shortage of radiotherapy services by country. By scaling up radiotherapy departments in lower-middle- income countries, we could potentially see more than 26.9 million life-years saved in low- and middle- income countries over the lifetime of the patients who received treatment [65].

A global call to action against cancer in low- and middle-income countries is desperately needed [66]. While there are organizations attempting to make a difference [67,68], much needs to be accomplished to stem this global crisis. Seven hundred and forty woman die **per day** from cervical cancer. The majority of these deaths are in relatively young woman and the deaths result in unmeasurable pain and suffering. The world health organization (WHO) has made safer motherhood a priority [69], the same urgency needs to be directed to cervical cancer. Vaccination programs are important, but we cannot ignore women that already have HPV. We, and others, implore the global woman's health movement to make the treatment of cervical cancer a priority [70].

Conclusions/Summary

Cervical cancer is one of the leading causes of cancer death among women [1], representing the fourth most common malignancy in women worldwide [3]. In order to tackle this complex problem there needs to be action on multiple fronts, including

primary and secondary prevention, improvements in treatment and access to care. The treatment of cervical cancer is a global health crisis that needs to be a call to action for the world health community. The GCIG through CCRN is bringing relevant and important trials to low and middle income countries. Our hope is that new attention can be brought to cervical cancer, especially among governmental and philanthropic agencies.

References

1. Global Cancer Statistics. Accessed on 13 May 2016. Available at: <http://www.cdc.gov/cancer/international/statistics.htm>.
2. Human papillomavirus (HPV) and cervical cancer. Accessed on 10 June 2016. Available at: <http://www.who.int/mediacentre/factsheets/fs380/en/>.
3. Global Cancer Facts & Figures 2nd Edition. Accessed on 10 June 2016. Available at: <http://www.cancer.org/acs/groups/content/@epidemiologysurveillance/documents/document/acspc-027766.pdf>.
4. SEER Stat Fact Sheets: Cervix Uteri Cancer. Accessed on 26 October 2016. Available at: <http://seer.cancer.gov/statfacts/html/cervix.html>.
5. Landoni F, Maneo A, Colombo A, et al. Randomised study of radical surgery versus radiotherapy for stage Ib-IIa cervical cancer. *Lancet* 1997;350:535-540.
6. Gadducci A, Sartori E, Maggino T, et al. The clinical outcome of patients with stage Ia1 and Ia2 squamous cell carcinoma of the uterine cervix: a Cooperation Task Force (CTF) study. *Eur J Gynaecol Oncol* 2003;24:513-516.
7. Sedlis A, Bundy BN, Rotman MZ, et al. A randomized trial of pelvic radiation therapy versus no further therapy in selected patients with stage IB carcinoma of the cervix after radical hysterectomy and pelvic lymphadenectomy: A Gynecologic Oncology Group Study. *Gynecol Oncol* 1999;73:177-183.
8. Morris M, Eifel PJ, Lu J, et al. Pelvic radiation with concurrent chemotherapy compared with pelvic and para-aortic radiation for high-risk cervical cancer. *N Engl J Med* 1999;340:1137-1143.
9. Rose PG, Bundy BN, Watkins EB, et al. Concurrent cisplatin-based radiotherapy and chemotherapy for locally advanced cervical cancer. *N Engl J Med* 1999;340:1144-1153.

10. Whitney CW, Sause W, Bundy BN, et al. Randomized comparison of fluorouracil plus cisplatin versus hydroxyurea as an adjunct to radiation therapy in stage IIB-IVA carcinoma of the cervix with negative para-aortic lymph nodes: a Gynecologic Oncology Group and Southwest Oncology Group study. *J Clin Oncol* 1999;17:1339-1348.
11. Eifel PJ, Winter K, Morris M, et al. Pelvic irradiation with concurrent chemotherapy versus pelvic and para-aortic irradiation for high-risk cervical cancer: an update of radiation therapy oncology group trial (RTOG) 90-01. *J Clin Oncol* 2004;22:872-880.
12. Rose PG, Ali S, Watkins E, et al. Long-term follow-up of a randomized trial comparing concurrent single agent cisplatin, cisplatin-based combination chemotherapy, or hydroxyurea during pelvic irradiation for locally advanced cervical cancer: a Gynecologic Oncology Group Study. *J Clin Oncol* 2007;25:2804-2810.
13. Keys HM, Bundy BN, Stehman FB, et al. Cisplatin, radiation, and adjuvant hysterectomy compared with radiation and adjuvant hysterectomy for bulky stage IB cervical carcinoma. *N Engl J Med* 1999;340:1154-1161.
14. Stehman FB, Ali S, Keys HM, et al. Radiation therapy with or without weekly cisplatin for bulky stage 1B cervical carcinoma: follow-up of a Gynecologic Oncology Group trial. *Am J Obstet Gynecol* 2007;197:503 e501-506.
15. Potter R, Georg P, Dimopoulos JC, et al. Clinical outcome of protocol based image (MRI) guided adaptive brachytherapy combined with 3D conformal radiotherapy with or without chemotherapy in patients with locally advanced cervical cancer. *Radiother Oncol* 2011;100:116-123.
16. Sturdza A, Potter R, Fokdal LU, et al. Image guided brachytherapy in locally advanced cervical cancer: Improved pelvic control and survival in RetroEMBRACE, a multicenter cohort study. *Radiother Oncol* 2016.
17. Perez CA, Breaux S, Madoc-Jones H, et al. Radiation therapy alone in the treatment of carcinoma of uterine cervix. I. Analysis of tumor recurrence. *Cancer* 1983;513-1402.
18. Eifel PJ, Thoms WW, Jr., Smith TL, et al. The relationship between brachytherapy dose and outcome in patients with bulky endocervical tumors treated with radiation alone. *Int J Radiat Oncol Biol Phys* 1994;28:113-118.
19. Viswanathan AN, Cormack R, Rawal B, et al. Increasing brachytherapy dose predicts survival for interstitial and tandem-based radiation for stage IIIB cervical cancer. *Int J Gynecol Cancer* 2009;19:1402-1406.
20. Tanderup K, Eifel PJ, Yashar CM, et al. Curative radiation therapy for locally advanced cervical cancer: brachytherapy is NOT optional. *Int J Radiat Oncol Biol Phys* 2014;88:537-539.

21. Sankaranarayanan R, Swaminathan R, Brenner H, et al. Cancer survival in Africa, Asia, and Central America: a population-based study. *Lancet Oncol* 2010;11:165-173.
22. Waggoner SE, Chernicky CL. "Molecular Biology of Cervical and Vulvar Carcinoma." In *Gynaecologic Cancer: Controversies in Management*. Gershenson, Gore, McGuire, Quinn, Thomas. Elsevier Science, Philadelphia, 2004
23. Website: Cervical Cancer Treatment (PDQ®)—Health Professional Version. Available at: http://www.cancer.gov/types/cervical/hp/cervical-treatment-pdq#link/_532_toc. Accessed 26 October 2016.
24. Fujiwara K, Monk B, Devouassoux-Shisheboran M. Adenocarcinoma of the uterine cervix: why is it different? *Curr Oncol Rep*. 2014 Dec;16(12):416. doi: 10.1007/s11912-014-0416-y.
25. Wright AA, Howitt BE, Myers AP, et al. Oncogenic mutations in cervical cancer: genomic differences between adenocarcinomas and squamous cell carcinomas of the cervix. *Cancer*. 2013 Nov 1;119(21):3776-83.
26. Ojesina AI, Lichtenstein L, Freeman SS, et al. Landscape of genomic alterations in cervical carcinomas. *Nature*. 2014 Feb 20;506(7488):371-5.
27. Bulkman NW, Berkhof J, Rozendaal L, et al. Human papillomavirus DNA testing for the detection of cervical intraepithelial neoplasia grade 3 and cancer: 5-year follow-up of a randomised controlled implementation trial. *Lancet*. 2007;370(9601):1764-72. Epub 2007/10/09.
28. Kitchener HC, Almonte M, Thomson C, et al. HPV testing in combination with liquid-based cytology in primary cervical screening (ARTISTIC): a randomised controlled trial. *Lancet Oncol*. 2009;10(7):672-82. Epub 2009/06/23.
29. Naucier P, Ryd W, Tornberg S, et al. Human papillomavirus and Papanicolaou tests to screen for cervical cancer. *N Engl J Med*. 2007;357(16):1589-97. Epub 2007/10/19.
30. Ronco G, Giorgi-Rossi P, Carozzi F, et al. Efficacy of human papillomavirus testing for the detection of invasive cervical cancers and cervical intraepithelial neoplasia: a randomised controlled trial. *Lancet Oncol*. 2010;11(3):249-57. Epub 2010/01/22.
31. Kitchener HC, Gilham C, Sargent A, et al. A comparison of HPV DNA testing and liquid based cytology over three rounds of primary cervical screening: extended follow up in the ARTISTIC trial. *Eur J Cancer*. 2011;47(6):864-71. Epub 2011/02/22.
32. Huh WK, Ault KA, Chelmow D, et al. Use of primary high-risk human papillomavirus testing for cervical cancer screening: interim clinical guidance. *Gynecol Oncol* 2015;136:178–82.

33. Zhao Q, Modis Y, High K, et al. Disassembly and reassembly of human papillomavirus virus-like particles produces more virion-like antibody reactivity. *Virology journal*. 2012;9:52. Epub 2012/02/24.
34. FUTURE II Study Group. Quadrivalent vaccine against human papillomavirus to prevent high-grade cervical lesions. *N Engl J Med*. 2007;356(19):1915-27.
35. Paavonen J, Naud P, Salmeron J, et al. Efficacy of human papillomavirus (HPV)-16/18 AS04-adjuvanted vaccine against cervical infection and precancer caused by oncogenic HPV types (PATRICIA): final analysis of a double-blind, randomised study in young women. *Lancet*. 2009;374(9686):301-14. Epub 2009/07/10.
36. 2015 HPV Academy Policy: American Academy of Pediatrics. Human Papillomaviruses. In: Kimberlin DW, Brady MT, Jackson MA, Long SS, eds. *Red Book: 2015 Report of the Committee on Infectious Diseases*. 30th ed. Elk Grove Village, IL: American Academy of Pediatrics; 2015: 576-583.
37. Dobson SR, McNeil S, Dionne M, et al. Immunogenicity of 2 doses of HPV vaccine in younger adolescents vs 3 doses in young women: a randomized clinical trial. *JAMA*. 2013;309(17):1793-802. Epub 2013/05/02.
38. Joura EA, Giuliano AR, Iversen OE, et al. A 9-valent HPV vaccine against infection and intraepithelial neoplasia in women. *N Engl J Med*. 2015;372(8):711-23. Epub 2015/02/19.
39. Van Damme P, Olsson SE, Block S, et al. Immunogenicity and Safety of a 9-Valent HPV Vaccine. *Pediatrics*. 2015;136(1):e28-39. Epub 2015/06/24.
40. Petrosky E, Bocchini JA Jr, Hariri S, et al. Use of 9-valent human papillomavirus (HPV) vaccine: updated HPV vaccination recommendations of the advisory committee on immunization practices. *MMWR Morb Mortal Wkly Rep* 2015;64:300-4.
41. Sanofi Pasteur Press Release. "Sanofi Pasteur MSD's new 9-valent HPV vaccine is now available in the UK." Accessed 18 December 2016. Available at: <http://www.spmsd.co.uk/sanofi-pasteur-msds-new-9-valent-hpv-vaccine-now-available-uk/>
42. Riethmuller D, Jacquard AC, St Guily JL, et al. Potential impact of a nonavalent HPV vaccine on the occurrence of HPV-related diseases in France. *BMC Public Health* 2015;15: 453.
43. Yang, DY. Update on the new 9-valent vaccine for human papillomavirus prevention. *Canadian Family Physician*, 2016;62:5, 399-402

44. Vici P, Pizzuti L, Mariani L, et al. Targeting immune response with therapeutic vaccines in premalignant lesions and cervical cancer: hope or reality from clinical studies. *Expert Rev Vaccines*. 2016; 9: 1-10.

45. Koh WJ, Greer BE, Abu-Rustum NR, et al. Cervical Cancer, Version 2.2015. *J Natl Compr Canc Netw*. 2015 Apr;13(4):395-404. PMID: 25870376

46. Gaffney DK, Du Bois A, Narayan K, et al. Practice patterns of radiotherapy in cervical cancer among member groups of the Gynecologic Cancer Intergroup (GCIG). *Int J Radiat Oncol Biol Phys*. 2007 Jun 1;68(2):485-90. Epub 2007 Mar 2. PMID:17336465

47. Chemoradiotherapy for Cervical Cancer Meta-Analysis Collaboration¹. Reducing uncertainties about the effects of chemoradiotherapy for cervical cancer: a systematic review and meta-analysis of individual patient data from 18 randomized trials. *J Clin Oncol*. 2008 Dec 10;26(35):5802-12.

48. Ryu SY, Lee WM, Kim K, et al. Randomized clinical trial of weekly vs. triweekly cisplatin-based chemotherapy concurrent with radiotherapy in the treatment of locally advanced cervical cancer. *Int J Radiat Oncol Biol Phys*. 2011 Nov 15;81(4):e577-81.

49. National Comprehensive Cancer Network Guidelines. Cervical Cancer. Available at: <http://www.tri-kobe.org/nccn/guideline/gynecological/english/cervical.pdf> Accessed August 2013.

50. Haie-Meder C, Morice P, Castiglione M, ESMO Guidelines Working Group. Cervical cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Annals of Oncol* 2010; 21:37-40.

51. Viswanathan A, Beriwal S, De Los Santos J, et. al. The American Brachytherapy Society Treatment Recommendations for Locally Advanced Carcinoma of the Cervix Part II: High Dose-Rate Brachytherapy. *Brachytherapy* 2012;11:47-52

52. Han KM, Fyles A, Pintilie M, Viswanathan AN. Trends in the utilization of brachytherapy in cervical cancer in the United States. *Int J Radiat Oncol Biol Phys* 2013;87:111-9.

53. Georg D, Kirisits C, Hillbrand M, Dimopoulos J, Potter R. Image-guided radiotherapy for cervix cancer: high-tech external beam therapy versus high-tech brachytherapy. *Int J Radiat Oncol Biol Phys* 2008;71:1272-8.

54. Patel MK, Cote ML, Ali-Fehmi R, Buckers T, Munkarah AR, Elshaikh MA. Trends in the utilization of adjuvant vaginal cuff brachytherapy and/or external beam radiation

treatment in stage I and II endometrial cancer: a surveillance, epidemiology, and end-results study. *Int J Radiat Oncol Biol Phys* 2012;83:178-84.

55. Nout RA, Smit VT, Putter H, et al. Vaginal brachytherapy versus pelvic external beam radiotherapy for patients with endometrial cancer of high-intermediate risk (PORTEC-2): an open-label, non-inferiority, randomised trial. *Lancet* 2010;375:816-23.

56. Montana GS, Fowler WC, Varia MA, Walton LA, Mack Y, Shemanski L. Carcinoma of the cervix, stage III. Results of radiation therapy. *Cancer* 1986;57:148-54.

57. Lanciano RM, Martz K, Coia LR, Hanks GE. Tumor and treatment factors improving outcome in stage III-B cervix cancer. *Int J Radiat Oncol Biol Phys* 1991;20:95-100.

58. Hanks GE, Herring DF, Kramer S. Patterns of care outcome studies. Results of the national practice in cancer of the cervix. *Cancer* 1983;51:959-67.

59. Corn BW, Hanlon AL, Pajak TF, Owen J, Hanks GE. Technically accurate intracavitary insertions improve pelvic control and survival among patients with locally advanced carcinoma of the uterine cervix. *Gynecologic Oncology* 1994;53:294-300.

60. Viswanathan AN, Moughan J, Small W, Jr., et al. The quality of cervical cancer brachytherapy implantation and the impact on local recurrence and disease-free survival in radiation therapy oncology group prospective trials 0116 and 0128. *International journal of gynecological cancer : official journal of the International Gynecological Cancer Society* 2012;22:123-31.

61. Erickson B, Eifel P, Moughan J, Rownd J, Iarocci T, Owen J. Patterns of brachytherapy practice for patients with carcinoma of the cervix (1996-1999): a patterns of care study. *Int J Radiat Oncol Biol Phys* 2005;63:1083-92.

62. Eifel PJ, Ho A, Khalid N, Erickson B, Owen J. Patterns of radiation therapy practice for patients treated for intact cervical cancer in 2005 to 2007: a quality research in radiation oncology study. *Int J Radiat Oncol Biol Phys* 2014;89:249-56.

63. Gill BS, Lin JF, Krivak TC, et al. National Cancer Data Base analysis of radiation therapy consolidation modality for cervical cancer: the impact of new technological advancements. *Int J Radiat Oncol Biol Phys* 2014;90:1083-90.

64. Maranga IO, Hampson L, Oliver AW, et al. Analysis of factors contributing to the low survival of cervical cancer patients undergoing radiotherapy in Kenya. *PloS one* 2013;8:e78411.

65. Atun R, Jaffray DA, Barton MB, et al. Expanding global access to radiotherapy. *The Lancet Oncology* 2015;16:1153-86.

66. Ferlay J, Shin HR, Bray F, et al. Estimates of worldwide burden of cancer in 2008. GLOBOCAN. *Int J Cancer*. 2010;127:2893–917

67. Einck JP, Hudson A, Shulman AC, et al. Implementation of a high-dose-rate brachytherapy program for carcinoma of the cervix in Senegal: a pragmatic model for the developing world. *Int J Radiat Oncol Biol Phys* 2014;89:462-7.

68. Radiating Hope (2015). Available at: <http://www.radiatinghope.org>. Accessed in 2015.

69. AbouZahr C. Safe motherhood: a brief history of the global movement 1947-2002. *Br Med Bull*. 2003, 67: 13-25.

70. Singhrao R, Huchko M, Yamey G. Reproductive and maternal health in the post-2015 era: cervical cancer must be a priority. *PLoS Med* 2013;10:e1001499.

Figure Legends:

Figure 1. Countries around the world have many GCIG members or are interested in joining GCIG or CCRN. Reproduced with permission from Gaffney, D.K., Suneja G., Ryu S.Y., McCormick M., Plante, M., Mileshekin, L., Small, Jr., W., Bacon, M., Stuart, G., Kitchener, H. The Cervix Cancer Research Network: A Global Outreach Effort on Behalf of the Gynecologic InterGroup, International Journal of Radiation Oncology*Biophysics, 2015, 92:506-508.

Figure 2. Estimated age-standardized rates (world) of deaths, females, cervical cancer, worldwide in 2012. Reproduced with permission from Ferlay J., Soerjomataram I., Ervik M., Dikshit R., Eser S., Mathers C., Rebelo M., Parkin D.M., Forman D., Bray, F. GLOBOCAN 2012 v1.0, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 11 [Internet]. Lyon, France: International Agency for Research on Cancer; 2013. Available from: <http://globocan.iarc.fr>, accessed on 21/12/2016.

Accepted

Table 1. Treatment capacity for different modalities based on setting. Reproduced with permission from Chuang et al., Management and Care of Women with Invasive Cervical Cancer: American Society of Clinical Oncology Resource-Stratified Clinical Practice Guideline, Journal of Global Oncology, 2(5):311-340.

Table 1 – Treatment Capacity.

Treatment	Setting			
	Basic	Limited	Enhanced	Maximal
Surgery	Simple (extrafascial) hysterectomy or more extensive hysterectomy can be performed*	Modified radical or radical hysterectomy	Capable of performing most major surgeries, including radical hysterectomy, radical trachelectomy,† pelvic and para-aortic LN sampling, and pelvic exenteration† Following are not available: PET scan, interventional radiology, sentinel node biopsy/IORT, or bevacizumab	Radical hysterectomy, radical trachelectomy, pelvic and para-aortic LN sampling, sentinel node biopsy, and pelvic exenteration; RT, chemotherapy, interventional radiology, palliative care service, and bevacizumab are all available
Chemotherapy	Availability of chemotherapy drugs is unpredictable	Chemotherapy may be available	Chemotherapy available; bevacizumab not available	Chemotherapy available; bevacizumab is available
RT	No RT available	Limited external RT with no brachytherapy available; in some areas where there is only brachytherapy and no external RT, this will be considered as basic level	RT including external beam and brachytherapy available; interventional radiology not available	RT including external beam and brachytherapy available; interventional radiology available
Pathology	Pathology services are not available; if there is a way to send pathology for review when needed, that should occur (Basic pathology may be available, but diagnosis is often delayed for more than 1 month; there are no frozen sections or pathology consultations in the region)	Pathology services in development (There are basic pathology and frozen section services; consultations are not readily available)	Pathology services in development or not always available (Pathology services including frozen sections are available; tumor registry and regular multidisciplinary conferences are not consistently available in the region)	Pathology available (Full pathology services including diagnosis, consultation, tumor registry, and multidisciplinary conferences are available)
Palliative care	Palliative care service is in development; basic palliative care, including pain and symptom management, should be provided‡	Pain and symptom management available; palliative care service is in development	Palliative care service not always available	Palliative care service available

NOTE. It is the view of the American Society of Clinical Oncology that health care providers and health care system decision makers should be guided by the recommendations for the highest stratum of resources available. This guideline is intended to complement but not replace local guidelines. Bold font indicates addition of a recommended action over a previous resource level (eg, in limited setting, a bold action is one that was not recommended in basic).

Abbreviations: IORT, intraoperative radiation therapy; LN, lymph node; PET, positron emission tomography; RT, radiotherapy.

*Where medical facilities exist to take care of women who are at high risk for postoperative complications.

†Can be performed at some enhanced levels.

‡Palliative care is multifaceted and in some contexts can be provided concurrently with tumor-directed therapy. Pain management and best supportive care are necessary but insufficient parts of palliative care in all settings. Women with advanced cervical cancer with or without access to tumor-directed therapy may have specific late-stage symptoms that require clinicians to perform or offer urogenital-specific interventions. See Special Commentary.

Figure 1

Accepted

Figure 2

Accepted A

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Small, W;Bacon, MA;Bajaj, A;Chuang, LT;Fisher, BJ;Harkenrider, MM;Jhingran, A;Kitchener, HC;Mileshkin, LR;Viswanathan, AN;Gaffney, DK

Title:

Cervical Cancer: A Global Health Crisis

Date:

2017-07-01

Citation:

Small, W., Bacon, M. A., Bajaj, A., Chuang, L. T., Fisher, B. J., Harkenrider, M. M., Jhingran, A., Kitchener, H. C., Mileshkin, L. R., Viswanathan, A. N. & Gaffney, D. K. (2017). Cervical Cancer: A Global Health Crisis. *CANCER*, 123 (13), pp.2404-2412. <https://doi.org/10.1002/cncr.30667>.

Persistent Link:

<http://hdl.handle.net/11343/292853>