DR. AMAYA LOPEZ BUSTINDUY (Orcid ID : 0000-0001-6131-4159)

Article type : Short Communication

Short communication

**Point-of-care tests for infectious diseases: barriers to
implementation across three London teaching hospitals**

Amaya L. Bustinduy^{1,2*}, Dakshika Jeyaratnam³, Sam Douthwaite⁴, Sarah Tonkin-Crine⁵,
Rebecca Shaw¹, Laura Hyrapetian⁶, Nick Sevdalis⁷, Simon Goldenberg⁴, Elizabeth Adams⁸,
Mike Sharland¹ (CLAHRC South London; Paediatric Infection Network).

¹ Paediatric Infectious Diseases Research Group, Infection and Immunity Institute, St George's
University of London, UK;

² Clinical Research Department, London School of Hygiene & Tropical Medicine, London, UK;

³ Department of Microbiology, King's College Hospital NHS Foundation Trust, London, UK;

⁴ Virology Department, Guy's and St Thomas' NHS trust, London, UK;

⁵ Nuffield Department of Primary Care Health Sciences, University of Oxford;

⁶ St George's NHS trust, London, UK;

⁷ Health Service & Population Research Department, King's College London, UK;

⁸ Aquarius Population Health, London, UK

Corresponding author:

*Amaya Bustinduy

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/APA.13867](https://doi.org/10.1111/APA.13867)

This article is protected by copyright. All rights reserved

27 Clinical Research Department

28 London School of Hygiene & Tropical Medicine

29 Keppel Street, WC1, London, UK

30 Amaya.Bustinduy@lshtm.ac.uk

31

32 Left running header: *Bustinduy and others*

33 Right running header: *Implementation barriers for POCT*

34

35 Word count: 1544

36 Figures: 1

37 Tables: 1

38

39

40

41 **List of abbreviations**

42 POCT- Point-of-care tests

43 CRP-C-reactive protein

44 RSV- Respiratory Syncytial Virus

45 NHS- National Health Service

46 ED- Emergency Department

47 KCH- King's College Hospital

48 SGH- St George's Hospital

49 GSTT- Guy's and St Thomas' Hospital

50 PCR- Polymerase Chain Reaction

AMR- Antimicrobial Resistance

Existing Point-of-care tests (POCT) to help identify infection-related causes of illness can complement diagnostic and disposition decisions in children attending emergency departments.(1) Evidence-based clinical algorithms can integrate such POCT to aid in the admission and discharge decision process. Paediatric studies validating these tools are scarce, with very few studies conducted in UK centres.(2-5) POCT can be based on host infection markers (e.g. finger prick tests for C-reactive protein (CRP) to help decide if the patient has a bacterial or viral infection) or pathogen detection tests (e.g. throat/nose swabs to rapidly diagnose viral infections such as RSV or influenza). The use of POCT may reduce time in the emergency department,(6) help rationalise antibiotic prescribing,(4) and reduce investigations in these children.(1, 5) On admission to the ward, POCT can also help with infection control procedures to reduce the risk of transmission of hospital-related infections. These benefits however have been mostly documented in adults (7).

We present the experience across of three paediatric tertiary centres in South London with different POCT implementation strategies, research driven, during one respiratory disease season (winter 2014 -spring 2015) with the aim of improving paediatric clinical process outcomes and potentially reduce antibiotic use, and primarily focus on the barriers encountered for implementation.

At St George's NHS trust, a service evaluation of febrile children presenting to Emergency Department (ED) was undertaken from October 2014 to March 2015. The aim of the study was to collect clinical information on febrile children and to develop an evidence-

based tool to reduce avoidable admissions in those with low-risk infections. Results from the main study are reported elsewhere (8) Outcome data included disposition of the patient, antibiotic use and re-attendance to the ED within 28 days. As a secondary outcome, ED staff were trained on the use of three POCT that were introduced by company provided training: Alere-Afinion™ CRP test (4 minutes), Alere i influenza A/B™ (10 minutes) and Alere BinaxNOW® RSV card (15 minutes). ED nurses and doctors were interviewed about their experience of managing febrile children, following national guidelines [9], and their perceptions on using POCT. Semi-structured interviews were carried out following an interview guide and interviews were audio-recorded and transcribed verbatim. At King's College Hospital (KCH, an observational study of children presenting with symptoms or signs of upper respiratory tract infection to was undertaken from January to July 2015. A ward-based diagnostic platform called BioFire Filmarray (BioMérieux, Marcy-l'Étoile, France) was used with company provided training. Ward staff were trained to perform the respiratory pathogen assay which detects seventeen respiratory viruses and three bacteria by polymerase chain reaction (PCR) in 70 minutes. A staff satisfaction questionnaire was used with a graded response 1-5 on ease of use and utility. Questionnaires were distributed trust wide. At Guy's and St Thomas' NHS trust – Evelina Children's Hospital (GSTT), a ward-based service evaluation was conducted from October 2014 to March 2015 at, using the Enigma® MiniLab™ FluAB-RSV PCR assay (Enigma Diagnostics Ltd, Salisbury, UK), a fully automated molecular platform able to detect influenza A and B and RSV in under 90 minutes. (9-11) The platform was introduced by company provided training of staff with a competency assessment then by the research nurse at which point staff got a barcode for the machine. Relevant staff identified by ward matron and interested clinical teams. Ward staff were trained to perform the test on any child with signs or symptoms of upper or lower respiratory tract infection. Duplicate samples were obtained in viral transport medium and tested in parallel with the standard laboratory-based assay (RVP Fast v2, Luminex, Austin, TX, USA). Diagnostic accuracy, error rate, turnaround time and use of hospital resources were measured and compared with the previous influenza season in which only the RVP was used. A staff satisfaction questionnaire was offered to all ward staff (trained or untrained in the use of the test) contacted through an electronic trust wide email distribution list, and they were asked to provide quantitative and qualitative feedback about their experience of using the Enigma test.

A total of 942 paediatric patients had a POCT performed across the three centres during October 2014 to July 2015. (**Table 1**) At St. George's ED, POCT uptake was 30% less than the laboratory counterpart test for CRP and viral respiratory panel, with 102 POCT compared to 341 laboratory-based tests. This was mainly for serum CRP samples, whereby

225 were sent to the laboratory compared to 41 tested by POCT. Fifteen medical ED staff including 3 consultants, 5 junior doctors, 4 senior registrars and clinical fellows and 3 nurses were interviewed to explore individual perceptions on the use of these tests. Overall, the concerns about using the CRP POCT were; being unsure about performance of the test, missing the 'odd sick child' and using the test in isolation. The advantages mentioned were; the immediateness of the result available in 4 minutes, the ease of use and being a blood sparing procedure. For the respiratory viruses POCT they were highly appreciated for the possibility of quickly cohorting patients, decreasing antibiotic use and being able to reassure parents. Downfalls mentioned included questions about performance of the test, error messages and sample duplication.

At KCH, 460 tests were performed on the ward. The BioFire platform was well received. Any concerns from staff members were shared with the PI. Thirty-four respondents, a mixture of nursing staff and paediatric consultants, completed the survey. Highlighted advantages included; fast results in 1 hour, good tests for infection control, early treatment and reduction of length of stay. Concerns included a limitation on space to house the equipment, needed for additional trained staff and high cost of the test.

In the GSTT paediatric ward, the ENIGMA platform POCT used had a good uptake among staff. More POCT were done compared to routine laboratory testing (462 patients tested by POCT vs 370 patients tested by routine laboratory). Results from the free text portion of the questionnaire from 34 staff who completed it were analysed for themes. Advantages included; parental anxiety reduction, better bed allocation, fast cohorting or isolation, increased safety and reduction in antibiotic use. Concerns raised by staff members included difficulty handling discrepant results (POCT-laboratory), more people needed to be trained and no changes to patient care observed.

We have described three hospitals across South London with different settings (ED and ward-based) and testing/clinical protocols but that all implemented a POCT to improve process outcomes and complement antibiotic stewardship programmes. In the two ward-based centres, although staff overall welcomed the POCT, the main issue was not having enough trained personnel in all shifts. As the uptake of their test increased beyond the ward where the machines were housed, excessive demands were placed on the trained staff to run tests from patients outside their ward, leading to a queue of tests to be run. The decision by the KCH and GSTT trusts to implement such a POCT will ultimately depend on impact on length of stay, antibiotic use and cost-effectiveness analysis (analysis awaited). In the ED-based study (St. George's ED), implementation barriers were centred around the clinician's uncertainty about the test and fear of under-investigating a potentially sick child. Other

problems included running a new technology within a busy service that was not yet integrated into the departmental clinical care pathways. Particular reticence was found with the implementation of CRP and the fear of a poor negative predictive value as a rule-out test, despite the growing evidence that this is not the case. (12) Overall, the ward-based implementation had a more positive feedback from the staff, however parallel sampling for laboratory testing makes this platform difficult to justify its cost.

This report is set within context of the global threat of antimicrobial resistance (AMR) and the growing recognition on the value of POCTs as valuable rapid diagnostic tools. (13, 14) Scaling up of these tests may help reduce unnecessary empiric antibiotic use and reduce invasive investigations often performed to overcome diagnostic uncertainty.(15) Implementing these tests, however, is challenging. (2) Behaviour changes are needed to modify long-established ways of providing clinical care and empirically prescribing antibiotics for febrile children. (16) New technology needs time to become established as part of routine care and to build evidence on its potential benefits.(17) The need to shift practice and integrate POCT into paediatric care as in adult settings is clear, and will require a multi-stage approach. Overcoming implementation barriers is paramount in order to succeed in this approach and to make local and national recommendations. (18)

We propose a stepwise framework to integrate POCT into paediatric clinical practice based on the three major areas identified from the different centres as summarised in **Figure 1**. 1) POCT RCT paediatric studies: multi-centred studies with cost-effective analysis to demonstrate improved outcomes. 2) Increasing staff knowledge and confidence to decrease uncertainty on the tests: Need for more comprehensive and teaching and training on the different devices, with particular emphasis on their evidence-based efficacy; 3) Integration: integrate POCT into existing care pathways and guidelines.

Qualitative assessment on the impact of interventions are helpful to understand implementation difficulties and can help pave the way for future integration of novel interventions such as POCT into paediatric practice.

Acknowledgments

We would like to thank the staff members at St George's Children's ED that contributed to the work presented. To Alere TM for the donation of the POCT and reagents for the study. Enigma diagnostics for funding the GSTT project and Evelina London Mountain ward staff, patients and parents for support and feedback. Thank staff and patients of the paediatric (especially Toni and Guy) wards and bioMérieux for the diagnostic platforms and tests.

186

187 **Competing interests and funding**

188 DJ project funding from bioMérieux.

189 SD project funding from Enigma diagnostics.

190 EA, CM and AV work for Aquarius who were funded by Enigma Diagnostics to perform the
191 study.192 **References**

- 193 1. Doan Q, Enarson P, Kissoon N, Klassen TP, Johnson DW. Rapid viral diagnosis for acute febrile
194 respiratory illness in children in the Emergency Department. *Cochrane Database Syst Rev* 2014;
195 9:CD006452
- 196 2. Bonner AB, Monroe KW, Talley LI, Klasner AE, Kimberlin DW. Impact of the rapid diagnosis of
197 influenza on physician decision-making and patient management in the pediatric emergency
198 department: results of a randomized, prospective, controlled trial. *Pediatrics* 2003; 112:363-7
- 199 3. Iyer SB, Gerber MA, Pomerantz WJ, Mortensen JE, Ruddy RM. Effect of point-of-care
200 influenza testing on management of febrile children. *Academic emergency medicine : official journal*
201 *of the Society for Academic Emergency Medicine* 2006; 13:1259-68
- 202 4. Doan QH, Kissoon N, Dobson S, Whitehouse S, Cochrane D, Schmidt B, et al. A randomized,
203 controlled trial of the impact of early and rapid diagnosis of viral infections in children brought to an
204 emergency department with febrile respiratory tract illnesses. *The Journal of pediatrics* 2009;
205 154:91-5
- 206 5. Poehling KA, Zhu Y, Tang YW, Edwards K. Accuracy and impact of a point-of-care rapid
207 influenza test in young children with respiratory illnesses. *Archives of pediatrics & adolescent*
208 *medicine* 2006; 160:713-8
- 209 6. Yen K, Gorelick MH. Strategies to improve flow in the pediatric emergency department.
210 *Pediatr Emerg Care* 2007; 23:745-9; quiz 50-1
- 211 7. Schuetz P, Muller B, Christ-Crain M, Stolz D, Tamm M, Bouadma L, et al. Procalcitonin to
212 initiate or discontinue antibiotics in acute respiratory tract infections. Evidence-based child health : a
213 *Cochrane review journal* 2013; 8:1297-371
- 214 8. Bustinduy AL, Chis Ster I, Shaw R, Irwin A, Thiagarajan J, Beynon R, et al. Predictors of fever-
215 related admissions to a paediatric assessment unit, ward and reattendances in a South London
216 emergency department: the CABIN 2 study. *Arch Dis Child* 2017; 102:22-8
- 217 9. Goldenberg SD, Edgeworth JD. The Enigma ML FluAB-RSV assay: a fully automated molecular
218 test for the rapid detection of influenza A, B and respiratory syncytial viruses in respiratory
219 specimens. *Expert Rev Mol Diagn* 2015; 15:23-32

10. Douthwaite ST, Walker C, Adams EJ, Mak C, Vecino Ortiz A, Martinez-Alier N, et al. Performance of a Novel Point-of-Care Molecular Assay for Detection of Influenza A and B Viruses and Respiratory Syncytial Virus (Enigma MiniLab) in Children with Acute Respiratory Infection. *J Clin Microbiol* 2016; 54:212-5
11. <http://filmarray.com/the-panels/>.
12. Nijman RG, Moll HA, Vergouwe Y, de Rijke YB, Oostenbrink R. C-Reactive Protein Bedside Testing in Febrile Children Lowers Length of Stay at the Emergency Department. *Pediatr Emerg Care* 2015; 31:633-9
13. Jones CH, Glogowska M, Locock L, Lasserson DS. Embedding new technologies in practice - a normalization process theory study of point of care testing. *BMC Health Serv Res* 2016; 16:591
14. Kaman WE, Andrinopoulou ER, Hays JP. Perceptions of point-of-care infectious disease testing among European medical personnel, point-of-care test kit manufacturers, and the general public. *Patient Prefer Adherence* 2013; 7:559-77
15. O'Neill J. Rapid diagnostics: stopping unnecessary use of antibiotics. The review on antimicrobial resistance. UK: *Wellcome Trust; HM Government* 2015
16. Byington CL, Castillo H, Gerber K, Daly JA, Brimley LA, Adams S, et al. The effect of rapid respiratory viral diagnostic testing on antibiotic use in a children's hospital. *Archives of pediatrics & adolescent medicine* 2002; 156:1230-4
17. Jennings LC, Skopnik H, Burckhardt I, Hribar I, Del Piero L, Deichmann KA. Effect of rapid influenza testing on the clinical management of paediatric influenza. *Influenza Other Respir Viruses* 2009; 3:91-8
18. CLAHRC SL. <http://www.clahrc-southlondon.nihr.ac.uk/>.

Table 1: Number of patients tested within the study period by POCT and in the reference laboratory.

	POCT	Main Laboratory
St George's NHS trust (Children's ED)	102	341
King's College Hospital (Paediatric ward)	460	520
Guy's and St Thomas' trust (Paediatric ward)	462	370

Author Manuscript

Figure 1: Proposed step-wise process for the full integration of POCT into paediatric clinical practice. Randomised control trials (RCT) with integrated cost effectiveness analysis. Training of medical staff to increase uptake leading to practice change.

apa_13867_f1.tif

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Bustinduy, AL;Jeyaratnam, D;Douthwaite, S;Tonkin-Crine, S;Shaw, R;Hyrapetian, L;Sevdalis, N;Goldenberg, S;Adams, EJ;Sharland, M

Title:

Point-of-care tests for infectious diseases: barriers to implementation across three London teaching hospitals

Date:

2017-07-01

Citation:

Bustinduy, A. L., Jeyaratnam, D., Douthwaite, S., Tonkin-Crine, S., Shaw, R., Hyrapetian, L., Sevdalis, N., Goldenberg, S., Adams, E. J. & Sharland, M. (2017). Point-of-care tests for infectious diseases: barriers to implementation across three London teaching hospitals. ACTA PAEDIATRICA, 106 (7), pp.1192-1195. <https://doi.org/10.1111/apa.13867>.

Persistent Link:

<http://hdl.handle.net/11343/292887>