**Fetal brain injury in complicated monochorionic pregnancies: diagnostic yield of prenatal MRI following surveillance
ultrasound and influence on prognostic counselling**

Running Head: Prenatal MRI for brain injury in complicated monochorionic gestations

Authors: Alice Robinson¹, Mark Teoh¹, Andrew Edwards^{1,2}, Michael Fahey^{3,4}, Stacy Goergen^{5,6}

¹ Fetal Diagnostic Unit, Monash Health, 246 Clayton Road, Clayton, Victoria 3168, Australia

² Department of Obstetrics and Gynaecology, Monash University, 246 Clayton Road, Clayton, Victoria 3168, Australia

³ Department of Paediatrics, Monash University, 246 Clayton Road, Clayton, Victoria 3168, Australia

⁴ Paediatric Neurology Unit, Monash Health, 246 Clayton Road, Clayton, Victoria 3168, Australia

⁵ Departments of Medical Imaging and Surgery, Monash University, Southern Clinical School, 246 Clayton Rd, Victoria 3168,

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1002/pd.5059](https://doi.org/10.1002/pd.5059)

Australia

⁶ Monash Imaging, Monash Health, 246 Clayton Rd, Clayton, Victoria 3168, Australia

Corresponding Author:

Stacy K Goergen MBBS (Hons), MCLinEpi, FRANZCR

E: Stacy.Goergen@monashhealth.org

T: +613 9594 2200

M: +61409190660

246 Clayton Rd.

Clayton, Victoria 3168

Australia

The authors have no conflicts of interest or funding sources to declare.

Word Count:

Main text = 4101 (Abstract = 190, References = 1036)

Tables – 4; Figures – 9

What is already known about this topic:

- Complicated monochorionic pregnancies are associated with an increased risk of adverse neurodevelopmental outcome with some injuries occurring prenatally, and prematurity likely to be a significant contributing factor to these adverse outcomes. A small number of case series have reported increased sensitivity and specificity of magnetic resonance imaging (MRI) compared with ultrasound for fetal brain injuries following twin – twin transfusion syndrome and co-twin demise.

What does this study add:

- Fetuses with normal cranial ultrasound (US) who were survivors of co-twin demise either before or after twin – twin transfusion (TTTS) were more likely to have abnormal brain magnetic resonance imaging (MRI) findings that changed prognostic counselling than were co-survivors of TTTS, selective intrauterine growth restriction, or twin anemia - polycythemia sequence.
- Addition of fetal MRI to the routine surveillance of complicated monochorionic gestations (cMCGs) with US may help to refine neurodevelopmental prognostication, and therefore pregnancy management, and postnatal care of children who are

survivors of co-twin demise. A prospective evaluation of the incorporation of fetal MRI, including brain volumetry, into the routine surveillance of cMCGs is required to inform clinical practice guidelines regarding the routine use of MRI in the imaging surveillance of these pregnancies.

ABSTRACT:

Objective: To determine the additional diagnostic information provided by prenatal (fetal) magnetic resonance imaging (pMRI) following tertiary ultrasound (US) for fetal cranial abnormalities in complicated monochorionic gestations (cMCGs)

Methods: Women with cMCGs complicated by twin – twin transfusion syndrome (TTTS), co-twin demise (CD), selective intrauterine growth restriction (sIUGR), and / or twin anemia – polycythemia sequence (TAPS) who were referred for pMRI after tertiary US were included. Primary outcome: additional diagnostic information by pMRI that changed prognostic counselling.

Results: 33 women with 48 live fetuses had pMRI at a median of 25 weeks (range: 21 – 29). 3/10 survivors of spontaneous CD, 1/8 survivors of CD after TTTS and 1/30 co-survivors had diagnostic information added by pMRI that altered counselling; US was normal in 2 and in the other 3 under- represented parenchymal injury ($5/33 = 15\%$; 95% CI $\pm 0.27 - 0.03$). Additional findings included occipital lobe infarction, hemispheric injury, dural sinus thrombosis, ischemia producing polymicrogyria, and intraventricular hemorrhage. Another 8 /33 women had additional information provided by pMRI that did not alter counselling.

Conclusion: pMRI resulted in changed prognostic counselling in 5/33 pregnancies. Evaluation of incorporation of pMRI into routine surveillance of complicated MCGs is needed.

Introduction

Monochorionic twins represent 20% of all multiple gestations, and of these, 15 - 30% will develop complications related to placental sharing such as twin – twin transfusion syndrome (TTTS), twin anemia – polycythaemia sequence (TAPS), selective growth restriction (sIUGR), and fetal death in utero. These complications, and their treatments (including fetoscopic surgery, intrauterine transfusion, and preterm delivery) confer increased risk of neurodevelopmental impairment to the survivors.

Survivors of intrauterine co-twin death (CD) have a greater than 20% average risk of neurodevelopmental impairment (NDI)¹⁻³.

Although fetoscopic laser photocoagulation of placental anastomosing vessels (FLPAV) is now the treatment of choice for sonographically diagnosed TTTS of Quintero stage II or greater⁴, between 3 and 16% of survivors have brain abnormalities that are demonstrable on postnatal cranial ultrasonography⁵⁻⁷ and 6–25% suffer from NDI⁸⁻¹². In a recent evaluation of children participating in the Eurofoetus trial, 12% of survivors of TTTS treated with FLPAV were found to have neurological morbidity, either owing to prematurity or as a result of hemodynamic changes associated with the disease and its treatment¹³. However, survivors of CD have an even higher risk of NDI especially when the demise occurs prior to or without FLPAV treatment for TTTS¹.

An uncertain proportion of the total cerebral injuries that occur in complicated monochorionic gestations (cMCGs) are prenatal, and

identification of such injuries has the potential to change pregnancy counselling and management. Improved prenatal diagnosis of brain injuries would allow prognostic refinement for the individual fetus that goes beyond informing families about the general risk of NDI known to be associated with specific complications of monochorionicity. While ultrasound (US) is the primary imaging modality used to monitor cMCGs for complications, more recently, prenatal MRI (pMRI) has been utilized to evaluate the intracranial structures in fetuses affected by TTTS and / or death of a monochorionic co-twin¹⁴⁻²⁴. Brain abnormalities have been identified with pMRI in between 11 and 100% of these cases, in series ranging from 1 – 27 pregnant subjects, when US was normal. However, the effect of pMRI following tertiary US, with regard to prognostic counselling of women with a range of monochorionic gestation complications, including fetal co-survivors of TTTS, fetuses with sIUGR and TAPS as well as co-twin demise, has previously not been evaluated.

The aim of this study was to determine the additional diagnostic information provided by pMRI in cMCGs when pMRI is performed following tertiary prenatal US, and to assess the effect on prognostic counselling.

Patients and Methods

A waiver of the requirement for a full ethics submission was provided by our institutional review board (Project number 16230Q) on the basis of the study protocol satisfying criteria for quality assurance / audit and in compliance with the provisions of the Declaration of Helsinki.

This was a single-institution retrospective study of patients referred to our institution between January 2007 and January 2016 who were referred for management of complications in a monochorionic pregnancy. Our institution is a tertiary and quaternary referral center for fetal diagnosis, intervention, and pMRI. During the study period, an average of 24 women per year (range 14 – 35) with complicated monochorionic gestations were referred to our fetal diagnostic unit. All women who had tertiary prenatal US performed at our center as well as pMRI for the same pregnancy and who had a diagnosis of either:

- i) Quintero Stage I²⁵ or greater TTTS treated with FLPAV followed by
 - a. Co-survivorship of both twins
 - b. Demise of one of the twins or
- ii) Spontaneous CD

- iii) sIUGR (defined as estimated fetal weight below the 10th percentile²⁶)
- iv) spontaneous or post-FLPAV TAPS

were included in this study. Women were referred for pMRI at the discretion of the fetal diagnostic unit based on abnormal US findings and / or increased risk of fetal NDI based on complications of monochorionicity. Timing of pMRI was determined on an individual basis, usually after 24 weeks gestation, and greater than 2 weeks after CD or FLPAV. The identification of included study subjects from our fetal diagnostic unit and pMRI databases and radiology information system (RIS) is illustrated in Figure 1. Women with MCGs without one or more of TTTS, CD, sIUGR, or TAPS were not referred for pMRI and thus not included in the study. Diagnosis of cMCG was often made in an external facility or institution but verified in all cases with US following referral to our institution.

MRI Technique:

pMRI was performed on a 1.5T Avanto MRI unit (Siemens, Erlangen, Germany) using an 8-channel torso array and 20-channel spine array provided by the scanner manufacturer. No sedation was used and patients were not asked to fast. The scan was

performed in the left side down decubitus position unless the patient preferred to lie completely supine. The field of view was 250 – 300 mm and matrix 256 X 256. Pulse sequences used to image the fetal brain included: T2 HASTE images with 3mm slice thickness and no gap in three orthogonal planes and in addition, manually triggered single images were acquired during maternal breath holding as necessary if fetal motion was producing substantial motion degradation. In all cases, axial, coronal and sagittal T2 HASTE and axial and coronal fat suppressed T1 weighted images were acquired through the fetal neck, chest, abdomen and pelvis, as is our routine for all pMRI studies. However, the abnormal cranial findings in particular are the subject of this study. Echoplanar gradient echo and diffusion weighted (DW) acquisitions ($b = 0, 1000$), as well as T1 – weighted axial images, and sagittal and axial True FISP sequences were acquired. Only one fetus (subject 11) studied in 2007 prior to our routine performance of DWI, was evaluated without this sequence. Duration of the fetal cranial study was approximately 25 minutes, but occasionally longer if additional acquisitions were required due to motion degradation.

Image analysis

Qualitative interpretation of all studies was undertaken at the time of the study by either one of two radiologists with several years' experience in pMRI performance and interpretation and subspecialisation in paediatric radiology or paediatric neuroradiology. This interpretation occurred with knowledge of the preceding ultrasound findings. In all cases, a report template was used to standardize data collection / reporting. We also performed quantitative assessment of the fetal brain by measuring the cerebral hemispheres (fronto-occipital cerebral, biparietal cerebral, and biparietal bone diameter), corpus callosum length, vermis (superior – inferior height, anterior – posterior width at the fastigial point), cerebellum (transverse cerebellar diameter) and comparing these measurements with published gestational age normative data^{27,28}.

All ultrasound studies were performed and interpreted by one of three obstetrician sonologists with specific training and qualification in obstetric and gynecological ultrasound and fetal diagnosis. All studies were performed using Philips iU-22 (Koninklijke Philips Electronics NV, Eindhoven, The Netherlands), or GE Voluson E8 Expert (GE Medical Systems, Kretztechnik GmbH & Co, Zipf, Austria) ultrasound machines. Examinations were performed transabdominally using C5-1, C5-2, and C9-4 MHz transducers on Philips machines or C4-8 and C1-5 MHz transducers on GE machines. Where satisfactory views could not be obtained and the fetus was cephalic, transvaginal ultrasound was performed using 5-9 MHz endovaginal transducers.

Ultrasound analysis included fetal biometry (head circumference, bone biparietal diameter from outer to inner of the fetal skull, abdominal circumference, femur length, and estimated fetal weight), which was compared to normative data²⁹⁻³², as well as triplanar neurosonography according to the International Society of Ultrasound in Obstetrics and Gynaecology Guidelines³³ (including subjective assessment of extra-axial space, cerebral hemispheres, interhemispheric fissure, sylvian fissure, cavum septum pellucidum and corpus callosum, thalami, brainstem, lateral ventricles, cerebellar hemispheres and vermis, and objective measurements of the atrium of the lateral ventricles, cerebellar diameter³⁴ and cisterna magna). During the 9 years of the study, increasing experience enabled more detailed visualization of sulcation patterns by ultrasound.

The gestational age at the time of the latest ultrasound preceding the first pMRI was recorded and the time elapsed between the two studies calculated. In women who had more than one pMRI, only the first pMRI date and result were used to determine interval between US and pMRI and only the findings on the first pMRI of the series is presented in the tables.

Data Collection

The following data were compiled from our institutional databases:

- a. the nature of the complication of monochorionicity and any treatment provided such as FLPAV or intrauterine transfusion
- b. gestational age at the time of the latest US preceding pMRI and the pMRI
- c. nature of US and pMRI findings
- d. pregnancy outcome
- e. post mortem results where available
- f. postnatal imaging findings

Data Analysis

For each case, an assessment was made jointly by the three co-authors who are subspecialists in obstetric and gynaecological ultrasound regarding the contribution of pMRI to the diagnostic evaluation of the fetal head for structural abnormalities by scoring their subjective interpretation of the added value of pMRI in following manner:

1. Normal (with a preceding normal US)
2. Normal (with a preceding abnormal fetal brain US)

3. Abnormal (with a preceding abnormal or normal fetal brain US) but no clinically important added information provided by pMRI
4. Abnormal (with a preceding abnormal fetal brain US) and clinically important information provided by pMRI.
5. Abnormal (with a preceding normal fetal brain US) and clinically important information provided by pMRI.

The number and proportion of pregnancies with additional findings on pMRI and changed prognostic counselling were measured as were the number of cases with additional findings on pMRI that did not alter counselling. Time elapsed between US and pMRI for cases with and without additional findings that altered counselling was compared.

The chi-square test was used to test for differences in the proportion of cases in which counselling changed as a result of pMRI for pregnancies with co-twin demise and co-twin survivorship. The Wilcoxon signed rank test was used to assess for differences in non-parametric continuous data, and the paired t-test for differences in normally distributed continuous data.

Results

During the study period, 216 women with MCGs were referred to the fetal diagnostic and therapy unit of our institution due to suspected complications of MCGs including TTTS, CD, TAPS and sIUGR. 33 of these women, with 48 live fetuses, had a pMRI with preceding tertiary US (Figure 1) and it was these women and their fetuses who were our study subjects, as summarized in Tables 1, 2, and 3. The rate of brain abnormalities overall was 15% (7/48) by ultrasound and 27% (13/48) by pMRI. In the group with TTTS, there was a 14% (3/22) rate of abnormalities detected by ultrasound, and 18% (4/22) by pMRI. In the small group of fetuses with TAPS or sIUGR, there was respectively a 25% (1/4) and 0% (0/4) rate of brain abnormality by ultrasound, and a 25% (1/4) and 50% (2/4) rate of brain abnormality by pMRI. Finally, in the group with CD, the rate of brain abnormalities was 20% (2/10) by ultrasound and 30% (3/10) by pMRI in those with spontaneous CD, and 13% (1/8) by ultrasound and 38% (3/8) by pMRI in those with CD post-FLPAV.

Thirty-one of the 33 women had pregnancy outcome data available. One autopsy was performed out of 6 perinatal deaths in the series and is presented in Table 1. Systematic postnatal paediatric and neurology follow-up was not available as patients did not always attend our institution for follow-up. From 37 live births, only 2 had neonatal follow-up data available up to 12 months of age.

The median gestational age at pMRI was 25 weeks with a range of 21 to 29 weeks. There was no difference in the median gestational age at pMRI between those with an abnormal US (7 fetuses, 7 pregnancies) and those with a normal US (41 fetuses, 26 pregnancies); 25 weeks (range 22 to 28 weeks) versus 25 weeks (range 21 to 29 weeks) respectively.

Ten subjects had serial pMRI studies (9 subjects had 2 pMRI, 1 subject had 3 pMRI), and only the first pMRI is presented in the tables. None of the follow-up pMRI studies showed additional findings, and one showed resolution of diffusion restriction (Subject 24, described below and in figure 9).

I. Additional Findings on pMRI resulting in clinically important additional findings and change to counselling (figures 2 – 6)

Five of 33 women (0.15; 95% confidence interval (CI): 0.027 – 0.03) with five fetuses, 4 of 18 of whom were survivors of co-twin demise and 1 of 30 who was a co-survivor at the time of pMRI, had additional findings that changed prognostic counselling. The difference in proportion of additional clinically important findings on pMRI for co-twin survivors compared with survivors of co-twin demise was statistically significant (chi square test statistic = 4.48, $p = 0.03$, at 0.05 level of significance). These additional findings included: bilateral occipital lobe infarction with evidence of severe bilateral basal ganglia ischemic infarction at post mortem (subject

4, figure 2a-c); dural sinus thrombosis (subject 12, figure 3a-b); asymmetric widening of the extra-axial space overlying one cerebral hemisphere with associated reduced size of the underlying cerebral hemisphere especially affecting the intermediate zone and subplate without evidence of cortical abnormality (subject 10, figure 4a-d); diffuse bilateral hemispheric injury with loss of zonal anatomy and deficient primary sulcal development that evolved to generalized polymicrogyria in the postnatal period (subject 22, figure 5a-f); and intraventricular haemorrhage (subject 3).

II. Additional Findings on pMRI that did not result in change to prognostic counselling (figures 7 – 9)

In eight women, and eight of their fetuses, additional findings that did not alter prognostic counselling were found. In 7 cases, the affected twin was one of two co-surviving twins of the 15 co-surviving pairs ($7/30 = 0.23$; 95%CI: 0.38 – 0.08) and in 1 case, one of the 18 survivors of co-twin demise ($1/18 = 0.06$; 95% CI: 0.05 – 0.16). This difference was not statistically significant (chi square test statistic = 1.74, p value – 0.19 at 0.05 level of significance). These additional findings were: Two cases of extensive hemispheric encephalomalacia with moderate ventriculomegaly demonstrated on US (subjects 19 and 21, figure 6a-c); a germinal matrix infarct in a fetus who represented as a child with gross motor and speech delay at 16 months of age (subject 14, figure 7);

generalized extra-axial space enlargement in two ex-recipients of TTTS with normal brain biometry and no structural abnormality (subjects 23 and 25, figure 8a-c); diffusion restriction in the parieto-occipital region at 22 weeks gestation (19 days post FLPAV) of an ex-donor fetus affected by sIUGR that resolved and did not evolve to local parenchymal ischemic injury on two subsequent pMRI studies over 5 weeks (subject 24, figure 9a-e); two cases of brain biometry on the 5th to 10th percentile on pMRI performed 7 and 21 days after US demonstrating fetal cranial biometry close to the 50th percentile (Subjects 31 and 33).

In one case (Table 3, subject 24) a true diffusion restriction abnormality not only resolved but failed to progress to hemispheric infarction or polymicrogyria on follow-up studies.

III. Time elapsed between US and pMRI

There was no clinically important difference in the time elapsed between US and pMRI in the fetuses with additional findings on pMRI that changed counselling compared to those without (Table 4). However, the small number of subjects in the former group precluded application of the Wilcoxon signed rank test to evaluate this difference for statistical significance.

IV. Gestational age at event or intervention and time elapsed before pMRI

In the 40 fetuses with a known event (co-twin death) or intervention (FLPAV), the median gestational age of the event or intervention was 21 weeks. In those with an abnormality on pMRI, the median gestational age of pMRI was 24 weeks. In those without an abnormality on pMRI, the median gestational age at pMRI was 25 weeks. Overall, the median time elapsed prior to pMRI was 32 days (range 6-55 days). In those with any abnormality on pMRI, the median time elapsed was 30 days, and in those with no abnormality on pMRI, the time elapsed was 34 days, a difference which is unlikely to be of clinical significance.

Discussion

Ultrasound is routinely used to monitor MCGs for specific complications, including TTTS, TAPS, and sIUGR. Pulsed Doppler US plays a key role in diagnosing and monitoring haemodynamic changes in TTTS and TAPS, and placental function in sIUGR.

Specifically, the detection of abnormal umbilical artery, middle cerebral artery and ductus venosus flow velocity waveforms in late stage TTTS and fetal compromise in sIUGR; and the detection of a large discordance in fetal middle cerebral artery peak systolic velocities in the diagnosis of TAPS. FLPAV is technically feasible from 16 weeks gestation onwards. An alternative treatment of fetal intrauterine blood transfusion (IUT) of the anemic fetus is sometimes performed in pregnancies complicated with TAPS. It is

the modality of choice to monitor fetal growth and placental function in these pregnancies in order to plan delivery, particularly in the setting of sIUGR.

While it is clear that US is the method of choice for fetal haemodynamic and growth assessment to enable detection and treatment of the complications of monochorionic twinning, our study has shown that although it frequently identifies “markers” of fetal brain injury, such as ventriculomegaly, periventricular echogenicity, and subarachnoid space enlargement, it may not be the optimal imaging modality in isolation to definitively diagnose abnormalities such as ischemia, haemorrhage, malformations, or to distinguish uncomplicated ventricular or subarachnoid space enlargement from underlying brain injury. In addition, in 4 of 33 of the pregnant women who had fetal prognosis altered by pMRI, US was normal. Although the conventional metrics of diagnostic test performance (sensitivity, specificity and likelihood ratios) cannot be generated for pMRI compared with US, due to lack of a postnatal gold standard, our findings generally support the work of others that has demonstrated the greater accuracy of pMRI for the diagnosis of fetal brain abnormalities, in the setting of co-twin demise and TTTS in particular. We have also demonstrated a lower incidence of prognosis changing additional diagnoses as a result of pMRI when it is performed in co-surviving twins with one or more of: TTTS

treated with FLPAV; sIUGR; or TAPS when compared with the occurrence of brain injury in fetuses who have survived co-twin demise.

There is currently no evidence to guide the timing of pMRI (relative to the occurrence of monochorionicity complications) in order to optimize the detection of abnormalities, and thus to be of most benefit in prognostic counselling. The median gestational age of pMRI in this retrospective study was 25 weeks, with a large range from 21 to 29 weeks. pMRI was performed at a median of 32 days after a known event or intervention (in the 40 cases where this was known). Earlier pMRI may enable earlier diagnosis which is particularly important in jurisdictions with more limitations on access to late termination of pregnancy. Performing pMRI too soon after the demise of a co-twin or diagnosis of advanced stage TTTS may result in evolving ischaemic injury and malformations of cortical development being underdiagnosed, although use of diffusion – weighted imaging offers the potential for acute ischaemic injury to be identified before volume loss, cystic encephalomalacia or malformations of cortical development become evident.

Prospective evaluation of the natural evolution of diffusion restriction within fetal brain parenchyma is required before this information can be used for prognostic counselling when standard T2-weighted gradient echo and single shot T2 weighted spin echo imaging of the fetal brain shows no abnormality.

Quarello et al³⁵ who studied US and MRI for the detection of brain injuries in fetuses affected by TTTS, performed routine prenatal ultrasound surveillance and added pMRI at 32 weeks; pMRI was performed earlier only in the event of an abnormal fetal brain ultrasound examination. Although they stated that “fetal cerebral MRI proved reliable when performed after 28 weeks” they performed imaging using a 1 Tesla MRI scanner and published their study nearly a decade ago, raising questions about the applicability of this conclusion given technological advances, including higher field strength, higher slew rates, diffusion-weighted imaging and parallel processing, that have occurred in the interim. These have resulted in faster scanning and improved image quality. In addition, the retrospective nature of their study, and their performance of early MRI contingent upon the finding of US abnormalities, significantly limits our ability to draw conclusions from their work about the accuracy of early MRI (in other words prior to 24 – 25 weeks). This question is important, both because of variable jurisdictional restrictions on late termination of pregnancy and the need to inform optimal health services utilization for pregnancy surveillance.

Major limitations of the current study include its retrospective nature, non-consecutive referred population (which may bias our conclusions about the proportion of MCGs in whom MRI reveals additional brain abnormalities), lack of postnatal follow-up data, and non-standardized timing of performance of MRI in relation to key events such as diagnosis and treatment of TTTS, recognition

of CD, and development of brain abnormality(ies) diagnosed with US. The median gestational age of pMRI in this study was 25 weeks, and it is possible that some studies may have been performed too early to detect an abnormality. Long term follow up of neurodevelopmental outcome would be the gold standard in determining the accuracy of both US and pMRI in the antenatal diagnosis of major neurological abnormalities in MCGs. It would also help immensely in allowing more precise prenatal counselling in these complicated pregnancies. In addition, long term follow up data for fetuses with apparently minor neurological abnormalities, such as subject 14, who had a small germinal matrix cyst but significant motor and speech delay prior to the age of 2 years, is urgently required to inform prognostic counselling for these patients. In fact, our ability to demonstrate these small abnormalities with contemporary pMRI techniques now surpasses our ability to apply this information. It would be simplistic and inappropriate to extrapolate outcome data from preterm infants with similar abnormalities, such as haemorrhagic or ischaemic infarction.

Dedicated imaging of the entire placenta was not performed in our study. Further evaluation of MRI in the identification of placental complications such as infarction and haemorrhage, and the prognostic importance of such identification in the setting of MCGs and sIUGR, is warranted.

Righini et al²², Hoffman et al²³, and Weisz et al²⁴ observed progression from cerebral hemispheric parenchymal DW abnormality to infarction in 4 fetuses with abnormal hemispheric signal on DW following CD or TTTS, but we did not observe this in our single case that presented with DW imaging abnormality (subject 24). While it is possible that our observation of apparent diffusion restriction was artefactual in this case, due to motion, it is our view that further consecutive patients with CD or post treatment for TTTS need to be studied before the inevitability of completed brain infarction can be assumed when restricted diffusion is identified within the fetal brain in this setting.

Future work should include development and prospective evaluation of a standardized MCG fetal monitoring protocol that involves pMRI (including DWI), as well as US, before and after treatment for TTTS and after CD. Assessment of the relationship between specific types of brain abnormalities demonstrated on pMRI and postnatal and clinical outcome is required to inform parental counselling. Jatzko et al¹⁷ have examined this in 11 survivors of SCD and found that 3 of 6 survivors with prenatally diagnosed cerebral lesions had adverse neurodevelopmental outcome whereas 5 of 5 survivors with normal pMRI had normal

neurodevelopmental outcomes. This requires validation in a larger series with pMRI being performed at standard times during the pregnancy for all subjects.

Fetal brain MRI volumetry and diffusion tensor imaging (DTI) scalar measurements require further evaluation at specific points in pregnancy with regard to their prognostic value in complicated MCGs, including fetuses with sIUGR, especially when qualitative assessment of the fetal brain is normal. Tarui et al³⁶ demonstrated differential growth trajectories for fetal hemispheric parenchyma and cerebella of donor, recipient, and normal control fetuses but used linear measurements and not volumetry to demonstrate this. It is expected that volumetry may be more sensitive in demonstrating a delay in brain growth and thus may be more accurate not only to predict NDI but also to assess the effect of neuroprotective interventions in complicated MCGs.

Current clinical practice guidelines relating to the US surveillance of monochorionic gestations should incorporate recommendations regarding the desirability of centers undertaking treatment of complicated monochorionic gestations having access to pMRI services to enable optimal diagnosis of fetal brain injuries³⁷.

Conclusion

The use of pMRI as an adjunct to ultrasound in the assessment of complicated monochorionic pregnancies resulted in a change to prognostic counselling in 5 out of 33 pregnancies in this study. Evaluation of incorporation of pMRI into routine surveillance of complicated MCGs at specific and appropriate points in gestation is needed to inform practice.

REFERENCES

1. van Heteren CF, Nijhuis JG, Semmekrot BA, *et al.* Risk for surviving twin after fetal death of co-twin in twin – twin transfusion syndrome. *Obstet Gynecol* 1998; 92:215–219.
2. Grether JK, Nelson KB, Cummins SK. Twinning and cerebral palsy: experience in four northern California counties, births 1983 through 1985. *Pediatrics* 1993; 92:854–858.
3. Pharoah PO, Adi Y. Consequences of in-utero death in a twin pregnancy. *Lancet* 2000; 355:1597–1602.
4. Senat MV, Deprest J, Boulvain M, *et al.* Endoscopic laser surgery versus serial amnioreduction for severe twin-to-twin transfusion syndrome. *N Engl J Med.* 2004; 351:136–44.
5. Lopriore E, Wezel-Meijler G, Middeldorp JM, *et al.* Incidence, origin, and character of cerebral injury in twin-to-twin transfusion syndrome treated with fetoscopic laser surgery. *Am J Obstet Gynecol* 2006; 194:1215–1220.
6. Spruijt M, Steggerda S, Rath M, *et al.* Cerebral injury in twin – twin transfusion syndrome treated with fetoscopic laser surgery. *Obstet Gynecol* 2012; 120:15–20.
7. van Klink JM, Koopman HM, van Zwet EW, *et al.* Cerebral injury and neurodevelopmental impairment after amnioreduction

- versus laser surgery in twin – twin transfusion syndrome: a systematic review and meta-analysis. *Fetal Diagn Ther* 2013; 33:81–89.
8. Simpson LL. Twin – twin transfusion syndrome. *Am J Obstet Gynecol* 2013; 208:3–18.
 9. Graef C, Ellenrieder B, Hecher K, *et al.* Long-term neurodevelopmental outcome of 167 children after intrauterine laser treatment for severe twin – twin transfusion syndrome. *Am J Obstet Gynecol* 2006; 194:303–308.
 10. Hack KE, Koopman-Esseboom C, Derks JB. Long-term neurodevelopmental outcome of monochorionic and matched dichorionic twins. *PLoS One* 2009; 4: e6815.
 11. Lopriore E, Middeldorp JM, Sueters M. Long-term neurodevelopmental outcome in twin-to-twin transfusion syndrome treated with fetoscopic laser surgery. *Am J Obstet Gynecol* 2007; 196:231–234.
 12. van Klink JM, Koopman HM, Oepkes D. Long-term neurodevelopmental outcome in monochorionic twins after fetal therapy. *Early Hum Dev* 2011; 87:601–606.
 13. Salomon LJ, Ortqvist L, Aegerter P. Long-term developmental follow-up of infants who participated in a randomized clinical trial of amniocentesis vs laser photocoagulation for the treatment of twin-to-twin transfusion syndrome. *Am J Obstet Gynecol*

2010; 203:444–447.

14. Merhar SL, Kline-Fath BM, Meinzen-Derr J, *et al.* Fetal and postnatal brain MRI in premature infants with twin – twin transfusion syndrome. *Journal of Perinatology* 2013; 33:112-118 (doi: 10.1038/jp.2012.87).
15. Kline-Fath BM, Calvo-Garcia MA, O'Hara SM, *et al.* Twin – twin transfusion syndrome: cerebral ischemia is not the only fetal MR imaging finding. *Pediatr Radiol* 2007; 37:47-56.
16. Jelin AC, Norton ME, Barth A, *et al.* Intracranial magnetic resonance imaging findings in the surviving fetus after spontaneous monochorionic co-twin demise. *Am J Obstet Gynecol* 2008; 199:398.e1-398.
17. Jatzko B, Rittenschober-Böhm J, Mailath-Pokorny M, *et al.* Cerebral Lesions at Fetal Magnetic Resonance Imaging and Neurologic Outcome After Single Fetal Death in Monochorionic Twins. *Twin Res Hum Genet.* 2015; 18:606-12. (doi: 10.1017/thg.2015.49).
18. Griffiths PD, Sharrack S, Chan KL, *et al.* Fetal brain injury in survivors of twin pregnancies complicated by demise of one twin as assessed by in utero MR imaging. *Prenat Diagn.* 2015; 35:583-91 (doi: 10.1002/pd.4577).

19. Goergen SK, Batt M, Teoh M, *et al.* Structural abnormalities of the fetal brain in complicated monochorionic twin gestations: What does MRI add? Ultrasound Meets Magnetic Resonance, oral presentation abstract, Rome, April 2015.
20. Glenn OA, Norton ME, Goldstein RB, *et al.* Prenatal Diagnosis of Polymicrogyria by Fetal Magnetic Resonance Imaging in Monochorionic Cotwin Death. *J Ultrasound Med.* 2005; 24:711-716.
21. Righini A, Salmona S, Bianchini E, *et al.* Prenatal magnetic resonance imaging evaluation of ischemic brain lesions in the survivors of monochorionic twin pregnancies: report of 3 cases. *J Comput Assist Tomogr.* 2004; 28:87-92.
22. Righini A, Kustermann A, Parazzini C, *et al.* Diffusion-weighted magnetic resonance imaging of acute hypoxic-ischemic cerebral lesions in the survivor of a monochorionic twin pregnancy: case report. *Ultrasound Obstet Gynecol.* 2007; 29:453–456 (doi: 10.1002/uog.3967).
23. Hoffmann C, Weisz B, Yinon Y, *et al.* Diffusion MRI findings in monochorionic twin pregnancies after intrauterine fetal death. *Am J Neuroradiol.* 2013; 34:212-6.
24. Weisz B, Hoffmann C, Ben-Baruch S, *et al.* Early detection by diffusion-weighted sequence magnetic resonance imaging of severe brain lesions after fetoscopic laser coagulation for twin – twin transfusion syndrome. *Ultrasound Obstet Gynecol.*

- 2014; 44:44-9 (doi: 10.1002/uog.132830).
25. Quintero RA, Morales WJ, Allen MH, *et al.* Staging of twin – twin transfusion syndrome. *J Perinatol.* 1999; 19:550-555.
26. Gratacós E, Lewi L, Muñoz B, *et al.* A classification system for selective intrauterine growth restriction in monochorionic pregnancies according to umbilical artery Doppler flow in the smaller twin. *Ultrasound Obstet Gynecol.* 2007; 30:28-34.
27. Parazzini C, Righini A, Rustico M, *et al.* Prenatal magnetic resonance imaging: brain normal linear biometric values below 24 gestational weeks. *Neuroradiology.* 2008; 50: 877-83 (doi: 10.1007/s00234-008-0421-7).
28. Tilea B, Alberti C, Adamsbaum C, *et al.* Cerebral biometry in fetal magnetic resonance imaging: new reference data. *Ultrasound Obstet Gynecol.* 2009; 33: 173-81 (doi: 1002/uog.6276).
29. Chitty LS, Altman DG, Henderson A, Campbell S. Charts of fetal size: 2. Head measurements. *Br J Obstet Gynaecol.* 1994; 101:35-43.
30. Chitty LS, Altman DG, Henderson A, Campbell S. Charts of fetal size: 3. Abdominal measurements. *Br J Obstet Gynecol.* 1994; 101:125-131.
31. Chitty LS, Altman DG. Charts of fetal size: limb bones. *Br J Obstet Gynaecol.* 2002; 109:919-929.

32. Hadlock FP, Harrist RB, Martinez-Poyer J. In utero analysis of fetal growth: a sonographic weight standard. *Radiol.* 1991; 181:129-133.
33. Paladini D for ISUOG Education Committee. Sonographic examination of the fetal central nervous system: guidelines for performing the 'basic examination' and the 'fetal neurosonogram'. *Ultrasound Obstet Gynecol.* 2007; 29:109-116.
34. Snijders RJM, Nicolaides KH. Fetal biometry at 14-40 weeks gestation. *Ultrasound Obstet Gynecol.* 1994; 4:34-48.
35. Quarello E, Molho M, Ville Y. Incidence, mechanisms, and patterns of fetal cerebral lesions in twin-to-twin transfusion syndrome. *J Mat-Fetal and Neonatal Med.* 2007; 20:589-597.
36. Tarui T, Khwaja OS, Estroff JA, *et al.* Altered fetal cerebral and cerebellar development in twin – twin transfusion syndrome. *Am J Neuroradiol.* 2012; 33:1121-6 (doi: 10.3174/ajnr.A2922).
37. Khalil A, Rodgers M, Baschat A, *et al.* ISUOG Practice Guidelines: role of ultrasound in twin pregnancy. *Ultrasound Obstet Gynecol.* 2016; 47:247–263 (doi:10.1002/uog.15821).
38. Slaghekke F, Kist WJ, Oepkes D, *et al.* Twin anemia-polycythemia sequence: diagnostic criteria, classification, perinatal management and outcome. *Fetal Diagn Ther.* 2010; 27:181-90 (doi: 10.1159/000304512).

PD_5059_F2a.jpg

ript

PD_5059_F2b.jpg

Aut

PD_5059_F2c.jpg

PD_5059_F3ab.jpg

PD_5059_F4ab.jpeg

PD_5059_F4c.jpg

PD_5059_F4d.jpg

PD_5059_F5abcd.jpeg

PD_5059_F5e f.jpg

PD_5059_F6abc.jpeg

PD_5059_F7.jpg

PD_5059_F8ab.jpg

PD_5059_F8c.jpg

PD_5059_F9ab.jpg

PD_5059_F9cde.jpg

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Robinson, A;Teoh, M;Edwards, A;Fahey, M;Goergen, S

Title:

Fetal brain injury in complicated monochorionic pregnancies: diagnostic yield of prenatal MRI following surveillance ultrasound and influence on prognostic counselling

Date:

2017-06-01

Citation:

Robinson, A., Teoh, M., Edwards, A., Fahey, M. & Goergen, S. (2017). Fetal brain injury in complicated monochorionic pregnancies: diagnostic yield of prenatal MRI following surveillance ultrasound and influence on prognostic counselling. *PRENATAL DIAGNOSIS*, 37 (6), pp.611-627. <https://doi.org/10.1002/pd.5059>.

Persistent Link:

<http://hdl.handle.net/11343/292927>