Title Page

The measurement properties of the Spence Children's Anxiety Scale- Parent version in a large international pooled sample of young people with Autism Spectrum Disorder

Word Count (manuscript/ text)

5,301 words

Final Author Names Listed (in order of authorship with and without qualifications)

MAGIATI, Iliana^{1*}, Ph.D., D.Clin.Psy; LERH, Jian Wei^{1*}, BSocSci; HOLLOCKS, Matthew J.^{2,3}, Ph.D.; ULJAREVIC, Mirko^{4,5}, Ph.D.; RODGERS, Jacqui⁶, Ph.D.; McCONACHIE, Helen⁷, Ph.D.; OZSIVADJIAN, Ann⁸, D.Clin.Psy; SOUTH, Mikle⁹, Ph.D.; VAN HECKE, Amy¹⁰, Ph.D.; HARDAN, Antonio¹¹, M.D.; LIBOVE, Robin¹¹, B.Sc.; LEEKAM, Susan¹², Ph.D.; & SIMONOFF, Emily, MD, FRCPsych

MAGIATI, Iliana^{1*}; LERH, Jian Wei^{1*}; HOLLOCKS, Matthew J.^{2,3}; ULJAREVIC, Mirko^{4,5}; RODGERS, Jacqui⁶; McCONACHIE, Helen⁷; OZSIVADJIAN, Ann⁸; SOUTH, Mikle⁹; VAN HECKE, Amy¹⁰; HARDAN, Antonio¹¹; LIBOVE, Robin¹¹; LEEKAM, Susan¹²; & SIMONOFF, Emily

(underlined are the authors' last names)

****These two authors have contributed substantially and equally to the conceptualization, development, analyses and writing up of this paper and are both joint first authors.***

Authors' Professional Affiliations

¹ Department of Psychology, National University of Singapore, 9 Arts Link, Singapore 117570, Singapore;

² Department of Child and Adolescent Psychiatry, Institute of Psychiatry, Psychology & Neuroscience, King's College London, DeCrespigny Park, Denmark Hill, London, SE5 8AF, UK;

³ Department of Clinical Psychology, Faculty of Medicine and Health Sciences, Norwich Medical School, University of East Anglia, Norwich, UK;

⁴ Olga Tennison Autism Research Centre, School of Psychological Science, La Trobe University, Victoria 3086, Australia.

⁵ Cooperative Research Centre for Living with Autism (Autism CRC), Long Pocket, Brisbane, Queensland, Australia

⁶Clinical Psychology, Ridley Building, Institute of Neuroscience, Newcastle University, Newcastle, NE1 7RU, UK;

⁷Institute of Health and Society, Newcastle University, Sir James Spence Institute, Royal Victoria Infirmary, Queen Victoria Road, Newcastle upon Tyne; NE1 4LP, UK

⁸Evelina London Children's Hospital, St Thomas' Hospital, London, UK;

⁹Department of Psychology, College of Family, Home and Social Sciences, Brigham Young University, Provo, USA;

¹⁰Department of Psychology, College of Arts and Sciences, Marquette University, Cramer Hall, Milwaukee, WI 53201-1881, USA

¹¹Department of Psychiatry and Behavioral Sciences, Stanford University School of Medicine, Stanford University, Stanford, CA 94305, USA

¹²Wales Autism Research Centre, School of Psychology, Cardiff University, Cardiff, CF10 3AT, UK;

Corresponding Author's Address

Dr. Iliana Magiati, Department of Psychology, National University of Singapore, 9 Arts Link, #02-24, Block AS4. Singapore 117570.

Tel. +65 65161143.

Email: psyim@nus.edu.sg

Running title: Measurement properties of the SCAS-P in ASD

Number of manuscript/ text pages: 24

Number of tables: 9

Number of figures: 1

Grant Information: see Acknowledgements for a breakdown of funding for all studies in this pooled international dataset.

The measurement properties of the Spence Children's Anxiety Scale- Parent version in a large international pooled sample of young people with Autism Spectrum Disorder

Lay Abstract

Young people with ASD are often affected by anxiety. Assessing and measuring anxiety in ASD reliably and accurately is challenging, as research has yet to identify which existing anxiety measures are most useful and relevant when used with youth with ASD. The present study examined the measurement properties and factor structure of the Spence Children's Anxiety Scale-Parent Version (SCAS-P) in a large international pooled sample of youth with ASD. Data from 870 participants from 12 studies in the UK, USA and Singapore were pooled. The accuracy of the existing SCAS-P full scale and its subscales was as good as that reported in typically developing children. The subscale measuring fear of animals, insects, environment and doctors, however, had poor accuracy. Thirty items out of the existing 38 SCAS-P items were identified to measure anxiety more accurately than the existing SCAS-P, but the way these items were organized together into subtypes was different to the anxiety subtypes commonly found in typically developing children. Moreover, these subscales were not consistent in one half compared to the other half of this large sample. The limitations of the present study, the use of SCAS-P to screen for anxiety problems in ASD and future research directions are discussed.

Scientific Abstract

Anxiety-related difficulties are common in ASD, but measuring anxiety reliably and validly is challenging. Despite an increasing number of studies, there is no clear agreement on which existing anxiety measure is more psychometrically sound and what is the factor structure of anxiety in ASD. The present study examined the internal consistency, convergent, divergent and discriminant validity, as well as the factor structure of the Spence Children's Anxiety Scale-Parent Version (SCAS-P), in a large international pooled sample of 870 caregivers of youth with ASD from 12 studies in the UK, USA and Singapore who completed the SCAS-P. Most were community recruited, while the majority had at least one measure of ASD symptomatology and either cognitive or adaptive functioning measures completed. Existing SCAS-P total scale and subscales had excellent internal consistency and good convergent, divergent and discriminant validity similar to or better than SCAS-P properties reported in typically developing children, except for the poorer internal consistency of the physical injury subscale. Confirmatory Factor Analysis (CFA) of the existing SCAS-P six-correlated factor structure was a poor fit for this pooled database. Principal component analysis using half of the pooled sample identified a 30-item five correlated factor structure, but a CFA of this PCA-derived structure in the second half of this pooled sample revealed a poor fit, although the PCA-derived SCAS-P scale and subscales had stronger validity and better internal consistency than the original SCAS-P. The study's limitations, the use of the SCAS-P to screen for DSM-derived anxiety problems in ASD and future research directions are discussed.

Key Words: autism spectrum disorder, anxiety, parent report, measurement, assessment, reliability, validity, factor structure.

The measurement properties of the Spence Children's Anxiety Scale- Parent version in a large international pooled sample of young people with Autism Spectrum Disorder

Introduction

Individuals with ASD have significantly higher rates of clinically elevated anxiety symptoms (10-84%) or diagnosed anxiety disorders (about 40%) compared to individuals without ASD or with other conditions (van Steensel et al., 2011; White et al., 2009). Anxiety-related difficulties can significantly interfere with and negatively impact development, functioning and quality of life in ASD (Davis III et al., 2014; Pellecchia et al., 2016). However, the identification and measurement of anxiety in ASD is often complex and challenging.

Emerging evidence suggests people with ASD experience both “typical” anxiety (i.e. worries about separation, achievement or social evaluation, common specific phobias) as well as more idiosyncratic anxiety presentations more specifically relating to ASD (i.e., worries about change, specific sensory related fears; social anxiety without fear of negative evaluation; see Ozsivadjian et al., 2012; Kerns et al., 2014; Trembath et al., 2012; Rodgers et al., 2016). Furthermore, there are often significant difficulties in disentangling anxiety from ASD symptoms (Lecavalier et al., 2014). For instance, worries about change may be part of generalized worry or a feature of restricted, repetitive behaviors (Kerns & Kendall, 2012). Nevertheless, a small number of factor analytic studies have been able to distinguish anxiety from ASD (i.e., White et al., 2012; Renno & Wood, 2013), suggesting that these *can* likely be disentangled both statistically and clinically (see also Kerns et al., 2016).

Children with ASD also vary considerably in their intellectual and verbal abilities, insight, emotional understanding and expression, and physiological symptoms (Cook et al., 2013; Didehbani et al., 2012; Mazefsky et al., 2011; Ozsivadjian et al., 2012), which can impact their ability to rate or describe their emotional states (Grondhuis & Aman, 2012; Lecavalier et al., 2014). This often necessitates obtaining multi-informant reports from caregivers or significant others. Although anxiety rating agreement between caregivers and children with ASD has ranged from poor (i.e. Kaat & Lecavalier, 2015; Kerns et al., 2015; Renno & Wood, 2013) to moderately good (i.e. Magiati et al., 2014; Blakeley-Smith et al., 2012; Ozsivadjian et al., 2014; van Steensel et al., 2012), Storch et al. (2012) found that parental reports contributed significantly to the clinical diagnosis of anxiety. Thus, parent report remains an important way of assessing anxiety in youth with ASD.

Another current consideration is whether adapted or modified anxiety measures are required, or whether existing measures developed for typically developing children have adequate reliability and validity for youth with ASD (Kerns & Kendall, 2012; Ollendick & White, 2012). There is, to date, no anxiety-specific informant measure developed specifically for ASD¹, with the exception of the Anxiety Scale for Children – Autism Spectrum Disorder (ASC-ASD; Rodgers et al., 2016) which showed promising psychometric properties. A small, but growing, number of studies have examined the psychometric properties of existing caregiver-reported anxiety measures when used with 7-18 year old individuals with ASD and

¹ A small number of measures of broader psychopathology, including but not specifically focusing on anxiety, have been developed and tested in people with ASD. A modified structured caregiver interview piloted with children with ASD (Autism Comorbidity Interview-Present and Lifetime; ACI-PL; Leyfer et al., 2006) is currently not available for researchers or clinicians to use. The ASD-Comorbid for Children (Matson & Wilkins, 2008) and the Baby and Infant Scale for children with Autistic Traits (BISCUIT; Matson et al., 2009) are informant rating scales with evidence of satisfactory psychometric properties, but which assess only a small number of anxiety symptoms as part of mixed worry/ depressed or anxiety/ repetitive behaviour subscales or which have a very limited age range.

verbal or cognitive functioning standard scores above 60 (see Table 1 for summary of caregiver reported measures; see also reviews by Lecavalier et al., 2014 and Wigham & McConachie, 2014). The parent measures examined tended to have satisfactory to excellent internal consistencies, and superior convergent and divergent validity than child self-reports. Few studies have examined alternative clinical cut-offs or have identified a consistent anxiety factor structure in ASD (see Table 1; White et al., 2015).

The present study: rationale and research aims/ questions

Despite the recent availability of a promising anxiety specific measure based on ASD-relevant anxiety factors (Rodgers et al., 2016), there is still a need to further examine the factor structure of “typical” anxiety in ASD (White et al., 2015) and to identify a psychometrically sound measure to assess traditional DSM-derived anxiety symptoms in ASD in order to elucidate the common anxiety features across different populations.

The present study therefore examined the psychometric properties and factor structure of the Spence Children’s Anxiety Scale-Parent version (SCAS-P). The SCAS-P provides a total, as well as six DSM-oriented subscale, scores and covers a wide range of anxiety symptoms, including a number of common specific phobia (the most commonly reported anxiety disorder in ASD; van Steensel et al., 2011) and social anxiety items assessing fear of negative evaluation (which may help in distinguishing caregiver-reported social avoidance due to fear of negative evaluation or due to ASD-related social communication challenges). It was selected in this study as it (a) has strong psychometric properties with typically developing children from diverse backgrounds (i.e. Spence, 1999; Nauta et al., 2004; Whiteside & Brown, 2008; Wang et al., 2015; Arendt et al., 2014); (b) is designed to parallel

DSM anxiety disorders' criteria (albeit DSM-IV-TR criteria, as it was developed before the release of DSM-5 in 2013); (c) it is one of the most frequently used anxiety measures in ASD research (Grondhuis & Aman, 2012; Wigham & McConachie, 2014) and (d) it is a freely available informant measure officially translated in more than 20 languages, making it cost-effective and easily accessible internationally in resource-limited clinical and research settings.

In terms of its measurement properties, Zainal and colleagues (2014) reported that the SCAS-P full scale and subscales had acceptable to good internal consistency ($\alpha = .60$ to $.88$), and satisfactory sensitivity ($.75$) and specificity ($.71$) against the K-SADS clinical interview using a cut-off score of one standard deviation above the SCAS-P normative mean (Nauta et al., 2004) in a preliminary study of 32 youth with ASD. It has shown good preliminary evidence of moderate to good parent-child agreement ($ICC = .59$ to $.69$; Magiati et al., 2014; Ozsivadjian et al., 2014), although May and colleagues (2015) found poor parent-child agreement in a younger sample of cognitively able children with ASD. To our knowledge, no study has yet examined the measurement properties of the SCAS-P for use in children with ASD with a sufficiently large sample, nor has its factor structure been explored in ASD. The present study therefore aimed to investigate the psychometric properties (internal consistency, convergent, divergent and discriminant validity) and factor structure of the SCAS-P in a large, international, multi-site pooled sample of children with ASD.

Methods

Participants

Data were pooled from 12 different studies from the UK (N=9 studies; 465 participants), Singapore (N=1 study; 241 participants) and the USA (N=2 studies; 164 participants; see Table 2 for participant characteristics and details of original studies).

Caregivers of 870 children (763 males, 87.7%) aged 5.58- 18.67 years old ($M = 11.6$, $SD = 2.77$) participated. Most of the informants were mothers (75%; Table 2). All 870 participants had a professional clinical diagnosis of Autism or ASD (727; 83.7%), Asperger's (111; 12.8%) or Pervasive Developmental Disorder-Not Otherwise Specified (32; 3.7%). In all three countries, community professional diagnoses were based on national guidelines for ASD diagnosis involving the use of DSM-IV-TR or ICD-10 diagnostic criteria and a number of validated diagnostic assessments, including semi-structured caregiver interviews and child observations by qualified professionals (see Table 2). In addition, a number of studies also included a measure of autism symptom severity ($n=479$ participants; 55%; see Measures), of which 393 / 82.3% scored above clinical cut off on the respective caregiver-reported ASD measure or the clinician-rated Autism Diagnostic Observation Schedule (ADOS; Lord et al., 2001²).

Twenty-one participants (2.4%) from Study 3, who had a caregiver-reported co-

² 130 of 130 (100%) met SRS cut-off (Constantino & Gruber, 2005) in studies 6, 11 and 12; 102 of 122 (83.6%) met SCQ cut-off (Rutter et al., 2003) in studies 1, 9 and 10; 162 of 238 (68.1%) met the DBC-ASA cut-off of 14 in Study 7 (Steinhausen & Metzke, 2004); and 118 of 124 (95.2%) met ADOS autism-spectrum cut-off (Lord et al., 2001) in studies 3, 4, 11 and 12.

occurring clinical diagnosis of anxiety disorder and who participated in Study 3 because they were seeking treatment for anxiety related concerns, constituted the clinical subsample in the present pooled database. All other participants were the “unselected” community sample, recruited from non-help seeking for anxiety settings (i.e. national or local autism research databases, special or mainstream schools, parent support groups, clinics where referral was for ASD but not for anxiety, etc.).

Measures

Anxiety. The *Spence Children's Anxiety Scale-Parent Version* (SCAS-P; Spence, 1999; Nauta et al., 2004) is a caregiver-completed DSM-IV-TR derived anxiety measure comprising 38 items rated on a 4-point scale (from 0 to 3; higher scores=more anxiety) assessing symptoms in six subscales: separation (6 items; score range 0-18), social (6 items; 0-18), generalized (6 items; 0-18), panic/agoraphobia (9 items; 0-27), physical injury/specific phobias (5 items; 0-15) and obsessive compulsive disorder (6 items; 0-18). A SCAS-P total score (range 0-114) one SD or more above the normative mean (mean 14.2; SD=9.7 in Nauta et al., 2004) is considered to be clinically elevated (Spence, personal communication, October 2012). In typically developing children, the SCAS-P has a factor invariance of six factors across age, gender and different countries, excellent convergent and divergent validity, acceptable to excellent internal consistency, and good discriminant validity between anxiety disordered and non-clinically anxious groups (Nauta et al., 2004; Whiteside & Brown, 2008; Li et al., 2011; Zhao et al., 2012). It also appears to have promising psychometric properties in youth with ASD in studies in the UK, Singapore and Australia (Zainal et al., 2014; Magiati et al., 2014; Magiati et al., 2016; Ozsivadjian et al., 2014;

Russell & Sofronoff, 2005).

Autism Symptom Severity. The caregiver or teacher-reported *Social Responsiveness Scale* (SRS; Constantino & Gruber, 2005; n=130), the 29-item *Developmental Behavior Checklist* (Einfeld & Tonge, 2002) *Autism Screening Algorithm score* (DBC-ASA; n=238)³, the *Social Communication Questionnaire* (SCQ; Rutter et al., 2003; n=122) or the *Autism Diagnostic Observation Schedule* (ADOS; Lord et al., 2001; n=124) were used to establish the number of participants scoring above cut-off scores as a way to confirm caregiver reported clinical diagnosis of ASD and to measure autism symptom severity in some of the studies (see Table 2).

IQ/ Adaptive Functioning. The *Wechsler Intelligence Scale for Children–Third Edition* (WISC-III; Wechsler, 1991; n=20), the *Wechsler Abbreviated Scale of Intelligence* (WASI; Wechsler, 1999; n=161), the *Stanford Binet Intelligence Scales–Fifth Edition* (SB5; Roid, 2003; n=54), the *Kaufman Brief Intelligence Test–Second Edition* (KBIT-2; Kaufman & Kaufman, 2005; n=43) or the *Scales of Independent Behavior–Revised Short Form* (SIB-R Short; Bruininks et al., 1996; n=239) were used as a measure of overall IQ or adaptive functioning in in some of the studies (see Table 2). All these measures have a normative mean of 100 and a standard deviation of 15.

Procedure

All studies were approved by their respective institutional ethics committees. The caregivers and youth were recruited according to the studies' approved research protocols

³ The Developmental Behavior Checklist (Einfeld & Tonge, 2002) *Autism Screening Algorithm score* (DBC-ASA) has been found to discriminate well between children with disabilities with and without ASD (Brereton et al., 2002; Steinhausen & Metzke, 2004).

and completed the measures summarized above and in Table 2.

Statistical Analytical Plan

Missing data. No participant was excluded, and there was no more than 10% missing data for any one particular measure (Bennett, 2001). All missing data were managed and replaced according to manual/ measure guidelines⁴.

Data Harmonization. Although pulling data together has many advantages in terms of increasing power, allowing the creation of larger and likely more representative of the population samples and maximizing the use of smaller-scale data (Griffith et al., 2015; Hussong et al., 2014), integration of data from different studies often employing different measures is also very challenging. Careful consideration therefore needs to be given to how the datasets can be harmonized as much as possible. Integrating data from intellectual, adaptive and autism symptom severity measures was complex, as different measures and informants were employed. To address this, and following methods discussed by Griffith and colleagues (2015), Hussong et al., (2013) and Schaap et al. (2011), we harmonized cognitive and adaptive functioning data, all of which had the same normative mean of 100 and SD of 15 by creating an ordinal “approximate level of functioning” classification variable with scores ranging from 1 (=standard score <40 corresponding to severe/ profound ID range as per DSM) to 8 (SS $\geq$ 120, corresponding to superior functioning as per the Wechsler scales’ classification)⁵. To ensure higher levels of consistency, we harmonized only caregiver-

⁴ Average caregivers’ ratings were used to tabulate the subscale and total scores for eight participants in Study 7 for whom SCAS-P data were given by both caregivers.

⁵ 8= Standard Score $\geq$ 120 (superior); 7= 110-119 (high average); 6= 90-109 (average); 5= 80-89 (low average); 4= 70-79 (borderline); 3= 55-69 (mild ID range); 2= 40-54 (moderate ID range); and 1= < 40 (severe or profound ID range); categories based on the Wechsler scales classification and DSM-5 ID ranges. All cognitive

reported autism symptom severity measures⁶ by converting the raw scores from the different autism screening measures to a 0 to 1 metric (dividing the total raw score of each participant by each scale's maximum score).

Statistical Analyses. To examine their fit in our ASD sample, a series of SCAS-P factor models identified in the literature (see Results) were constructed in CFA models within a general structural equation model framework with a robust maximum likelihood (MLR) estimator in MPLUS v5 (Muthén & Muthén, 2012) to generate model fit indices. The MLR estimator was used due to non-normally distributed data. The *Satorra-Bentler* Scaled Chi-Square test was used to compare models to account for the scaling due to the use of the robust estimator. As the Chi-Square statistic tends to underestimate goodness of fit in large sample sizes (Bollen, 1989), Hu and Bentler (1999) recommend examining multiple indices $RMSEA \leq .06$; $CFI \geq .95$; and $TLI \geq .95$ to determine the model fit.

Participants ($n=849^7$) were then randomly assigned into two groups ($n_1=425$; $n_2=424$) with similar rates of participants from the three countries. Each group was deemed sufficiently large to explore the underlying factor structure using Principal Component Analysis (PCA) and validating the identified factor structure using CFA (Norris & Lecavalier 2010; Tabachnick & Fidell, 2013; Yong & Pearce, 2013). The two subgroups were not

and adaptive measures employed in the different studies correlate very highly with each other in data presented in their respective manuals. We acknowledge that different measures assess different components of intellectual/adaptive functioning and similar, but also different, underlying constructs, which is why we did not use the actual scores, but rather the broader classification categories. This approach maximizes use of all available data, while still acknowledging that the scores obtained are highly correlated, but not directly comparable – this is the reason we did not use the continuous standard scores.

⁶ In Study 12, 43 participants had ADOS, SRS, and SCQ data; the SRS scores were used for these participants, because its scoring method provides greater score range.

⁷ The clinical subsample of 21 participants from Study 3 were not included in these analyses, as they had a known identified anxiety disorder and were recruited into the research study due to specifically seeking intervention for anxiety related difficulties.

significantly different in terms of chronological age, gender, ASD symptomatology, functioning, and SCAS-P anxiety total score ($p > .05$; all were small effect size differences, except harmonized functioning which was small-to-medium). The PCA (with direct oblimin rotation, as the factors were theoretically and empirically correlated; Spence, 1998) was run with the first group. Parallel analysis and scree plots determined the number of factors. The PCA derived model was then tested for model fit in the second subsample.

The reliability and validity of the existing SCAS-P and PCA-derived SCAS-P was examined in the full sample. Pearson's, biserial, Spearman's Rho and Cramer's V correlations examined the relationship between SCAS-P total and subscale scores with demographic variables, harmonized autism symptomatology, and functioning classification.

Welch ANOVAs with Bonferroni corrections, Chi square tests, and Games–Howell Post-Hoc tests were used to examine differences (i) among the three countries; (ii) between those who met ASD cut-off in the screening measures ($n=394$) and those who had professional diagnoses of ASD but did not meet ASD cut-off or did not have measures of autism symptomatology completed ($n=476$); and (iii) between community ($n=849$) and clinically referred for anxiety ($n=21$) participants. Effect sizes were interpreted as per Cohen (1988; e.g. $r < .30$ small; $.30-.49$ medium; $> .50$ large; $d < .49$ small; $.50-.79$ medium; large $> .80$).

Results

Preliminary analyses: differences between samples/ countries and between subsamples with and without ASD screening data

Excluding the clinically referred for anxiety subsample from Study 3 ($n=21$), the three country subsamples differed statistically significantly in chronological age, SCAS-P total score, ASD symptomatology, functioning classification, and gender rates with small to large effect size differences (Table 3). Post-hoc Games–Howell tests showed that the UK subsample had higher SCAS-P total score and ASD symptomatology compared to the US and Singaporean participants with medium-to-large effect size differences. The UK and US participants were also significantly older and higher functioning than those from Singapore with a small and large effect size difference respectively (Table 3). However, the country differences in SCAS-P total scores were no longer significant after controlling for age, gender, functioning, and ASD symptomatology ($F(2, 355)=1.91, p=.15, d=.049$).

Also excluding the clinically referred subsample, there were statistically significant differences in chronological age, SCAS-P (total, social and generalized anxiety scores only), mean functioning classification, and gender between the participants with professional diagnoses of ASD who scored above clinical cut-off in one of the different screening measures employed in the pooled studies ($n=376$) and the subsample with professional diagnoses who scored below recommended clinical cut-offs in the screening measures administered or who had no ASD symptom ratings available ($n=473$), but effect sizes of these differences were mostly small ($p<.05, .007 \leq \text{effect sizes} \leq .19$; Table 4), with the exception of functioning classification, where the participants not meeting cut-off or not having ASD

screening data available had somewhat higher functioning rankings (Table 4). The difference in SCAS-P anxiety total score between the two subsamples remained significant when age, gender, and functioning classification were controlled for, but the effect size of the difference was very small (Table 4). Because of the small effect size differences between those with and without ASD screening measure data, and as all studies had recruited participants with valid professional diagnoses following established national procedures, we proceeded with full sample analyses.

Comparison with SCAS-P norms

The pooled unselected subsample had significantly higher mean SCAS-P total and subscale scores than published norms ($p < .001$; Nauta et al., 2004; Table 5). Using the suggested cut-off score of >24 , 76.2% (16/ 21) participants in the clinical subsample as compared to 46.8% (397/ 849) from the unselected subsample scored above cut-off.

Item analyses

All the items had at least .33 corrected item-total correlations, except item 16 (“my child needs to keep checking that s/he has done things right”; $r = .13$). The average corrected item-total correlation for the 38 SCAS-P items was .52 ($SD = .11$). The average corrected item-subscale correlation was .53 ($SD = .11$) and the average corrected subscale-total correlation was .69 ($SD = .10$).

Internal Consistency

Internal consistency for the total score items was excellent at $\alpha = .93$. For the subscales, all Cronbach's alphas were $> .75$, except physical injury which was suboptimal (Table 5).

Validity

Convergent validity. The SCAS-P total and subscale scores had moderate to strong positive correlations with the DBC-anxiety subscale ($r(236) = .64$ for total; $r(236) = .32-.56$ for subscales; all $p < .001$; $n = 238$ from Study 7 only in which used the DBC).

Divergent validity. SCAS-P had non-significant or small correlations ($< .20$) with age, gender, and overall functioning classification, with the exception of a positive small-to-medium effect size relationship between age, functioning classification and social phobia, and between overall functioning and social and generalized anxiety (Table 6). ASD symptomatology had significant medium-to-large positive correlations with SCAS-P total and subscale scales, except for the physical injury subscale which had a small correlation (Table 6). Using data from Study 7 only, the SCAS-P had small-to-moderate correlations with the DBC-disruptive/antisocial subscale ($r(236) = .47$ for total; $r(236) = .23-.44$ for subscales; all $p < .001$).

Discriminant validity. The clinical subsample from Study 3 ($n = 21$) had significantly higher SCAS-P total scores, generalized and social anxiety scores than the unselected subsample (Table 7).

Confirmatory Factor analysis

Four different factor models examined by Nauta et al. (2004) in the normative SCAS-P sample were explored (Table 8). The one factor model provided a better fit for the data than the null model, but the six uncorrelated factor model did not improve the model fit. The six correlated factor model provided a better fit than six uncorrelated factors. Lastly, the 5-factor model with Generalized Anxiety as a second-order factor provided a significantly better fit than the previous models. Overall, however, the indices suggested that none of these models

provided adequate fit for our ASD sample. A PCA was thus conducted to explore the SCAS-P factor structure in our sample⁸.

Revised Factor Structure Analyses

Principal Component Analysis. PCA was first run on the 38 SCAS-P items with the randomly selected first half of the sample's participants ($n_1=425$), with direct oblimin (delta=0) rotation. The Kaiser-Meyer-Olkin test of sampling adequacy was excellent at .93 (Field, 2009). Bartlett's test of sphericity was also significant ($\chi^2(703, n=425) = 7891.76, p < .001$) indicating that the inter-item correlations were adequate for PCA. The points of inflexion on the Scree plot suggested one, three, or five components (Figure 1), while eight factors had eigenvalues >1 . Parallel analysis (O'Connor, 2000) suggested five components.

Extracting for five components and suppressing correlation coefficients $<.38$ to ensure that the selected items had minimal relation with the component, the initial five components accounted for 51.1% of the variance. As several items had communalities significantly below .40 (Costello & Osborne, 2005) or factor loadings below .38, they were removed sequentially and the PCA was re-run each time until a simple structure was achieved (Thurstone, 1947). Items 16, 23, 29, and 34 were removed sequentially because of low communality, followed by items 3 and 36 because of low factor loadings $<.38$, and lastly items 17 and 38 due to cross-loading. A simple factor structure of five components was obtained with the remaining 30 items (see Table 9) accounting for a total of 57.9% of the variance⁹.

⁸ We also ran the CFA with the participants scoring above cut-off in the autism screening measures only and the results were consistent with the CFA for the full sample, therefore we report the full sample analyses.

⁹ We first ran the PCA following the same steps and criteria only with the participants who scored above cut-off in one of the three caregiver-reported screening measures employed in the different studies and found the same 5-factor structure explaining very similar variance (54.8%), the only difference being that two more SCAS-P

The five components had significant positive moderate-to-high inter-correlations ($r \geq .38$). The first component, Social/Generalized Anxiety symptoms, accounted for 35.2% of the variance and its nine items were mainly on social anxiety, but with some generalized and separation anxiety symptoms. The second component accounted for 7.3% of the variance and contained 5 items, measuring mostly separation anxiety symptoms but also fear of darkness and feeling scared. The third component contained eight items measuring somatic or panic symptoms accounting for 6.5% of the variance. The fourth factor contained four items all assessing obsessive-compulsive symptoms and accounted for an additional 4.7% of variance. The last component was a Specific Phobia factor containing four items on medical/dental phobia and fear of public places or using public transport and accounting for 4.2% of the variance.

Item analyses and reliability of revised PCA-derived SCAS-P. The item means and SDs, corrected item-total, corrected item-subscale and corrected subscale-total correlations of the revised 30-item SCAS-P are presented in Table 5. All items had at least .34 corrected item-total correlations. The average corrected item-total correlation, item-subscale correlation and subscale-total correlations were .54 ($SD=0.09$), .59 ($SD=0.08$) and .62 ($SD=0.06$) respectively. Internal consistency for the 30-SCAS-P items was .93 and $\alpha \geq .69$ for the subscales (Table 5).

Validity of the PCA-derived SCAS-P. The PCA-derived SCAS-P total and its subscales had good convergent validity with the DBC-anxiety subscale in Study 7 (r (236) = .65 for total; .38-.55 for subscales; all $p < .001$). In the same study subsample, the revised

items were included in the final solution. Given the consistent findings, we report on the analyses for the full sample, as the strength of this pooled database lies primarily in the large, diverse sample.

SCAS-P total and subscale scores were positively correlated with the DBC-disruptive/antisocial subscale ($r(236) = .26-.46, p < .001$) with small to medium effect size correlations. In the full sample, the PCA-derived SCAS-P total and subscales scores had non-significant or significant small correlations with age, gender and functioning classification, indicating divergent validity (with the exception of small-to-medium effect size correlations between mixed generalized/social anxiety subscale and age, functioning classification, and ASD symptomatology; Table 6). The SCAS-P total and the mixed generalized/ social anxiety subscale scores were significantly higher in the clinical subsample than the unselected participants, providing some preliminary evidence of discriminant validity, although effect sizes were small (Table 7). The original SCAS-P full-scale correlated $r(868) = .99$ with the PCA-revised SCAS-P full scale. The original SCAS-P subscales were also positively and highly correlated with their corresponding revised subscales (i.e., $r = .88-.95$), except for the PCA-derived Physical Injury/ Specific Phobia subscale which was moderately correlated with the original SCAS-P Physical Injury subscale ($r(868) = .57$).

Confirmatory Factor Analysis. A CFA was then run in the other half of the pooled sample ($n_2 = 424$) to test the goodness of fit of the PCA derived model summarized above (and in Table 8). The model fit indices (CFI, TFI, RMSEA) suggested that the PCA derived model did not fit the other half of our sample adequately (Table 8).

Discussion

Measurement properties of the original 38-item SCAS-P

Overall, the original SCAS-P full scale and all but one subscales had excellent internal consistencies and convergent, divergent and discriminant validity similar to or better than SCAS-P data reported in typically developing (TD) children (Whiteside & Brown, 2008; Li et al., 2011; Nauta et al., 2004). The Physical Injury subscale had a low Cronbach's alpha of .55, consistent with previous findings (Li et al., 2011; Magiati et al., 2016; Nauta et al., 2004) suggesting that this particular subscale, although commonly endorsed in our sample, may be inconsistent psychometrically.

However, the SCAS-P total correlated .47 with the DBC antisocial/ aggressive subscale, suggesting some overlap between anxiety and externalizing behaviors in our sample. This is consistent with other qualitative and quantitative studies reporting that often anxiety in ASD may come across or be expressed as irritability or aggression (Ozsivadjian et al., 2012; Mazefsky et al., 2012; Lydon et al., 2015). The positive association of age and functioning with the SCAS-P social anxiety subscale is developmentally expected - increasing social anxiety symptoms are often observed with age in typically developing children, while studies in ASD have also shown that with increases in age and ability comes increased awareness of social isolation and difference, a risk factor for developing anxieties about social situations (i.e. Kerns et al., 2016; Sukhodolsky et al., 2008). At the same time, the original SCAS-P performed less well in the present study in terms of structural and discriminant validity as compared to other published studies with typically developing children (Nauta et al., 2004; Whiteside & Brown, 2008).

Factor structure of the original SCAS-P in ASD

The SCAS-P factor models in the existing literature (see Nauta et al., 2004) did not provide an adequate fit for our data, suggesting that the underlying conceptual and structural basis of the SCAS-P anxiety symptoms may to some extent be different in ASD. Other studies exploring factor structures of existing anxiety measures in ASD have also reported similarly poor fits - Renno & Wood (2013) and White et al. (2015) using the Multidimensional Anxiety Scale for Children-Parent version (MASC-P; March et al., 1997); and Stern et al. (2014) using the Screen for Child Anxiety Related Disorders (SCARED-P; Birmaher et al., 1997). White et al. (2015) found a mixed factor of separation anxiety and panic items and two separate social anxiety factors (i.e., evaluation and performance focused) using the MASC-P, while Stern et al. (2014) observed a mixed panic and generalized anxiety factor comprising a mixture of items from the other subscales, such as school phobia, using the SCARED-41-P. It is possible that the lack of fit of the factor structures of existing measures in different samples of participants with ASD may be explained by the fact that anxiety symptoms in ASD may be manifestations of broader and more complex underlying pathways and processes, such as emotional dysregulation, alexithymia, hyper-arousal, intolerance of uncertainty and/ or sensory sensitivities (e.g. Kerns et al., 2016; White et al., 2014; Mazefsky et al., 2012; South & Rodgers, 2017). These broader underlying ASD-related vulnerabilities may also explain the findings of “mixed” factors in the earlier and the present studies in ASD. Others, however, have reported preliminary evidence of a good fit (i.e., Hallett et al., 2013a, using the CASI-Anxiety). Of the “traditional” anxiety factors, separation and social anxiety subscales, and panic subscale (physiological symptoms), to a lesser extent, were most consistently replicated in the studies mentioned earlier.

Measurement properties of the revised PCA-derived 30-item SCAS-P

The PCA suggested an alternative five-factor SCAS-P structure, with a number of items loading on factors not commonly suggested in the existing literature (Nauta et al., 2004). For example, social and generalized anxiety items loaded together on one “mixed” factor. As discussed earlier, it is possible that, compared to normative samples, some individual SCAS-P items load onto different anxiety factors because of different, and ASD-distinct, underlying pathways explaining these fears and anxieties (White et al., 2015; Kerns et al., 2016; South & Rodgers, 2017). For example, two items relating to fears of using public toilets or being in crowds load onto the social anxiety and panic/ agoraphobia factors respectively in the normative SCAS-P. However, these items load onto the specific phobias factor in the PCA-derived SCAS-P in this study, which could be because these fears may be related to specific overwhelming sensory experiences in toilets and crowds for those with ASD, rather than to “traditional” social anxiety concerns (see Kerns, 2016 for more on the clinical implications of this in the differential diagnosis of anxiety in ASD).

Compared to the original SCAS-P, the revised PCA-derived SCAS-P had similar corrected internal consistencies (with the exception of the specific phobia subscale, which was better in the PCA-derived SCAS-P factor structure), item-total, item–subscale, and subscale-total correlations, divergent validity and discriminant validity, but somewhat stronger convergent validity, as shown by higher correlations with the DBC anxiety subscale. ASD symptomatology was less strongly associated with the PCA-derived SCAS-P subscales, with the exception of a medium association with mixed GAD/ social subscale. This could potentially mean that PCA helped remove overlapping items that tap on both anxiety and ASD severity.

The moderate similarities in content and the high positive correlations between the original SCAS-P and the corresponding revised subscales suggest that the underlying anxiety factors may generally be similar. Nevertheless, the lack of a consistent, replicable, and adequate factor structure in the two randomly selected large subsamples in the present pooled dataset is difficult to interpret. Clearly, differences in recruitment and sample characteristics may partially explain the lack of consistency, as we were able to compare the two randomly selected subsamples on only some characteristics and measures. The large heterogeneity of ASD may contribute to less consistent structures, or possibly to different underlying structures for different subgroups. Furthermore, the items to which the caregivers responded were all enquiring about typical anxiety presentations and there was no further clarification or elaborations requested (i.e. whether what parents were describing related specifically to anxiety anticipation or to ASD symptoms and associated distress more broadly; (see Kerns, 2016). It is possible that measures which will include both “traditional” as well as more ASD-distinct anxiety presentations and symptoms may derive more comprehensive, and thus more consistent, structures across different samples. Currently, however, due to the concerns with lack of consistency, it is likely premature to use the PCA-derived SCAS-P scale and factor structure for clinical or research purposes.

Comparing the measurement properties of the original/ revised SCAS-P to other existing anxiety measures in ASD

Both the original and revised SCAS-P versions and the SCARED-71-P have demonstrated promising evidence of discriminant validity in terms of significantly higher means in the clinical help-seeking for anxiety sample compared to the unselected community-recruited sample, although this needs to be confirmed with studies employing

clinical diagnostic interviews to establish sensitivity and specificity of the measure. Good sensitivity and specificity for the SCARED-41-P and SCARED-71-P has been reported in a clinically anxious help-seeking sample (Van Steensel et al., 2013), but much poorer accuracy was reported for a number of other parent and self-report anxiety measures developed for typically developing children when compared against the augmented for ASD Anxiety Disorder Interview Schedule Child/ Parent (ADIS C/P; Kerns et al. 2015; Stern et al., 2014; van Steensel et al., 2012).

Both original and revised PCA-derived SCAS-P versions and the other anxiety measures examined in the literature so far had poor structural validity, except for the CASI-Anxiety which showed promising evidence of adequate model fit (Hallett et al., 2013a; Renno & Wood, 2013; Stern et al., 2014; White et al., 2015). Overall, with regards to existing anxiety measures, both the original SCAS-P and the SCARED-71-P and CASI-Anxiety appear to be promising in their use with children and adolescents with ASD, but future research needs to examine how to improve their structural validity, sensitivity and specificity for use as a screening tool with this population. It is likely that such improvements in the structural validity will come from including and piloting in existing measures addendum subscales measuring more ASD-specific anxiety presentations (i.e., see Kerns et al., 2017; Bearss et al., 2015; Rodgers et al., 2016).

Strengths and Limitations of the present study

This large pooled ASD sample which includes participants from 12 studies from three countries made it possible to examine structural validity and factor structure of, as well as to test the adequacy of the new structure in, a large diverse group of young people with ASD. As most of the participants were recruited through community settings, this allowed the

investigation of the use of the SCAS-P in non-clinically referred young people for anxiety with ASD.

At the same time, however, pooling participants from multiple studies and countries also presented several challenges and harmonization of data from different sources and measures is inherently difficult as there was no common measure for ASD severity, cognitive or adaptive functioning. Some loss in the richness and range of data available for harmonization is inevitable (in the present study, this was the case for scores from intellectual/ adaptive functioning and autism symptomatology which were pooled together in ordinal categories, rather than used continuously). It is also possible that pooling data from 12 studies in three countries might have masked other potential differences, such as sociocultural differences, beyond the identified examined differences in age, gender, overall functioning, or ASD symptom severity. Furthermore, only some of the pooled studies had additional child report anxiety measures or clinician diagnostic interviews available, thus we were not able to investigate informant agreement nor to establish which participants met clinician-rated diagnostic criteria for an anxiety disorder. Lastly, although participants across all levels of intellectual/ verbal/ adaptive functioning were included, our pooled sample comprised more children and adolescents with ASD who were functioning within the non-impaired range in terms of cognitive/ adaptive skills.

Possible implications

For Clinical Practice. Following further replication, the SCAS-P appears to be a reasonably good choice as a freely available first line screening tool for typical DSM-derived anxiety in ASD, provided that it is employed alongside other multi-informant and multi-

method information gathering approaches (i.e. using the ASC-ASD by Rodgers et al., 2016 to also examine more ASD-related anxiety difficulties; or the ADIS C/P ASD addendum clinical interview by Kerns and colleagues, 2014; see also Vasa et al., 2016). If the original SCAS-P is used, we recommend the use of the total score as a general screen for anxiety, and not of the separate subscales, as the underlying subscale/ factor structure remains inconsistent and unclear and requires further exploration. We specifically caution against the use of the original SCAS-P Physical Injury subscale, as some of its items may not be tapping on a latent specific phobia factor. Furthermore, clinicians should not simply invite caregivers to “tick” symptoms off the checklist; instead, we encourage them to explore the informants’ responses further, in order to establish the precise content of anxiety concerns and to elucidate whether they may be best explained or understood as anxiety involving anticipation or as a distressed reaction as a consequence of their ASD symptoms (e.g., Kerns & Kendall, 2012; see Kerns et al., 2014; 2016 for evidence and recommendations differentiating ASD and anxiety symptomatology using systematic interviewing procedures).

For Future Research. Several unanswered questions remain. In our summary in Table 1, it appears that measures developed for typically developing youth generally do not have strong evidence for discriminant validity, sensitivity or specificity, or good factor structure fits when employed in youth with ASD. CFAs often reveal that the measures’ original structures have inadequate fits in ASD samples, although studies using EFAs have identified relatively similar factor structures consistently using different measures (i.e., Hallett et al., 2013a; Renno & Wood, 2013; Stern et al., 2014; White et al., 2015; this study).

Future research should thus look into (i) modifying, adapting or enhancing existing measures for individuals with ASD, while gathering normative ASD data; and/ or (ii)

developing and improving ASD-specific anxiety measures (Kerns et al., 2015; Sterling et al., 2015; White et al., 2015; Rodgers et al., 2016). Children's cognitive profile and verbal skills also appear to affect anxiety symptom endorsement to some extent (Hallett et al., 2013b; Witwer & Lecavalier, 2007) and hence, it will be important to validate the measures, establish norms and cut-offs separately for ASD youths of different levels of ability.

Furthermore, youth with ASD also present with more idiosyncratic ASD-related anxiety (see Bearss et al., 2015; Kerns et al., 2014; Ozsivadjian et al., 2012; for a review, see Magiati et al., 2017). Future research should examine how ASD-related anxiety symptoms relate to traditional DSM-derived anxiety symptoms in young people with ASD (Kerns et al., 2014) and the underlying structural validity of measures including both traditional and ASD-related anxiety symptoms. Researchers can use existing work with focus group methodologies which have generated candidate anxiety items to develop an ASD-specific anxiety measure (Bearss et al., 2015; Rodgers et al., 2016) or augment modified measures of traditional DSM-defined anxiety symptoms with an ASD-addendum subscale (Ozsivadjian et al., 2014; Kerns et al., 2014).

Research efforts to better assess, understand and treat anxiety in children and adolescents with ASD have intensified and have considerably improved our understanding of anxiety in ASD. The present study contributes to this growing body of literature by examining a widely used and easily accessible caregiver-report anxiety measure. Future research efforts should focus on improving our understanding of the structure of, and relationship between, traditional and more ASD-related presentations of anxiety and our ability to reliably and validly identify these in individuals with ASD.

Acknowledgements

We would like to thank all the caregivers and families, schools and clinics, participants and staff involved in all studies and countries. Not in any particular order, Study 1 includes data from independent research commissioned by the UK National Institute for Health Research (NIHR) under the Research for Patient Benefit programme (PB-PG-0408-16069) and data collected by Dr. Janes and Dr. Connolly as part of doctoral theses for the qualification of Doctor of Clinical Psychology. Studies 2 and 5 are grateful to the Database of Children with ASD Living in the North East (www.daslne.org) for assistance with recruitment. Daslne is funded by the UK autism research charity Autistica and by The Children's Foundation. Studies 4 & 5 were supported by funding provided by Northumbria, Tyne and Wear NHS Foundation Trust awarded to Dr. Jacqui Rodgers. Study 3 was commissioned by the National Institute for Health Research (NIHR) under the Research for Patient Benefit programme (PB-PG-0408-16069). Study 6 was supported through a PhD studentship awarded to Dr. N. Darus by the Ministry of Health in Malaysia. Study 7 was supported by a Start-up grant awarded to Dr. I. Magiati from the National University of Singapore, Faculty of Arts and Social Sciences. Study 8 was funded by the NIHR Biomedical Research Centre (BRC) in Mental Health (grant code: PCCKASA) and the South London and Maudsley Charitable Funds (awarded to Prof. E. Simonoff). Study 9 was supported by the Charles Hawkins Fund. Study 10 was supported by a PhD studentship from the Wales Office of Research and Development for Health and Social Care, National Institute for Social Care and Health Research, awarded to Dr. M. Uljarevic and a project grant from the Baily Thomas Charitable Fund awarded to Prof. S. Leekam. The data collection was carried out at Cardiff University by Dr. M. Uljarevic and by Dr. J. Lidstone. Study 11 was supported by grants from Simons Foundation Autism Research Initiative, the Mosbacher Family Fund for Autism Research, The Escher Fund at the Silicon Valley Community Foundation, and Stanford University's Child Health Research Institute. Study 12 was supported by grants from the Autism Society of Southeastern Wisconsin and the National Center for Advancing Translational Sciences, National Institutes of Health (NIH), through grant number 8KL2TR000056 to Dr. A. Van Hecke. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIH, NHS, NIHR or Department of Health. The authors report no conflicts of interest.

References

- American Psychiatric Association (APA; 2013). Diagnostic and statistical manual of mental disorders (5th ed.) (DSM-5). Washington, DC: Author.
- Arendt, K., Hougaard, E., & Thastum, M. (2014). Psychometric properties of the child and parent versions of Spence Children's Anxiety Scale in a Danish community and clinical sample. *Journal of anxiety disorders*, 28(8), 947-956.
- Bearss, K., Taylor, C.A., Aman, M.G., Whittemore, R., Lecavalier, L., Miller, J., et al. (2015). Using qualitative methods to guide scale development for anxiety in youth with autism spectrum disorder. *Autism*. Published online before print. doi:10.1177/1362361315601012
- Bennett, D. A. (2001). How can I deal with missing data in my study? *Australian and New Zealand Journal of Public Health*, 25(5), 464-469.
- Birmaher, B., Khetarpal, S., Brent, D., Cully, M., Balach, L., Kaufman, J., et al. (1997). The screen for child anxiety related emotional disorders (SCARED): Scale construction and psychometric characteristics. *Journal of the American Academy of Child & Adolescent Psychiatry*, 36(4), 545-553.
- Blakeley-Smith, A., Reaven, J., Ridge, K., & Hepburn, S. (2012). Parent-child agreement of anxiety symptoms in youth with autism spectrum disorders. *Research in Autism Spectrum Disorders*, 6(2), 707-716.
- Brereton, A. V., Tonge, B. J., Mackinnon, A. J., & Einfeld, S. L. (2002). Screening young

- people for autism with the developmental behavior checklist. *Journal of the American Academy of Child & Adolescent Psychiatry*, 41(11), 1369-1375.
- Bollen, K. A. (1989). *Structural equations with latent variables*. New York: Wiley.
- Boulter C., Freeston, M., South, M. & Rodgers, J. (2014) Intolerance of Uncertainty as a Framework for Understanding Anxiety in Children and Adolescents with Autism Spectrum Disorders. *Journal of Autism and Developmental Disorders*, 44(6), 1391–1402.
- Bruininks, R. H., Woodcock, R. W., Weatherman, R. F., & Hill, B. K. (1996). *SIB-R: Scales of independent behavior-Revised*. Itasca, IL: Riverside Publishing.
- Chamberlain, P.D., Rodgers, J., Crowley, M.J., White, S.E., Freeston, M.H. & South, M. (2013). A potentiated startle study of uncertainty and contextual anxiety in adolescents diagnosed with Autism Spectrum Disorder. *Molecular Autism*, 4(1), 31-41.
- Chorpita, B. F., Yim, L., Moffitt, C., Umemoto, L. A., & Francis, S. E. (2000). Assessment of symptoms of DSM-IV anxiety and depression in children: A revised child anxiety and depression scale. *Behaviour Research and Therapy*, 38(8), 835-855.
- Constantino, J. N., & Gruber, C. P. (2005). *The social responsiveness scale manual*. Los Angeles, CA: Western Psychological Services.
- Cook, R., Brewer, R., Shah, P., & Bird, G. (2013). Alexithymia, not autism, predicts poor recognition of emotional facial expressions. *Psychological science*, 24(5), 723-732.
- Costello, A. B., & Osborne, J. W. (2005). Best practices in exploratory factor analysis: four

recommendations for getting the most from your analysis. Practical Assessment

Research & Evaluation. Retrieved from <http://pareonline.net/pdf/v10n7.pdf>

Darus, N. (2016). *Anxiety in Autism Spectrum Disorder (ASD): The Influence of Executive and Sensory Processing Dysfunctions*. Unpublished Ph.D. thesis, Newcastle University, UK.

Davis III, T. E., White, S. W., & Ollendick, T. H. (2014). *Handbook of Autism and Anxiety*. Springer.

Didehbani, N., Shad, M. U., Kandalaft, M. R., Allen, T. T., Tamminga, C. A., Krawczyk, D. C., et al. (2012). Brief report: Insight into illness and social attributional style in Asperger's syndrome. *Journal of autism and developmental disorders*, 42(12), 2754-2760.

Einfeld, S. L., & Tonge, B. J. (2002). *Manual for the Developmental Behaviour Checklist (2nd ed.)*. Melbourne and Sydney: Monash University Centre for Developmental Psychiatry and Psychology, and School of Psychiatry, University of South Wales.

Field, A. (2009). *Discovering statistics using SPSS (3rd eds.)*. Sage publications.

Fletcher-Watson, S., Leekam, S.R., Connolly, B., Collis, J.M., Findlay, J.M., McConachie, H., et al. (2012). Attenuation of change blindness in children with Autism Spectrum Disorders. *British Journal of Developmental Psychology*, 30(3), 446-458.

Griffith, L.E., van den Heuvel, E., Fortier, I., Sohel, N., Hofer, S.M., et al. (2015). Statistical approaches to harmonize data on cognitive measures in systematic reviews are rarely reported. *Journal of Clinical Epidemiology*, 68(2), 154-162.

Grondhuis, S. N., & Aman, M. G. (2012). Assessment of anxiety in children and adolescents with autism spectrum disorders. *Research in Autism Spectrum Disorders*, 6(4), 1345-1365.

Hallett, V., Ronald, A., Colvert, E., Ames, C., Woodhouse, E., Lietz, S., et al. (2013b). Exploring anxiety symptoms in a large-scale twin study of children with autism spectrum disorders, their co-twins and controls. *Journal of Child Psychology and Psychiatry*, 54(11), 1176-1185.

Hallett, V., Lecavalier, L., Sukhodolsky, D. G., Cipriano, N., Aman, M. G., McCracken, J. T., et al. (2013a). Exploring the manifestations of anxiety in children with autism spectrum disorders. *Journal of autism and developmental disorders*, 43(10), 2341-2352.

Hollocks, M.J., Howlin, P., Papadoupoulos, A., Khondoker, M., & Simonoff, E. (2014). Differences in HPA-axis and Heart Rate Responsiveness to Psychosocial Stress in Children with Autism Spectrum Disorders with and without co-morbid anxiety. *Psychoneuroendocrinology*, 46, 32-45.

Hollocks, M.J., Ozsivadjian, A., Matthews, C.E., Howlin, P., & Simonoff, E. (2013). The relationship between attentional bias and anxiety in children and adolescents with autism spectrum disorders. *Autism Research* 6(4):237-247.

Hu, L., & Bentler, P. (1999). Cutoff criteria for fit indices in covariance structure analyses: Conventional criteria versus new alternatives. *Structural Equation Modeling*, 6(1), 1-55.

- Hussong, A.M., Curran, P.J., & Bauer, D.J. (2013). Integrative Data Analysis in Clinical Psychology Research. *Annual Review of Clinical Psychology*, 9: 61-89.
- Kaat, A. J., & Lecavalier, L. (2015). Reliability and Validity of Parent-and Child-Rated Anxiety Measures in Autism Spectrum Disorder. *Journal of autism and developmental disorders*, 45(10), 3219-3231.
- Kaufman, J., Birmaher, B., Brent, D., Rao, U., Flynn, C., Moreci, P., et al. (1996). Schedule for affective disorders and schizophrenia for school-age children—present and lifetime version (K-SADSPL): Initial reliability and validity data. *Journal of the American Academy of Child and Adolescent Psychiatry*, 36(7), 980–988.
- Kaufman, A. S., & Kaufman, N. L. (2005). *Kaufman brief intelligence test—second edition (K-BIT-2)*. Circle Pines: American Guidance Service.
- Kerns, C. M., & Kendall, P. C. (2012). The presentation and classification of anxiety in autism spectrum disorder. *Clinical Psychology: Science and Practice*, 19(4), 323-347.
- Kerns, C. M., Kendall, P. C., Berry, L., Souders, M. C., Franklin, M. E., Schultz, R. T., et al. (2014). Traditional and atypical presentations of anxiety in youth with autism spectrum disorder. *Journal of autism and developmental disorders*, 44(11), 2851-2861.
- Kerns, C. M., Maddox, B. B., Kendall, P. C., Rump, K., Berry, L., Schultz, R. T., et al. (2015). Brief measures of anxiety in non-treatment-seeking youth with autism spectrum disorder. *Autism*, 19(8), 969-979.
- Kerns, C.M., Renno, P., Kendall, P.C., Wood, J.J., & Storch, E.A. (2017). Anxiety Disorders

Interview Schedule-Autism Addendum: Reliability and Validity in Children with Autism Spectrum Disorder. *Journal of Clinical Child & Adolescent Psychology*, 46, 88-100.

Kerns, C.M., Rump, K., Worley, J., Kratz, H., McVey, A., et al. (2016). The Differential Diagnosis of Anxiety Disorders in Cognitively Able Youth with Autism. *Cognitive and Behavioral Practice*, 23 (4), 530-547.

Lecavalier, L., Wood, J. J., Halladay, A. K., Jones, N. E., Aman, M. G., Cook, E. H., et al. (2014). Measuring anxiety as a treatment endpoint in youth with autism spectrum disorder. *Journal of autism and developmental disorders*, 44(5), 1128-1143.

Leyfer, O. T., Folstein, S. E., Bacalman, S., Davis, N. O., Dinh, E., Morgan, J., et al. (2006). Comorbid psychiatric disorders in children with autism: Interview development and rates of disorders. *Journal of Autism and Developmental Disorders*, 36(7), 849-861.

Li, J. C. H., Lau, W. Y., & Au, T. K. F. (2011). Psychometric properties of the Spence Children's Anxiety Scale in a Hong Kong Chinese community sample. *Journal of anxiety disorders*, 25(4), 584-591.

Lidstone, J., Uljarevic, M., Sullivan, J., Rodgers, J., McConachie, H., Freeston, M., et al. (2014). Relations among restricted and repetitive behaviours, anxiety and sensory features in children with autism spectrum disorders. *Research in Autism Spectrum Disorders*, 8(2), 82-92.

Lord, C., Rutter, M., DiLavore, P., & Risi, S. (2001). *Autism Diagnostic Observation Schedule (ADOS) manual*. Los Angeles, CA: Western Psychological Services.

- Lydon, S., Healy, O., O'Callaghan, O., Mulhern, T., & Holloway, J. (2015). A systematic review of the treatment of fears and phobias among children with Autism Spectrum Disorders. *Review Journal of Autism and Developmental Disorders*, 2, 141-154.
- Magiati, I., Chan, J. Y., Tan, W. L. J., & Poon, K. K. (2014). Do non-referred young people with Autism Spectrum Disorders and their caregivers agree when reporting anxiety symptoms? A preliminary investigation using the Spence Children's Anxiety Scale. *Research in Autism Spectrum Disorders*, 8(5), 546-558.
- Magiati, I., Ong, C., Lim, X. Y., Tan, J. W. L., Ong, A. Y. L., Patricia, F., et al. (2016). Anxiety symptoms in young people with autism spectrum disorder attending special schools: Associations with gender, adaptive functioning and autism symptomatology. *Autism*, 20(3), 306-320.
- Magiati, I., Ozsivadjian, A., & Kerns, C. (2017). Phenomenology and presentation of anxiety in Autism Spectrum Disorder. In C. Kerns, P. Renno, E. Storch, P. Kendall, & J. Wood (Eds.), *Anxiety in Children and Adolescents with Autism Spectrum Disorder*, 1st edition, p. 33-54. Elsevier Academic Press, US.
- March, J. S., Parker, J. D., Sullivan, K., Stallings, P., & Conners, C. K. (1997). The Multidimensional Anxiety Scale for Children (MASC): factor structure, reliability, and validity. *Journal of the American Academy of Child & Adolescent Psychiatry*, 36(4), 554-565.
- Maskey, M., Warnell, F., Parr, J., Le Couteur, A.S. & McConachie, H. (2013). Emotional and behavioural problems in children with ASD. *Journal of Autism and Developmental*

Disorders, 43(4), 851-859.

Matson, J. L., Wilkins, J., Sevin, J. A., Knight, C., Boisjoli, J. A., & Sharp, B. (2009).

Reliability and item content of the Baby and Infant Screen for Children with aUtism Traits (BISCUIT): Parts 1-3. *Research in Autism Spectrum Disorders*, 3, 336-344.

Matson, J. L., & Wilkins, J. (2008). Reliability of the autism spectrum disorders-comorbid for children (ASD-CC). *Journal of Developmental and Physical Disabilities*, 20(4), 327-336.

Matson, J. L., LoVullo, S. V., Rivet, T. T., & Boisjoli, J. A. (2009). Validity of the autism spectrum disorder-comorbid for children (ASD-CC). *Research in Autism Spectrum Disorders*, 3(2), 345-357.

May, T., Cornish, K., & Rinehart, N. J. (2015). Parent-Child Agreement Using the Spence Children's Anxiety Scale and a Thermometer in Children with Autism Spectrum Disorder. *Autism research and treatment*, 2015, 1-9.

Mazefsky, C. A., Pelphrey, K. A., & Dahl, R. E. (2012). The need for a broader approach to emotion regulation research in autism. *Child Development Perspectives*, 6, 92-97.

Mazefsky, C. A., Kao, J., & Oswald, D. P. (2011). Preliminary evidence suggesting caution in the use of psychiatric self-report measures with adolescents with high-functioning autism spectrum disorders. *Research in Autism Spectrum Disorders*, 5(1), 164-174.

McConachie, H., Barry, R., Spencer, A., Parker, L., Le Couteur, A.S. & Colver, AF. (2009). Daslⁿe: The challenge of developing a regional database for Autism Spectrum Disorder. *Archives of Disease in Childhood*, 94(1), 38-41.

- McConachie, H., McLaughlin, E., Grahame, V., Taylor, H., Honey, E., Tavernor, L., et al. (2013). Group therapy for anxiety in children with autism spectrum disorder. *Autism*, 18(6), 723-732.
- Mikita, N., Hollocks, M. J., Papadopoulos, A. S., Aslani, A., Harrison, S., Leibenluft, E., et al. (2015). Irritability in boys with autism spectrum disorders: an investigation of physiological reactivity. *Journal of Child Psychology and Psychiatry*, 56(10), 1118-1126.
- Muthén, L.K. & Muthén, B.O. (2012). Mplus User's Guide. Seventh Edition. Los Angeles, CA: Muth'en & Muth'en.
- Nauta, M. H., Scholing, A., Rapee, R. M., Abbott, M., Spence, S. H., & Waters, A. (2004). A parent-report measure of children's anxiety: Psychometric properties and comparison with child report in a clinic and normal sample. *Behaviour Research and Therapy*, 42(7), 813-839.
- Norris, M., & Lecavalier, L. (2010). Evaluating the use of exploratory factor analysis in developmental disability psychological research. *Journal of autism and developmental disorders*, 40(1), 8-20.
- O'Connor, B. P. (2000). SPSS and SAS programs for determining the number of components using parallel analysis and Velicer's MAP test. *Behavior Research Methods, Instrumentation, and Computers*, 32(3), 396-402.
- Ollendick, T. H., & White, S. W. (2012). The presentation and classification of anxiety in autism spectrum disorder: Where to from here?. *Clinical Psychology: Science and*

Practice, 19(4), 352-355.

Ozsivadjian, A., Hibberd, C., & Hollocks, M. J. (2014). Brief report: the use of self-report measures in young people with autism spectrum disorder to access symptoms of anxiety, depression and negative thoughts. *Journal of autism and developmental disorders*, 44(4), 969-974.

Ozsivadjian, A., Knott, F., & Magiati, I. (2012). Parent and child perspectives on the nature of anxiety in children and young people with autism spectrum disorders: a focus group study. *Autism*, 16(2), 107-121.

Parker, K. J., Garner, J. P., Libove, R. A., Hyde, S. A., Hornbeak, K. B., Carson, D. S., et al. (2014). Plasma oxytocin concentrations and OXTR polymorphisms predict social impairments in children with and without autism spectrum disorder. *Proceedings of the National Academy of Sciences*, 111(33), 12258-12263.

Pellecchia, M., Connell, J. E., Kerns, C. M., Xie, M., Marcus, S. C., & Mandell, D. S. (2016). Child characteristics associated with outcome for children with autism in a school-based behavioral intervention. *Autism*, 20(3), 321-329.

Renno, P., & Wood, J. J. (2013). Discriminant and convergent validity of the anxiety construct in children with autism spectrum disorders. *Journal of autism and developmental disorders*, 43(9), 2135-2146.

Rieske, R. D., Matson, J. L., Davis III, T. E., Konst, M. J., Williams, L. W., & Whiting, S. E. (2013). Examination and validation of a measure of anxiety specific to children with autism spectrum disorders. *Developmental neurorehabilitation*, 16(1), 9-16.

- Rodgers, J., Glod, M., Connolly, B., McConachie, H. (2012). The relationship between anxiety and repetitive behaviours in Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*, 42(11), 2404–2409.
- Rodgers, J., Riby, D.M., Janes, E., Connolly, B., McConachie, H. (2012). Anxiety and Repetitive Behaviours in Autism Spectrum Disorders and Williams syndrome: A Cross-syndrome Comparison. *Journal of Autism & Developmental Disorders*, 42(2) 175-180.
- Rodgers, J., Wigham, S., McConachie, H., Freeston, M., Honey, E., & Parr, J. R. (2016). Development of the anxiety scale for children with autism spectrum disorder (ASC-ASD). *Autism Research*. Advance online publication. DOI: 10.1002/aur.1603
- Roid, G. H. (2003). *Stanford-binet intelligence scales (SB5)*. Rolling Meadows, IL: Riverside.
- Russell, E., & Sofronoff, K. (2005). Anxiety and social worries in children with Asperger syndrome. *Australian and New Zealand journal of psychiatry*, 39(7), 633-638.
- Rutter, M., Bailey, A., & Lord, C. (2003). *Manual for the Social Communication Questionnaire*. Los Angeles, CA: Western Psychological Services.
- Schaap, L.A., Peeters, G.M.E.E., Dennison, E.M., Zambon, S., Nikolaus, T., et al. (2011). European Project on Osteoarthritis (EPOSA): methodological challenges in harmonization of existing data from five European population-based cohorts on aging. *BMC Musculoskeletal Disorders*, 12: 272. doi: 10.1186/1471-2474-12-272

- Schohl, K., Karst, J., Dolan, B., Stevens, S., Carson, A.M., & Vaughan Van Hecke, A. (2014). A replication and extension of the PEERS intervention: Examining effects on social skills and social anxiety in adolescents with autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 44, 532-545. DOI: 10.1007/s10803-013-1900-1
- South, M., Chamberlain, P. D., Wigham, S., Newton, T., Gray, L., Freeston, M., et al. (2014). Enhanced decision making and risk avoidance in young people with high-functioning Autism Spectrum Disorder. *Neuropsychology*, 28(2), 222-228, DOI 10.1037/neu0000016.
- South, M., & Rodgers, J. (2017). Sensory, Emotional and Cognitive Contributions to Anxiety in Autism Spectrum Disorders. *Frontiers in Human Neuroscience*, Volume 11, <https://doi.org/10.3389/fnhum.2017.00020>.
- Spence, S. H. (1997). Structure of anxiety symptoms among children: a confirmatory factor-analytic study. *Journal of abnormal psychology*, 106(2), 280-297.
- Spence, S. H. (1998). A measure of anxiety symptoms among children. *Behaviour research and therapy*, 36(5), 545-566.
- Spence, S. H. (1999). Spence Children's Anxiety Scale (Parent version). Brisbane: University of Queensland. Available at www.scaswebsite.com
- Spence, S. H., Barrett, P. M., & Turner, C. M. (2003). Psychometric properties of the Spence Children's Anxiety Scale with young adolescents. *Journal of anxiety disorders*, 17(6), 605-625.

van Steensel, F. J., Bögels, S. M., & Perrin, S. (2011). Anxiety disorders in children and adolescents with autistic spectrum disorders: A meta-analysis. *Clinical child and family psychology review*, 14(3), 302-317.

van Steensel, F. J., Deutschman, A. A., & Bögels, S. M. (2013). Examining the screen for child anxiety-related emotional disorder-71 as an assessment tool for anxiety in children with high-functioning autism spectrum disorders. *Autism*, 17(6), 681-692.

Vaughan Van Hecke, A., Stevens, S., Carson, A.M., Dolan, B., Karst, J., Schohl, K., et al. (2015). Measuring the plasticity of social approach: A randomized controlled trial of the effects of the PEERS intervention on EEG asymmetry in adolescents with autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 45(2), 316-335. DOI 10.1007/s10803-013-1883-y

Steinhausen, H. C., & Metzke, C. W. (2004). Differentiating the behavioural profile in autism and mental retardation and testing of a screener. *European Child & Adolescent Psychiatry*, 13(4), 214-220.

Stern, J. A., Gadgil, M. S., Blakeley-Smith, A., Reaven, J. A., & Hepburn, S. L. (2014). Psychometric properties of the SCARED in youth with autism spectrum disorder. *Research in autism spectrum disorders*, 8(9), 1225-1234.

Sterling, L., Renno, P., Storch, E. A., Ehrenreich-May, J., Lewin, A. B., Arnold, E., et al. (2015). Validity of the Revised Children's Anxiety and Depression Scale for youth with autism spectrum disorders. *Autism*, 19(1), 113-117.

Storch, E. A., Ehrenreich May, J., Wood, J. J., Jones, A. M., De Nadai, A. S., Lewin, A. B.,

- et al. (2012). Multiple informant agreement on the anxiety disorders interview schedule in youth with autism spectrum disorders. *Journal of Child and Adolescent Psychopharmacology*, 22(4), 292-299.
- Tabachnick, B. G., & Fidell, L. S. (2013). *Using Multivariate Statistics (6th eds.)*. Boston : Pearson.
- Trembath, D., Germano, C., Johanson, G., & Dissanayake, C. (2012). The experience of anxiety in young adults with autism spectrum disorders. *Focus on Autism and Other Developmental Disabilities*, 27(4), 213-224.
- Thurstone, L. L. (1947). *Multiple factor analysis*. Chicago, IL: University of Chicago Press.
- Vasa, R. A., Mazurek, M. O., Mahajan, R., Bennett, A. E., Bernal, M. P., Nozzolillo, A. A., et al. (2016). Assessment and treatment of anxiety in youth with Autism spectrum disorders. *Pediatrics*, 137(Supplement 2), S115-S123.
- Wang, M., Meng, Q., Liu, L., & Liu, J. (2015). Reliability and Validity of the Spence Children's Anxiety Scale for Parents in Mainland Chinese Children and Adolescents. *Child Psychiatry & Human Development*, 1-10.
- Wechsler, D. (1991). *Manual for the Wechsler Intelligence Scale for Children—Third Edition*. San Antonio, TX: The Psychological Corporation.
- Wechsler, D. (1999) *Wechsler Abbreviated Scale of Intelligence*. San Antonio: Harcourt Brace.
- White, S. W., Bray, B. C., & Ollendick, T. H. (2012). Examining shared and unique aspects

of social anxiety disorder and autism spectrum disorder using factor analysis. *Journal of autism and developmental disorders*, 42(5), 874-884.

White, S. W., Lerner, M. D., McLeod, B. D., Wood, J. J., Ginsburg, G. S., Kerns, C., et al. (2015). Anxiety in youth with and without autism spectrum disorder: examination of factorial equivalence. *Behavior therapy*, 46(1), 40-53.

White, S. W., Oswald, D., Ollendick, T., & Scahill, L. (2009). Anxiety in children and adolescents with autism spectrum disorders. *Clinical Psychology Review*, 29(3), 216-229.

Whiteside, S. P., & Brown, A. M. (2008). Exploring the utility of the Spence Children's Anxiety Scales parent-and child-report forms in a North American sample. *Journal of Anxiety Disorders*, 22(8), 1440-1446.

Wigham, S., & McConachie, H. (2014). Systematic review of the properties of tools used to measure outcomes in anxiety intervention studies for children with autism spectrum disorders. *PloS one*, 9(1).

Wigham, S., Rodgers, J., South, M., McConachie, H., & Freeston M.H. (2015). The interplay between sensory processing abnormalities, intolerance of uncertainty, anxiety and restricted and repetitive behaviours in Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*, 45(4), 943-52. DOI 10.1007/s10803-014-2248-x

Witwer, A. N., & Lecavalier, L. (2007). Autism screening tools: an evaluation of the social communication questionnaire and the developmental behaviour checklist–autism screening algorithm. *Journal of Intellectual and Developmental Disability*, 32(3), 179-

187.

Yong, A. G., & Pearce, S. (2013). A beginner's guide to factor analysis: Focusing on exploratory factor analysis. *Tutorials in Quantitative Methods for Psychology*, 9(2), 79-94.

Zainal, H., Magiati, I., Tan, J. W. L., Sung, M., Fung, D. S., & Howlin, P. (2014). A preliminary investigation of the Spence Children's Anxiety Parent Scale as a screening tool for anxiety in young people with autism spectrum disorders. *Journal of autism and developmental disorders*, 44(8), 1982-1994.

Zhao, J., Xing, X., & Wang, M. (2012). Psychometric properties of the Spence Children's Anxiety Scale (SCAS) in Mainland Chinese children and adolescents. *Journal of anxiety disorders*, 26(7), 728-736.

of measurement properties of caregiver-report anxiety measures in studies of young people with ASD

Studies	ASD Participants	Reliability	Validity
Matson & Wilkins (2008)	113-177 2-17 year olds with ASD; clinical and community recruited	$\alpha=.74$ Mean item inter-rater (familiar family member): $\alpha=.46$	<i>Structural</i> : EFA identified one worry/depressed factor among other internalising and externalising symptoms <i>Convergent</i> : Large correlations (i.e. .52 to .68 with BASC-2 somatization, depression, anxiety) <i>Divergent</i> : Zero to moderate correlations (i.e. .07 to .32 with BASC-2 attentional problems, withdrawal, hyperactivity, aggression, conduct problems)
Matson et al. (2009)	Mixed functioning (some with ID, Down syndrome)	Mean item test-retest of 2 weeks: $\alpha=.55$	
Rieske et al. (2013)	53 2-16 year olds with ASD from clinical setting; some with ID		<i>Convergent</i> : Large correlations (i.e. .70 to .74 with BASC-2 depression, anxiety, internalizing total) <i>Divergent</i> : non-significant or small correlations (i.e. -.05 to .14 with BASC-2 attentional problems, daily living)
Hallett et al. (2013a)	415 4-17 year olds with ASD from clinical settings; subgroups of $IQ \geq 70$ and < 70	Total items $\alpha=.87$ Subscales $\alpha=.65$ to .85	<i>Structural</i> : Promising evidence from EFA and CFA on 4-factor structure – GAD, SAD, Over-arousal, SoP <i>Divergent</i> : Small-to-moderate correlations (i.e. .08 to .31 with OCD measure, Vineland adaptive behaviour, Aberrant Behavior Checklist irritability, social withdrawal, stereotypy and inappropriate speech, PDD measure)
White et al. (2012)	30 12-17 year olds from clinical setting Verbal $IQ \geq 70$	Total $\alpha=.85$ Inter-informant (caregiver/child): $r=.45$	<i>Convergent</i> : Large (.78 with MASC-P); .47 with depressive symptoms <i>Divergent</i> : Small non-significant correlations (-.20 to .15 with ADOS and Verbal IQ)
Storch et al. (2012)	72 7-17 year olds with ASD from clinical settings $IQ \& VIQ > 70$	Total $\alpha = .59$ Inter-rater $r=.86$ Test-retest after 26 days $r=.83$	<i>Convergent</i> : .40 to .62 with MASC-P, CBCL anxiety and internalizing symptoms, clinician rated severity of anxiety <i>Divergent</i> : .03 to .47 with ADOS and CBCL externalizing behaviours subscales
Kerns et al (2015)	54 7-17 year olds unselected/ community sample; $IQ > 60$	total $\alpha = .90$	<i>Convergent</i> : .46 to .60 with SCARED-P and BASC-2 Parent Poor sensitivity (=53) & strong specificity (=95) using original cut-off; stronger specificity (=93), but weaker specificity (=71) with alternative cut-off
Kaat & Lecavalier (2015)	46 8-16 year olds with ASD with > 50% seeking treatment $IQ \geq 55$	Total $\alpha = .91$ Subscales $\alpha = .57$ to .88 <i>Test-retest</i> after 2 to 3 weeks: total = .82, subscales = .76 to .87 <i>Parent-child agreement</i> : Total= .23; Subscales= .08 to .27	<i>Convergent</i> : Large correlations (i.e. .64 to .85 with MASC-P, CSI anxiety) <i>Divergent</i> : Non-significant correlations (i.e. -.20 to .29 with child variables, emotion recognition, ADOS, CSI externalizing behaviour subscales, but .69 with CSI depressive disorders)
Renno & Wood (2013)	88 7-11 year olds with ASD and comorbid anxiety	Total score $\alpha = .86$ Subscales $\alpha = .77$ to .88	<i>Discriminant/Structural</i> : Anxiety symptom severity can be distinguished from ASD symptom severity with CFA, but poor discrimination between separation and social anxiety

1		disorder	<i>Parent-child agreement:</i>	<i>Convergent:</i> $r=.18$ to $.58$ with semi-structured interview ratings
2		IQ>70	Total= .17; Subscales= .06	<i>Divergent:</i> Non-significant or moderate correlations (i.e. $-.04$ to $.40$
3			to .50	with parent rated and semi-structured interview ASD symptom severity
4	White et al.	30 12-17 year olds	Total: $\alpha = .90$	<i>Convergent:</i> $.78$ with CASI
5	(2012)	with ASD and		<i>Divergent:</i> Non-significant or moderate correlations ($-.26$ to $.385$
6		comorbid anxiety	<i>Parent-child agreement:</i>	with ADOS, VIQ, depressive symptoms)
7		disorder; VIQ ≥ 70	Total=.36	<i>Discriminant:</i> Poor (i.e. not significantly higher than a published
8			Subscales=.11 to .52	unselected ASD sample; only 23.3% met cut-off)
9	White et al.	465 7-17 year old	<i>Parent-child agreement</i>	<i>Structural:</i> Poor model fit; EFA suggested alternative 4-factor
10	(2015)	with ASD and	for Total =.38	structure (SAD/Panic, SoP evaluation focused, somatic symptoms,
11		comorbid anxiety		SoP performance focused)
12		disorder;	Inter-item subscale	
13		VIQ>70	correlations:	
14			.33 to .34, except .18 for	
15			harm avoidance	
16	Kaat &	46 8-16 year olds	Total $\alpha = .92$;	<i>Convergent:</i> $.33$ to $.85$ with RCADS-P, CSI anxiety; $.65$ with CSI
17	Lecavalier	with ASD; >50%	Subscales $\alpha = .67$ to $.95$	depressive disorders
18	(2015)	seeking treatment for	<i>Test-retest</i> after 2 to 3	<i>Divergent:</i> Non-significant correlations (i.e. $-.21$ to $.26$ with child
19		anxiety;	weeks: Total=.83;	variables, emotion recognition, social cognition, ADOS, CSI
20		IQ ≥ 56	subscales = $.71$ to $.90$	externalizing behaviour subscales
21			<i>Parent-child agreement:</i>	
22			Total= .23; subscales= .0	
23			to .45	
24	Blakeley-	129 7-18 year olds	Total score $\alpha = .90$	<i>Structural:</i> Items loaded on their original factors generally in PCA
25	Smith et al.	with ASD/ clinical	Subscales $\alpha = .73$ to $.89$	(Panic, SAD, SoP), except GAD and school phobia items
26	(2012)	sample; VIQ ≥ 70		
27	Stern et al.		<i>Parent-child agreement</i>	<i>Convergent:</i> small-to-moderate correlations (i.e. total $r=.44$ with
28	(2014)		(in a sample of 63 8-14	clinician global clinical severity rating; Subscales $r=.28$ to $.46$ with
29			year olds with VIQ ≥ 80):	semi-structured interview ratings
30			total= .52; subscales=.27	
31			to .71	Good Sensitivity: total = $.71$, subscales = $.70$ to 1.00
32	Kerns et al.	54 7-17 year olds	Total score $\alpha = .91$	Acceptable Specificity: total = $.67$, subscales = $.33$ to $.83$
33	(2015)	with ASD;		<i>Convergent:</i> Large correlations (i.e. total $r= .60$ to $.73$ with BASC2-P
34		unselected, but 37%	<i>Parent-child agreement:</i>	and PARS)
35		had another anxiety	Total =.33	
36		disorder;		Sensitivity ($.58$) and specificity ($.86$) with original cut-off; better
37		IQ>60		sensitivity ($.95$), but poorer specificity ($.62$) with alternative cut-off
38	Van Steensel	115 7-18 year olds	Total score $\alpha = .94$ to $.95$	<i>Convergent:</i> $.42$ to $.75$ with semi-structured parent interview ratings
39	et al. (2013)	with ASD &	Subscales $\alpha = .70$ to $.88$	or another SCARED-71 caregiver report
40		comorbid anxiety		<i>Discriminant:</i> Good (i.e. significant differences in subscales scores
41		disorder	<i>Parent-child agreement:</i>	between disordered and no disorder, except for OCD subscale
42			Total score $r=.39$ to $.41$	Sensitivity (total =.95; subscales = $.78$ to $.90$); Specificity
43		High functioning		(subscales= $.39$ to $.67$) with original cut-off

1			<i>Parent-child agreement</i> on	Better specificity (.67 to .71) but lower sensitivity (.71 to .80) with
2			diagnosis of anxiety	alternative cut-off for separation, social & generalized anxiety
3			disorder: .52 to .85	
4	Zainal et al.	32 6-18 year olds	Total score α = .88	<i>Convergent:</i> Moderate-to-large correlations (i.e. .48 to .61 with semi-
5	(2014)	with ASD	Subscales α = .60 to .78	structured interview rating and DBC-Anxiety)
6		unselected;		<i>Discriminant:</i> Good (i.e. significant difference between anxiety
7		NVIQ $\geq$ 70		disordered and no disorder groups on total score, social phobia
8				subscale and panic subscale)
9				Promising Sensitivity (.71), specificity (.76), NPV (.90), but poor
10	Magiati et al.	38 8-18 year old	Total score α = .88	PPV (.45)
11	(2014)	unselected;	Subscales α = .51 to .80	
12		NVIQ >70	<i>Parent-child agreement:</i>	
13			Total=.69; Subscales=.42	
14			to .78; Mean item=-.01 to	
15			.69	
16	Magiati et al.	241 5-17 year olds	Total score α = .87	<i>Divergent:</i> -.01 to .48 with age, gender, adaptive functioning and
17	(2016)	with ASD;	Subscales α = .47 to .76	DBC autism symptom severity
18		unselected sample;		
19		Mixed level of IQ		
20		and/or adaptive		
21		functioning (SIB-R		
22		SS=58.8 (40.4))		
23	Ozsvadjian et	30 10-16 year olds	Total score α =.92	<i>Divergent:</i> non-significant correlation with full scale IQ, but small
24	al. (2014)	with ASD from	<i>Parent-child agreement:</i>	correlation with autism symptom severity (.38)
25		special school;	Total score =.59	
26		IQ $\geq$ 70		
27	May et al.	44 8-13 year olds	<i>Parent-child agreement:</i>	
28	(2015)	with ASD from	Total score =.25	
29		clinical sample;	Subscales = .11 to .31	
30		IQ $\geq$ 70		

ability; EFA = Exploratory Factor Analysis; BASC= Behaviour Assessment System for Children; PDD= Pervasive Developmental disorder; ADOS=Autism Diagnostic Observation Schedule; CFA= confirmatory factor analysis; GAD= generalized anxiety disorder; SAD= Separation Anxiety Disorder; SoP= Social Phobia; IQ= Intelligence Quotient; PCA= principal component analysis; NVIQ = Non-verbal Intelligence Quotient; CBCL= Child Behavior Checklist; PPV= Positive Predictive Value; DBC= Developmental Behavior Checklist; ADIS=Anxiety Disorders Interview Schedule.

1
 2 **Characteristics in current pooled database and for each dataset**

3 Sample description	4 Recruitment & inclusion- 5 exclusion criteria	6 Diagnosis	7 Autism Severity 8 measure	9 IQ or Adaptive 10 functioning & 11 other measures	12 SCAS-
13 465 (53.4%) UK 14 164 (18.9%) US 15 241 (27.7%) SG 16 Informants: 17 Mothers (651, 74.8%) 18 Fathers (77, 8.9%), 19 Both parents (8, 0.9%), 20 Grandparents (1, .1%), 21 Others (4, 0.5%) 22 Not reported (129, 23 14.8%). 24 19.5% in mainstream 25 schools, 31.8 % in 26 special schools, 0.1% 27 home-schooled, 58.5% 28 information not 29 available	30 Recruited through database, 31 information sent to suitable 32 families. 33 Inclusion: diagnosis of ASD, 34 8-16 years old 35 Exclusion: taking medication 36 for repetitive behaviours	37 727 (83.6%) Autism/ASD 38 111 (12.8%) Asperger's 39 syndrome 40 32 (3.7%) PDD-NOS 41 Caregiver report or research- 42 based; all made by qualified 43 professionals based on DSM- 44 IV, DSM-IV-TR or ICD-10 45 criteria according to standard 46 national procedures for 47 diagnosis of ASD in the 48 respective countries typically 49 including interviews with 50 caregivers and child 51 observations/ assessment.	52 SRS (n=130; 53 M=159.2; 54 SD=19.13) 55 SCQ (n=122; 56 M=21.33, 57 SD=6.56) 58 DBC-ASA 59 (n=238; 60 M=18.39; SD=9.33) Harmonized pooled item mean caregiver- reported autism symptom severity score (range 0-1; n=393; M=0.41; SD=0.20)	61 IQ scores (M= 62 96.58, SD= 20.36, 63 Range=13-144, 64 n=312) 65 Adaptive 66 functioning SS 67 scores (n=239; 68 M=58.8; SD=40.4) 69 Overall 70 "functioning" 71 classification mean 72 ranking score 73 (M=4.73, 74 SD=2.203, 75 (1=lower to 8= 76 superior; n=551)	77 M=2 78 SD=1 79 Range
80 UK community 81 sample, not help- 82 seeking for anxiety 83 8-16 years old (M= 84 53.45 months, 85 SD=24.86); 80% Male	86 Recruited through database, 87 information sent to suitable 88 families. 89 Inclusion: diagnosis of ASD, 90 8-16 years old 91 Exclusion: taking medication 92 for repetitive behaviours	93 Diagnosed through 94 multidisciplinary team 95 assessment following guidelines 96 of the UK National Autism Plan 97 for Children (Le Couteur, 98 2003); confirmed with SCQ 99 100% Autism/ASD diagnosis	100 SCQ 101 (M=24.80, SD= 102 5.07, Range=15- 103 31) 104 100% met cut-off	105 WISC-III short 106 form full IQ 107 (M=93.20, 108 SD=18.27, 109 Range= 69-133)	110 M=4 111 SD=2 112 Range
113 UK Community 114 sample, not help- 115 seeking for anxiety 116 9-15 years old (M= 117 47.37 months, 118 SD=24.00); 86.4%	119 Families given information 120 about joining research 121 database by diagnosing 122 clinicians, other service 123 providers, or voluntary 124 organisations	125 Multi-professional diagnostic 126 teams (as Study 1); Parent 127 report of diagnosis received and 128 validated with information 129 collected from professionals; 130 most children known to local 131 services and diagnosis	132 No additional 133 measure 134 administered as 135 part of the study	136 Not measured as 137 part of this study	138 M=3 139 SD=1 140 Range

Male 1 2 3 4 5	Inclusion: diagnosis of ASD; Any age to 18 years.	confirmed by professionals 75.4% Autism/ASD 24.6% Asperger's syndrome			
UK clinical sample help-seeking for anxiety 9 10 11 12 13 14	Approached by member of CAMHS team Inclusion: diagnosis of ASD, at least 2 anxiety disorders; 9 to 13 years 11 months; Full IQ>69; Exclusion: untreated ADHD or oppositional behaviors	Multi-professional diagnostic team, confirmed by ADOS Module 3 and SCQ. 100% Autism/ASD	ADOS diagnostic algorithm score (n= 20; M=10.35, SD=3.79, range=5– 19) 85% (17 out of 20) met cut-off	WASI full score (n=19; M=103.74, SD = 15.23, Range= 70-121)	M=4 SD=1 Range=
US research sample, not help-seeking for anxiety 15 16 17 18 19 20 21 22 23	Inclusion: Research diagnosis of ASD & FSIQ > 80; aged 8- 18 years	Research diagnosis of ASD, and above cut-off in SRS 100% Autism/ASD	ADOS diagnostic algorithm score (n= 62; M= 11.68, SD=3.57, Range= 7-20) SRS Total Score (n= 59), M=104, SD= 23.18, Range=55-149	WASI (n=43; M=110.53, SD=13.78, Range= 84-140)	M=2 SD=1 Range=
UK Community and research database, not help seeking for anxiety 24 25 26 27 28 29 30 31 32	Recruited through DASLne database Inclusion: 8-18 years old with ASD or Asperger's syndrome	As in Study 1 100% Autism/ASD	No formal measure administered as part of this study	WASI (n=14; M=101.93, SD=11.55, Range=83-119)	M=2 SD=1 Range=
UK community sample, not help seeking for anxiety 33 34 35 36 37 38 39 40	Recruited from mainstream and special schools Inclusion: age 8-16 years old with ASD or Asperger's syndrome	Based on SRS and parent reported diagnosis 58.8% Autism 41.2% Asperger's syndrome	SRS (n=32; M=162.55, SD= 20.19, Range = 116-202) 100% (32 out of 32) met cut-off	WASI (M=83.29, SD= 12.42, Range= 64- 116)	M=3 SD=1 Range=
Singapore Unselected community sample; 41 42 43	Recruited from special schools	Parent- report and special school formal entry criteria; all professional diagnosis;	DBC-Autism Screening Algorithm score	Adaptive behaviour SIB-R (n=239; M=58.83,	M=1 SD=1 Range=

5-17 years old (M=123.58 months, SD=35.88); 81.7% Male	Inclusion: age range 6-18 years old; diagnosis of Asperger's Syndrome, ASD, Autism/ Autistic disorder, PDD-NOS by professional	91.7% Autism; 6.2% Asperger's syndrome; 2.1% PDD-NOS	(n=238; M=16.23, SD=8.34, Range=0-42)	SD=40.37, Range= 0-171)	
		DBC-Autism Algorithm (M=18.4, SD=9.3, range=0-41)	68.1% (162 out of 238) > cut-off of 14	DBC-Anxiety subscale (n=238; M=4.66, SD=2.96, Range=0- 15)	
				DBC- disruptive/antisocia l subscale (n=238; M=12.54, SD=7.76, Range=0-34)	
Recruited from CAMHS	IQ $\geq$ 70 on WASI and with good verbal ability	ASD diagnosed by Multi- disciplinary Team in CAMHS services; 31 out of 52 based on ADOS/ADI used by local team, otherwise clinical diagnoses confirmed by SCQ; 100% ASD	No scores available for the study, but clinical diagnoses used ADOS/ ADI or SCQ.	WASI (M=101.42, SD=13.20, Range=76- 138)	M=3 SD=1 Range
10-16 years old (M=154.19 months, SD=22.99); 100% Male					
Special schools	IQ $\geq$ 70 on WASI and a score of $\geq$ 70 on the WIAT	SCQ; 100% Autism	SCQ (M=22.78, SD=7.50, Range=6- 35)	WASI (M=96.19, SD=13.65, Range=73- 122)	M=2 SD=1 Range
10-17 years old (M=157.78 months, SD=16.65); 100% Male			87.5% (28 out of 32) met cut-off		
Recruited through local schools with specialist autism provision and parent support groups.	Clinical diagnosis of ASD established by multidisciplinary clinical team.	ASD diagnosed by multidisciplinary diagnostic team. Clinicians from two health boards involved in diagnosing children confirmed DSM-IV-TR diagnosis –ADOS routinely used in these services.	SCQ (n=28;M=21.35, SD=4.12, Range=12- 31)	No formal measures; Parent report of language level based on the Diagnostic Interview for Social and Communication Disorders (DISCO) expressive language scale	M=2 SD=1 Range
6 to 18 years old (M=137.17 months, SD=40.90); 98.9% Male	Exclusions: brain injury, cerebral palsy, any neuro/ musculo/ skeletal disorder/ malformation that would seriously limit ability to walk without help or a known genetic condition (e.g., Fragile X, Down syndrome).	63.2% Autism; 36.8% Asperger's syndrome	92.9% (26 out of 28) met cut-off		
	N= 63 recruited from Ireland and N= 32 from South Wales				
Children with ASD recruited through the	3-12 year old children with ASD who met ADOS and	ASD diagnosis based on ADOS, ADI-R and expert	SRS (n=55; M=154.27,	Stanford Binet 5 th Ed. (n=54;	M=1 SD=1

Autism and Developmental Disorders Research Registry and by flyers posted in the Autism and Developmental Disorders Clinic at Stanford University 8 9 to 12 years old 10 (M=114.79 months, 11 SD=23.43); 80.4% 12 Male	ADI-R diagnostic criteria. All participants were (i) pre- pubertal, (ii) in good medical health, and (iii) willing to provide a blood sample. Participants included if they had a full-scale IQ >50 and no genetic conditions	clinical opinion. 51.8% Autism; 48.2% PDD- NOS	SD=18.64, Range=100-205) ADOS diagnostic algorithm score (n=51, M= 14.51, SD=5.21, Range=3-24) 100% (55) met SRS cut-off, but 94% (48) met ADOS cut-off	M=79.33, SD=26.12, Range=13– 128; N=54)	Range
13 14 RCT Community 15 sample recruited via in- 16 house waiting list for 17 social skills treatment 18 at University Autism 19 Clinic (not selected for 20 anxiety); assessed at 3 21 time points: before and 22 after intervention, and 23 at 6-months follow-up 24 to 16 years old 25 (M=161.02 months, 26 SD=16.82); 83.7% 27 Male	(a) aged between 11 and 16 years old; (b) fluent in English; (c) no history of adolescent major mental illness, such as bipolar, schizophrenia, or psychosis; (d) no history of hearing, visual, or physical impairments; (e) adolescent wants to learn to make and keep friends; (f) KBIT Verbal IQ > 70	Established community diagnosis by professional of either Autism, Asperger, or Pervasive Developmental Disorder—NOS; and meeting criteria for ASD or Autism on the ADOS-G Module 4 100% ASD on ADOS	ADOS diagnostic algorithm score (M=10.16, SD=3.27, Range=7– 18) SCQ (n=42; M=17.90, SD=6.38, Range=3-30) SRS (M=163.16, SD=18.17, Range = 125 – 199) 100% met ADOS and SRS cut-off	Kaufman Brief Intelligence Test- Second Edition (M=105.79, SD=19.28, Range=71– 144)	M=2 SD=1 Range

Communication Questionnaire (Rutter et al., 2003); SRS = Social Responsiveness Scale (Constantino & Gruber, 2005); ADOC = Autism Diagnostic Observation
Diagnostic Interview – Revised; DBC = Developmental Behaviour Checklist; WISC-III = Wechsler Intelligence Scale for Children Third Edition; WASI =
of Intelligence; KBIT = Kaufman Brief Intelligence Test; FSIQ= Full Scale Intelligence Quotient; SIB-R = Scales of Independent Behavior-Revised; Das
m Spectrum Disorder Living in the North East; SLaM = South London and Maudsley NHS Foundation Trust; CAMHS = Child and Adolescent Mental H

1

2 Statistics for variables and comparison among country subsamples (n=849; excluding the 21 participants from the clinically anxious subsample)

3

4	Mean (SD) / Mode [#]			Welch ANOVA / Chi-	Effect	<i>p</i>	Post-hoc	Effect
5				square [#]	size, η		analyses	size, d
6	UK	US	Singapore					
7								
8								
9								
10								
11								
12								
13								
14	146.10 (29.75)	142.58 (32.66)	123.58 (35.88)	F (2, 381.43)=34.93	0.29	<.001***	UK=US>SG	.11-.68
15								
16								
17	31.89 (18.99)	22.64 (15.89)	18.23 (11.23)	F(2, 425.50)=69.96	0.35	<.001***	UK>US>SG	.32-.88
18								
19								
20	162.28 (19.91)	158.17 (18.87)	-		-			
21	n=32	n=98						
22	23.12 (5.94)	17.90 (6.34)	-		-			
23								
24	n=80	n=42						
25								
26	-	-	18.29 (9.33)	-	-	-	-	-
27			n=238					
28	0.59 (0.13)	0.45 (0.16)	1.31(0.16)	F(2, 109.10)= 147.46	0.63	<.001***	UK>US>SG	0.86-
29								
30	n=112	n=42	n=238					1.90
31								
32								
33	5.71 (1.18)	5.73 (1.79)	[adaptive	χ^2 (2, N = 532) = 192.60	0.60	<.001***	US=UK>SG	0.013-
34	N=153	N=140	functioning]					1.28
35			3.40 (2.28)					
36			N=239					
37								
38	Male	Male	Male	χ^2 (2, N = 849) = 13.27	0.13	.001***	-	
39								
40								

41 square test for gender; ⁺ see Methods, Statistical plan; lower score indicates less severe symptoms; range 0-1; ⁺⁺ see Methods, Statistical Plan; functioning] to 8 (superior functioning); * p<.05; ** p<.01; *** p<.001. SRS=Social Responsiveness Scale; SCQ=Social Communication Questionnaire.

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

1 ve statistics for variables and comparison among participants meeting and not meeting ASD cut-off (n=849; excluding the 21 help-seeking for an
2

Mean (SD) / Mode [#]					
Meeting ASD		Not meeting ASD		Welch ANOVA /	Effect size, η
screening cut-off	(n=376)	screening cut-off/ no	ASD screening score		
		available (n=473)	Chi-square [#]		<i>p</i>
	132.57 (34.19)	144.16 (32.22)	F(1, 781.71)= 25.34	0.17	<.001 ^{***}
	23.96 (15.60)	28.02 (18.88)	F(1, 845.74)= 11.78	0.11	.001 ^{**}
	3.14 (3.46)	3.58 (4.08)	F(1, 843.49)= 2.97	0.058	.085
	4.34 (3.60)	4.87 (3.90)	F(1, 828.43)= 4.21	0.070	.040
	4.42 (2.90)	4.46 (3.09)	F(1, 823.94)= 0.40	0.0068	.841
	4.37 (4.02)	6.12 (4.75)	F(1, 843.55)= 33.61	0.19	<.001 ^{***}
	3.46 (3.19)	3.96 (3.83)	F(1, 845.16)= 4.30	0.070	.038
	4.24 (3.20)	5.04 (3.68)	F(1, 840.01)= 11.37	0.11	.001
classification	4.24 (2.28)/	5.49 (1.8)/	χ^2 (1, N = 532) = 50.50	0.31	<.001 ^{***}
	6 “average”	6 “average”			
	[n=347]	[n=185]			
	1.16 (.36)	1.10 (.30)	χ^2 (1, N = 849) = 6.35	0.086	.012 [*]
	Male	Male			

29 i-square test for gender and Overall Functioning classification; ⁺⁺ see Methods, Statistical Plan; classification range 1 (lower functioning) to 8 (superior fu
30 * $p \leq .001$.
31

32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Statistics, subscale-total correlations and internal consistency of original and PCA-derived SCAS-P full scale and subscales

	Pooled sample's [#] Mean (SD)	Pooled sample's Item Mean (SD)	Corrected subscale-total correlations	Cronbach's α	Normative means (SD) from Nauta et al. (2004)	One sample t-test comparison of study's mean with norms	Effect size, d
SCAS-P	3.38 (3.82)	0.38 (0.43)	.76	.83	1.0 (1.6)	t(848)=18.19*	0.81
10	4.63 (3.78)	0.78 (0.63)	.74	.76	2.6 (2.8)	t(848)=15.67*	0.61
11	4.44 (3.01)	0.89 (0.60)	.49	.55	2.6 (2.3)	t(848)=17.83*	0.69
12	5.34 (4.52)	0.90 (0.76)	.67	.84	4.2 (2.8)	t(848)=7.36*	0.30
13	3.74 (3.57)	0.63 (0.60)	.67	.80	1.1 (1.7)	t(848)=21.55*	0.94
14	4.68 (3.50)	0.79 (.59)	.82	.82	2.7 (2.0)	t(848)=16.50*	0.70
15							
16	26.22 (17.61)	0.70 (0.47)	-	.93	14.2 (9.7)	t(848)=19.90*	0.85
17							
18							
19							
20							
21	2.60 (3.52)	0.33 (0.44)	.71	.86			
22	4.54 (3.34)	0.92 (0.68)	.58	.76			
23	2.61 (2.50)	0.66 (0.63)	.62	.69			
24	8.17 (6.29)	0.92 (0.70)	.66	.90			
25							
26	2.07 (2.43)	0.52 (0.61)	.53	.75			
27							
28	19.99 (14.17)	0.68 (0.48)	-	.93			

* if $p < (.05/13) = .0038$; [#] all participants without clinical subsample (n=849).

1

Table 2 Correlations between child characteristics and participants' original SCAS-P scores and PCA-derived 30 item SCAS-P score

3

[illegible]

functioning classification and other continuous variables; %original SCAS-P subscale; ^SPCA-derived SCAS-P subscale. The values above the diagonal line correspond to the child characteristics and original SCAS-P total and subscales scores, while the values below the diagonal line correspond to the corrections between PCA-derived SCAS-P total and subscales scores.

seen in the total raw scores of the caregiver reported autism symptom measures and the SCAS-P total score were also similar to the relationship reported for CQ3= .35; DBC-ASA $r=.38$, all $p<.01$.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

on between referred (n=21) and unselected subsamples

Mean (SD) for clinical subsample (n=21)	Mean (SD) for unselected subsample (n=849)	Welch ANOVA	Effect size, d	p
3.57 (4.57)	3.38 (3.82)	F (1, 20.70)=4.73	0.52	.041
4.51 (4.48)	4.63 (3.78)	F(1, 20.71)=4.44	0.50	.047
5.71 (3.38)	4.44 (3.01)	F(1, 20.79)=2.93	0.40	.102
8.76 (4.07)	5.34 (4.52)	F(1, 21, 24)=14.36	0.79	.001*
6.00 (3.46)	3.74 (3.57)	F(1, 21.06)=8.72	0.64	.008
8.14 (3.61)	4.68 (3.50)	F(1, 20.94)=18.86	0.97	<.001*
40.90 (18.57)	26.22 (17.61)	F(1, 20.90)=12.85	0.81	.002*
4.43 (4.13)	2.60 (3.52)	F(1, 20.73)=4.03	0.48	.058
8.86 (4.52)	4.54 (3.34)	F(1, 20.54)=5.47	0.58	.030
3.57 (3.20)	2.61 (2.50)	F(1, 20.61)=2.76	0.33	.187
13.19 (4.93)	8.17 (6.29)	F(1, 21.64)=20.99	0.89	<.001*
2.95 (2.40)	2.07 (2.43)	F(1, 21.03)=2.76	0.36	.111
34.00 (14.69)	19.99 (14.17)	F(1, 20.93)=11.53	0.76	.003*

cant if $p < (.05/13) = <.0038$.

1
2
3 of Confirmatory Factor Analyses for the SCAS-P

	X ²	df	CFI	TFI	RMSEA (90% CI)	Model Comparison*
	11574	703	-	-	-	-
	4055	665	0.69	0.67	0.077 (0.074 - 0.079)	X ² (df) = 3470 (38) , p ≤ .01
	5151	665	0.59	0.56	0.101 (0.099 – 0.103)	N.S.
	2952	650	0.79	0.77	0.064 (0.061 – 0.066)	X ² (df) =1103 (15) , p ≤ .01
	3127	661	0.77	0.76	0.065 (0.063 – 0.068)	X ² (df) =175 (11) , p ≤ .01
	4242	663	0.67	0.65	0.079 (0.077 - 0.081)	-

calculated χ^2 vs. the previous model; CFI=Comparative Fit Index, TFI=
index; RMSEA=Root Mean Square Error of Approximation.

1

Items derived from Principal Component Analysis (PCA)

Items	Components				
	Mixed Social/ GAD	Separation anxiety	Somatic / Panic symptoms	OCD	Specific Phobias
people think of	.840				
make a fool of	.796				
people					
will do badly at school	.785				
(s)he has to talk in	.664				
has to take a test	.650				
something awful will	.603				
our family	.579				
something bad will happen	.452				
school in the mornings	.397				
to sleep on his/her		.789			
		.735			
is/her own at home		.683			
away from us / me		.492			
g afraid		.443			
her heart suddenly			-.759		
ckly for no reason					
her heart beating			-.756		
feeling as if (s)he			-.755		
g dizzy or faint			-.702		
shake			-.687		
			-.626		
care for no reason at			-.490		
will suddenly get a					
ere is nothing to be			-.476		
ngs over and over				.800	
ings in just the right				.797	
from happening					

(s)he has done things	.620	
1		
2		
3	.596	
4		.684
5		.621
6		.545
7		.541
8		

conducted using half of the unselected subsample randomly selected (n=425).

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Accepted Article

Scree plot of the Principal Components Analysis (PCA) for the SCAS-P.

Figure 1. Scree Plot of eigenvalues of components for Principal Component Analysis. Three arrows and a line were added to show the points of inflexion at component 2, 4 and 6.

19x14mm (300 x 300 DPI)

Accepted A

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Magiati, I;Lerh, JW;Hollocks, MJ;Uljarevic, M;Rodgers, J;McConachie, H;Ozsivadjian, A;South, M;Van Hecke, A;Hardan, A;Libove, R;Leekam, S;Simonoff, E

Title:

The measurement properties of the spence children's anxiety scale-parent version in a large international pooled sample of young people with autism spectrum disorder

Date:

2017-10-01

Citation:

Magiati, I., Lerh, J. W., Hollocks, M. J., Uljarevic, M., Rodgers, J., McConachie, H., Ozsivadjian, A., South, M., Van Hecke, A., Hardan, A., Libove, R., Leekam, S. & Simonoff, E. (2017). The measurement properties of the spence children's anxiety scale-parent version in a large international pooled sample of young people with autism spectrum disorder. *AUTISM RESEARCH*, 10 (10), pp.1629-1652. <https://doi.org/10.1002/aur.1809>.

Persistent Link:

<http://hdl.handle.net/11343/292994>