Relationships between self-reported and observed parenting behaviour, adolescent disordered eating attitudes and behaviours, and the 5-HTTLPR polymorphism: Data from the Australian Temperament Project.

Vanja Rozenblat^{a*}, Joanne Ryan^b, Eleanor Wertheim^c, Ross King^e, Craig A. Olsson^{e,d,b}, Primrose Letcher^d and Isabel Krug^a.

^a The University of Melbourne, Psychological Sciences, Faculty of Medicine, Dentistry and Health Sciences, Parkville, 3010 Victoria, Australia

^b Murdoch Children's Research Institute, Royal Children's Hospital Melbourne, Parkville 3010, Victoria, Australia

^c La Trobe University, School of Psychology and Public Health, Faculty of Health, Bundoora 3086, Victoria, Australia

^d The University of Melbourne, Department of Paediatrics, The Royal Children's Hospital Melbourne, Faculty of Medicine, Dentistry and Health Sciences, Parkville 3010, Victoria, Australia

^e Deakin University Geelong, Centre for Social and Early Emotional Development, School of Psychology, Faculty of Health, 3220, Victoria, Australia

*Corresponding author. School of Psychological Sciences, The University of Melbourne, Level 12 Redmond Barry Building, Parkville 3010, Australia. *E-mail address:* vanja@rozenblat.net (V.Rozenblat)

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1002/erv.2530](https://doi.org/10.1002/erv.2530)

This work was supported by an Early Career Researcher Grant (1350035), an Australian Research Council Senior Research Fellowship (DP 130101459), and the Australian Postgraduate Award.

ABSTRACT

This study examined whether self-reported and observationally-measured parental behaviours were associated with disordered eating, and investigated possible moderation by a serotonin transporter polymorphism (*5-HTTLPR*). Study 1 included 650 adolescents from the Australian Temperament Project (ATP) who completed the EDI-2 Drive for Thinness and Bulimia scales at 15/16 years and were genotyped for *5-HTTLPR*. Parents completed an ATP-devised measure of parental warmth and harsh punishment. Study 2 included a sub-group of 304 participants who also engaged in a video-recorded family interaction, with observed parental warmth and hostility coded by the Iowa Family Interaction Rating Scale. Greater self-reported parental warmth was associated with lower bulimia scores. Conversely, observationally-measured parental warmth was associated with lower drive for thinness, but not bulimia. Self-reported parental harsh punishment was associated with bulimia only, with observed parental hostility associated with neither outcome. *5-HTTLPR* genotype did not moderate the relationship between parent behaviours and adolescent disordered eating.

Keywords: disordered eating; gene environment interactions; *5-HTTLPR*; parenting behaviours; observational measurement.

Relationships between self-reported and observed parenting behaviour, adolescent disordered eating attitudes and behaviours, and the *5-HTTLPR* polymorphism: Data from the Australian Temperament Project.

The biopsychosocial approach to eating disorder (ED) aetiology proposes that risk factors for EDs range from those relating to the individual, such as genes and psychological traits, to those that form part of the environment, including relationships with parents and peers, significant life events, exposure to the 'thin ideal', and numerous other factors (Culbert, Racine, & Klump, 2015; Stice, 2002; Trace, Baker, Penas-Lledo, & Bulik, 2013). Risk of developing an ED is believed to be associated with an interplay between these factors. For example, individuals exposed to 'risky' environments may develop ED symptoms only if they also carry certain 'risky' psychological or genetic factors (Stice, Marti, & Durant, 2011). In particular, following a lack of findings of direct genetic effects (Root et al., 2011), a growing area of ED research involves examining how individual genetic differences may moderate the impact of certain environmental variables on ED symptoms (Rozenblat et al., 2017).

Most research to date has approached investigation of gene x environment (GxE) interactions through the diathesis-stress lens, investigating how genetic 'vulnerability' may increase susceptibility to stressful environments (e.g., Stoltenberg, Anderson, Nag, & Anagnopoulos, 2012).

However, GxE investigations across other fields increasingly support an alternative model, under which certain genetic factors conceptualised as conferring 'risk' may be better conceptualised as 'plasticity' factors, that are associated with better outcomes under positive or neutral environmental conditions (Belsky et al., 2009). The current study will examine possible genetic plasticity in disordered eating attitudes and behaviours by examining adolescent exposure to both 'positive' and 'negative' parenting behaviours, using a multi-method multi-source approach.

The concept of genetic plasticity has received some early support in the depression field (Belsky & Pluess, 2009; Uher & McGuffin, 2008). Belsky and Pluess (2009) identified a number of studies in which participants with the short (s) allele of the serotonin transporter *5-HTTLPR* polymorphism, typically considered a 'risk' allele, exhibited lower depression under conditions of few or no stressful life events, compared to those homozygous for the long (l) allele (Brummett et al., 2008; Eley et al., 2004). However, recent meta-analyses cast some doubt over these earlier findings of plasticity (e.g., soon to be published results of a large meta-analysis described by Culverhouse et al., 2013).

5-HTTLPR has been the most heavily studied polymorphism in investigations of GxE interactions, with one or two copies of the s-allele resulting in reduced serotonin transcription (Heils et al., 1996). The s-allele has been implicated in changes in appetite, mood, and the stress-response system (Gotlib, Joormann, Minor, & Hallmayer, 2008; Leibowitz & Alexander, 1998; Ruhé, Mason, & Schene, 2007), and thus is of direct relevance to ED aetiology. As such, most GxE investigations in eating pathology have focussed on *5-HTTLPR*, with a recent meta-analysis finding significant interactions between *5-HTTLPR* and traumatic life events, as well as sexual and physical abuse, in

predicting EDs and bulimia nervosa (BN), respectively (Rozenblat et al., 2017). However, no study in the ED field has examined GxE 'risk' from a plasticity perspective.

Parenting factors, such as parental behaviour and child-parent relationship quality, have been implicated in EDs (Quiles, Quiles, Pamies, & Treasure, 2013; Tetzlaff, Schmidt, Brauhardt, & Hilbert, 2016). As parenting practices range from more positive to more negative, they provide an excellent opportunity for investigating the plasticity hypothesis of GxE interactions in relation to eating pathology. This may involve analysing two dimensionally opposite parenting behaviours, such as warmth and hostility (Schaefer, 1959), which have some theoretical and empirical support for a role in disordered eating. For example, compared to controls, questionnaire-based studies investigating families of ED patients have found families tend to be characterised by lower warmth and fewer positive family bonds (Calam, Waller, Slade, & Newton, 1990; Cunha, Relvas, & Soares, 2009; Vidovic, Juresa, Begovac, Mahnik, & Tocijl, 2004), but greater family conflict and parental control (Calam et al., 1990; Canetti, Kanyas, Lerer, Latzer, & Bachar, 2008). Furthermore, parenting styles (Baumrind, 1971) conceptualised as low on warmth and high on control and hostility (i.e., authoritarian parenting) have been associated with greater disordered eating compared to parenting styles not low on warmth (Lobera, Ríos, & Casals, 2011; Zubatsky, Berge, & Neumark-Sztainer, 2015). Parental warmth is, therefore, a possible protective factor for disordered eating while hostility or harsh punishment may constitute risk factors, although limited research specifically investigates the latter.

Given the importance of accurate measurement of environmental stimuli in GxE research (Moffitt, Caspi, & Rutter, 2005), a second approach may involve investigating parenting behaviours using observational techniques. Unlike idiographic self-report measurement (Margolin et al., 1998),

observational measures assess families using the same metric, and may overcome issues such as limitations in self-awareness (Aspland & Gardner, 2003). Other than studies specifically examining parental expressed emotion (Vaughn & Leff, 1976), very few studies have thus far investigated the relationship between (non-mealtime) parental behaviours and eating pathology using observational techniques, and most have not analysed parental warmth or hostility (these include: Blair, Freeman, & Cull, 1995; Humphrey, 1989; Kog & Vandereycken, 1989; Lattimore, Wagner, & Gowers, 2000; Ratti, Humphrey, & Lyons, 1996; Stasch & Reich, 2000; Thomas, Hoste, & Le Grange, 2012). Across these studies, findings regarding parental warmth and hostility have been mixed (e.g., Humphrey, 1989, cf. Lattimore et al., 2000). This inconsistency is unsurprising given that most existing observational studies are limited by use of poor, unestablished, or inappropriate instruments to code the observational data (e.g., Kog & Vandereycken, 1989; Lattimore et al., 2000), have used small to medium sample sizes (largest $N = 74$, in Humphrey, 1989), and have analysed female-only samples. Furthermore, no studies have examined the relationship between observed parent behaviours and sub-clinical eating pathology. There remains a need to establish solid conclusions regarding how parental warmth and hostility are related to disordered eating using reliable and consistent observational approaches, alongside self-report measures and use of large sample sizes. One further possibility is that genetics may, to an extent, account for some discrepancies in the literature.

To our knowledge, two studies (Karwautz et al., 2011; van Strien, Snoek, van der Zwaluw, & Engels, 2010) have analysed parenting within a GxE framework in the eating disorder field. One study ($N = 265$) found an interaction between problematic parenting styles, in particular parental control, and the s-allele of *5-HTTLPR* in predicting AN diagnosis in a female-only discordant AN sister-

pair sample (Karwautz et al., 2011). The second study ($N = 279$) found an interaction between parental psychological control and the A1 allele of the Taq1A polymorphism on *DRD2* in predicting greater emotional eating in a mixed-gender sample (van Strien et al., 2010). However, neither study explicitly tested for genetic plasticity. Furthermore, both studies focussed largely on parental control and did not shed light on the role of other parenting styles, including positive parent behaviours.

The present investigation aimed to assess the relationship between self-reported and observed parental behaviours and adolescent disordered eating, and to investigate whether *5-HTTLPR* moderated this relationship, using the largest sample to date. This is an important improvement on previous ED research, given the critical nature of sample size in genetic research (Duncan & Keller, 2011). The first study included self-reported measures of parenting behaviour to examine the direct effects of parental warmth and use of harsh punishment on disordered eating attitudes and behaviours, as well as possible moderation by *5-HTTLPR*. The second study aimed to replicate these analyses with the use of observationally-measured parenting behaviours, to provide insight into how observed parenting behaviours are related to adolescent disordered eating using a sample size multiple times larger than previously assessed, as well as aiming to provide further support for any GxEs identified in Study 1. These studies represent an advancement both in research relating to the role of parenting factors in EDs, which has included predominantly self-report measures, as well as GxE interactions in the ED field, which has focused on presence or absence of negative outcomes only. This is the first investigation of observed parenting within a GxE framework in the ED field, and inclusion of a mixed-gender community sample allows for results to inform ED prevention initiatives.

STUDY 1

METHOD

Participants

Australian Temperament Project participants were initially recruited in infancy (4-8 months) in Victoria in 1983 via maternal and child health centres in urban and rural locations. The first survey included 2,443 infants (48.0% female), with 16 surveys completed to date. The present study involved a sub-set of 650 participants (50.2% female), who had completed the 11th survey at age 15-16 years and self-identified as Caucasian. Participants provided information on disordered eating attitudes and behaviours, as well as a saliva sample for genotyping. One parent of each participant completed a parent-reported measure of parenting behaviours. To increase sample size for the present study, in 2015, 196 participants who had completed the survey but had not provided saliva for DNA extraction were invited to provide a saliva sample. Of 196 participants, 107 participants were successfully contacted and 83 individuals agreed to provide a sample. Parents and adolescents provided written informed consent for each survey wave and for the collection of saliva samples. The data collection was approved by the Australian Institute of Family Studies Ethics Review and carried out in accordance with the latest version of the Declaration of Helsinki. The final sample included a greater proportion of participants from the highest socio-economic (SES) quartile compared to the original sample in 1983 (see Table 1 for participant characteristics).

----- Include Table 1 about here -----

Measures

Disordered eating attitudes and behaviours. The Eating Disorder Inventory-2 (EDI-2; Garner, 1991) Drive for Thinness and Bulimia subscales assessed adolescent disordered eating. The Drive for Thinness subscale consists of seven items measuring participants' desire to lose weight or fear of weight gain (e.g., "I am preoccupied with the desire to be thinner") with Cronbach's $\alpha = .92$ in the current sample. The Bulimia subscale consists of 8 items measuring bulimic behaviours, including bingeing and purging (e.g., "I stuff myself with food"), with a somewhat lower internal consistency of $\alpha = .74$ in the present sample. Responses ranged from *Never* to *Always*, and were scored on the original 1 – 6 scale, as recommended for non-clinical samples (Schoemaker, van Strien, & van der Staak, 1994), and then averaged with higher mean scores reflecting greater disordered eating attitudes and behaviours. Wording of the EDI-2 was modified somewhat for an Australian audience, as detailed in Krug et al. (2016).

Parent behaviours: The ATP-devised Parenting Practices Scale (Prior, Sanson, Smart, & Oberklaid, 2000) was used to assess parental behaviours, with parents responding to five questions investigating parental warmth (e.g., "In general, how easy is it to spend time with your teenager?"), with $\alpha = .77$, and 10 questions regarding parental punishment (e.g., "I use threats of punishment to control him/her"), also $\alpha = .77$. Responses were recorded on a 5-point scale from *Always/Almost Always* to *Never*, and re-coded so higher scores reflected greater warmth or greater punishment.

5-HTTLPR genotype: DNA was isolated using Qiagen QIAamp kits from buccal epithelial cells via cotton swabs, and genotyping performed as described previously Jorm et al. (2000). The additional 83 saliva samples obtained for Study 2 were collected via Oragene saliva tubes, with DNA extracted and genotyped at the Australian Genomics Research Facility (AGRF), Adelaide, Australia. In

both cases, *5-HTTLPR* genotype was coded into s-present (s/s or s/l genotype) or s-absent (l/l genotype) groups.

Sociodemographics: Adolescent age, height in centimetres, and weight in kilograms were self-reported, allowing for calculation of participant body mass index (BMI; kg/m²). Socioeconomic status (SES) was calculated on the basis of parent-reported maternal and paternal education and employment status.

Data analyses

Using IBM SPSS Version 21, four multiple linear regression models separately assessed the direct effects of parental warmth and parental harsh discipline on participant EDI-2 Drive for Thinness and Bulimia scores, controlling for BMI and gender. Additional models tested for moderation by the *5-HTTLPR* polymorphism by including the gene x environment interaction terms. These models also controlled for the potential confounding effects of gender and BMI as per Keller (2014), by including all the covariate x gene and covariate x parenting style interaction terms in the regression models. Prior to analyses, missing data (23.5%) for the BMI variable were imputed using multiple imputation, with no systematic patterns of missingness observed.

Power analysis

Power analysis was conducted using Quanto (<http://biostats.usc.edu/Quanto.html>), with the following specifications: Continuous outcome, independent individuals design, and a dominant model with s-allele frequency of 43% (based on allele frequencies below). To detect a large to medium sized GxE interaction (R-sq = .02) at an alpha level of .05, a sample of 389 was required,

compared to the present sample of 650. However, this sample was not powered to detect small effects (i.e. $R^2 < .01$).

RESULTS

Descriptive statistics for participants in Study 1 are featured in Table 2. Participant *5-HTTLPR* genotype distribution (l/l = 197, s/l = 341, and s/s = 112) met the Hardy-Weinberg Equilibrium, $\chi^2 = 2.96$, $df = 1$, $p > .05$. Tables featuring results of all regression models are in Supplementary Materials.

----- Include Table 2 about here -----

Across all regression models examining main effects, there was a direct effect of female gender and BMI on Drive for Thinness scores, with the former also predicting Bulimia scores (all $p < .001$). There was a significant relationship between Bulimia, but not Drive for Thinness, and self-reported parental warmth ($\beta = .11$, $p = .005$) and harsh punishment ($\beta = .11$, $p = .003$). There was also an interaction between gender and harsh punishment in predicting Drive for Thinness ($\beta = .30$, $p = .038$), with scores for males lower under conditions of higher harsh punishment, while they remained largely unchanged for females. There were no main effects of *5-HTTLPR*, or any significant interaction effects between *5-HTTLPR* and parent behaviours.

STUDY 2

METHOD

Participants

Study 2 consisted of a sub-sample of 304 participants (53.0% female) from Study 1 who were selected to participate in an observational family interaction task based on responses to the 11th survey, as part of an earlier investigation into factors contributing to risk and resilience for

adolescent adjustment. Approximately three-quarters of those invited consented to participate, and were characterised into a problem (16.7%), high risk (21.7%), or low risk (55.2%) group. Participants in the problem group either exhibited elevated levels of depressed mood, as indicated by endorsing five or more DSM-III (APA, 1980) symptoms of depression, reported frequent substance use, or exhibited anti-social behaviour, determined by endorsing four or more delinquent acts (e.g., stealing, fighting, driving a car without permission). Based on a number of factors that differentiated the problem group from the main ATP sample (such as earlier behaviour problems, school adjustment difficulties, and peer relationships), a group of problem free yet 'high risk' participants was identified. A gender-balanced low risk group was drawn randomly from the remaining sample. Participants completed the EDI-2 and provided saliva samples as described in Study 1. The final sample for Study 2 included a lower proportion of participants from the lowest SES quartile compared to the original sample. Demographic information is featured in Table 1.

Measures

Disordered eating attitudes and behaviours, BMI, and age were measured as in Study 1.

Observational measure of parent behaviours: Trained interviewers performed home visits during which participants and one of their parents (usually the mother) engaged in a video-recorded 15-minute discussion based on a set of cards containing 14 questions about family life (e.g., teenager's accomplishments and disappointments, parental rules and fairness). Parenting behaviours were coded by a team who had undergone extensive training and were blind to adolescent group membership, using the Iowa Family Interaction Rating Scale (IFIRS; Melby et al., 1998), a macro-level observational coding system that was designed to measure behaviours and emotions in family discussions with adolescents. This is considered a valid tool for measuring

behaviours in a variety of dyadic interactions and has been validated against self and other reports from family members (Melby & Conger, 2001). Two scales were selected for the present study, parental warmth and parental hostility.

Warmth measured parental expressions of care, concern, and support directed at their child, including expressions of approval, affectionate physical contact, and building upon or reciprocating warmth displayed by their child. It was rated on a 9-point scale from *no warmth* to *frequent warmth*. Average intraclass correlation to assess inter-rater reliability for this scale was .80, based on cross examination of nine videos.

Hostility measured the extent to which parents engaged in hostile behaviours directed to their child, including rejection, active ignoring, showing contempt, or expressing complaints or critical remarks. It was rated on a 9-point scale from *no hostility* to *frequent hostility*. The average intraclass correlation for this scale was .75, which compares favourably to previous studies using the IFIRS and is deemed acceptable (Ge, Best, Conger, & Simons, 1996).

Data analyses

Data were analysed using IBM SPSS Version 21, as in Study 1, via four linear regression models examining the direct effects of observed parental warmth and observed parental hostility on EDI-2 Drive for Thinness and Bulimia scales. Additional models examined possible moderation by 5-HTTLPR, again controlling for gender and BMI. In the analyses investigating parental warmth and 5-HTTLPR, covariate x gene and covariate x environment contrast terms were included, as recommended by Keller (2014). However, this was not possible in the investigation of parental hostility and 5-HTTLPR due to issues of excessive collinearity (further discussed in Results section).

Finally, to ensure that any differences between Studies 1 and 2 reflected measurement effects as opposed to sample effects, the direct effects of self-reported parental warmth and self-reported use of harsh punishment were also tested in the Study 2 sub-sample. GxE interactions were not included in this analysis to avoid multiple testing issues. Missing data for BMI (28.29%) were imputed via multiple imputation, with no systematic patterns of missingness observed.

Power analysis

Power analysis was conducted using Quanto (<http://biostats.usc.edu/Quanto.html>), as for Study 1, with s-allele frequency of 43% (based on allele frequencies for Study 2 participants). The current sample was powered to detect a large to medium sized GxE interaction, $R^2 = .03$, which requires 258 participants at an alpha level of .05, but was not powered to detect effects smaller than this.

RESULTS

Descriptive statistics for participants in Study 2 are featured in Table 3. *5-HTTLPR* genotype distribution (l/l = 90, s/l = 154, and s/s = 60) met the Hardy-Weinberg Equilibrium, $\chi^2 = .16$, $df = 1$, $p > .05$. Results from all regression models are located in the Supplementary Materials.

----- Insert Tables 5 about here -----

Female gender predicted greater Drive for Thinness and Bulimia scores (all $p < .001$), while BMI was associated with the former only. There were also significant direct effects of greater

observed parental warmth on reduced Drive for Thinness ($\beta = -.10, p = .029$), but not Bulimia, and no direct effects of observed parental hostility. No significant main effects of *5-HTTLPR* or interaction between *5-HTTLPR* and observed parenting behaviours were observed.

The models investigating moderation by *5-HTTLPR* in the relationship between observed parental hostility and disordered eating do not include gene x covariate and environment x covariate contrast terms due to issues with collinearity. Although Keller (2014) argues collinearity functions to control for alternate explanations of any GxE interaction, the gene x environment interaction showed minimal change regardless of whether the additional contrast terms were included in the models. Therefore, the simpler model less affected by collinearity is presented.

The follow-up analysis investigating self-reported parental warmth and use of harsh punishment in the present sub-sample revealed main effects of the warmth and punishment on Bulimia scores ($\beta = -.14, p = .013$ and $\beta = .14, p = .011$, respectively) but not Drive for Thinness scores. These results mirrored those of the larger sample in Study 1.

DISCUSSION

We adopted a multi-method assessment approach in two studies to investigate the relationship between parental behaviours (warmth, harsh punishment, and hostility) and disordered eating attitudes and behaviours. We also investigated whether the relationship between parenting factors and eating pathology was moderated by the serotonin transporter *5-HTTLPR* polymorphism. Results indicated that greater parent-reported warmth was associated with lower bulimia symptoms, and in concordance, parent-reported use of harsh punishment techniques was associated with greater bulimia symptoms. Conversely, greater observed parental warmth was

associated with lower adolescent drive for thinness, while greater observed parental hostility did not predict either disordered eating outcome. No moderation by *5-HTTLPR* was identified in any of these relationships.

Parental warmth and harsh punishment

Findings of a relationship between parental warmth and use of harsh punishment techniques with disordered eating attitudes and behaviours supports the results of past studies that have assessed these parental behaviours via questionnaires. For example, parental warmth and care have been identified as lower in families of patients with AN and BN compared to controls (Tetley, Moghaddam, Dawson, and Rennoldson, 2014), while parental harsh punishment has been associated with BN in a number of studies (Rorty, Yager, & Rossotto, 1995; Stuart et al., 1990), with these patterns of findings also reflected in studies of child development more broadly (e.g., Bender et al., 2007; Yap, Pilkington, Ryan, & Jorm, 2014).

However, few studies have investigated these associations using observed parenting behaviours, as we have here, and a novel pattern of results was identified compared with research based on questionnaires. Across both samples, the two self-reported measures of parenting behaviours were clearly associated with adolescent bulimia symptoms, but neither was related to drive for thinness. Conversely, the observed parenting behaviours tended to be associated with drive for thinness but not bulimia. Other studies have also found limited correspondence between parent responses on questionnaires and their actual observed behaviours (e.g., Kallstrom-Fuqua, 2004). Such discrepancies are not unexpected for a number of reasons, including socially desirable or idiosyncratic responding to a questionnaire, or issues of ecological validity during observational tasks. It is also possible that the two measures of parental warmth used in the present study tapped

into slightly different constructs. The observational measurement may have had a stronger focus on caring and warm behaviour displayed by the parent, irrespective of the child's behaviour, while the questionnaire measure perhaps reflected more greatly the level of general positive interaction between parent and child, albeit from the parent's perspective. This may suggest that the precise relationship between parental behaviours and adolescent disordered eating is highly nuanced and varies according to the specific element of parenting or disordered eating measured.

Gene x Environment Interactions

The lack of significant GxE interactions identified in the present study contrasts with Karwautz et al. (2011) ($N = 256$), who found that problematic parenting styles interacted with the s-allele of 5-HTTLPR to predict AN diagnosis in a discordant sister-pair sample. One possibility is that the influence of 5-HTTLPR may only be evident when clinical-level disordered eating pathology, such as AN, is considered. Another consideration is that Karwautz et al. (2011) examined parental control, a different parental behaviour to those examined in the present study. The other study identifying a possible interaction between parenting and genes ($p = 0.06$) in the ED field also investigated parental control (van Strien et al., 2010) ($N = 276$). One possibility therefore is that parental control but not other parenting behaviours may interact with genetic factors to predict eating pathology. However, it remains possible that discrepant findings between the current study and previous investigations are due to low power in both Karwautz et al. (2011) and van Strien et al. (2010), which increases the chances of false positive findings (Duncan, Pollastri, & Smoller, 2014). Sample sizes required for genetic association studies are very large due to the small effect sizes purportedly involved, and the present sample size ($N = 650$) was more than double that of the previous two studies identifying GxE interactions involving parenting in EDs. Similarly, numerous past studies that have identified

plasticity related to the *5-HTTLPR* s-allele in other fields, and indeed have been cited as early evidence of this effect, have been vastly underpowered. A large portion of these studies included fewer than one hundred and fifty participants (e.g., Kim, 2010; Pluess, Belsky, Way, & Taylor, 2010; Taylor et al., 2006; Wilhelm et al., 2006).

The lack of moderation by *5-HTTLPR* in the relationship between parental warmth and bulimic behaviours and drive for thinness supports the view that this polymorphism does not act as a plasticity allele in the presence of family environments typically considered 'protective'. This is also reflected in the fact that, to the authors' knowledge, no prior study investigating GxEs in the ED field has reported differential susceptibility in response to positive environments (distinct from absence of negative stressors). In other fields, parental warmth and *5-HTTLPR* have not been heavily studied within a GxE framework, with the few existing studies presenting a mix of results (e.g., Hankin et al. 2011, Kochanska, Kim, Barry, & Philibert, 2011). Given the large-scale publication bias that purportedly affects studies of GxE interactions (Duncan & Keller, 2011), it cannot be ascertained how many other unpublished investigations have failed to find GxE effects involving parental warmth.

Strengths and Limitations

The present paper contributes to research examining parenting factors in disordered eating by analysing a large-scale community sample, with a focus on replication across two samples using multi-method multi-source assessment. In contrast, past studies have largely adopted questionnaire-based measures, often retrospective, with limited investigation using observational measurement of parenting behaviours. A few limitations of the present study must also be noted. The present study did not examine comorbid psychopathology, so was unable to assess whether this factor may possibly mediate the identified relationships between particular parenting behaviours and

disordered eating outcomes. The present study also examined the biallelic model of *5-HTTLPR*. There is some evidence to suggest that examination of a triallelic model may better reflect *5-HTTLPR* activity (Wendland et al., 2006), with support from some studies in the ED field (Steiger et al., 2009; Stoltenberg, Anderson, Nag, & Anagnopoulos, 2012 but not Richardson et al., 2008), although more broadly findings have been tentative (e.g., Uher & McGuffin, 2008). In the present study data were not available to examine the triallelic model, which may yet have resulted in identification of a GxE interaction in the present sample. Furthermore, the sample size, particularly in Study 2, was relatively underpowered for genetic association studies (although it was above the mean $N = 288$ of GxE studies in the ED field; Rozenblat et al., 2017). Conversely, the sample in Study 1 constitutes the third largest investigation of its kind in the field (following Akkermann et al. 2012, $N = 765$, and the combined-samples analyses in Rozenblat et al., 2017, $N > 1,000$). This is an important strength. Furthermore, assessing disordered eating attitudes and behaviours dimensionally increases power and sensitivity compared to case-control designs, in which the 'clinical' group may represent heterogeneous underlying patterns of eating pathology or genetic influences (Abbott, 2008). Assessing disordered eating symptoms in a population-based sample also increases generalisability of results and utility for informing prevention and early intervention initiatives, although may limit the applicability of results to clinical populations. Finally, as with all studies of candidate genes, *5-HTTLPR* represents only one polymorphism in a system of genetic factors likely to be involved in ED pathology, and thus such investigations constitute stepping stones towards increasing our understanding of the genetic underpinnings of EDs.

Clinical Implications and Future Directions

Results support a clear relationship between parent behaviours and adolescent disordered eating, across both genders, independent of individual genetic predisposition. Findings can inform parenting programs targeting ED prevention or psychoeducation for 'at risk' families. For example, one systematic review of parent-focussed ED prevention initiatives stated early positive findings required support from additional research regarding the family setting (Hart, Cornell, Damiano, & Paxton, 2015). Adoption of longitudinal designs would also help establish whether parental behaviours constitute protective/risk factors, or are better conceptualised as correlates of eating pathology. Furthermore, investigations into genetic plasticity could also expand beyond *5-HTTLPR* to examine cumulative plasticity across numerous polymorphisms testing multiple positive environmental factors. Selection of positive eating-related outcomes, such as positive body image (Webb, Wood-Barcalow, & Tylka, 2015), as opposed to absence of eating-related pathology, may also be more appropriate for reflecting the relationship between genetic plasticity and environmental factors. However, it is also possible that regardless of precise measurement techniques, the role of *5-HTTLPR* in EDs may prove to be very minor at best, such that continued investigation into parenting and other environmental factors may constitute a more fruitful approach to untangling the complex aetiology of eating pathology.

Acknowledgements: The ATP study is located at The Royal Children's Hospital Melbourne and is a collaboration between Deakin University, The University of Melbourne, the Australian Institute of Family Studies, The University of New South Wales, The University of Otago (NZ), and the Royal Children's Hospital; further information available at www.aifs.gov.au/atp. The views expressed in this paper are those of the authors and may not reflect those of their organisational affiliations, nor of other collaborating individuals or organisations. We acknowledge all collaborators who have contributed to the Australian Temperament Project, especially Professors Ann Sanson, Margot Prior, Frank Oberklaid, John Toumbourou and Ms Diana Smart. We would also like to sincerely thank the participating families for their time and invaluable contribution to the study. This paper forms part of Vanja Rozenblat's PhD with publication undertaken at The University of Melbourne.

Conflicts of interest: The authors declare no conflicts of interest.

References

- Abbott, A. (2008). Psychiatric genetics: The brains of the family. *Nature*, 454(7201), 154-157.
- Akkermann, K., Kaasik, K., Kiive, E., Nordquist, N., Orelund, L., & Harro, J. (2012). The impact of adverse life events and the serotonin transporter gene promotor polymorphism on the development of eating disorder symptoms. *Journal of Psychiatric Research*, 46, 38 – 43.
- American Psychiatric Association (APA). (1980). Diagnostic and statistical manual of mental disorders. 3rd. Washington, DC: Author.
- Aspland, H., & Gardner, F. (2003). Observational measures of parent-child interaction: An introductory review. *Child and Adolescent Mental Health*, 8(3), 136-143.
- Baumrind, D. (1971). Current patterns of parental authority. *Developmental Psychology*, 4(1p2), 1.
- Belsky, J., Jonassaint, C., Pluess, M., Stanton, M., Brummett, B., & Williams, R. (2009). Vulnerability genes or plasticity genes? *Molecular Psychiatry*, 14(8), 746-754. doi:10.1038/mp.2009.44
- Belsky, J., & Pluess, M. (2009). Beyond diathesis stress: differential susceptibility to environmental influences. *Psychological Bulletin*, 135(6), 885-908. doi:10.1037/a0017376
- Bender, H. L., Allen, J. P., McElhaney, K. B., Antonishak, J., Moore, C. M., Kelly, H. O. B., & Davis, S. M. (2007). Use of harsh physical discipline and developmental outcomes in adolescence. *Development and Psychopathology*, 19(01), 227-242.
- Blair, C., Freeman, C., & Cull, A. (1995). The families of anorexia nervosa and cystic fibrosis patients. *Psychological Medicine*, 25(05), 985-993.
- Brummett, B. H., Boyle, S. H., Siegler, I. C., Kuhn, C. M., Ashley-Koch, A., Jonassaint, C. R., . . . Williams, R. B. (2008). Effects of environmental stress and gender on associations among symptoms of depression and the serotonin transporter gene linked polymorphic region (5-HTTLPR). *Behavior Genetics*, 38(1), 34-43.

- Calam, R., Waller, G., Slade, P., & Newton, T. (1990). Eating disorders and perceived relationships with parents. *International Journal of Eating Disorders*, 9(5), 479-485.
- Canetti, L., Kanyas, K., Lerer, B., Latzer, Y., & Bachar, E. (2008). Anorexia nervosa and parental bonding: the contribution of parent–grandparent relationships to eating disorder psychopathology. *Journal of Clinical Psychology*, 64(6), 703-716.
- Culbert, K. M., Racine, S. E., & Klump, K. L. (2015). Research Review: What we have learned about the causes of eating disorders - a synthesis of sociocultural, psychological, and biological research. *Journal of Child Psychology and Psychiatry*, 56(11), 1141-1164.
- Culverhouse, R. C., Bowes, L., Breslau, N., Nurnberger Jr, J. I., Burmeister, M., Fergusson, D. M., . . . Bierut, L. J. (2013). Protocol for a collaborative meta-analysis of 5-HTTLPR, stress, and depression. *BMC Psychiatry*, 13(1), 304.
- Cunha, A. I., Relvas, A. P., & Soares, I. (2009). Anorexia nervosa and family relationships: Perceived family functioning, coping strategies, beliefs, and attachment to parents and peers. *International Journal of Clinical and Health Psychology*, 9(2), 229-240.
- Deas, S., Power, K., Collin, P., Yellowlees, A., & Grierson, D. (2011). The relationship between disordered eating, perceived parenting, and perfectionistic schemas. *Cognitive Therapy and Research*, 35(5), 414-424.
- Duncan, L. E., & Keller, M. C. (2011). A Critical Review of the First 10 Years of Candidate Gene-by-Environment Interaction Research in Psychiatry. *American Journal of Psychiatry*, 168, 1041 - 1049.
- Duncan, L. E., Pollastri, A. R., & Smoller, J. W. (2014). Mind the gap: why many geneticists and psychological scientists have discrepant views about gene-environment interaction (GxE) research. *American Psychologist*, 69(3), 249-268.

- Eley, T. C., Sugden, K., Corsico, A., Gregory, A. M., Sham, P., McGuffin, P., Plomin, R., & Craig, I. W. (2004). Gene–environment interaction analysis of serotonin system markers with adolescent depression. *Molecular Psychiatry*, 9(10), 908-915.
- Garner, D. M., L. (1991). *Eating Disorders Inventory-2 Professional Manual*. Odessa, F: Psychological Assessment Resources, Inc.
- Ge, X., Best, K. M., Conger, R. D., & Simons, R. L. (1996). Parenting behaviors and the occurrence and co-occurrence of adolescent depressive symptoms and conduct problems. *Developmental Psychology*, 32(4), 717.
- Gotlib, I. H., Joormann, J., Minor, K. L., & Hallmayer, J. (2008). HPA axis reactivity: a mechanism underlying the associations among 5-HTTLPR, stress, and depression. *Biological Psychiatry*, 63(9), 847-851.
- Hankin, B., Nederhof, E., Oppenheimer, C., Jenness, J., Young, J., Abela, J., . . . Oldehinkel, A. (2011). Differential susceptibility in youth: evidence that 5-HTTLPR x positive parenting is associated with positive affect ‘for better and worse’. *Translational Psychiatry*, 1(10), e44.
- Hart, L. M., Cornell, C., Damiano, S. R., & Paxton, S. J. (2015). Parents and prevention: a systematic review of interventions involving parents that aim to prevent body dissatisfaction or eating disorders. *International Journal of Eating Disorders*, 48(2), 157-169.
- Heils, A., Teufel, A., Petri, S., Stöber, G., Riederer, P., Bengel, D., & Lesch, K. P. (1996). Allelic variation of human serotonin transporter gene expression. *Journal of Neurochemistry*, 66(6), 2621-2624.
- Humphrey, L. L. (1989). Observed family interactions among subtypes of eating disorders using structural analysis of social behavior. *Journal of Consulting and Clinical Psychology*, 57(2), 206.

- Jáuregui Lobera, I., Bolaños Ríos, P., & Garrido Casals, O. (2011). Parenting styles and eating disorders. *Journal of Psychiatric and Mental Health Nursing*, 18(8), 728-735.
- Jorm, A., Prior, M., Sanson, A., Smart, D., Zhang, Y., & Easteal, S. (2000). Association of a functional polymorphism of the serotonin transporter gene with anxiety-related temperament and behavior problems in children: a longitudinal study from infancy to the mid-teens. *Molecular Psychiatry*, 5(5), 542-547.
- Kallstrom-Fuqua, A. C. (2004). Examining parenting outcomes of childhood sexual abuse survivors utilizing observation and self-report methods. *Doctoral Dissertation, The University of North Texas*.
- Karwautz, A. F., Wagner, G., Waldherr, K., Nader, I. W., Fernandez-Aranda, F., Estivill, X., Holliday, J., Collier, D. A., & Treasure, J. L. (2011). Gene-environment interaction in anorexia nervosa: relevance of non-shared environment and the serotonin transporter gene. *Molecular Psychiatry*, 16(6), 590-592.
- Keller, M. C. (2014). Gene x environment interaction studies have not properly controlled for potential confounders: the problem and the (simple) solution. *Biological Psychiatry*, 75(1), 18-24.
- Kim, S. (2010). *Children's Genotypes Interact with Maternal Responsive Care in Predicting Competencies and Problems: Diathesis-Stress or Differential Susceptibility?* Paper presented at the 57th Annual Meeting (October 26-31, 2010).
- Kochanska, G., Kim, S., Barry, R. A., & Philibert, R. A. (2011). Children's genotypes interact with maternal responsive care in predicting children's competence: Diathesis-stress or differential susceptibility? *Development and Psychopathology*, 23(2), 605.

- Kog, E., & Vandereycken, W. (1989). Family interaction in eating disorder patients and normal controls. *International Journal of Eating Disorders*, 8(1), 11-23.
- Krug, I., King, R. M., Youssef, G. J., Sorabji, A., Wertheim, E. H., Le Grange, D., Hughes, E. K., Letcher, P., & Olsson, C. A. (2016). The effect of low parental warmth and low monitoring on disordered eating in mid-adolescence: Findings from the Australian Temperament Project. *Appetite*, 105, 232-241.
- Lattimore, P. J., Wagner, H. L., & Gowers, S. (2000). Conflict avoidance in anorexia nervosa: An observational study of mothers and daughters. *European Eating Disorders Review*, 8(5), 355-368.
- Leibowitz, S. F., & Alexander, J. T. (1998). Hypothalamic serotonin in control of eating behavior, meal size, and body weight. *Biological Psychiatry*, 44(9), 851-864.
- Margolin, G., Oliver, P. H., Gordis, E. B., O'hearn, H. G., Medina, A. M., Ghosh, C. M., & Morland, L. (1998). The nuts and bolts of behavioral observation of marital and family interaction. *Clinical Child and Family Psychology Review*, 1(4), 195-213.
- Melby, J., Conger, R., Book, R., Rueter, M., Lucy, L., Repinski, D., & Scaramella, L. (1998). *The Iowa Family Interaction Rating Scales*. Ames: Iowa State University. Institute for Social and Behavioral Research.
- Melby, J. N., & Conger, R. D. (2001). *The Iowa Family Interaction Rating Scales: Instrument summary*. Mahwah, NJ, US: Lawrence Erlbaum Associates Publishers.
- Moffitt, T. E., Caspi, A., & Rutter, M. (2005). Strategy for investigating interactions between measured genes and measured environments. *Archives of General Psychiatry*, 62(5), 473-481.

- Pluess, M., Belsky, J., Way, B. M., & Taylor, S. E. (2010). 5-HTTLPR moderates effects of current life events on neuroticism: differential susceptibility to environmental influences. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 34(6), 1070-1074.
- Prior, M., Sanson, A., Smart, D., & Oberklaid, F. (2000). Pathways from infancy to adolescence: Australian Temperament Project 1983–2000. *Melbourne, Australia: Australian Institute of Family Studies*.
- Quiles, M. Y., Quiles, S. M., Pamies, A. L., J. B. A., & Treasure, J. (2013). Peer and family influence in eating disorders: A meta-analysis. *European Psychiatry*, 28(4), 199-206.
- Ratti, L. A., Humphrey, L. L., & Lyons, J. S. (1996). Structural analysis of families with a polydrug-dependent, bulimic, or normal adolescent daughter. *Journal of Consulting and Clinical Psychology*, 64(6), 1255.
- Richardson, J., Steiger, H., Schmitz, N., Joober, R., Bruce, K. R., Israel, M., Gauvin, L., Anestin, A. S., Dandurand, C., Howard, H., & de Guzman, R. (2008). Relevance of the 5-HTTLPR polymorphism and childhood abuse to increased psychiatric comorbidity in women with bulimia-spectrum disorders. *The Journal of Clinical Psychiatry*, 69(6), 981-990.
- Root, T. L., Szatkiewicz, J. P., Jonassaint, C. R., Thornton, L. M., Pinheiro, A. P., Strober, M., ... & Bulik, C. M. (2011). Association of candidate genes with phenotypic traits relevant to anorexia nervosa. *European Eating Disorders Review*, 19(6), 487-493.
- Rorty, M., Yager, J., & Rossotto, E. (1995). Aspects of childhood physical punishment and family environment correlates in bulimia nervosa. *Child Abuse & Neglect*, 19(6), 659-667.
- Rozenblat, V., Ong, D., Fuller-Tyszkiewicz, M., Akkermann, K., Collier, D., Engels, R. C., . . . Karwautz, A. (2017). A systematic review and secondary data analysis of the interactions between the

- serotonin transporter 5-HTTLPR polymorphism and environmental and psychological factors in eating disorders. *Journal of Psychiatric Research*(84), 62 - 72.
- Ruhé, H. G., Mason, N. S., & Schene, A. H. (2007). Mood is indirectly related to serotonin, norepinephrine and dopamine levels in humans: a meta-analysis of monoamine depletion studies. *Molecular Psychiatry*, 12(4), 331-359.
- Schaefer, E. S. (1959). A circumplex model for maternal behavior. *The Journal of Abnormal and Social Psychology*, 59(2), 226.
- Schoemaker, C., van Strien, T., & van der Staak, C. (1994). Validation of the Eating Disorders Inventory in a nonclinical population using transformed and untransformed responses. *International Journal of Eating Disorders*, 15(4), 387-393.
- Stasch, M., & Reich, G. (2000). Interpersonal relation pattern in families with a bulimic member- interaction analysis. *Praxis der Kinderpsychologie und Kinderpsychiatrie*, 49(3), 157-175.
- Steiger, H., Richardson, J., Schmitz, N., Joober, R., Israel, M., Bruce, K. R., Gauvin, L., Dandurand, C., & Anestis, A. (2009). Association of trait-defined, eating-disorder sub-phenotypes with (biallelic and triallelic) 5HTTLPR variations. *Journal of Psychiatric Research*, 43(13), 1086-1094.
- Stice, E. (2002). Risk and maintenance factors for eating pathology: a meta-analytic review. *Psychological bulletin*, 128(5), 825 - 848.
- Stice, E., Marti, C. N., & Durant, S. (2011). Risk factors for onset of eating disorders: Evidence of multiple risk pathways from an 8-year prospective study. *Behaviour Research and Therapy*, 49(10), 622-627.

- Stoltenberg, S. F., Anderson, C., Nag, P., & Anagnopoulos, C. (2012). Association between the serotonin transporter triallelic genotype and eating problems is moderated by the experience of childhood trauma in women. *Int J Eat Disord*, 45(4), 492-500.
- Stuart, G. W., Laraia, M. T., Ballenger, J. C., & Lydiard, R. B. (1990). Early family experiences of women with bulimia and depression. *Archives of Psychiatric Nursing*, 4(1), 43-52.
- Taylor, S. E., Way, B. M., Welch, W. T., Hilmert, C. J., Lehman, B. J., & Eisenberger, N. I. (2006). Early family environment, current adversity, the serotonin transporter promoter polymorphism, and depressive symptomatology. *Biological Psychiatry*, 60(7), 671-676.
- Tetley, A., Moghaddam, N. G., Dawson, D. L., & Rennoldson, M. (2014). Parental bonding and eating disorders: A systematic review. *Eating Behaviors*, 15(1), 49-59.
- Tetzlaff, A., Schmidt, R., Brauhardt, A., & Hilbert, A. (2016). Family functioning in adolescents with binge-eating disorder. *European Eating Disorders Review*, 24(5), 430-433.
- Thomas, S. A., Hoste, R. R., & Le Grange, D. (2012). Observed connection and individuation: Relation to symptoms in families of adolescents with bulimia nervosa. *International Journal of Eating Disorders*, 45(7), 891-899.
- Trace, S. E., Baker, J. H., Penas-Lledo, E., & Bulik, C. M. (2013). The genetics of eating disorders. *Annual Review of Clinical Psychology*, 9, 589-620.
- Uher, R., & McGuffin, P. (2008). The moderation by the serotonin transporter gene of environmental adversity in the aetiology of mental illness: review and methodological analysis. *Molecular Psychiatry*, 13(2), 131-146.
- van Strien, T., Snoek, H. M., van der Zwaluw, C. S., & Engels, R. C. (2010). Parental control and the dopamine D2 receptor gene (DRD2) interaction on emotional eating in adolescence. *Appetite*, 54(2), 255-261.

- Vaughn, C., & Leff, J. (1976). The measurement of expressed emotion in the families of psychiatric patients. *British Journal of Clinical Psychology*, 15(2), 157-165.
- Vidovic, V., Juresa, V., Begovac, I., Mahnik, M., & Tocji, G. (2004). Perceived family cohesion, adaptability and cohesion in eating disorders. *European Eating Disorders Review*, 13(1), 19-28.
- Volkow, N. D., Wang, G.-J., Maynard, L., Jayne, M., Fowler, J. S., Zhu, W., Logan, J., Gatley, S. J., Ding, Y. S., Wong, C., & Pappas, N. (2003). Brain dopamine is associated with eating behaviors in humans. *International Journal of Eating Disorders*, 33(2), 136-142.
- Wendland, J. R., Martin, B. J., Kruse, M. R., Lesch, K. P., & Murphy, D. K. (2006). Simultaneous genotyping of four functional loci of human *SLC6A4*, with a reappraisal of 5-HTTLPR and rs25531. *Molecular Psychiatry*, 11, 224 – 226.
- Webb, J. B., Wood-Barcalow, N. L., & Tylka, T. L. (2015). Assessing positive body image: Contemporary approaches and future directions. *Body Image*, 14, 130-145.
- Wilhelm, K., Mitchell, P. B., Niven, H., Finch, A., Wedgwood, L., Scimone, A., Blair, I. P., Parker, G., Schofield, P. R. (2006). Life events, first depression onset and the serotonin transporter gene. *The British Journal of Psychiatry*, 188(3), 210-215.
- Yap, M. B. H., Pilkington, P. D., Ryan, S. M., & Jorm, A. F. (2014). Parental factors associated with depression and anxiety in young people: A systematic review and meta-analysis. *Journal of Affective Disorders*, 156, 8-23.
- Zubatsky, M., Berge, J., & Neumark-Sztainer, D. (2015). Longitudinal associations between parenting style and adolescent disordered eating behaviors. *Eating and Weight Disorders-Studies on Anorexia, Bulimia and Obesity*, 20(2), 187-194.

Table 1.

Sociodemographic details of participants included in Study 1 and Study 2.

	Full Sample			Females		Males	
	<i>N</i>	<i>N</i>	%	<i>N</i>	%	<i>N</i>	%
Study 1							
Sex	650						
Males		312	49.8				
Females		338	50.2				
SES Quartile	632						
Highest		231	36.6	116	36.6	115	36.5
Medium-High		197	31.2	103	32.5	94	29.8
Medium-Low		129	20.4	59	18.6	70	22.2
Lowest		75	11.9	39	12.3	36	11.4
Parent Marital Status	645						
Married/Defacto		538	83.4	264	81.5	274	85.4
Separated/Divorced		73	11.3	42	13.0	31	9.7
Single/Widowed		22	3.4	10	3.1	12	3.7
Remarried		12	1.8	8	2.5	4	1.2
Study 2							
Sex	304						
Males		143	47.0				
Females		161	53.9				
SES Quartile	296						
Highest		103	34.8	53	33.3	50	36.5
Medium-High		100	33.8	55	34.6	45	32.8
Medium-Low		61	20.6	31	19.5	30	21.9
Lowest		32	10.8	20	12.6	12	8.8
Parent Marital Status	299						
Married/Defacto		242	81.0	126	80.3	116	81.7

Separated/Divorced	37	12.4	17	10.8	20	14.1
Single/Widowed	12	4.0	8	5.1	4	2.8
Remarried	8	2.7	2	3.8	2	1.4

Author Manuscript

Table 2.

Descriptive statistics for key variables investigated in Study 1.

Variable	Full Sample	Mean (<i>SD</i>)	
		Females	Males
Age (years)	15.72 (0.16)	15.74 (0.14)	15.72 (0.17)
Parental warmth	4.22 (0.63)	4.25 (0.65)	4.19 (0.61)
Parental punishment	2.00 (0.56)	1.94 (0.55)	2.06 (0.57)
EDI-2 Bulimia	1.78 (0.65)	1.94 (0.73)	1.62 (0.53)
EDI-2 Drive for Thinness	2.23 (1.15)	2.81 (1.22)	1.64 (0.70)
BMI	21.27 (3.28)	21.20 (3.10)	21.34 (3.44)

Table 3.

Descriptive statistics for key variables investigated in Study 2.

Variable	Mean (<i>SD</i>)	Mean (<i>SD</i>)	
		Females	Males
Age (years)	15.73 (0.15)	15.73 (0.13)	15.72 (0.10)
Parental warmth	4.38 (1.93)	4.57 (1.92)	4.16 (1.92)
Parental hostility	1.92 (1.53)	2.06 (1.65)	1.76 (1.37)
EDI-2 Bulimia	1.89 (0.73)	2.08 (0.80)	1.68 (0.58)
EDI-2 Drive for Thinness	2.34 (1.25)	2.99 (1.28)	1.61 (0.67)
BMI	21.55 (3.59)	21.65 (3.40)	21.45 (3.77)

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Rozenblat, V;Ryan, J;Wertheim, E;King, R;Olsson, CA;Letcher, P;Krug, I

Title:

Relationships Between Self-Reported and Observed Parenting Behaviour, Adolescent Disordered Eating Attitudes and Behaviours, and the 5-HTTLPR Polymorphism: Data From the Australian Temperament Project

Date:

2017-09-01

Citation:

Rozenblat, V., Ryan, J., Wertheim, E., King, R., Olsson, C. A., Letcher, P. & Krug, I. (2017). Relationships Between Self-Reported and Observed Parenting Behaviour, Adolescent Disordered Eating Attitudes and Behaviours, and the 5-HTTLPR Polymorphism: Data From the Australian Temperament Project. EUROPEAN EATING DISORDERS REVIEW, 25 (5), pp.381-388. <https://doi.org/10.1002/erv.2530>.

Persistent Link:

<http://hdl.handle.net/11343/293167>