DR. MONICA WONG (Orcid ID : 0000-0003-2510-8854)

Article type : Review Article

Use of time and adolescent health-related quality of life/wellbeing:

A scoping review

Authors: Monica Wong (MD),^{1,2} Kate Lycett (PhD),² Tim Olds (PhD),^{2,3} Lisa Gold (PhD),^{2,4} Melissa Wake (MD),^{1,2,4}

Affiliations: ¹The University of Melbourne, Melbourne, Australia; ²Murdoch Childrens Research Institute, Melbourne, Australia; ³University of South Australia, Adelaide, Australia; ⁴Deakin University, Geelong, Australia; ⁵Department of Paediatrics and the Liggins Institute, The University of Auckland, New Zealand.

Short title:

Time use and adolescent quality of life

Correspondence to:

Professor Melissa Wake, Department of Paediatrics: Child & Youth Health, Auckland City Hospital, Grafton, Auckland, NEW ZEALAND. Phone: +64 9 923 7690,

Email: melissa.wake@auckland.ac.nz

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/apa.13929](https://doi.org/10.1111/apa.13929)

This article is protected by copyright. All rights reserved

1 ABSTRACT

2 Time use could profoundly affect adolescents' health-related quality of life (HRQL). Ideally, overall
3 time use patterns would be considered, because activities within a 24-hour day are inherently
4 correlated (more in one activity means less in another). This review focused on the associations of 1)
5 overall time use patterns and 2) components of time use patterns with HRQL in adolescents.

6 Conclusion:

7 ~~More~~ physical activity, less screen time and more/adequate sleep, in isolation, are associated with
8 better profile-based HRQL subscales. Greater understanding of adolescents' overall time use patterns
9 and HRQL is a priority for policy development.

10 **Keywords:** Adolescent, lifestyle, quality of life, scoping review, time use

11 KEYNOTES

- 12 • Overall patterns of time use could profoundly affect adolescents' current and future health-related
- 13 quality of life (HRQL).
- 14 • This review identified many studies examining specific aspects of time use and HRQL, with
- 15 better outcomes associated with more physical activity, less screen time and more sleep/absence
- 16 of sleep deprivation.
- 17 • Large-scale representative surveys are needed that incorporate valid measures of all major time
- 18 use components simultaneously alongside utility- and profile-based HRQL.

19 INTRODUCTION

20 The 2016 Lancet commission has centred global attention on the health and wellbeing of adolescents
21 (10-19 years) (1). Adolescence is typically considered a time of optimal health, yet it directly accounts
22 for 35% of the global burden of disease, mostly from non-communicable diseases such as obesity and
23 poor mental health (2). It is also a crucial transition phase (2, 3) when health patterns are established
24 and track into adulthood (4).

25 How adolescents spend their time has been shown to impact on certain aspects of health (5). "Health"
26 encompasses health status and functional status, as well as quality of life (6). Encompassed within
27 quality of life is health-related quality of life (HRQL), i.e., quality of life in the context of one's health
28 and/or illness (7). HRQL can be used as an indicator to guide healthcare resource allocation (8), as it

can be measured at the population level across all life stages. Therefore, it is important to understand whether potentially-modifiable time use patterns (i.e., general nature and timing of daily activities (5)) in adolescence are key determinants of HRQL.

Time use potentially has a bidirectional relationship with health: health influences time use, while time use can improve or reduce health. Current health promotion strategies recommend well-balanced lifestyles in children for optimal health (9, 10), but what a well-balanced lifestyle looks like for an adolescent remains unclear. A comprehensive insight into time use seems an essential prerequisite to designing targeted intervention programs to promote health-enhancing lifestyles (11).

To understand relationships between time use patterns and HRQL, both need to be reliably measured. Fortunately, such measures now exist. There are two main types of HRQL instruments: health profile and utility measures. Health profile measures have been widely available for the last two decades (12). They span multiple subdomains with good precision, but do not allow for calculation of quality-adjusted life years (QALYs) (6). Newer utility measures do produce QALYs but are shorter and have fewer items than health profile measures, reducing their sensitivity (13-15).

The two main types of time use measures are stylised analogues of time use diaries and 24-hour diaries. Stylised questions ask respondents to recall the habitual duration of relevant activities; they are widely used but often only cover a section of the day and are generally inaccurate (16). A 24-hour diary which records all activities over a 24-hour day either prospectively or retrospectively, is more accurate as it reduces social desirability bias and recall errors (17), and may consider activities' co-existence and displacement (5, 18). For example, high levels of physical activity and sedentary behaviour can co-exist, typically in "techno-active" boys who spend relatively large proportions of time on physical activity and screen time (19, 20).

Both types of measures can be analysed as components of time use in isolation (variable-centred) or in combination (person-centred) (21-23). A person-centred approach captures all activities an individual undertakes in a specified time period and explores them as a whole (21, 24). This requires complex data treatment and analysis to acknowledge the boundedness of time, i.e., that any increase in time spent in one activity necessarily decreases time in another (21, 24). In contrast, a variable-centred approach investigates one or more discrete activities in isolation, which are then typically analysed in simpler linear models, but does not acknowledge the finite nature of time and the necessary trade-offs (21, 24).

1 Adolescent time use patterns may present an opportunity to transform future health status (24-26).
 2 Thus, this scoping review aims to evaluate the associations of 1) person-centred time use and 2)
 3 variable-centred time use with HRQL in population-based studies involving 10 to 19 year olds.

4 **METHODS**

5 A preliminary scoping review in late 2015 informed the choice of search terms, databases and
 6 restrictions to be used. The search strategy was developed with a librarian expert in this area at the
 7 Royal Children's Hospital, Melbourne. Medline (Ovid), PsycINFO and Embase (Ovid) databases were
 8 searched in February 2016 using a common set of MeSH terms, thesaurus and/or keywords. PubMed
 9 was searched to retrieve yet-to-be indexed papers. Appendix 1 shows the Medline search strategy,
 10 which was then adapted for the other databases. Due to limited resources and the relative novelty of
 11 standardised time use and HRQL measurement in children, we restricted the review to papers
 12 published in English in the last twenty years.

13
 14 Table 1 lists the inclusion and exclusion criteria. Data were extracted by a single author (MoW).
 15 Additional relevant articles were found by hand-searching reference lists of the identified articles and
 16 from experts in the field. From our prior knowledge, we did not expect a meta-analysis to be possible,
 17 so planned a scoping review (27). No formal quality rating was undertaken.

18 **RESULTS**

19 Figure 1 shows the progress to final inclusion of 29 articles. Only one article examined overall time
 20 use (i.e. person-centred) against a validated health profile HRQL measure (28); none used a utility
 21 HRQL measure. Therefore, we broadened the inclusion criteria for person-centred time use to include
 22 outcomes of wellbeing and functioning, identifying four further articles. An additional 24 articles
 23 examining individual activities (i.e. variable-centred) and HRQL were identified. As anticipated, time
 24 use instruments, HRQL constructs, data treatment and methodological decisions differed widely.
 25 Tables 2 and 3 present key information of the included studies, including type of time use measures
 26 (stylised vs 24-hour diary).

28 **Person-centred time use (Table 2)**

29 *Relationships between HRQL and person-centred time use*

30 Hunt's (28) cross-sectional community sample of 667 15-19 year olds was the only study to use a
 31 validated HRQL profile measure (the KIDSCREEN health profile measure (28, 29)) and a 24-hour
 32 time diary (table 2). Using latent profile models with their 31-item diary, they identified three
 33 probabilistic time use profiles in boys: i) 'productive', ii) 'high leisure', and iii) 'all-rounder'. Two
 34 profiles emerged in girls: i) 'higher study/lower leisure', and ii) 'moderate study/higher leisure'. There

was no association between time use and HRQL in boys. In girls, 40% of those in the ‘moderate study/higher leisure’ group scored >0.5 standard deviations (SD) above the internal mean for HRQL, compared to 18% in the ‘higher study/lower leisure’ group ($p=0.006$). Of the many analyses conducted, only five associations were seen between HRQL subdomains and time use, and none remained significant after adjustment for multiple testing.

Relationships between wellbeing/functioning and person-centred time use

A further four American studies investigated person-centred time use against various questionnaires to measure different aspects of wellbeing/functioning (e.g. internalising and externalising problems, academic results, see Table 2) (30-33). Nelson’s study (32) was the only one to measure an extensive list of activities using a validated stylised questionnaire with a large national sample, while the other smaller studies used study-designed time use questionnaires (30, 33) and/or had a limited list of activities (30, 31). In Nelson’s study (32), boys and girls were analysed together to identify five profiles: i) ‘social’, ii) ‘all-around’, iii) ‘unstructured recreation’, iv) ‘employed’ and v) ‘study’. The profiles differed by sex, race/ethnicity, socioeconomic status, school type and location. The ‘study’ group consistently showed the best perception of school climate, academic achievements, and the lowest levels of truancy and victimisation, while the ‘unstructured recreation’ group showed the worst outcomes for wellbeing/functioning.

The three remaining studies each examined different aspects of wellbeing/functioning (30, 31, 33). For example, Bartko (30) demonstrated that adolescents who participated in pro-social activities had the best psychosocial functioning. In Metzger’s study (31), participation in organised activities was also associated with positive outcomes; however, in certain profiles (e.g. ‘school and community’) it was associated with negative outcomes (e.g. delinquency, suspension). Zarrett’s (33) six identified profiles differed in terms of ‘positive youth development’, ‘contribution’ and ‘depression’, but not in ‘risk behaviour’, which was consistent across profiles.

Variable-centred time use

Table 3 summarises the 24 studies examining variable-centred time use. A wide range of mainly short measures was used to estimate habitual time spent in specific activities, which generally comprised physical activity, sedentary behaviour (often screen time specifically) and sleep. Similarly, various HRQL measures were used, with only two studies employing utility measures (34, 35).

Relationships between HRQL and variable-centred time use

Of the five studies stratifying by sex (23, 36-39), two found different results for boys and girls where the associations were only observed in boys (36, 38). The participants’ ages ranged from 9-19 years

and most studies were conducted in wealthy European countries, Australia, United States, Canada and Japan.

15 studies (11 cross-sectional, four longitudinal) investigated physical activity and HRQL, perhaps reflecting the global obesity epidemic (40). All used stylised questions to estimate frequency and/or duration time in physical activity; most used previously-reported questions (23, 34, 35, 37, 38, 41-47), while the remainder used study-designed measures (22, 36, 37, 48). Many physical activity measures comprised one to three items. Of the 15 studies, 13 showed that more time in moderate-vigorous physical activity was associated with better HRQL. The strongest association was observed in Omorou's (44) longitudinal national study of 1445 French 14-18 year olds, where the mean HRQL of adolescents who met vs did not meet physical activity recommendations over a two-year period was 0.3SD higher on the Duke Health Profile (sample mean 71.0, SD 15.5). The other 12 studies showed similar but weaker associations. Thus, more physical activity, when assessed in isolation, is associated with somewhat better HRQL.

Sedentary behaviour and HRQL has also been widely studied (13 studies; 10 cross-sectional, three longitudinal). Of the 13 studies, 11 examined screen time alone (22, 23, 34, 39, 41, 46, 47, 49-52), one examined reading and music listening on top of screen time (38), and one examined overall sedentary behaviour (duration of sitting) (44). Most studies (12 of 13) showed that more sedentary behaviour was associated with marginally worse HRQL. Mathers' (52) cross-sectional community survey of 925 Australian 13-19 year olds was the most robust in using a validated 24-hour time diary; all others used stylised questionnaires. Children with <2 hours/day of screen time scored 3.35 (95% confidence interval (CI): 1.23; 5.47) units higher on the PedsQL (range 0-100; higher scores indicate better HRQL) and 1.43 (95% CI: 0.07; 2.80) units higher on the KIDSCREEN (range 0-100; higher scores indicate better HRQL) than those with ≥ 4 hours/day (52). Ten other studies showed comparable results, indicating that more screen time, when assessed in isolation, is associated with marginally worse HRQL. Interestingly, in Jalali-Farahani's study (38), HRQL was not associated with music listening and reading -- suggesting that the type of sedentary behaviour is important.

Sleep and HRQL was investigated in seven studies with stylised questionnaires (five cross-sectional, two longitudinal), mostly in the form of sleep duration (six out of seven). Most showed that longer sleep duration or an absence of sleep deprivation, was associated with slightly better HRQL (22, 34, 53-56). For example, in Chen's (22) longitudinal community survey of 7794 Japanese 9-13 year olds, children with <8 hours/day of sleep were 1.25 times (95% CI: 1.01; 1.54) more likely to have worse HRQL (dichotomised) than those with ≥ 9 hours/day of sleep. Similarly, Paiva's (56) cross-sectional national study of 3195 Portuguese 12-19 year olds showed that children whose weekday and weekend

1 sleep duration differed by ≥ 3 hours (i.e. suggesting sleep deprivation on certain days) had 0.81 units
 2 (0.2SD) lower HRQL (mean 38.6 SD 5.8, higher scores indicate better HRQL, $p=0.003$).

3 **DISCUSSION**

4 **Principal findings**

5 Due to a dearth of research, we could not determine how time use patterns are associated with utility-
 6 based HRQL. However, we found some limited evidence from five studies that overall time use is
 7 associated with health profile HRQL and wellbeing/functioning. Thus, a well-balanced lifestyle, i.e.
 8 high participation in multiple types of activities (e.g. ‘all-rounder’, ‘productive’) was associated with
 9 better academic performance and social support and less problematic behaviours, while high
 10 involvement in leisure activities alone (e.g. ‘unstructured recreation’) was associated with the worst
 11 academic involvement, problematic behaviour and psychological functioning. Regarding aspects of
 12 time examined in isolation from each other (an inherently flawed approach, as outlined in the
 13 Introduction), frequent participation in physical activity, less screen time and adequate/more sleep all
 14 consistently showed small to moderate associations with somewhat better HRQL.

16 **Interpretation of findings**

17 *Person-centred time use studies*

18 Contrary to expectations, the one extant study (28) suggested that a ‘well-balanced’ adolescent
 19 lifestyle was not associated with better HRQL in boys and, although there was some evidence of an
 20 association in girls for overall HRQL, this did not play out as differences in the more detailed
 21 dimensions underlying the global measure. This could be the truth — one interpretation being that
 22 cumulative effects of health-enhancing and health-compromising activities may not become
 23 subjectively apparent until adulthood. Alternatively, these broadly null findings might reflect study
 24 limitations, including low response rate (52%), relatively small sample size, homogeneous sample
 25 (adolescents in full time education and living at home), blunt time diary (31 activities in 15 minute
 26 blocks; not validated for this age group) and/or blunt HRQL measure (the 10-item KIDSCREEN).
 27 While well conducted, this study does not suffice to definitively address the question. Nevertheless, it
 28 provided more insight than any other studies included.

30 Interpreting the findings of the four studies that examined overall time use and wellbeing/ functioning
 31 is challenging due to the heterogeneity of exposure and outcome measures. All studies utilised stylised
 32 time use questionnaires with limited items (6 to 19) and diverse measures of wellbeing/functioning
 33 (e.g. Child Behavior Checklist, Profiles of Student Life -- Attitudes and Behaviors Survey). Residual
 34 confounding is another possible issue as, for example, none adjusted for body mass index (a strong
 35 correlate of HRQL in adolescents) (57). Low response rate leading to selection bias is a potential issue

in three studies (30, 32, 33). Due to this variability we have interpreted results cautiously and cannot draw definitive conclusions.

Variable-centred time use

The findings that frequent participation in physical activity, less screen time and adequate/more sleep are consistently associated with somewhat better HRQL was more compelling. Individual associations appeared to be small to moderate in size and therefore of uncertain public policy significance; it is not known whether, when combined, these would ‘add up’ to larger overall beneficial associations with HRQL. Limitations include utilisation of study-designed measures (22, 34, 36, 37, 39, 48, 49), stylised questionnaires without specific types of activities (e.g. sports, dance, athletics for physical activity) (22, 23, 34, 36-38, 41-45, 47, 48) and crude time use measures without actual duration spent on the activity (22, 23, 36, 42). These are major issues, because such measures are known to be extremely inaccurate when compared with actigraphy (58). Ceiling effects for HRQL were observed in many studies, restricting the ranges of differences that were possible, and residual confounding was likely (e.g. 12 studies did not adjust for body mass index). However, the major weakness in these studies is that activities were analysed in isolation, neglecting the compositional nature of time use, so that attributing cause and effect to any one activity type is innately flawed.

We identified, but did not discuss, one relevant systematic review that did not meet our selection criteria (though four of its included papers did (22, 23, 41, 46) and were included). Suchert investigated sedentary behaviour and mental health and similar to our findings, found that in 10 of 14 studies more screen time was associated with worse HRQL and/or wellbeing/functioning (59).

Strengths and limitations

Strengths of this review include a comprehensive search strategy of a wide range of databases going back 20 years, and *a priori* inclusion and exclusion criteria. We included papers published in peer-reviewed journals only. Limitations include restriction to English language publications, and we were unable to conduct dual study screening, selection and data extraction. Moreover, almost all studies included were from wealthy developed countries (except for two Iranian studies) (38, 49), limiting our conclusions to these nationalities and socioeconomic statuses. Meta-analysis was not possible due to the high heterogeneity in methodological designs between studies, most of which do not account for the finite, bounded nature of time. Furthermore, a scoping review such as this does not provide formal assessment of methodological quality of the studies included.

1 **Implications and future research**

2 This review critically appraises the evidence regarding relationships between time use and HRQL in
 3 largely-healthy adolescents. It strongly suggests that time use patterns may be an important modifiable
 4 determinant of HRQL, wellbeing and functioning in this age group. However, quantity and quality
 5 was extremely limited, as we could locate only one empirical cross-sectional person-centred study on
 6 time use in relation to HRQL. Among the other studies reviewed we identified a lack of standardised,
 7 validated time use instruments; methodological variability in time use analyses and selection of
 8 covariates; and failure to account for the interconnected nature of activities across the full 24-hour day
 9 using appropriate techniques. We concur with Hunt's (26) and Ferrar's (24) call for valid age-
 10 appropriate instruments to investigate time use and HRQL and wellbeing in adolescents.

11
 12 Large-scale representative surveys are now needed that use valid measures incorporating all major
 13 types of activities alongside validated, widely-used (and preferably utility based) HRQL measures.
 14 This would allow researchers to chart complex interactions between time use and HRQL in
 15 adolescence or even across lifespans, and assist clinicians, researchers and policymakers to identify
 16 clusters of individuals at risk. This could in turn assist in a more tailored intervention approach in
 17 developing adolescent policy that considers the multiple ways in which adolescents spend their time.

18 19 **ACKNOWLEDGEMENTS**

20 Professor Wake is supported by Australian National Health & Medical Research Council Senior
 21 Research Fellowship 1046518. Research at the Murdoch Childrens Research Institute is supported by
 22 the Victorian Government's Operational Infrastructure Support Program.

23 24 **CONFLICT OF INTEREST**

25 The authors declared no potential conflicts of interest, including no specific financial interests relevant
 26 to the subject of this manuscript.

27 **ABBREVIATIONS**

- 28 • CI: Confidence intervals
- 29 • HRQL: Health-related quality of life
- 30 • PedsQL: Paediatric Quality of Life Inventory
- 31 • QALY: Quality-adjusted life years
- 32 • SD: Standard deviation

1 REFERENCES

- 2 1. Patton GC, Sawyer SM, Santelli JS, Ross DA, Afifi R, Allen NB, et al. Our future: a
3 Lancet commission on adolescent health and wellbeing. *The Lancet* 2016; 387:
4 2423-78.
- 5 2. World Health Organisation. Adolescent health epidemiology [Internet]. World
6 Health Organisation; 2014 [Available
7 from: http://www.who.int/maternal_child_adolescent/epidemiology/adolescence/en/.
- 8 3. World Health Organisation. Health for the world's adolescents [Internet]. World
9 Health Organisation; 2014 [Available from: [http://apps.who.int/adolescent/second-](http://apps.who.int/adolescent/second-decade/section3)
10 [decade/section3](http://apps.who.int/adolescent/second-decade/section3).
- 11 4. Juonala M, Magnussen CG, Berenson GS, Venn A, Burns TL, Sabin MA, et al.
12 Childhood adiposity, adult adiposity, and cardiovascular risk factors. *New Engl J*
13 *Med* 2011; 365: 1876-85.
- 14 5. Larson RW, Verma S. How children and adolescents spend time across the world:
15 work, play, and developmental opportunities. *Psychol Bull* 1999; 125: 701.
- 16 6. Guyatt GH, Feeny DH, Patrick DL. Measuring health-related quality of life. *Ann Intern*
17 *Med* 1993; 118: 622-9.
- 18 7. Ware JJ, Dewey J. Health status and outcome assessment tools. *Int J Health Educ*
19 2000; 3: 138-48.
- 20 8. Centers for Disease Control and Prevention. Measuring Healthy Days: Population
21 Assessment of Health-Related Quality of Life. Atlanta, GA: CDC; 2000.
- 22 9. Government of South Australia. Eat well be active healthy weight strategy for South
23 Australia 2006-2010. In: Department of Health, editor. Adelaide 2006.
- 24 10. National Health Service Choices. Get healthy as a family: NHS choices; 2015
25 [Available from: [http://www.nhs.uk/Livewell/childhealth6-15/Pages/Involve-the-whole-](http://www.nhs.uk/Livewell/childhealth6-15/Pages/Involve-the-whole-family-in-your-childs-weight-management.aspx)
26 [family-in-your-childs-weight-management.aspx](http://www.nhs.uk/Livewell/childhealth6-15/Pages/Involve-the-whole-family-in-your-childs-weight-management.aspx).
- 27 11. Kreuter MW, Wray RJ. Tailored and targeted health communication: strategies for
28 enhancing information relevance. *Am J Health Behav* 2003; 27 Suppl 3: S227-32.
- 29 12. Varni JW, Burwinkle TM, Seid M. The PedsQL 4.0 as a school population health
30 measure: feasibility, reliability, and validity. *Qual Life Res* 2006; 15: 203-15.
- 31 13. Feeny D. A utility approach to the assessment of health-related quality of life. *Med*
32 *care* 2000; 38: 1151-4.

- 1 14. Griebisch I, Coast J, Brown J. Quality-adjusted life-years lack quality in pediatric care:
2 a critical review of published cost-utility studies in child health. *Pediatrics* 2005;
3 115: e600-14.
- 4 15. Stevens K, Ratcliffe J. Measuring and valuing health benefits for economic evaluation
5 in adolescence: an assessment of the practicality and validity of the Child Health
6 Utility 9D in the Australian adolescent population. *Value Health* 2012; 15: 1092-9.
- 7 16. United Nations Economic Commission for Europe. Guidelines for Harmonising Time
8 Use Surveys. Luxembourg: United Nations Economic Commission for Europe; 2013.
- 9 17. Kelly P, Fitzsimons C, Baker G. Should we reframe how we think about physical
10 activity and sedentary behaviour measurement? Validity and reliability
11 reconsidered. *Int J Behav Nutr Phys Act* 2016; 13: 1-10.
- 12 18. Mekary RA, Willett WC, Hu FB, Ding EL. Isotemporal substitution paradigm for
13 physical activity epidemiology and weight change. *Am J Epidemiol* 2009; 170: 519-
14 27.
- 15 19. Ferrar K, Olds T, Maher C. More than just physical activity: time use clusters and
16 profiles of Australian youth. *J Sci Med Sport* 2013; 16: 427-32.
- 17 20. Telama R, Nupponen H, Piéron M. Physical activity among young people in the
18 context of lifestyle. *Eur Phys Educ Rev* 2005; 11: 115-37.
- 19 21. Hagell A, Peck S, Zarrett N. Trends in adolescent time-use in the United Kingdom. In:
20 Hagell A, editor. Changing adolescence: social trends and mental health. UK: Policy
21 Press Scholarship Online; 2012. p. 47-73.
- 22 22. Chen X, Sekine M, Hamanishi S, Yamagami T, Kagamimori S. Associations of lifestyle
23 factors with quality of life (QOL) in Japanese children: a 3-year follow-up of the
24 Toyama Birth Cohort Study. *Child Care Health Dev* 2005; 31: 433-9.
- 25 23. Lacy KE, Allender SE, Kremer PJ, de Silva-Sanigorski AM, Millar LM, Moodie ML, et al.
26 Screen time and physical activity behaviours are associated with health-related
27 quality of life in Australian adolescents. *Qual Life Research* 2012; 21: 1085-99.
- 28 24. Ferrar K, Chang C, Li M, Olds TS. Adolescent time use clusters: a systematic review. *J*
29 *Adolesc Health* 2013; 52: 259-70.
- 30 25. Prochaska JO. Multiple Health Behavior Research represents the future of
31 preventive medicine. *Prev Med* 2008; 46: 281-5.
- 32 26. Hunt E, McKay EA. What can be learned from adolescent time diary research. *J*
33 *Adolesc Health* 2015; 56: 259-66.

- 1 27. Peters MD, Godfrey CM, Khalil H, McInerney P, Parker D, Soares CB. Guidance for
2 conducting systematic scoping reviews. *International journal of evidence-based*
3 *healthcare* 2015; 13: 141-6.
- 4 28. Hunt E, McKay EA, Dahly DL, Fitzgerald AP, Perry IJ. A person-centred analysis of the
5 time-use, daily activities and health-related quality of life of Irish school-going late
6 adolescents. *Qual Life Res* 2015; 24: 1303-15.
- 7 29. Ravens-Sieberer U, Gosch A, Rajmil L, Erhart M, Bruil J, Duer W. KIDSCREEN-52
8 quality-of-life measure for children and adolescents. *Expert Rev Pharmacoecon*
9 *Outcomes Res* 2005; 5: 353-64.
- 10 30. Bartko WT, Eccles JS. Adolescent participation in structured and unstructured
11 activities: a person-oriented analysis. *J Youth Adolesc* 2002; 32: 233-41.
- 12 31. Metzger A, Crean HF, Forbes-Jones EL. Patterns of organized activity participation in
13 urban, early adolescents: associations with academic achievement, problem
14 behaviors, and perceived adult support. *J Early Adolesc* 2009; 29: 426-42.
- 15 32. Nelson IA, Gastic B. Street ball, swim team and the sour cream machine: a cluster
16 analysis of out of school time participation portfolios. *J Youth Adolesc* 2009; 38:
17 1172-86.
- 18 33. Zarrett N, Fay K, Li Y, Carrano J, Phelps E, Lerner RM. More than child's play:
19 variable- and pattern-centered approaches for examining effects of sports
20 participation on youth development. *Dev Psychol* 2009; 45: 368-82.
- 21 34. Chen G, Ratcliffe J, Olds T, Magarey A, Jones M, Leslie E. BMI, health behaviors, and
22 quality of life in children and adolescents: a school-based study. *Pediatrics* 2014;
23 133: e868-74.
- 24 35. Boyle S, Jones G, Walters S. Physical activity, quality of life, weight status and diet in
25 adolescents. *Qual Life Res* 2010; 19: 943-54.
- 26 36. Liu J, Sekine M, Tatsuse T, Fujimura Y, Hamanishi S, Lu F, et al. Outdoor physical
27 activity and its relation with self-reported health in Japanese children: results from
28 the Toyama birth cohort study. *Child Care Health Dev* 2015; 41: 920-7.
- 29 37. Galan I, Boix R, Medrano MJ, Ramos P, Rivera F, Pastor-Barriuso R, et al. Physical
30 activity and self-reported health status among adolescents: a cross-sectional
31 population-based study. *BMJ Open* 2013;3: 1-10.

- 1 38. Jalali-Farahani S, Amiri P, Chin YS. Are physical activity, sedentary behaviors and
2 sleep duration associated with body mass index-for-age and health-related quality
3 of life among high school boys and girls? *Health Quality Life Outcomes* 2016; 14: 30.
- 4 39. Herman KM, Hopman WM, Sabiston CM. Physical activity, screen time and self-rated
5 health and mental health in Canadian adolescents. *Prev Med* 2015; 73: 112-6.
- 6 40. World Health Organisation. Report of the commission on ending childhood obesity.
7 Geneva: World Health Organisation; 2016.
- 8 41. Dalton WT, Schetzina KE, Pfortmiller DT, Slawson DL, Frye WS. Health behaviors
9 and health-related quality of life among middle school children in Southern
10 Appalachia: data from the winning with wellness project. *J Pediatr Psychol* 2011; 36:
11 677-86.
- 12 42. Sanchez-Lopez M, Salcedo-Aguilar F, Solera-Martinez M, Moya-Martinez P, Notario-
13 Pacheco B, Martinez-Vizcaino V. Physical activity and quality of life in schoolchildren
14 aged 11-13 years of Cuenca, Spain. *Scand J Med Sci Sports* 2009; 19: 879-84.
- 15 43. Spengler S, Woll A. The more physically active, the healthier? The relationship
16 between physical activity and health-related quality of life in adolescents: The
17 MoMo study. *J Phys Act Health* 2013; 10: 708-15.
- 18 44. Omorou AY, Langlois J, Lecomte E, Briancon S, Vuillemin A. Cumulative and
19 bidirectional association of physical activity and sedentary behaviour with health-
20 related quality of life in adolescents. *Qual Life Res* 2016; 25: 1169-78.
- 21 45. Breslin G, Gossrau-Breen D, McCay N, Gilmore G, MacDonald L, Hanna D. Physical
22 activity, gender, weight status, and wellbeing in 9- to 11-year-old children: a cross-
23 sectional survey. *J Phys Act Health* 2012; 9: 394-401.
- 24 46. Gopinath B, Hardy LL, Baur LA, Burlutsky G, Mitchell P. Physical activity and
25 sedentary behaviors and health-related quality of life in adolescents. *Pediatrics*
26 2012; 130: e167-74.
- 27 47. Finne E, Bucksch J, Lampert T, Kolip P. Physical activity and screen-based media use:
28 cross-sectional associations with health-related quality of life and the role of body
29 satisfaction in a representative sample of German adolescents. *Health Psychol Behav*
30 *Med* 2013; 1: 15-30.
- 31 48. Petracci E, Cavrini G. The effect of weight status, lifestyle, and body image
32 perception on health-related quality of life in children: a quantile approach. *Qual*
33 *Life Res* 2013; 22: 2607-15.

- 1 49. Dolatabadi NK, Eslami AA, Mostafavi F, Hassanzade A, Moradi A. The relationship
2 between computer games and quality of life in adolescents. *J Educ Health Promot*
3 2013; 2: 20.
- 4 50. Goldfield GS, Cameron JD, Murray M, Maras D, Wilson AL, Phillips P, et al. Screen
5 time is independently associated with health-related quality of life in overweight
6 and obese adolescents. *Acta Paediatr* 2015; 104: e448-e54.
- 7 51. Kantor RM, Grimes GR, Limbers CA. Physical activity, sedentary behaviors, and
8 health-related quality of life in rural Hispanic youth. *Trans Issues Psychol Sci* 2015;
9 1: 239-49.
- 10 52. Mathers M, Canterford L, Olds T, Hesketh K, Ridley K, Wake M. Electronic media use
11 and adolescent health and well-being: cross-sectional community study. *Acad*
12 *Pediatr* 2009; 9: 307-14.
- 13 53. Tzischinsky O, Shochat T. Eveningness, sleep patterns, daytime functioning, and
14 quality of life in Israeli adolescents. *Chronobiol Int* 2011; 28: 338-43.
- 15 54. Gustafsson ML, Laaksonen C, Aromaa M, Asanti R, Heinonen OJ, Koski P, et al.
16 Association between amount of sleep, daytime sleepiness and health-related quality
17 of life in school children. *J Adv Nurs* 2016; 72: 1263-72.
- 18 55. Segura-Jiménez V, Carbonell-Baeza A, Keating X, Ruiz J, Castro-Piñero J. Association
19 of sleep patterns with psychological positive health and health complaints in
20 children and adolescents. *Qual Life Res* 2015; 24: 885-95.
- 21 56. Paiva T, Gaspar T, Matos MG. Sleep deprivation in adolescents: correlations with
22 health complaints and health-related quality of life. *Sleep Med* 2015; 16: 521-7.
- 23 57. Williams J, Wake M, Hesketh K, Maher E, Waters E. Health-related quality of life of
24 overweight and obese children. *JAMA* 2005; 293: 70-6.
- 25 58. Adamo KB, Prince SA, Tricco AC, Connor-Gorber S, Tremblay M. A comparison of
26 indirect versus direct measures for assessing physical activity in the pediatric
27 population: a systematic review. *Int J Pediatr Obes* 2009; 4: 2-27.
- 28 59. Suchert V, Hanewinkel R, Isensee B. Sedentary behavior and indicators of mental
29 health in school-aged children and adolescents: a systematic review. *Prev Med*
30 2015; 76: 48-57.
- 31

TABLES

Table 1. Inclusion and exclusion criteria

Inclusion criteria	Exclusion criteria
Published in 1996-2016	Primarily focused on children aged less than 10 years or adults
Primary focus on healthy individuals aged 10-19 years	Clinical or special populations (e.g. children with autism spectrum disorder)
Sample size more than 100	Theoretical or methodological papers or literature reviews
Exposure variable reported as actual time spent on activities	Theses or book chapters
Outcomes measured using validated tools	Time use expressed as metabolic equivalents or energy expenditure
English language	
Published in peer-reviewed journals	

Table 2. Summary of included studies on person-centred time use. All time use measures were subjective; only studies with validated measures of HRQL/wellbeing included.

Author (Ref) year	Sample; design	N (% boys); response; Age (yr)	Measures ⁱ	Main findings ⁱⁱ	Limitations
-------------------	----------------	--------------------------------	-----------------------	-----------------------------	-------------

Author (Ref) year	Sample; design	N (% boys); response; Age (yr)	Measures ⁱ	Main findings ⁱⁱ	Limitations
Hunt (28) 2014	Community based, Ireland. Cross-sectional	667 (47% boys) 52% response 15-19yr	<ul style="list-style-type: none"> - Time use: 31 time-diary items in 15-minute categories; 1 weekday and 1 weekend day; adapted from Irish Economic and Social Research Institute time use survey for adults 2005 - HRQL: KIDSCREEN-52; 10 domain scores + 10-item KIDSCREEN-10 global HRQL score - Stratification: sex 	<p>Time use profiles derived for boys and girls separately using latent profile models</p> <ul style="list-style-type: none"> • Boys, 3 profiles: 'productive' (more time in study, paid work, housework, and less time in leisure), 'high leisure' (more time on leisure but less time on study, paid work, housework, sleep, self-care and school), and 'all-rounder' (average amount of time in sleep, self-care, school, slightly above-average leisure time, and more time in study and housework). • Girls, 2 profiles: 'moderate study/high leisure' (less time in sleep, moderate in study, more in leisure); 'moderate study/low leisure' (more time in sleep, housework and study, less in leisure). • Boys: no association of time profiles with global HRQL. Girls: 40% in the 'moderate study/higher leisure' group scored >0.5 SD above the mean for global HRQL, vs 18% in the 'higher study/lower leisure' group (p=0.006, association remained after adjustment for multiple testing). • Time profiles vs HRQL subscales: none significant after adjustment for multiple testing. 	<ul style="list-style-type: none"> - A non-utility measure for HRQL - All adolescents were in school and living at home, which may have reduced variability
Bartko (30) 2002	Community based, US. Cross-sectional	918 (50% boys) - 16-17yr	<ul style="list-style-type: none"> - Time use: 11 stylised items; study designed - Wellbeing/functioning: GPA, problem behaviour (15-item), Children's Depression Inventory, Harter's Global Self-Worth scale, Child Behavior Checklist - Confounder: sex, SES 	<ul style="list-style-type: none"> • Out of 7 wellbeing scales analysed, time use profiles were associated with 6 wellbeing scales. • Externalising behaviour differed the most between time profiles: the 'uninvolved' group had the highest (i.e. worst) score (mean 8.51 SD 9.27) while the 'high involved' group had the lowest score (mean 4.91 SD 5.35). 	<ul style="list-style-type: none"> - Limited list of activities - Study designed time use instrument that has not been validated. - Crude measures of time use
Metzger (31) 2009	Community based, US. Cross-sectional	2495 (49% boys) 99% response 13yr	<ul style="list-style-type: none"> - Time use: 6 stylised items; Adapted from the Search Institute's Profiles of Student Life: Attitudes and Behaviors Surveys - Wellbeing/functioning: GPA, school attendance, suspension histories, Children Institute's Assessment of Adolescent Behaviors and Attitudes Towards School and Community, 	<ul style="list-style-type: none"> • Time profiles were associated with all nine aspects of wellbeing measured. • The most significant association was in the 'negative affect' score where the 'uninvolved' group reported the worst while the 'multiply involved' showed the best score (i.e. least negative affect) (marginal mean 2.20 and 1.89 respectively). • Effect size could not be assessed as the standard deviation was not provided. 	<ul style="list-style-type: none"> - Limited list of activities

Author (Ref) year	Sample; design	N (% boys); response; Age (yr)	Measures ⁱ	Main findings ⁱⁱ	Limitations
Nelson (32) 2009	National sample, US. Cross-sectional	6338 (51% boys) 41% response 10 th grade	<ul style="list-style-type: none"> - Confounder: sex, age, ethnicity - Time use: 18 stylised items; based on the Educational Longitudinal Study of 2002 - Wellbeing/functioning: Educational Longitudinal study of 2002; 6-item survey on recognitions at school - Confounder: sex, age, ethnicity, school 	<ul style="list-style-type: none"> • Time profiles were associated with all aspects of wellbeing measured. • Truancy and delinquency differed the most between time profiles: the ‘unstructured recreation’ profile (less time in academic and enrichment activities outside of school and social activities but more screen time) reported the highest levels of truancy (mean 1.73 SD 0.56) while the ‘study’ group (more time in homework, reading, school clubs and sports but less time on social activities) had the lowest truancy score (mean 1.49 SD 0.39). 	<ul style="list-style-type: none"> - Different scales were used for each aspect of wellbeing making the results challenging to interpret.
Zarrett (29) 2009	National sample, US. Cross-sectional	1357 (48% boys) 39% 11yr	<ul style="list-style-type: none"> - Time use: 19 stylised items; study designed - Wellbeing/functioning: PYD, Profiles of Student Life–Attitudes and Behaviors Survey, Teen Assessment Project Survey Question Bank, CES-D, Monitoring the Future questionnaire - Confounder: sex, ethnicity, SES, youth goal orientation 	<ul style="list-style-type: none"> • In examining the association between six time use profiles and four wellbeing scales, the ‘high engaged’ profile was associated with the best PYD while the ‘nonsports’ profile showed the worst PYD score (mean (SE) 74.84 (1.32) and 68.00 (0.57) respectively, $p < 0.05$). Similar results were found in the ‘contribution’ score. • The ‘high engaged’ and ‘nonsports’ groups had the worst ‘depression’ score while the ‘sports only’ group had the best score (mean (SE) 14.60 (1.06), 14.07 (0.45) and 11.78 (0.69) respectively, $p < 0.05$). • Risk behaviour did not differ between the six time use profiles. 	<ul style="list-style-type: none"> - Study designed time use instrument that has not been validated.

ⁱ CES-D: Center for Epidemiological Studies Depression Scale, GPA: Grade point average, PYD: Positive youth development, SES: Socioeconomic status

ⁱⁱ SD: Standard deviation, SE: Standard error

Table 3. Summary of included studies on variable-centred time use. All time use measures were subjective.

Author (Ref) year	Sample; design	N (%boys); response; Age (yr)	Measures ⁱ	Main findings	Limitations
Boyle (35) 2010	Community based, England. Cross-sectional	1771 (50% boys) 33% response 11-15yr	<ul style="list-style-type: none"> - PA: 24 stylised items; Western Australian CAPANS - HRQL: PedsQL, EQ-5D - Confounder: sex, age, ethnicity, SES 	<ul style="list-style-type: none"> • There was no difference in HRQL between adolescents meeting vs not meeting recommended PA guidelines. 	

Author (Ref) year	Sample; design	N (%boys); response; Age (yr)	Measures ⁱ	Main findings	Limitations
			- Stratification: season		
Sánchez-López (42) 2008	Community based, Spain. Cross-sectional	1073 (48% boys) 76% response 11-13yr	- PA: 2 stylised items; activity type unspecified; actual duration unspecified; CHIP-CE - HRQL: CHIP-CE - Confounder: sex, age, BMI	<ul style="list-style-type: none"> Physically active adolescents had higher HRQL scores except in risk avoidance domain. Satisfaction, comfort, resilience and achievement domain scores were higher by 0.26, 0.12, 0.51, 0.29 scores respectively (P<0.001) on a scale of 1-5. Risk avoidance was higher in those that are sedentary by 0.18 (P<0.001). Effect size of CHIP-CE: 0.20=small, 0.50=moderate, 0.80=large. 	<ul style="list-style-type: none"> PA measured using two items of CHIP-CE. Crude measures of time use
Spengler (43) 2013	National sample, Germany. Cross-sectional	1828 (51% boys) 40% response 11-17yr	- PA: 3 stylised items; activity type unspecified; MoMo-PAQ - HRQL: KINDL-R - Confounder: SES	<ul style="list-style-type: none"> An hour increase in PA per week raised total HRQL score by 0.42 points (P<0.001, SD 10.3) on a 0-100 scale. Physical activity explained 2.4% of variance of HRQL by a linear regression (P<0.001). 	
Liu (36) 2015	Community based, Japan. Longitudinal	5238 (48% boys) 50% response 5-7; 11-13yr	- PA: 1 stylised item; activity type unspecified; actual duration unspecified; study designed - HRQL: COOP - Confounder: breakfast eating, sleep duration, headache, abdominal pain - Stratification: sex	<ul style="list-style-type: none"> Adolescents with frequent PA at follow-up were more likely to report better HRQL (86.4% vs 79.7% good self-reported health in boys, p<0.001; 82.5% vs 78.7% in girls, p=0.015). There was no significant difference at baseline (aged 5-7 years). Boys who reported persistent (i.e. at baseline and follow-up) in participation in PA were more likely to have better HRQL at age 12 years (OR (95% CI) 1.47 (1.14; 1.89)). No difference was found in girls. 	<ul style="list-style-type: none"> Study designed time use instrument that has not been validated. Crude measures of time use Follow-up bias
Petracci (48) 2013	Community based, Italy. Cross-sectional	4338 (50% boys) 95% response 13-14yr	- PA: 1 stylised item; activity type unspecified; study designed ^{ii,iii,iv,v} - HRQL: EQ-VAS - Confounder: no	<ul style="list-style-type: none"> Adolescents with <2 hours/week of PA had lower HRQL with small effect size. There was no significant difference between adolescents who exercised 2-11 hours/week and those who exercised >11 hours/week. 	<ul style="list-style-type: none"> Study designed time use instrument that has not been validated.
Galan (37) 2013	National sample, Spain. Cross-sectional	21811 (-) 78% response 11-18yr	- PA: 1 stylised item; activity type unspecified; study designed - HRQL: KIDSCREEN	<ul style="list-style-type: none"> Increasing PA was associated with higher HRQL especially when increasing from 5 to 7 days/week of moderate-vigorous PA (unstandardised β (95% CI) 3.0 (1.7; 4.3), and 5.1 (3.7; 6.4) respectively; mean 50 SD 10). 	<ul style="list-style-type: none"> Number of each sex was not stated. Study designed time

Author (Ref) year	Sample; design	N (%boys); response; Age (yr)	Measures ⁱ	Main findings	Limitations
			<ul style="list-style-type: none"> - Confounder: age, BMI, ethnicity, SES, numerous others - Stratification: sex 		use instrument that has not been validated.
Breslin (45) 2012	Community based, Northern Ireland. Cross-sectional	1424 (45% boys) 73% response 9-11yr	<ul style="list-style-type: none"> - PA: 2 stylised items; activity type unspecified; adapted from PASS projectⁱⁱ ^{iv} - HRQL: CHIP-CE, KIDSCREEN - Confounder: sex, BMI 	<ul style="list-style-type: none"> • Adolescents with high PA had significantly higher HRQL (P=0.003) with small effect size ($\eta_p^2 \approx 0.01$). 	
Dalton (41) 2010	Community based, US. Cross-sectional	152 (46% boys) 29% response 11-12yr	<ul style="list-style-type: none"> - PA: 2 stylised items; activity type unspecified; adapted from Glasgow's Measures of Patient Health Behaviors 2005 - ScrT: subjective; 2 items; adapted from Glasgow's Measures of Patient Health Behaviors 2005ⁱⁱ - HRQL: PedsQL - Confounder: SES, BMI 	<ul style="list-style-type: none"> • Being physically active was associated with a higher level of total HRQL score on a scale of 0-100 (standardised $\beta=0.24$ p<0.004). • A longer duration of screen time was associated with a lower total HRQL score (standardised $\beta=-0.45$ p<0.001). 	- Small sample size
Gopinath (46) 2012	Community based, Australia. Longitudinal	169 (44% boys) 75%; 52% response 12; 18yr	<ul style="list-style-type: none"> - PA: 10 stylised items; Adolescent Physical Activity Recall Questionnaire - ScrT: 5 items; Adolescent Sedentary Activity Questionnaire - HRQL: PedsQL - Confounder: sex, age, ethnicity, SES, BMI 	<ul style="list-style-type: none"> • Adolescents with the highest level of PA (≥ 2.57 vs ≤ 1.21 hours/day) had higher overall HRQL on a scale of 0-100 (mean (SE) 82.3 (0.9) and 79.15 (0.9) respectively). • Those who frequently participated in PA over 5 years (≥ 2.57 vs ≤ 1.21 hours/day) had a higher HRQL (mean (SE) 83.2 (1.6) and 79.57 (1.4) respectively). • Adolescents with high screen time (≥ 3.93 vs ≤ 2.50 hours/day) had lower physical, psychosocial, emotional, school and overall HRQL (mean (SE) 78.30 (0.9) and 81.83 (0.9) respectively) • Those with high screen time (≥ 3.93 vs ≤ 2.50 hours/day) over 5 years had lower HRQL (mean (SE) 75.87 (1.5) and 82.21 (1.5) respectively, p<0.001). 	- Follow-up bias

Author (Ref) year	Sample; design	N (%boys); response; Age (yr)	Measures ⁱ	Main findings	Limitations
Lacy (24) 2012	Community based, Australia. Cross-sectional	3040 (56% boys) 51% response 11-18yr	<ul style="list-style-type: none"> - PA: 5 stylised items; activity type unspecified; actual duration unspecified; adapted from the 2002 National Children's Nutrition Survey - ScrT: 8 stylised items; adapted from the 2002 National Children's Nutrition Survey - HRQL: PedsQL - Confounder: SES, BMI - Stratification: sex, age 	<ul style="list-style-type: none"> • Boys who were most physically active at recess, lunchtime and after school every day had higher HRQL by 5.31 (95% CI 3.56; 7.07, $p<0.001$), 8.11 (95% CI 6.16; 10.1, $p<0.001$) and 6.37 (95% CI 3.24; 9.50, $p<0.001$) units on a 0-100 scale. Similar results were found in girls. • An hour increase in screen time was related to 1.30-unit reduction in HRQL in boys (95% CI -1.70; -0.89, $p<0.001$) and 1.82-unit reduction in HRQL in girls (95% CI -2.42; -1.23, $p<0.001$). 	<ul style="list-style-type: none"> - Crude measures of time spent on PA recorded.
Omorou (44) 2015	National sample, France. Longitudinal	1445 (43% boys) - 14-18yr	<ul style="list-style-type: none"> - PA: 25 stylised items; activity type unspecified; IPAQ - SB: 2 stylised items; IPAQ - HRQL: Duke Health Profile - Confounder: sex, age, SES, BMI 	<ul style="list-style-type: none"> • The cumulative level of meeting PA guidelines was associated with an increase in overall HRQL (standardised $\beta=5.5\text{-unit} \pm 1.0$, $p<0.001$), social (8.7 ± 1.2, $p<0.001$), physical (3.8 ± 1.2, $p=0.001$), and mental (5.2 ± 1.4, $p<0.001$) scores on a scale of 0-100 than not meeting PA guidelines. • Cumulative high SB (>7 hours/day) was correlated with lower HRQL (standardised $\beta=-3.1 \pm 1.2$, $p=0.003$). 	<ul style="list-style-type: none"> - Response rate not provided - Follow-up bias
Finne (47) 2013	National sample, Germany. Cross-sectional	6813 (51% boys) 67% response 11-17yr	<ul style="list-style-type: none"> - PA: 1 stylised item; activity type unspecified; adapted from KiGGS survey - ScrT: 1 stylised item; adapted from KiGGS survey - HRQL: KINDL-R - Confounder: age, SES, ethnicity, BMI, smoking, alcohol, sleep duration 	<ul style="list-style-type: none"> • High PA was associated with higher levels of HRQL. There were small to moderate effect sizes for daily vs. no regular PA for both sexes. Emotional and social wellbeing were affected the most in boys (moderate effects: $d=0.56\text{-}0.59$) and physical then social wellbeing in girls (small to moderate effects: $d=0.43$ and $d=0.32$). • High screen time was related to lower HRQL. The largest effect was found in school domain ($d=-0.37$). In boys, only physical wellbeing and school were significantly affected by high screen time ($p<0.001$), with the largest effects for school ($d=-0.31$) 	<ul style="list-style-type: none"> - Crude measures of time use
Chen (23) 2005	Community based, Japan. Longitudinal	7794 (50% boys) 90%; 75% response 9-13yr	<ul style="list-style-type: none"> - PA: 1 stylised item; activity type unspecified; actual duration unspecified; study designed - ScrT: 2 stylised items; study designed - Sleep: 2 stylised items; study designedⁱⁱ 	<ul style="list-style-type: none"> • Adolescents who participated in PA 'seldom' or almost never' were more likely to have poorer HRQL than those with 'very often' at baseline (OR (95% CI) 1.58 (1.38; 1.81) and 1.92 (1.29; 2.87) respectively). Those who reduced PA and kept low PA were associated with poorer HRQL (OR 2.10 (1.84; 2.39) and 2.21 (1.03; 4.09) respectively). 	<ul style="list-style-type: none"> - Study designed time use instrument that has not been validated. - Crude measures of

Author (Ref) year	Sample; design	N (%boys); response; Age (yr)	Measures ⁱ	Main findings	Limitations
			<ul style="list-style-type: none"> - HRQL: COOP - Confounder: sex, age, BMI 	<ul style="list-style-type: none"> • Adolescents who increased screen time (<2 vs ≥2 hours/day) were associated with poor HRQL (OR 1.24 (1.10; 1.39)). Those with ≥ 3h television viewing were associated with poorer HRQL (OR 1.20 (1.10; 1.39)). • Those with less sleep (<8 vs ≥9 hours/day) were associated with poorer HRQL (OR 1.25 (1.01; 1.54)). 	<ul style="list-style-type: none"> - time use for PA - Follow up bias
Chen (34) 2014	Community based, Australia. Cross-sectional	3353 (49% boys) - 9-15yr	<ul style="list-style-type: none"> - PA: 1 stylised item; activity type unspecified; adapted from the Youth Risk Behavior Survey - ScrT: 2 stylised items; adapted from the Youth Risk Behavior Survey - Sleep: 5 stylised items; study designed^{iv,v} - HRQL: CHU9D - Confounder: sex, age, SES 	<ul style="list-style-type: none"> • An increase in the number of physically active days was correlated with an increase in HRQL by 0.004-unit (SE=0.001, P<0.001 and SE=0.002, P = 0.05 for primary and high school children respectively) on a scale of 0-1. • Each additional hour of screen time was associated with a decreased adjusted mean utility of -0.008 and -0.009 (SE=0.001, P <0.001 both) for primary and high school students respectively. • An extra hour of sleep was significantly associated with an additional adjusted mean utility of 0.010 (SE=0.003, P<0.001) and 0.014 (SE=0.014, P= 0.001) for primary and high school students respectively. 	<ul style="list-style-type: none"> - Response rate was unknown. - Inclusion criteria: disadvantaged communities. - Study designed time use instrument for sleep that has not been validated.
Jalali-Farahani (38) 2016	Community based, Iran. Cross-sectional	465 (51% boys) 96% response 14-17yr	<ul style="list-style-type: none"> - PA: 3 stylised items; activity type unspecified; QAPACE - ScrT: 2 stylised items; QAPACE - Music/reading: 2 stylised items; QAPACE - Sleep: 1 stylised item; QAPACE^{iv,v} - HRQL: PedsQL - Stratification: sex, school/vacation 	<ul style="list-style-type: none"> • PA was positively related to HRQL during school period. • Screen time was negatively associated with school functioning during school period. • There was no association between sleep and HRQL. • Time spent on music listening/reading was positively correlated to HRQL during school period. During vacation period, they were negatively associated with emotional functioning in boys. 	
Kantor (51) 2015	Community based, U.S. Cross-sectional	448 (51% boys) 96% response 7-12yr	<ul style="list-style-type: none"> - ScrT: 2 stylised items; SPAN^{vi} - HRQL: PedsQL - Confounder: sex, ethnicity, grade 	<ul style="list-style-type: none"> • Greater duration on television was associated with lower HRQL (unstandardised β=-0.97-unit, SE 0.41, p<0.05) on a 0-100 scale. 	<ul style="list-style-type: none"> - Inclusion criteria: Hispanic children who could read in English or Spanish.
Dolatabadi (49) 2013	Community based, Iran.	444 (52% boys) 90% response	<ul style="list-style-type: none"> - Computer games: 1 stylised item; study designed 	<ul style="list-style-type: none"> • There was a positive relationship between duration of computer games playing and HRQL (p=0.01, r=0.11), but no relationship between playing computer games 	<ul style="list-style-type: none"> - Study designed time use instrument that

Author (Ref) year	Sample; design	N (%boys); response; Age (yr)	Measures ⁱ	Main findings	Limitations
	Cross-sectional	12-15yr	- HRQL: WHOQOL-BREF - Confounder: SES	(regardless of duration) and HRQL.	has not been validated.
Herman (39) 2015	National sample, Canada. Cross-sectional	7725 (51% boys) 68% response 12-17yr	- ScrT: 3 stylised items; study designed ^{vi} - HRQL: SRH, SRMH - Confounder: age, BMI, ethnicity, SES, lifestyle - Stratification: sex	• Adolescents with high screen time were more likely to have worse HRQL (boys: OR (95% CI) 1.28 (1.08, 1.52); girls: 2.06 (1.74, 2.44)), and worse mental health (boys: 1.34 (1.11, 1.62); girls: 1.52 (1.28, 1.80)).	- SMRH not previously validated in adolescents. - Study designed time use instrument that has not been validated.
Mathers (52) 2009	Community based, Australia. Cross-sectional	925 (50% boys) 56% response 13-19yr	- ScrT: 259 time-diary items; MARCA - HRQL: CHQ, KIDSCREEN, PedsQL, K-10, SDQ - Confounder: SES, sex, age, BMI	• Adolescents with the highest screen time (≥ 255 vs < 121 minutes/day) had poorer KIDSCREEN scores (unstandardised β (95% CI) -1.43 (-2.80; -0.07), $P=0.04$) and PedsQL scores by 3.35 on a scale of 0-100 (95% CI -5.47; -1.23, $P=0.002$).	
Goldfield (50) 2015	Community based, Canada. Cross-sectional	358 (27% boys) - 14-18yr	- ScrT: 3 stylised items; study designed - HRQL: PedsQL - Confounder: age, sex, SES, ethnicity, BMI, PA intensity, PA duration, caloric intake	• Screen time duration was associated with lower HRQL (adjusted $r=-0.16$, standardised $\beta=-0.16$, $p=0.009$). • Computer time and video games were associated with lower HRQL.	- Inclusion criteria: overweight and obese adolescents - Study designed time use instrument that has not been validated.
Segura- Jimenez (55) 2014	Community based, Spain. Cross-sectional	684 (53% boys) 95% response 6-18yr	- Sleep: 1 stylised item; HBSC - HRQL: HBSC - Confounder: age, sex, SES	• Sleep durations of >10 hours/day in children <12 years old and >9 hours/day in adolescents were associated with reporting no health complaints (OR 2.3, $P=0.005$ and OR 1.7, $P=0.047$ respectively).	
Paiva (56) 2015	National sample, Portugal. Cross-sectional	3195 (46% boys) 90% response 12-19yr	- Sleep: 3 stylised items; HBSC - HRQL: KIDSCREEN - Confounder: no	• Adolescents with sleep deprivation had lower HRQL but the effect size was small ($\eta^2=0.003$; $p=0.003$).	- Limited information about actual sleep duration

Author (Ref) year	Sample; design	N (%boys); response; Age (yr)	Measures ⁱ	Main findings	Limitations
Tzischinsky (53) 2011	Community based, Israel. Cross-sectional	470 (50% boys) - 12-16yr	- Sleep: 2 stylised items; SSHS - HRQL: PedsQL - Confounder: no	• Weekday sleep duration was significantly correlated with HRQL ($r=0.17$, $p<0.001$ and $r=-0.32$, $p<0.001$ respectively).	
Gustafsson (54) 2016	Community based, Finland. Longitudinal	568 (47% boys) 63-67% response 10-15yr	- Sleep: 4 stylised items; HBSC - HRQL: PedsQL - Stratification: age	• The association between sleep and HRQL was strongest in the 15-year olds, weaker in the 12-year olds and no association in the 10-year-olds.	- Follow-up bias

ⁱ BMI: Body mass index, CAPANS: Child and Adolescent Physical Activity and Nutrition Survey, CHIP-CE: Child Health and Illness Profile, CHQ: Child Health Questionnaire, CHU9D – Child Health Utility 9D, COOP: Cooperative Functional Assessment Charts, EQ-5D: EuroQoL five dimensions questionnaire, EQ-VAS: EuroQoL Visual Analogue Scale, HBSC: Health Behaviour in School-aged Children, IPAQ: International Physical Activity Questionnaire, K-10: Kessler Psychological Distress Scale, KiGGS: German Health Interview and Examination Survey, MoMo-PAQ: MoMo physical activity questionnaire, PA: Physical Activity, PASS: Physical Activity in Scottish Schoolchildren, PedsQL: Pediatric Quality of Life Inventory, QAPACE: Quantification de l'Activite Physique en Altitude Chez les Enfants, SB: Sedentary Behaviour, ScrT: Screen Time, SDQ: Strengths and Difficulties Questionnaire, SES: Socioeconomic status, SPAN: School Physical Activity and Nutrition, SRH: Self-Rated Health, SRMH: Self-Rated Mental Health, SSHS: School Sleep Habits Survey, WHOQOL-BREF: WHO Quality of Life-BREF

ⁱⁱ Dietary pattern

ⁱⁱⁱ Weight status

^{iv} SES: Socioeconomic Status

^v Sex

^{vi} Physical activity (not time-related)

FIGURES

Figure 1. Progress through the stages of study selection

Appendix 1. Search history in Medline

1. ((time adj3 (use*1 or usage)) or leisure or chore*1 or (social adj interaction*1)).tw,kf,hw
2. (sitting or sedentary or screentime or (screen adj time) or TV or television or computer*1 or videogame* or (video adj game*1)).tw,kf,hw.
3. (play or sport*1 or physical activit* or exercise).tw,kf,hw.
4. sleep.tw,kf.
5. *sleep/ or *sleep deprivation/ or exp *sleep stages/
6. exp *Exercise/ or exp Internet/
7. *Sedentary Lifestyle/
8. *"Quality of Life"/
9. (quality adj3 life).tw,kf.
10. (diary or diaries or (self adj report*) or scale*1 or instrument*1 or survey*1 or questionnaire*).tw,kf.
11. exp "Surveys and Questionnaires"/
12. (child* or adolescen* or youth*1 or pediatric* or paediatric*).tw,kf,hw.
13. px.fs.
14. health status/
15. (1 or 2 or 3 or 4 or 5 or 6 or 7) and (8 or 9) and (10 or 11) and 12 and (13 or 14)
16. limit 15 to (english language and last 10 years) **Results: 1094**

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Wong, M;Lycett, K;Olds, T;Gold, L;Wake, M

Title:

Use of time and adolescent health-related quality of life/well-being: ascoping review

Date:

2017-08-01

Citation:

Wong, M., Lycett, K., Olds, T., Gold, L. & Wake, M. (2017). Use of time and adolescent health-related quality of life/well-being: ascoping review. ACTA PAEDIATRICA, 106 (8), pp.1239-1245. <https://doi.org/10.1111/apa.13929>.

Persistent Link:

<http://hdl.handle.net/11343/293169>