Effects of Australian Summer Monsoon on Sea Surface Temperature Diurnal Variation over the Australian North-Western Shelf

Xiao Hua Wang^{a, b, *} and Haifeng Zhang^{a, b}

^a *The Sino-Australian Research Centre for Coastal Management, UNSW Canberra, Canberra, ACT, 2600, Australia*

^b *School of Physical, Environmental and Mathematical Sciences, UNSW Canberra, Canberra, ACT, 2600, Australia*

^{*} *Corresponding author: x.h.wang@unsw.edu.au*

Abstract

Five-year (2010 to 2014) sea surface temperature (SST) data from the Advanced Very High Resolution Radiometer (AVHRR) are used to study the effects of the Australian summer monsoon on SST diurnal variation (DV) over the Australian North-Western shelf (NWS, defined as 105°E-125°E, 25°S-10°S). Strong DV events identified with amplitude of 0.5-2K were observed over the NWS. A double-peak seasonal pattern of DV is obtained, with the strongest DV occurring in February/March and October/November. This seasonal DV pattern over the NWS is largely due to the Australian summer monsoon, which reduces the easterly trade wind during the summer monsoonal period, favouring the development of SST DV. Since the monsoon region is distributed globally in the tropical oceans, in the western and eastern North Pacific, as well as in the southern Indian Ocean, we anticipate that strong SST DV may also exist in these parts of coastal oceans.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1002/2017GL075008](https://doi.org/10.1002/2017GL075008)

Key words: AVHRR; Australian North-Western Shelf; SST diurnal variation; Australian summer monsoon

Author Manuscript

1. Introduction

The Southern Hemisphere trade winds are the easterly to southeasterly winds that affect tropical and subtropical Australia in the austral winter. During the austral summer, the temperature gradient between the landmass in northern Australia and the surrounding oceans generates moist northwesterly wind from the Indian Ocean, which replaces the easterly trade winds. This northwesterly wind over the northern part of western Australia is often referred to as the Australian summer monsoon [e.g., *Gentilli, 1971; Suppiah, 1992*]. The westerly of Australian summer monsoon, which typically begins in October/November and lasts until March/April each year, has a strong effect on both meteorological conditions and ocean state over the Australian North-Western Shelf (NWS, defined as 105°E-125°E, 25°S-10°S) and along the Australian west coast [e.g., *Wyrki, 1962; Godfrey and Ridgway, 1985; Weller et al., 2012; Jourdain et al., 2013*].

Both the monsoon and the ocean state in the Australian NWS and west coast may present significant interannual variability due to the El Niño-Southern Oscillation (ENSO) and the Indian Ocean Dipole (IOD). During La Niña, a strong Leeuwin Current is driven by higher coastal sea level, which advects more warm water southwards. During El Niño, the Leeuwin Current becomes weaker and the water temperature is generally cooler [*Feng and Meyers, 2003; Zinke et al., 2014*]. In particular, during La Niña in 2010/2011 a strong marine heat wave occurred along the west coast, with the highest sea surface temperature (SST) recorded in the region. This heat wave was caused by the combination of high sea level in the western equatorial Pacific associated with La Niña condition and unusually high air-sea heat flux into the ocean [*Pearce and Feng, 2013*]. Furthermore, in recent years an increasing trend in SST

in this area has been revealed, which is associated with global warming, and appears to be more significant than that in the Indian Ocean [Caputi *et al.*, 2009; Pearce and Feng, 2007].

SST diurnal variation (DV) is generally defined as the daily temperature variation in the upper few meters of the ocean. A diurnal warm layer typically has a temperature difference relative to the water body below on the order of 0-3K; but in some cases, it can reach values up to 5-8K [e.g., Gentemann *et al.*, 2008; Karagali and Høyer, 2014]. Past studies on SST DV ranged from the determination and characterization of DV events [e.g., Sverdrup *et al.*, 1942; Soloviev and Lukas, 1997; Clayson and Weitlich, 2005; Kawai and Wada, 2007; Gentemann *et al.*, 2008; Eastwood *et al.*, 2011; Karagali and Høyer, 2014; and Zhang *et al.*, 2016a], to the roles of DV events in air-sea interaction [e.g., Halpern and Reed, 1976; Li *et al.*, 2001; Clayson and Bogdanoff, 2013; Marullo *et al.*, 2016], and to their impacts on long-term climate patterns [e.g., Masson *et al.*, 2012].

The significance of SST DV has been widely recognized. Many studies indicate that a better understanding of DV events is essential to better represent the air-sea interaction in weather and climate models [e.g., Fairall *et al.*, 1996; Clayson and Bogdanoff, 2013]. Clayson and Bogdanoff [2013] found that excluding SST DV effects can, to different extents in different regions, underestimate the air-sea heat flux. In Brunke *et al.* [2008], after adding an SST DV scheme, the Community Atmosphere Model (CAM3.1) was better at predicting the diurnal cycle in air temperature and that in precipitation. According to some research, incorporating SST DV signal can enhance the performance of a climate model on long term scale. For instance, Masson *et al.* [2012] showed that in their general circulation model, the prediction of ENSO amplitude, frequency, and skewness is significantly better if SST DV effects are

taken into account. Over the Australian NWS region, it is particularly important to study the DV effects given this domain's role in the tropical climate system and in marine heat wave investigation. The SST DV in this region may also have potential impact on intraseasonal and interannual oscillations such as the Madden-Julian Oscillation (MJO), ENSO evolution, and air temperature prediction over the continent. Finally, the formation of thermal stratification has been accepted as an essential physical condition for the occurrence of phytoplankton bloom in the open waters [e.g., *Riley*, 1957; *Nelson and Smith*, 1991; *Tett and Walne*, 1995; *Ruardij et al.*, 1997] and coastal waters [*Azumaya et al.*, 2001]. As a vast coastal region, the NWS contains shallow waters that are the habitats of temperature sensitive ecosystems such as coral reefs [*Zhu et al.*, 2014]. Therefore, our understanding of SST DV will greatly benefit the studies of the Australian NWS marine ecosystems and their coastal bio-physical dynamics.

In this research, we investigate the effects of the Australian summer monsoon on SST DV over the Australian NWS using a five-year (2010 through 2014) Advanced Very High Resolution Radiometer (AVHRR) SST data set produced by the Australian Bureau of Meteorology (Bureau) and the outputs of the Australian Community Climate and Earth System Simulator-Regional (ACCESS-R) numerical weather prediction model. The organization of the rest of this paper is as follows. In section 2, we give details of the data sets used and a simple physical DV model. In section 3, we present the effects of the Australian monsoon on SST DV. Summary of the study is provided in section 4.

2. Data and Model

Five-year (1st January 2010 to 31st December 2014) AVHRR SST data are used to study the effects of the Australian summer monsoon on SST DV events over the Australian NWS, together with the hourly meteorological data from ACCESS-R 24-hour forecast. A brief description of both data sets and a physical DV model as an example is presented in this section.

2.1. Data sets

2.1.1. SST data

The SST data used in this research are NOAA-19 High-Resolution Picture Transmission AVHRR data version 2 (“fv02”), produced by the Bureau as a contribution to the Integrated Marine Observing System (IMOS; <http://www.imos.org.au/>). Four types of AVHRR SST products, namely, the Level 2 Pre-processed (L2P), Level 3 Un-collated (L3U), Level 3 Collated (L3C), and Level 3 Super-collated (L3S), have been consistently reprocessed for each satellite platform over the region around Australia (70°E-170°W, 70°S-20°N) from 1992 to 2015 [Beggs *et al.*, 2010, 2013]. IMOS AVHRR SST products are compliant with the Group for High Resolution Sea Surface Temperature (GHRSSST) Data Specification (GDS) 2.0 revision 5 (r5) in the Network Common Data Form (NetCDF) file format [GHRSSST Science Team, 2010]. Gridding the best quality L2P data for each pass generates the L3U data at $0.02^\circ \times 0.02^\circ$ resolution. The L3C products are composed of multiple passes of the same satellite sensor over a fixed time period, typically a single daytime period and a single night-time period. The data set was selected due to its high spatial resolution and SST algorithms designed to minimise day-night SST bias [Zhang *et al.*, 2016b]. The L3U files are

used for validation. For SST DV analyses, to avoid repeated merging of the L3U files within a single daytime or night-time period, daily L3C files are adopted.

The validation results using in-situ data by *Zhang et al.* [2016b] showed that the AVHRR SST data set is of overall good quality, although the daytime measurements have an average of 0.19K underestimation. Readers are referred to *Zhang et al.* [2016b] for detailed discussion on the AVHRR SST data quality control method and validation. We also wish to note that since AVHRR SST data are only available under cloud-free conditions, our analysis will unavoidably have this sampling bias, so it may not represent typical SST variation in regions that are frequently obscured by clouds.

2.1.2. Meteorological data

Daily maximum solar shortwave insolation ($SSI_{\max}$), wind speed and direction at 10-m height above sea level are obtained using the ACCESS-R weather prediction model hourly outputs. The original ACCESS climate model is a fully coupled Earth System Model, developed by the Bureau in collaboration with the Commonwealth Scientific and Industrial Research Organisation (CSIRO). In ACCESS, the ocean module, the Modular Ocean Model version 4 (MOM4), and the atmosphere module, the Met Office Unified Model (UM), are coupled via the Ocean-Atmosphere-Sea Ice-Snowpack (OASIS) coupler. The observation processing system uses most routinely available in-situ and remotely sensed data. The frequency of data assimilation is 6-hourly for base times of 00, 06, 12, and 18 UTC using four-dimensional variational algorithms (4dVAR). However, as a transformed and regional version, ACCESS-R weather prediction model is no longer coupled to the ocean module. The SST, which is

used as the boundary condition to ACCESS-R, is derived from observations. ACCESS-R became fully operational since September 2009, and can forecast a variety of surface parameters, including the sensible and latent heat fluxes, shortwave and longwave insolation, and winds [Puri *et al.*, 2013]. The model spatial resolution was increased from $0.375^\circ \times 0.375^\circ$ prior to 17th April 2013 to $0.11^\circ \times 0.11^\circ$ on 17th April 2013 and kept it so. Preliminary comparison showed an increase in the forecast skill relative to the original coarser resolution model [National Meteorological and Oceanographic Centre Operations Bulletin Number 98, 2013]. Therefore, the higher resolution wind data from the model are used for the period from 17th April 2013 onwards.

The ACCESS-R $SSI_{\max}$ and wind speed over the NWS are compared with the European Centre for Medium-Range Weather Forecasts (ECMWF) reanalysis $SSI_{\max}$ data and Cross-Calibrated Multi-Platform (CCMP) sea surface wind data for the same five-year period, respectively (Figure 1). The CCMP data sets are the composite of cross-calibrated satellite winds obtained from the Remote Sensing Systems (RSS) to produce a high-resolution ($0.25^\circ \times 0.25^\circ$) gridded analysis [Atlas *et al.*, 2011]. The two sets of data provide high correlations with $R = 0.85$ and 0.88 for $SSI_{\max}$ and wind speed, respectively. The ACCESS-R underestimates both $SSI_{\max}$ and wind speed slightly with biases of -27.4 Wm^{-2} and -0.33 ms^{-1} , respectively.

Figure 1. Joint distribution of ACCESS-R and ECMWF/CCMP data for (a) SSI_{max} and (b) wind speed over the Australian NWS (105°E-125°E, 25°S-10°S) from 2010 to 2014.

2.2. A simple DV model

The process of SST DV is concerned with solar radiation and wind mixing in the surface boundary layer. A number of numerical models are available to understand DV events and predict the variability of the DV process. They range from turbulence closure models [e.g., *Mellor and Yamada*, 1982; *Karagali et al.*, 2017], to statistical models [e.g., *Price et al.*, 1987; *Webster et al.*, 1996; *Gentemann et al.*, 2003], to air-sea coupled models [e.g., *Noh et al.*, 2011], and to physical-empirical hybrid models [e.g., *Gentemann et al.*, 2009; *Weihs and Bourassa*, 2014]. In essence, the balance between the net heat flux and wind-driven turbulent mixing over a daily cycle determines the magnitude of SST DV. As an example, this process can be represented by a simple physical model for a sub-surface boundary layer (under the cool skin layer) [e.g., *Zeng and Beljaars*, 2005; *Takaya et al.*, 2010] in the following equation:

$$\frac{\partial}{\partial t} \int_{-d}^{-\delta} T dz = \frac{Q + R_s - R(-d)}{\rho_w c_w} - K_w \frac{\partial T}{\partial z} \Big|_{z=-d} \quad (1)$$

where T is temperature, δ is the depth of the cool skin layer, d is the measurement depth at which the diurnal cycle can be omitted (set as 3 m in this study), Q is the sum of latent heat, sensible heat, and longwave radiation, R_s is shortwave radiation, ρ_w is seawater density, c_w is seawater heat capacity, and K_w is turbulent diffusion coefficient. Following *Large et al.* [1994],

$$K_w(z) = k u_{*w}(-z) / \phi_t \left(\frac{-z}{L} \right) \quad (2)$$

where $k = 0.4$ is the Von Karman constant, u_{*w} is the surface friction velocity, and z is negative in the ocean. The stability function is:

$$\phi_t \left(\frac{d}{L} \right) = \begin{cases} 1 + 5 \frac{d}{L} & \text{for } \frac{d}{L} \geq 0 \\ \left(1 - 16 \frac{d}{L} \right)^{-0.5} & \text{for } \frac{d}{L} < 0 \end{cases} \quad (3)$$

In (3), L is the Monin-Obukhov length calculated as $L = \rho_w c_w u_{*w}^3 / (k F_d)$, and $F_d = g \alpha_w [Q + R_s - R(-d)]$ where α_w is the thermal expansion coefficient and $R(-d)$ is shortwave insolation at depth d .

3. Results and Discussion

3.1. SST DV seasonal variability

The SST DV amplitude (also referred to as dSST) in this study is calculated as the daytime SST minus the night-time SST at the same location within the same day (local time). NOAA-19 crosses the local equator at ~14 local solar time (LST) in the day and ~2 LST at night. As

the minimum SST in the daily cycle is observed around dawn (5-7 LST), and the maximum SST, at 14-16 LST [e.g., Gentemann *et al.*, 2003; Karagali *et al.*, 2012; Zhang *et al.*, 2016a], the SST DV amplitude calculated in this study is likely to underestimate the actual maximum DV to some extent. However, it is believed that the general SST DV patterns should be captured, as demonstrated previously by SST DV studies using AVHRR data [e.g., Böhm *et al.*, 1991; Hawkins *et al.*, 1993; Gentemann *et al.*, 2003].

Figure 2 shows the five-year monthly-averaged DV, wind and SSI_{max} over the whole study domain of the NWS. The wind and SSI_{max} averages only include data from days on which DV data are available at that location. The time series of wind vectors over the NWS region shows all-year southerly wind with a strong southeasterly component in the non-monsoon season and a strong southwesterly component in the monsoon season (Figure 2a). A double-peak pattern is revealed for DV over the whole study domain (Figure 2b). The first peak in DV amplitude occurs in February/March, reaching a dSST amplitude of 0.6K. The second peak appears in October/November with a dSST of 0.5K. Both DV peaks coincide with the minimum wind speed and high SSI_{max} in those months. Minimum DV occurs in the austral winter (June/July) when wind speed and SSI_{max} reach maximum and minimum, respectively. Figure 2c shows the five-year monthly- and area-averaged DV but only for the strongest one-third DV events, with collocated wind speed and SSI_{max} . Although not as obvious as in Figure 2b, the double-peak DV pattern can be observed with a maximum amplitude reaching 1.4K in February/March.

Figure 2. (a) Five-year weekly- and area-averaged wind vector; (b) five-year monthly- and area-averaged DV, wind speed, and SSI_{max} over the whole study domain of the Australian NWS (105°E-125°E, 25°S-10°S). The wind and SSI_{max} averages only include data from days on which DV data are available at that location. (c) Same as (b), except for the strongest one-third DV events with the collocated wind and SSI_{max} .

3.2. Effects of the Australian summer monsoon on SST DV

Figures 3 and 4 show five-year monthly-averaged wind vector, dSST and counts of clear days (a proxy for cloud cover) for November, January, March, and August. The former three months represent the beginning, middle, and end of the Australian summer monsoon, respectively, whereas August is a typical non-monsoon season when the SST reaches the coldest of the year. Around October/November, westerly monsoon winds start to develop quickly and largely cancel the easterly trade winds (Figure 3a). At this stage, the resultant winds are very weak ($\sim 1\text{-}3 \text{ ms}^{-1}$). Also, the monsoon winds carry a large amount of water vapour; thus, clouds begin to form and the counts of clear days start to decline (Figure 4a). However, despite the decreasing solar radiation due to more frequent cloud cover, the SST

DV can be very large ($\sim 1\text{-}2\text{K}$ on average), indicating the dominant role of weak wind in determining DV over this region.

Figure 3. Five-year-averaged monthly wind vector and dSST for (a) Nov, (b) Jan, (c) Mar, and (d) Aug.

Figure 4. Spatial distributions of five-year-averaged monthly counts (for Nov, Jan, Mar, and Aug) of the days on which DV data are available, i.e., days on which there are both local daytime and night-time quality controlled SST measurements at the same location.

From January to February, the westerly monsoon winds have replaced the trade winds. Stable westerly winds, which are relatively weak ($3\text{--}5\text{ ms}^{-1}$), are observed over this region (Figure 3b). More clouds lead to the fewest clear days of the year in January (Figure 4b). Still, the SST DV is relatively strong, mainly due to the weak wind.

The monsoon winds begin to decay in March/April (Figure 3c). Before the easterly trade winds take over, there is another wind speed trough during this period (Figures 2b and c). Clear days are still rarely seen because it takes some time for the clouds to dissipate (Figure

4c). Expectedly, the SST DV sees another peak during this period, with a maximum dSST near the coast on the order of 2K.

The austral winter (May to September) marks the typical non-monsoon climate. Moderate to large easterly trade winds dominate in this period. Figure 3d shows consistent easterly trade winds reaching $5\text{--}6\text{ ms}^{-1}$ in August with a monthly-averaged SST DV close to 0K over the entire NWS region. As expected, most days in August are cloud free, with the counts of clear days on the order of 25 days (Figure 4d).

To further investigate the effects of monsoonal winds on the SST DV, daily-averaged surface turbulent diffusion coefficient K_w defined by Equation (2) is calculated (Figure 5). The heat fluxes used in the calculation are daily averages, and the depth d is set to 3 m. Small K_w represents weak wind mixing and strong stratification due to surface heating, which in turn produces a large SST DV. Due to reduced wind speed in November and March, K_w is the minimum in the region with a value around $0.1\text{ m}^2\text{s}^{-1}$. K_w in January reaches a maximum value of $0.35\text{ m}^2\text{s}^{-1}$ at the latitude of 24°S , corresponding to a band of strong trade wind deflected to southerly by the monsoon (Figure 3b). In August, persistent southeasterly trade winds produce moderate K_w over the NWS, despite of low K_w along the coast of the NWS.

In conclusion, large dSST over the NWS is produced by strong SSI_{max} and weak wind speed at the onset and termination of the Australian summer monsoon season in October/November and February/March, respectively. Two opposing wind systems of easterly trade winds and westerly monsoon winds weaken the wind mixing in the surface boundary layer, ensuring large SST DV events during these months. During the peak of the monsoon season (austral

winter), strong and persistent monsoon (trade) winds of northwesterly (southeasterly) dominate. Large wind mixing events prevent the development of large SST DV. Due to low solar radiation in the austral winter, dSST reaches its minimum in its seasonal variability.

Figure 5. Spatial distributions of the five-year-averaged monthly turbulent diffusion coefficient defined by Equation (2) for Nov, Jan, Mar, and Aug.

4. Summary

Five-year (2010 to 2014) AVHRR SST data produced by the Bureau were used to study the effects of the Australian monsoon on SST DV events over the Australian NWS, together with the hourly meteorological data from the ACCESS-R 24-hour forecast. For the five-year average, strong DV events with amplitude of 0.5-2K show a double-peak seasonal pattern over the NWS, where the strongest DV events occur in February/March and

October/November, and the weakest, in June and July. This strong seasonal DV pattern is largely due to the Australian summer monsoon, which has reduced the easterly trade winds at the beginning and end of the monsoonal period. As a result, the wind speed reaches its minimum in these months, causing large DV events due to the weak wind mixing at the sea surface. The strong DV events are further enhanced by the maximum $SSI_{\max}$ change from winter to summer at relatively higher latitudes of the NWS. Other factors, such as cloud cover, play a minor role in affecting SST DV of the region.

Since the monsoon region is distributed globally in the tropical oceans of the western and eastern North Pacific, as well as the southern Indian Ocean, the seasonal reversal from easterly trade winds to westerly monsoon winds over these regions can also be observed (e.g., the tropical Asia, West Africa and India in the Northern Hemisphere). Thus, a double-peak seasonal pattern of DV may also exist in these parts of coastal oceans. The effects on SST DV by these regional tropical monsoons may be a subject of our future study.

Acknowledgements

This is a Sino-Australian Research Centre for Coastal Management Publication Number 37. H. Zhang has been financially supported by the China Scholarship Council since August 2014 for his Ph.D. studies in Australia. The authors would like to thank the Bureau of Meteorology for providing the ACCESS-R forecast meteorological data and the AVHRR SST data. We also thank Anna Maggiorano for providing assistance in literature review on the Australian summer monsoon. If one would like to acquire the AVHRR SST and ACCESS-R data, please contact Dr. Helen Beggs (Helen.Beggs@bom.gov.au). The comments from three anonymous reviewers and the editor have improved the manuscript. We acknowledge the assistance of English editing by Dr Zuojun Yu.

References

- Atlas, R., R. N. Hoffman, J. Ardizzone, S. M. Leidner, J. C. Jusem, D. K. Smith, and D. Gombos (2011) A cross-calibrated, multiplatform ocean surface wind velocity product for meteorological and oceanographic applications. *Bull. Am. Meteorol. Soc.*, 92(2), 157-174, doi: 10.1175/2010BAMS2946.1.
- Azumaya, T., Y. Isoda, and S. Noriki(2001), Modeling of the spring bloom in Funka Bay, Japan. *Cont. Shelf Res.*, 21, 473-494, doi:10.1016/S0278-4343(00)00091-1
- Beggs, H., L. Majewski, G. Paltoglou, E. Schulz, I. Barton, and R. Verein (2010), Report to GHRST11 from Australia–BLUElink and IMOS, in Proceedings of the 11th GHRST Science Team Meeting, Lima, 21–25 June 2010. <https://www.ghrsst.org/documents/q/category/ghrsst-science-team-meetings/ghrsst-xi-lima-peru/>
- Beggs, H., L. Majewski, C. Griffin, R. Verein, P. Sakov, X. Huang, L. Garde, and C. Tingwell (2013). Report to GHRST14 from Australia - BlueLink and IMOS, in Proceedings of the 14th GHRST Science Team Meeting, Woods Hole, USA, 17–21 June 2013. <https://www.ghrsst.org/documents/q/category/ghrsst-science-team-meetings/ghrsst-xiv-woods-hole-ma-usa/>
- Böhm, E., S. Marullo, and R. Santoleri (1991), AVHRR visible-IR detection of diurnal warming events in the western Mediterranean Sea, *Int. J. Remote Sens.*, 12(4), 695-701, doi:10.1080/01431169108929686.
- Brunke, M. A., X. Zeng, V. Misra, and A. Beljaars (2008), Integration of a prognostic sea surface skin temperature scheme into weather and climate models, *J. Geophys. Res.*, 113(D21), doi:10.1029/2008JD010607.

Caputi, N., De Lestang, S., Feng, M. and Pearce, A. (2009), Seasonal variation in the long term warming trend in water temperature off the Western Australian coast. *Mar. Freshwater Res.*, 60(2), 129-139, doi: 10.1071/MF08199.

Clayson, C. A., and A. S. Bogdanoff (2013), The effect of diurnal sea surface temperature warming on climatological air-sea fluxes, *J. Clim.*, 26(8), 2546-2556, doi:http://dx.doi.org/10.1175/JCLI-D-12-00062.1.

Clayson, C. A., and D. Weitlich (2005), Diurnal warming in the tropical Pacific and its interannual variability, *Geophys. Res. Lett.*, 32(21), doi:10.1029/2005GL023786.

Eastwood, S., P. Le Borgne, S. Péré, and D. Poulter (2011), Diurnal variability in sea surface temperature in the Arctic, *Remote Sens. Environ.*, 115(10), 2594-2602, doi:10.1016/j.rse.2011.05.015.

Fairall, C., E. F. Bradley, J. Godfrey, G. Wick, J. B. Edson, and G. Young (1996), Cool-skin and warm-layer effects on sea surface temperature, *J. Geophys. Res.*, 101(C1), 1295-1308, doi:10.1029/95JC03190.

Feng, M. and G. Meyers, (2003), Interannual variability in the tropical Indian Ocean: a two-year time-scale of Indian Ocean Dipole. *Deep-sea Res. II*, 50(12), 2263-2284, doi:10.1016/S0967-0645(03)00056-0.

Gentemann, C. L., C. J. Donlon, A. Stuart-Menteth, and F. J. Wentz (2003), Diurnal signals in satellite sea surface temperature measurements, *Geophys. Res. Lett.*, 30(3), doi:10.1029/2002GL016291.

Gentemann, C. L., P. J. Minnett, and B. Ward (2009), Profiles of ocean surface heating (POSH): A new model of upper ocean diurnal warming, *J. Geophys. Res.*, 114(C7), doi:10.1029/2008JC004825.

Gentemann, C. L., P. J. Minnett, P. Le Borgne, and C. J. Merchant (2008), Multi-satellite measurements of large diurnal warming events, *Geophys. Res. Lett.*, 35, L22602, doi:10.1029/2008GL035730.

Gentilli, J. (1971), Australian climatic factors. In Gentilli, J., editor, *World survey of climatology*, Amsterdam, Elsevier, 35-52.

GHR SST Science Team (2010), The Recommended GHR SST Data Specification (GDS) 2.0, document revision 5, available from the GHR SST International Project Office, 2011, pp 123.

Godfrey, J.S. and K. R. Ridgway, (1985), The large-scale environment of the poleward-flowing Leeuwin Current, Western Australia: longshore steric height gradients, wind stresses and geostrophic flow. *J. Phys. Oceanogr.*, 15(5), 481-495, doi:[http://dx.doi.org/10.1175/1520-0485\(1985\)015<0481:TLSEOT>2.0.CO;2](http://dx.doi.org/10.1175/1520-0485(1985)015<0481:TLSEOT>2.0.CO;2)

Halpern, D., and R. K. Reed (1976), Heat budget of the upper ocean under light winds, *J. Phys. Oceanogr.*, 6(6), 972-975, doi:[http://dx.doi.org/10.1175/1520-0485\(1976\)006<0972:HBOTUO>2.0.CO;2](http://dx.doi.org/10.1175/1520-0485(1976)006<0972:HBOTUO>2.0.CO;2).

Hawkins, J., R. Clancy, and J. Price (1993), Use of AVHRR data to verify a system for forecasting diurnal sea surface temperature variability, *Int. J. Remote Sens.*, 14(7), 1347-1357, doi:10.1080/01431169308953961.

Jourdain, N. C., A. S. Gupta, A. S. Taschetto, C. C. Ummenhofer, A. F. Moise, and K. Ashok (2013), The Indo-Australian monsoon and its relationship to ENSO and IOD in reanalysis data and the CMIP3/CMIP5 simulations, *Clim. Dyn.*, 41(11-12), 3073-3102, doi: 10.1007/s00382-013-1676-1

Karagali, I., and J. Høyer (2014), Characterisation and quantification of regional diurnal SST cycles from SEVIRI, *Ocean Sci.*, 10, 745-758, doi:10.5194/os-10-745-2014.

Karagali, I., J. Høyer, and C. Hasager (2012), SST diurnal variability in the North Sea and the Baltic Sea, *Remote Sens. Environ.*, 121, 159-170, doi:10.1016/j.rse.2012.01.016.

Karagali, I., J. L. Høyer, and C. J. Donlon (2017), Using a 1-D model to reproduce the diurnal variability of SST, *J. Geophys. Res. Oceans*, 122, 2945-2959, doi:10.1002/2016JC012542.

Kawai, Y., and A. Wada (2007), Diurnal sea surface temperature variation and its impact on the atmosphere and ocean: A review, *J. Oceanogr.*, 63(5), 721-744, doi:10.1007/s10872-007-0063-0.

Large, W. G., J. C. McWilliams, and S. C. Doney (1994), Oceanic vertical mixing: A review and a model with a nonlocal boundary layer parameterization, *Rev. Geophys.*, 32(4), 363-403, doi:10.1029/94RG01872.

Li, W., H. Liu, and Y. Yu (1994), Impacts of Diurnal Cycle of SST on the Intraseasonal Variation of Surface Heat Flux over the Western Pacific Warm Pool, *Adv. Atmos. Sci.*, 5, 793-806.

Marullo, S., P. Minnett, R. Santoleri, and M. Tonani (2016), The diurnal cycle of sea surface temperature and estimation of the heat budget of the Mediterranean Sea, *J. Geophys. Res.*, 121(11), 8351-8367, doi:10.1002/2016JC012192.

Masson, S., P. Terray, G. Madec, J.-J. Luo, T. Yamagata, and K. Takahashi (2012), Impact of intradaily SST variability on ENSO characteristics in a coupled model, *Clim. Dyn.*, 39(3-4), 681-707, doi:10.1007/s00382-011-1247-2.

Mellor, G. L., and T. Yamada (1982), Development of a turbulence closure model for geophysical fluid problems, *Rev. Geophys.*, 20(4), 851-875, doi:10.1029/RG020i004p00851.

National Meteorological and Oceanographic Centre Operations Bulletin Number 98 (2013), *APSI upgrade of the ACCESS-R Numerical Weather Prediction system*, Australian Bureau of Meteorology, Available at <http://www.bom.gov.au/australia/charts/bulletins/apob98.pdf>

- Nelson, D. M., W. O. Smith, (1991), Sverdrup revisited: Critical depths, maximum chlorophyll levels, and the control of Southern Ocean productivity by the irradiance-mixing regime. *Limnol. Oceanogr.*, 36(8), 1650-1661, doi:10.4319/lo.1991.36.8.1650
- Noh, Y., E. Lee, D. H. Kim, S. Y. Hong, M. J. Kim, and M. L. Ou (2011), Prediction of the diurnal warming of sea surface temperature using an atmosphere-ocean mixed layer coupled model, *J. Geophys. Res.*, 116(C11), doi:10.1029/2011JC006970.
- Pearce, A. and M. Feng (2007), Observations of warming on the Western Australian continental shelf. *Mar. Freshwater Res.*, 58(10), 914-920, doi:10.1071/MF07082.
- Pearce, A. F., and M. Feng (2013). The rise and fall of the “marine heat wave” off Western Australia during the summer of 2010/2011. *J. Mar. Syst.*, 111, 139-156, doi:10.1016/j.jmarsys.2012.10.009.
- Price, J. F., R. A. Weller, C. M. Bowers, and M. G. Briscoe (1987), Diurnal response of sea surface temperature observed at the long - term upper ocean study (34° N, 70° W) in the Sargasso Sea, *J. Geophys. Res.*, 92(C13), 14480-14490, doi:10.1029/JC092iC13p14480.
- Puri, K., G. Dietachmayer, P. Steinle, M. Dix, L. Rikus, L. Logan, M. Naughton, C. Tingwell, Y. Xiao, and V. Barras (2013), Implementation of the initial ACCESS numerical weather prediction system, *Aust. Meteorol. Oceanogr. J.*, 63, 265-284.
- Riley, G. A. (1957), Phytoplankton of the North Central Sargasso Sea, 1950-52. *Limnol. Oceanogr.*, 35, 228-234, doi:10.1002/lno.1957.2.3.0252
- Ruardij, P., and W. V. Raaphorst, (1995). Benthic nutrient regeneration in the ERSEM ecosystem model of the North Sea. *Neth. J. Sea Res.*, 33, 453-483, doi: 10.1016/0077-7579(95)90057-8

Soloviev, A., and R. Lukas (1997), Observation of large diurnal warming events in the near-surface layer of the western equatorial Pacific warm pool, *Deep-Sea Res. I*, 44(6), 1055-1076, doi:10.1016/S0967-0637(96)00124-0.

Suppiah, R. (1992). The Australian summer monsoon: a review. *Prog. in Phys. Geogr.*, 16(3), 283-318, doi:10.1177/030913339201600302.

Sverdrup, H. U., M. W. Johnson, and R. H. Fleming (1942), *The Oceans: Their physics, chemistry, and general biology*, Prentice-Hall New York.

Takaya, Y., J.-R. Bidlot, A. C. M. Beljaars, and P. A. E. M. Janssen (2010), Refinements to a prognostic scheme of skin sea surface temperature, *J. Geophys. Res.*, 115, C06009, doi:10.1029/2009JC005985.

Tett, P., and A. Walne, (1995). Observations and simulations of hydrography, nutrients and plankton in the southern North Sea. *Ophelia*, 42, 371-416, doi: 10.1080/00785326.1995.10431514.

Webster, P. J., C. A. Clayson, and J. A. Curry (1996), Clouds, radiation, and the diurnal cycle of sea surface temperature in the tropical western Pacific, *J. Clim.*, 9(8), 1712-1730, doi:http://dx.doi.org/10.1175/1520-0442(1996)009<1712:CRATDC>2.0.CO;2.

Weihs, R. R., and M. Bourassa (2014), Modeled diurnally varying sea surface temperatures and their influence on surface heat fluxes, *J. Geophys. Res.*, 119(7), 4101-4123, doi:10.1002/2013JC009489.

Weller, E., M. Feng, H. Hendon, J. Ma, S.P. Xie, and N. Caputi, (2012), Interannual variations of wind regimes off the subtropical Western Australia coast during austral winter and spring. *J. Clim.*, 25(16), 5587-5599, doi: 10.1175/JCLI-D-11-00324.1.

Wyrtki, K. (1962), The upwelling in the region between Java and Australia during the south-east monsoon. *Mar. Freshwater Res.*, 13(3), 217-225, doi: 10.1071/MF9620217.

Zeng, X., and A. Beljaars (2005), A prognostic scheme of sea surface skin temperature for modeling and data assimilation, *Geophys. Res. Lett.*, 32, L14605, doi:10.1029/2005GL023030.

Zhang, H., H. Beggs, L. Majewski, H. Wang, and A. Kiss (2016a), Investigating Sea Surface Temperature Diurnal Variation over the Tropical Warm Pool Using MTSAT-1R Data, *Remote Sens. Environ.*, 183, 1-12, doi: 10.1016/j.rse.2016.05.002.

Zhang, H., H. Beggs, X. H. Wang, A. E. Kiss, and C. Griffin (2016b), Seasonal patterns of SST diurnal variation over the Tropical Warm Pool region, *J. Geophys. Res. Oceans*, 121, 8077–8094, doi:10.1002/2016JC012210.

Zhu, X., P. J. Minnett, R. Berkelmans, J. Hendee, and C. Manfrino (2014), Diurnal warming in shallow coastal seas: Observations from the Caribbean and Great Barrier Reef regions. *Cont. Shelf Res.*, 82, 85-98, doi:10.1016/j.csr.2014.03.002

Zinke, J., A. Rountrey, M. Feng, S.P. Xie, D. Dissard, K. Rankenburg, J.M. Lough, and M.T. McCulloch, (2014) Corals record long-term Leeuwin current variability including Ningaloo Niño/Niña since 1795. *Nat. Commun.*, 5, doi:10.1038/ncomms4607.

script

2017GL075008-f01-z.tif

Authc

cript

2017GL075008-f02-z-.tif

Auth

ct

A

2017GL075008-f03-z.tif

2017GL075008-f04-z-.tif

st

A

2017GL075008-f05-z-.tif

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Wang, XH;Zhang, H

Title:

Effects of Australian Summer Monsoon on Sea Surface Temperature Diurnal Variation Over the Australian North-Western Shelf

Date:

2017-10-16

Citation:

Wang, X. H. & Zhang, H. (2017). Effects of Australian Summer Monsoon on Sea Surface Temperature Diurnal Variation Over the Australian North-Western Shelf. *Geophysical Research Letters*, 44 (19), pp.9856-9864. <https://doi.org/10.1002/2017GL075008>.

Persistent Link:

<http://hdl.handle.net/11343/293551>