DR. MARK J HANLY (Orcid ID : 0000-0002-9279-7453)

Article type : Original Article

Gestational age and child development at age five in a population-based cohort of 97,989 Australian Aboriginal and non-Aboriginal children

Author list: Mark Hanly^a, Kathleen Falster^{a,b,c}, Georgina Chambers^{a,d}, John Lynch^{e,f}, Emily Banks^{b,g}, Nusrat Homaira^d, Marni Brownell^h, Sandra Eadesⁱ, Louisa Jorm^a

Author affiliations

^a Centre for Big Data Research in Health, UNSW Australia, Sydney, Australia

^b National Centre for Epidemiology and Population Health, Australian National University, Canberra, Australia

^c Centre for Social Research Methods, Australian National University, Canberra, Australia

^d School of Women's and Children's Health, UNSW Australia, Sydney, Australia

^e School of Public Health, University of Adelaide, Adelaide, Australia

^f School of Social and Community Medicine, University of Bristol, UK

^g The Sax Institute, Sydney, Australia

^h Manitoba Centre for Health Policy, Department of Community Health Sciences, Max Rady College of Medicine, Rady Faculty of Health Sciences, University of Manitoba, Winnipeg, Canada

ⁱ Baker Heart and Diabetes Institute, Melbourne, Australia

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/ppe.12426](https://doi.org/10.1111/ppe.12426)

This article is protected by copyright. All rights reserved

1 **Corresponding author**

2 Dr Mark Hanly

3 Centre for Big Data Research in Health

4 Level 1, AGSM Building (G27)

5 UNSW Sydney NSW 2052 Australia

6 m.hanly@unsw.edu.au

7

8 Short title Gestational age and child development in Australian children

Abstract

Background: Preterm birth and developmental vulnerability are more common in Australian Aboriginal compared with non-Aboriginal children. We quantified how gestational age relates to developmental vulnerability in both populations.

Methods: Perinatal datasets were linked to the Australian Early Development Census (AEDC), which collects data on five domains, including physical, social, emotional, language/cognitive, and general knowledge/communication development. We quantified the risk of developmental vulnerability on ≥ 1 domains at age 5, according to gestational age and Aboriginality, for 97,989 children born in New South Wales, Australia, who started school in 2009 or 2012.

Results: 7,079 children (7%) were Aboriginal. Compared with non-Aboriginal children, Aboriginal children were more likely to be preterm (5% versus 9%), and developmentally vulnerable on ≥ 1 domains (20% versus 36%). Overall, the proportion of developmentally vulnerable children decreased with increasing gestational age, from 44% at ≤ 27 weeks to 20% at 40 weeks. Aboriginal children had higher risks than non-Aboriginal children across the gestational age range, peaking among early term children (RD 19.0, 95% CI 16.3, 21.7; RR 1.91, 95% CI 1.77, 2.06). The relation of gestational age to developmental outcomes was the same in Aboriginal and non-Aboriginal children, and adjustment for socioeconomic disadvantage attenuated the risk differences and risk ratios across the gestational age range.

Conclusions: Although the relation of gestational age to developmental vulnerability was similar in Aboriginal and non-Aboriginal children, Aboriginal children had a higher risk of developmental vulnerability at all gestational ages, which was largely accounted for by socioeconomic disadvantage.

Keywords: Gestational age; Preterm birth; Early childhood development; Indigenous population; Linked administrative data.

1 Introduction

2 Birth before 39 weeks' gestation is a leading risk factor for adverse neonatal outcomes¹ and
3 impaired development during early childhood and beyond.^{2,3} By school age, children born preterm
4 have an increased risk of developmental deficits on multiple domains.⁴ The risk of adverse
5 development is highest for extremely preterm babies, and declines with each additional week of
6 gestational age at birth through to full term.⁵⁻⁸ Recent evidence suggests that postterm babies, born
7 after 41 weeks' gestation, also have an elevated risk of special educational needs⁷ and developmental
8 vulnerability,⁹ broadly similar to children born at 37 weeks' gestation.

9 A social gradient in the prevalence of preterm birth has been observed in numerous settings¹⁰
10 and using alternative measures of disadvantage.¹¹ There is also an association between disadvantage
11 and early childhood development.¹² Accordingly, socioeconomic disadvantage potentially confounds
12 the relationship between gestational age and child development. Socioeconomic disadvantage may
13 also moderate the effect of gestational age on child development—deprivation during the early
14 childhood years may exacerbate the effects of prematurity, while relative affluence may be
15 protective.^{1,13,14} These moderating effects may arise through differential access to resources¹⁵ or
16 parenting behaviours¹⁶ that promote positive development.

17 In Australia, Aboriginal children are more likely to be born preterm,¹⁷ and developmentally
18 vulnerable at school entry,¹⁸ compared with their non-Aboriginal peers. It is unknown, however,
19 whether the relationship between gestational age and early childhood development is similar in both
20 populations. Aboriginal children are more likely to experience disadvantage from the time they are in
21 utero;¹⁹ the effect of preterm birth on child development may therefore be more severe for Aboriginal
22 compared with non-Aboriginal children due to the potential modifying effects of disadvantage during
23 early childhood.^{13,14} Combined with higher rates of preterm birth in Aboriginal children, this may
24 explain some of the disparity in child development between Aboriginal and non-Aboriginal children,
25 and inform targets for intervention to reduce the Aboriginal – non-Aboriginal development gap.

26 To better understand whether the association between gestational age and child development
27 differs between Aboriginal and non-Aboriginal children, and to explore the contribution of
28 socioeconomic disadvantage to this relationship, we used linked health and education data from
29 population-level administrative datasets to quantify child development outcomes at age five,
30 according to gestational age at birth, in Australian Aboriginal and non-Aboriginal children.

31 Methods

32 *Data sources and study population*

1 This study used the Seeding Success data resource, a population-based cohort of children who
2 were born in New South Wales (NSW), Australia, and participated in the Australian Early
3 Development Census (AEDC) in 2009 or 2012. The source data and characteristics of the cohort, have
4 been described in detail elsewhere.^{20,21} Briefly, AEDC data were linked to administrative datasets in
5 NSW, including: Perinatal Data Collection (PDC), the Register of Births, Deaths and Marriages
6 (RBDM) birth registrations, Admitted Patients Data Collection (APDC), and Public-School
7 Enrolments.

8 The Seeding Success cohort (n=166,278) was restricted to singleton children without
9 medically diagnosed special needs, and with available gestational age and AEDC data (n=151,088).
10 The study population was further restricted to those enrolled in a NSW Public School in 2009 or 2012
11 (n=97,989), because socioeconomic measures were unavailable for children enrolled in non-
12 government schools.

13 *Gestational age and Aboriginality*

14 Gestational age was recorded in the PDC as the number of completed weeks, using the best
15 available clinical estimate based on last menstrual period, and early ultrasound, where available.
16 Gestational age has previously been found to be recorded with a high degree of reliability in NSW
17 perinatal data.²² For descriptive statistics of the whole study population, outcomes were analysed by
18 week of gestation, from 27 to 43 weeks, with children born prior to 28 weeks (n=123), or after 42
19 weeks (n=141) grouped as ≤ 27 weeks and ≥ 43 weeks, respectively. For analyses of Aboriginal and
20 non-Aboriginal children, gestational age was recoded into six categories, because of small numbers of
21 Aboriginal children at either extreme of the gestational age range. Gestational age categories
22 included: early/moderate preterm (≤ 33 weeks), late preterm (34-36 weeks), early term (37-38 weeks),
23 full term (39-40 weeks), late term (41 weeks) and postterm (≥ 42 weeks).²³

24 Aboriginality was defined using multiple data sources to enhance Aboriginal identification,
25 which is often under-reported in administrative datasets.²⁴ Children were classified as Aboriginal if
26 the child or either parent was recorded as Aboriginal and/or Torres Strait Islander on any birth record
27 from the PDC, RBDM or APDC, or their AEDC record at school age.

28 *Early childhood development outcome*

29 The outcome of interest was a binary indicator of developmental vulnerability on ≥ 1 domains
30 measured by the AEDC. The AEDC is a nationwide census of Australian school starters implemented
31 triennially since 2009, and includes a population measure of early childhood development on five
32 domains: physical health and wellbeing; social competence; emotional maturity; language and
33 cognitive skills; and communication and general knowledge.²⁵ The AEDC is completed in the second

term (May-July), for all children enrolled in the first year of school, by a teacher who has known the child for at least two months. Coverage is almost total in NSW, with 99.9% of school starters assessed in 2009 and 97.3% in 2012.¹⁸ Children were classified as developmentally vulnerable if they scored in the lowest decile on one or more of the five AEDC domains, based on the 2009 thresholds.²⁶

Explanatory variables

Potential explanatory variables were identified *a priori* using directed acyclic graphs (available on request) and included: child's sex, parity, maternal age at childbirth, maternal education, parental occupation, maternal region of birth, maternal partnership status at childbirth, private health insurance/patient status at childbirth, smoking during pregnancy, antenatal care attendance, maternal comorbidities during pregnancy (including pre-existing and gestational hypertension and/or diabetes), preschool attendance in the year before school, school starting age, and English as a Second Language (ESL). Area-level variables, assigned according to the mother's statistical local area of residence at the child's birth, included: geographical remoteness, measured by the Accessibility/Remoteness Index of Australia (ARIA+);²⁷ and area-level socioeconomic disadvantage, measured by the Australian Bureau of Statistics' Index of Relative Socioeconomic Advantage and Disadvantage.²⁸ Where available covariates were hypothesised to lie on the causal pathway between gestational age and development, they were excluded from the analysis (e.g. five minute Apgar score, admission to special care nursery/NICU at birth, and resuscitation at birth).

Missing data

Most variables used in the analysis had less than 3% missing data, apart from maternal education (8.5%), parental occupation (6.8%), and preschool attendance (6.5%) (Supplementary Table S1). Complete covariate information was available for 80% (n=82,089) of the study population. Aboriginal children with poor development were more likely to have incomplete data, therefore, multiple imputation was used to limit potential missing data bias. Imputations were derived using fully conditional specification, and each incomplete variable was modelled in turn using the outcome and covariates from the analysis model as predictors. The adjusted analyses presented here were performed by pooling estimates from the ten imputed datasets using standard rules.²⁹

Statistical methods

We estimated the proportion of children with developmental vulnerability for the study population, with 95% confidence intervals, for each week of gestational age from 27 to 43 weeks. We then estimated the proportion of Aboriginal and non-Aboriginal children with developmental vulnerability for each gestational age category. To assess the independent associations between child development and both gestational age and Aboriginality, we fitted crude and adjusted logistic

1 regression models to estimate the risk difference and relative risk of developmental vulnerability
2 within each gestational age category compared with children born at full term, and for Aboriginal
3 compared with non-Aboriginal children. The risk estimates were calculated from the fitted logistic
4 regression parameters, by aggregating individual probability estimates to derive estimates of risk for
5 subgroups of interest.³⁰ This approach performs comparably to other common approaches to risk ratio
6 estimation,³¹ and avoided the common issue of non-converging log-binomial models.

7 To test whether the relationship between gestational age and child development differed
8 statistically between Aboriginal and non-Aboriginal children, we refitted the model to include a set of
9 dummy variables for the interaction between Aboriginality and gestational age. The fitted model
10 parameters were used to estimate the risk difference and relative risk of developmental vulnerability
11 for Aboriginal compared with non-Aboriginal children within each gestational age group. The
12 presence of effect modification was assessed from the fully adjusted model on the additive scale by
13 calculating the relative excess risk due to interaction (RERI),³² and on the multiplicative scale, by
14 examining the statistical significance of the interaction coefficients.

15 To quantify how much of the inequality in developmental vulnerability between Aboriginal
16 and non-Aboriginal children could be explained by differences in measured socioeconomic,
17 demographic and perinatal characteristics, the interaction model was fitted repeatedly, adding sets of
18 covariates sequentially. Model 0 was unadjusted. Model 1 adjusted for family-level socioeconomic
19 indicators, including low maternal education, low-skilled parental occupation and private health
20 insurance/patient status. Model 2 included all remaining perinatal and demographic covariates from
21 Table 1.

22 For all models, standard errors were calculated using the delta method, and were inflated to
23 account for the clustering of similar children within schools. Analyses were carried out using Stata
24 (version 12).³³

25 *Sensitivity analyses*

26 To assess the impact of excluding children attending non-government schools from the main
27 analysis, we compared the unadjusted association between gestational age and developmental
28 vulnerability in the analysis sample and all children with complete exposure and outcome data. To
29 assess the consistency of findings across different development domains, we examined the
30 relationship between gestational age and developmental vulnerability by Aboriginality on the five
31 AEDC domains. To assess the effect of multiple imputation, we compared the distribution of the
32 study variables by Aboriginal status for children with complete covariate data and the imputed
33 sample. Because the children with special needs are not assigned to an AEDC developmental
34 vulnerability category, and gestational age may be associated with special needs, we conducted a

sensitivity analysis under a ‘worst-case scenario’ assumption whereby all children with special needs were assumed to be developmentally vulnerable.

Ethics approval

Ethical approval was obtained from the NSW Population Health Services and Research Ethics Committee (2014/04/523), the NSW Aboriginal Health and Medical Research Council Ethics Committee (1031/14) and the Australian National University Human Research Ethics Committee (2014/384).

Results

Study population

Of the total 97,989 children, 7,079 (7.2%) were Aboriginal. A higher proportion of Aboriginal children were born early preterm (0.3% versus 0.1%), moderate preterm (1.8% versus 1.0%), late preterm (6.5% versus 3.8%), and early term (22.2% versus 21.3%), compared with non-Aboriginal children (Table 1). Higher proportions of Aboriginal children had indicators of socioeconomic disadvantage compared with their non-Aboriginal counterparts, including low maternal education (19.9% versus 5.6%), low parental occupation level (59.3% versus 24.3%) and lower rates of private health insurance/patient status at the child’s birth (6.6% versus 32.9%) (Table 1).

Gestational age, Aboriginality and child development outcomes

Overall, 20.8% (95% CI 20.6, 21.1) of children were developmentally vulnerable at age five. The proportion of developmentally vulnerable children was highest among those born ≤ 27 weeks’ gestation, at 43.9% (95% CI 35.1, 52.1), decreasing with increasing gestational age to 19.8% (95% CI 19.4, 20.3) among children born at 40 weeks (Figure 1). Among children born at ≥ 43 weeks, 27.0% (95% CI: 19.6, 34.3) were developmentally vulnerable. The proportion of Aboriginal children who were developmentally vulnerable was 35.9% (95% CI: 34.8, 37.0) compared with 19.7% (95% CI: 19.4, 19.9) of non-Aboriginal children. A higher proportion of Aboriginal children were developmentally vulnerable across the gestational age range (Figure 2).

Modelling results

Compared to full term children, the adjusted risk difference for developmental vulnerability was 1.8 percentage points (95% CI 1.1, 2.4) for early term children, 3.9 percentage points (95% CI 2.6, 5.1) for late preterm children, and 7.5 percentage points (95% CI 5.1, 9.9) for early/moderate preterm children (Table 2). The adjusted relative risks, compared with full term children, were 1.09 (95% CI 1.05, 1.12) for early term children, 1.19 (95% CI 1.13, 1.26) for late preterm children, and

1 1.37 (95% CI 1.25, 1.49) for early/moderate preterm children, respectively. There was no difference
2 in risk, in absolute or relative terms, between late or postterm children, compared with full term
3 children.

4 The burden of developmental vulnerability between Aboriginal and non-Aboriginal children
5 differed by 16.3 percentage points (95% CI 14.7, 17.8) in absolute terms, and was 83% higher (RR
6 1.83, 95% CI 1.74, 1.91) in relative terms (Table 2). After adjusting for differences in measured
7 covariates, the Aboriginal to non-Aboriginal children risk difference was attenuated to 5.4 percentage
8 points (95% CI, 4.3, 6.6), and the relative risk to 1.27 (95% CI 1.21, 1.32).

9 There was slight variation in the magnitude of the Aboriginal to non-Aboriginal risk
10 differences and relative risks of developmental vulnerability across the gestational age range (Table
11 3). The smallest absolute difference was observed in postterm children (RD 12.5, 95% CI 3.8, 21.2;
12 RR 1.62, 95% CI 1.17, 2.07), and the largest difference was among early term children (RD 19.0,
13 95% CI 16.3, 21.7; RR=1.91, 95% CI 1.77, 2.06) (Table 3; Model 0). Adjustment for family-level
14 socioeconomic indicators attenuated the Aboriginal-to-non-Aboriginal risk differences and relative
15 risks across the gestational age range. For example, for full term children, the unadjusted risk
16 difference was 15.6 percentage points (95% CI 13.8, 17.4), reduced to 7.1 percentage points (95% CI
17 5.6, 8.5) after adjustment for family-level socioeconomic indicators; this equates to a 54% relative
18 reduction (Table 3; Model 1). After adjusting for all available covariates, there were no interactions
19 between gestational age and Aboriginality on the additive scale (assessed using the RERI statistic) or
20 the multiplicative scale (assessed using the coefficient of the interaction term), indicating the
21 association between gestational age and child development was similar in Aboriginal and non-
22 Aboriginal children (Table 3; Model 2).

23 *Sensitivity analyses*

24 The relation of gestational age to developmental vulnerability was similar in the
25 analysis sample (N=97,989) and the sample with complete outcome and exposure data (N=151,088)
26 (Supplementary Figure S1). The pattern of association between gestational age and developmental
27 vulnerability for Aboriginal and non-Aboriginal children was broadly similar on the five
28 developmental domains (Supplementary Figure S2), and the aggregate outcome (i.e. vulnerable on ≥ 1
29 domains). Compared to the sample with complete covariate data, the imputed sample had a higher
30 proportion of children with developmental vulnerability, low maternal education and low paternal
31 occupation, especially among Aboriginal children (Supplementary Table S2). Accordingly, multiple
32 imputation enabled us to include data for more vulnerable children in the analysis.

33 Because children with special needs did not have AEDC outcome data, 4,072 (4.3%) non-
34 Aboriginal children and 531 (7.0%) Aboriginal children were excluded from the main analysis. When

children with special needs were included in the developmental vulnerability group, the proportion of children classified as vulnerable increased to 40.4% [95% CI: 39.3, 41.5] among Aboriginal children, and 23.1% [95% CI: 22.8, 23.4] among non-Aboriginal children, but, consistent with the main analysis, the relationship between gestational age and child development was similar in Aboriginal and non-Aboriginal children (Supplementary Figure S3).

Comment

Principal findings

In this contemporary population-based cohort of almost 100,000 Australian children, preterm children had a higher risk of developmental vulnerability, with increasing risks at earlier gestational ages. Although the association between gestational age and child development at age five was similar in Aboriginal and non-Aboriginal children, Aboriginal children had a higher risk of developmental vulnerability across the gestational age range. Within each gestational age category, adjustment for available socioeconomic, demographic and perinatal factors accounted for a substantial proportion of the absolute and relative inequalities between Aboriginal and non-Aboriginal children.

Interpretation

The general finding that each additional week of gestational age was associated with a decrease in the risk of developmental vulnerability from ≤ 27 weeks through to 40 weeks' gestation is consistent with a study of the same population that did not explore outcomes for Aboriginal children⁵ and studies in Scotland and the USA that examined child outcomes beyond the perinatal period by week of gestational age among more than 100,000 children.^{7,13} We also found that children born at 43 weeks' gestation had an elevated risk of developmental vulnerability, with similar risks compared to children born at 37 weeks' gestation, although the confidence intervals were wide, and overlapped with the estimate for children born at 40 weeks. A similar U-shaped pattern of risk in the full-term period between 37-43 weeks' gestation was found for developmental vulnerability at age 5 among more than 12,000 South Australian children,⁹ and for special educational needs among more than 360,000 Scottish children.⁷

For the first time, this study quantifies at scale the relationship between gestational age and early childhood development in Aboriginal children, a disproportionately disadvantaged population group in Australia. Although it is well known that preterm birth is more common in Aboriginal compared with non-Aboriginal children,¹⁷ this study shows that Aboriginal children have a higher risk of developmental vulnerability across the whole gestational age range. We also established that the overall relation of gestational age to child development is similar in Aboriginal and non-Aboriginal children. Aboriginal children were more likely than non-Aboriginal children to be disadvantaged, as

1 indicated by maternal schooling, parental occupation, and private health insurance/patient status. We
2 have shown that differences in the distribution of these indicators of socioeconomic
3 advantage/disadvantage account for a substantial proportion of the absolute and relative gap in child
4 development outcomes between Aboriginal and non-Aboriginal children at all gestational ages.
5 Although Aboriginal children are more likely to be born preterm, and to experience a disproportionate
6 burden of developmental vulnerability, the absence of significant interaction terms in our study
7 suggests that the relationship of gestational age to child development at age five is broadly similar in
8 Aboriginal and non-Aboriginal children.

9 *Strengths of the study*

10 The strengths of this study include the high population coverage of the source data, and the large
11 sample size, which enabled us to quantify outcomes for Aboriginal children, a small and vulnerable
12 population group. Linkage of multiple cross-sectoral administrative datasets enabled enhanced
13 identification of Aboriginal children compared with a single data source.²⁴

14 *Limitations of the data*

15 We were limited to covariates available from the source data, which were collected primarily for
16 administrative purposes, rather than research. Because there is likely to be residual confounding,
17 adjusted risk estimates of the association between gestational age and child development in
18 Aboriginal and non-Aboriginal children also include the effect of unmeasured covariates, including
19 the intergenerational effects of discrimination, marginalisation and removal of children that are
20 pertinent to Australian Aboriginal children and their families.³⁴ As key family-level socioeconomic
21 variables were not available for children enrolled in non-government schools, the study population
22 was restricted to the 70% of children who attended NSW Public Schools. However, sensitivity
23 analyses illustrate that the relationship between gestational age and child development outcomes at
24 age five were consistent in the study population and the broader population of children enrolled in
25 both non-government and government schools who had complete exposure and outcome data. An
26 additional limitation is that children with special needs were excluded from the main analysis because
27 the AEDC is not validated for use in this population. This was potentially problematic because
28 preterm birth is associated with an increased risk of special educational needs,⁷ and Aboriginal
29 children were disproportionately excluded from the analysis due to a higher prevalence of special
30 needs. However, our sensitivity analysis, that added children with special needs to the group of
31 children classified as developmentally vulnerable, showed a similar relationship between gestational
32 age and child development outcomes in Aboriginal and non-Aboriginal children, and did not alter the
33 qualitative conclusions of the study.

34 *Conclusions*

1 This study adds to growing evidence that, at a population level, increases of gestational age
2 up to 39 weeks are associated with a reduced risk of adverse child development outcomes. This
3 association is similar in both Aboriginal and non-Aboriginal children. Early preterm children are
4 routinely monitored and supported during early childhood in Australia; our study suggests there may
5 also be unmet developmental needs among late preterm and early term children.

6 Although the relationship between gestational age and early childhood development was
7 similar for Aboriginal and non-Aboriginal children, Aboriginal children had an increased risk of
8 developmental vulnerability by age five, regardless of gestational age. Our findings suggest that much
9 of the absolute and relative gap in outcomes between Aboriginal and non-Aboriginal children across
10 the gestational age range can be accounted for by the disproportionate burden of disadvantage
11 experienced by Aboriginal children. Accordingly, policies that improve the social and economic
12 conditions for disadvantaged children, including Aboriginal children, are likely to improve child
13 development in these vulnerable populations.

Funding

This work was supported by an Australian National Health and Medical Research Council (NHMRC) Project Grant (#1061713). KF was supported by an NHMRC Early Career Fellowship (#1016475) and an NHMRC capacity building grant (#573122). EB was supported by an NHMRC Senior Research Fellowship (#1042717). JL is funded by a National Health and Medical Research Council of Australia Partnership Project Grant (1056888) and Centre of Research Excellence (1099422). SE was supported by an NHMRC Career Development Fellowship (#1013418). MB was supported by the Manitoba Centre for Health Policy Population-Based Child Health Research Award.

Acknowledgements

The authors would like to thank the Australian Government Department of Education, the NSW Ministry of Health, the NSW Register of Births, Deaths and Marriages, the NSW Department of Education and the NSW Department of Family and Community Services (FACS) for allowing access to the data. The authors also thank the NSW Centre for Health Record Linkage for conducting the linkage of data sources detailed in this paper, and policy colleagues from NSW Health and FACS, including Emily Klineberg, Jessica Stewart and Michael Henman, for participation in discussions about this research. The Seeding Success investigator team comprises Louisa Jorm, Kathleen Falster, Sandra Eades, John Lynch, Emily Banks, Marni Brownell, Rhonda Craven, Kristjana Einarsdóttir, Deborah Randall, Sharon Goldfeld, Alastair Leyland, Elizabeth Best and Marilyn Chilvers.

References

1. Saigal S, Doyle LW. An overview of mortality and sequelae of preterm birth from infancy to adulthood. *The Lancet*. 2008;371:261-269.
2. Cheong JL, Doyle LW, Burnett AC, et al. Association between moderate and late preterm birth and neurodevelopment and social-emotional development at age 2 years. *JAMA Pediatrics*. 2017.
3. D'Onofrio BM, Class QA, Rickert ME, Larsson H, Långström N, Lichtenstein P. Preterm birth and mortality and morbidity. *JAMA Psychiatry*. 2013;70(11):1231.
4. Bhutta AT, Cleves MA, Casey PH, Cradock MM, Anand KJS. Cognitive and behavioral outcomes of school-aged children who were born preterm: A meta-analysis. *JAMA: Journal of the American Medical Association*. 2002;288(6):728-737.
5. Bentley JP, Roberts CL, Bowen JR, Martin AJ, Morris JM, Nassar N. Planned birth before 39 weeks and child development: A population-based study. *Pediatrics*. 2016;138(6).
6. Poulsen G, Wolke D, Kurinczuk JJ, et al. Gestational age and cognitive ability in early childhood: A population-based cohort study. *Paediatric and Perinatal Epidemiology*. 2013;27:371-379.
7. Mackay DF, Smith GCS, Dobbie R, Pell JP. Gestational age at delivery and special educational need: Retrospective cohort study of 407,503 schoolchildren. *PLoS Medicine*. 2010;7(6):1-10.
8. Kerstjens JM, De Winter AF, Bocca-Tjeertes IF, Bos AF, Reijneveld SA. Risk of developmental delay increases exponentially as gestational age of preterm infants decreases: A cohort study at age 4years. *Developmental Medicine and Child Neurology*. 2012;54(12):1096-1101.
9. Smithers LG, Searle AK, Chittleborough CR, Scheil W, Brinkman SA, Lynch JW. A whole-of-population study of term and post-term gestational age at birth and children's development. *BJOG*. 2015;122(10):1303-1311.
10. Poulsen G, Strandberg-Larsen K, Mortensen L, Barros H, Cordier S, Correia S. Exploring Educational Disparities in Risk of Preterm Delivery: A Comparative Study of 12 European Birth Cohorts. *Paediatric and Perinatal Epidemiology*. 2015;29:172-183.
11. Petersen CB, Mortensen LH, Morgen CS, et al. Socio-economic inequality in preterm birth: A comparative study of the Nordic countries from 1981 to 2000. *Paediatric and Perinatal*

- Epidemiology*. 2009;23(1):66-75.
12. Nicholson JM, Lucas N, Berthelsen D, Wake M. Socioeconomic inequality profiles in physical and developmental health from 0-7 years: Australian National Study. *Journal of Epidemiology & Community Health*. 2012;66(1):81-87.
13. Richards JL, Chapple-McGruder T, Williams BL, Kramer MR. Does neighborhood deprivation modify the effect of preterm birth on children's first grade academic performance? *Social Science & Medicine*. 2015;132:122-131.
14. Gisselmann M, Koupil I, De Stavola BL. The combined influence of parental education and preterm birth on school performance. *Journal of Epidemiology & Community Health*. 2011;65(9):764-769.
15. Burger K. How does early childhood care and education affect cognitive development? An international review of the effects of early interventions for children from different social backgrounds. *Early Childhood Research Quarterly*. 2010;25(2):140-165.
16. Smith KE, Landry SH, Swank PR. The role of early maternal responsiveness in supporting school-aged cognitive development for children who vary in birth status. *Pediatrics*. 2006;117(5):1608-1618.
17. Centre for Epidemiology and Evidence. *NSW Mothers and Babies 2014*. Sydney: NSW Ministry of Health; 2016. <http://www.health.nsw.gov.au/hsnsw/Pages/Mothers-and-Babies-2014.aspx> [Accessed 24/09/17].
18. Australian Government. *A Snapshot of Early Childhood Development in Australia - AEDI National Report*. Canberra; 2013. <https://www.aedc.gov.au/resources/detail/national-report-2012> [Accessed 24/09/17].
19. Australian Institute of Health and Welfare. *The Health and Welfare of Australia's Aboriginal and Torres Strait Islander Peoples: 2015*. Canberra; 2015. <https://www.aihw.gov.au/reports/indigenous-health-welfare/indigenous-health-welfare-2015/contents/table-of-contents> [Accessed 24/09/17].
20. Falster K, Jorm L, Eades S, et al. What factors contribute to positive early childhood health and development in Australian Aboriginal children? Protocol for a population-based cohort study using linked administrative data (The Seeding Success Study). *BMJ Open*. 2015;5:1-13.
21. Falster K, Jorgensen M, Hanly M, et al. Data Resource Profile: Seeding Success: a cross-sectoral data resource for early childhood health and development research in Australian

- 1 Aboriginal and non-Aboriginal children. *International Journal of Epidemiology*. 2017.
2 <https://doi.org/10.1093/ije/dyx051> [Accessed 24/09/17].
- 3 22. Taylor L, Pym M, Bajuk B, Sutton L, Travis S, Banks C. *New South Wales Mothers and*
4 *Babies 1998*. Sydney; 2000. [http://www.health.nsw.gov.au/phb/Pages/NSW-mothers-babies-](http://www.health.nsw.gov.au/phb/Pages/NSW-mothers-babies-1998.aspx)
5 [1998.aspx](http://www.health.nsw.gov.au/phb/Pages/NSW-mothers-babies-1998.aspx) [Accessed 24/09/17].
- 6 23. Richards JL, Drews-Botsch C, Sales JM, Flanders D, Kramer R. Describing the shape of the
7 relationship between gestational age at birth and cognitive development in a nationally
8 representative U.S. birth cohort. *Paediatric and Perinatal Epidemiology*. 2016;30:571-582.
- 9 24. Gialamas A, Pilkington R, Berry J, et al. Identification of Aboriginal children using linked
10 administrative data: Consequences for measuring inequalities. *Journal of Paediatrics and*
11 *Child Health*. 2016;52:534-540.
- 12 25. Brinkman S, Gregory T, Goldfeld S, Lynch J, Hardy M. Data Resource Profile: The Australian
13 Early Development Index (AEDI). *International Journal of Epidemiology*. 2014;43(4):1089-
14 1096.
- 15 26. Brinkman S, Gregory T, Harris J, Hart B, Blackmore S, Janus M. Associations between the
16 early development instrument at age 5, and reading and numeracy skills at ages 8, 10 and 12:
17 A prospective linked data study. *Child Indicators Research*. 2013;6(4):695-708.
- 18 27. Australian Bureau Of Statistics. *Australian Statistical Geography Standard (ASGS): Volume 5*
19 *- Remoteness Structure*. Vol 5. Canberra; 2013.
- 20 28. Australian Bureau Of Statistics. *Socio-Economic Indexes for Areas (SEIFA)*. Canberra; 2011.
- 21 29. White IR, Royston P, Wood AM. Multiple imputation using chained equations: Issues and
22 guidance for practice. *Statistics in Medicine*. 2011;30(4):377-399.
- 23 30. Kleinman LC, Norton EC. What's the risk? A simple approach for estimating adjusted risk
24 measures from nonlinear models including logistic regression. *Health Services Research*.
25 2009;44(1):288-302.
- 26 31. Knol MJ, Le Cessie S, Algra A, Vandenbroucke JP, Groenwold RHH. Overestimation of risk
27 ratios by odds ratios in trials and cohort studies: alternatives to logistic regression. *Canadian*
28 *Medical Association Journal*. 2012;184(8):895-899.
- 29 32. Knol MJ, VanderWeele TJ. Recommendations for presenting analyses of effect modification
30 and interaction. *International Journal of Epidemiology*. 2012;41(2):514-520.

- 1 33. StataCorp. Stata Statistical Software: Release 12. 2011.
- 2 34. Brown N. History, Law, and Policy as a Foundation for Health Care Delivery for Australian
3 Indigenous Children. *Pediatric Clinics of North America*. 2009;56(6):1561-1576.

Figure legends

Figure 1. Percentage of children developmentally vulnerable on one or more AEDC domains, and number of births, by gestational age at birth.

Figure 2. Percentage of children developmentally vulnerable on one or more AEDC domains, by gestational age group and Aboriginality.

Supporting Information

Table S1. Number and percentage of missing observations for the study variables.

Table S2. Comparison of study variables by Aboriginal status for the sample with complete covariate data and the imputed sample.

Figure S1. Risk of developmental vulnerability by gestational age for the population of children with complete exposure and outcome data (N=151,088), the study population with exclusion criteria applied (N=97,989)

Figure S2. Percentage of children developmentally vulnerable on one or more AEDC domains, by gestational age group and Aboriginality for five developmental domains: physical health and wellbeing; social competence; emotional maturity; language and cognitive skills; and communication and general knowledge

- 1 **Figure S3.** Percentage of children developmentally vulnerable, or with medically diagnosed special
- 2 needs, on one or more AEDC domains, by gestational age group and Aboriginality.

Table 1. Child, mother and area-level characteristics at birth and school age, by Aboriginality, for the imputed study population.

	Non-Aboriginal		Aboriginal		Total	
	n	(%)	n	(%)	n	(%)
Characteristics of child, mother and area of residence						
<i>Child characteristics at birth</i>						
n (%)	90,910	100.0	7,079	100.0	97,989	100.0
Gestational age group						
Early preterm (22-27 weeks)	102	0.1	21	0.3	123	0.1
Moderate preterm (28-33 weeks)	866	1.0	128	1.8	994	1.0
Late preterm (34-36 weeks)	3,469	3.8	463	6.5	3,932	4.0
Early term (37-38 weeks)	19,381	21.3	1,570	22.2	20,951	21.4
Full term (39-40 weeks)	49,714	54.7	3,731	52.7	53,445	54.5
Late term (41 weeks)	15,590	17.1	1,046	14.8	16,636	17.0
Postterm (≥42 weeks)	1,788	2.0	120	1.7	1,908	1.9
Male sex	46,185	50.8	3,475	49.1	49,660	50.7
<i>Maternal characteristics at birth</i>						
Parity						
0	37,654	41.4	2,385	33.7	40,040	40.9
1	31,637	34.8	1,978	27.9	33,614	34.3

2+	21,619	23.8	2,716	38.4	24,335	24.8
Maternal age at childbirth						
<20	3,087	3.4	1,218	17.2	4,305	4.4
20-24	13,385	14.7	2,181	30.8	15,566	15.9
25-29	25,140	27.7	1,766	24.9	26,906	27.5
30-34	30,221	33.2	1,277	18.0	31,498	32.1
35+	19,077	21.0	637	9.0	19,714	20.1
Mother born in Australia	64,932	71.4	6,716	94.9	71,648	73.1
Mother has a partner	75,755	83.3	3,338	47.2	79,093	80.7
Private patient/insurance	29,921	32.9	470	6.6	30,391	31.0
Smoking during pregnancy	12,882	14.2	3,415	48.2	16,297	16.6
Antenatal care in first 20 weeks	81,106	89.2	5,702	80.5	86,807	88.6
Maternal comorbidity ^a	9,740	10.7	672	9.5	10,412	10.6
<i>Child characteristics at the start of school</i>						
Attended preschool/childcare	67,007	73.7	4,953	70.0	71,960	73.4
English as a second language	16,455	18.1	231	3.3	16,686	17.0
School starting age						
<5 years	31,196	34.3	2,485	35.1	33,681	34.4

5-5.5 years	36,946	40.6	2,763	39.0	39,709	40.5
≥5.5 years	22,768	25.0	1,831	25.9	24,599	25.1
AEDC census year						
2009	43,315	47.6	3,190	45.1	46,505	47.5
2012	47,595	52.4	3,889	54.9	51,484	52.5
<i>Maternal/family characteristics at the start of school</i>						
Low maternal education (≤9 years school)	5,048	5.6	1,407	19.9	6,455	6.6
Low skilled parental occupation(s) (Grade 4/unemployed)	22,118	24.3	4,199	59.3	26,317	26.9
<i>Area of residence</i>						
Geographic remoteness ^b						
Major City	60,033	66.0	2,701	38.2	62,734	64.0
Inner Regional	22,951	25.2	2,462	34.8	25,413	25.9
Outer Regional	7,397	8.1	1,495	21.1	8,892	9.1
Remote/Very Remote	529	0.6	422	6.0	950	1.0
<i>Area-level advantage/disadvantage^c</i>						
Q1 (Most disadvantaged)	7,885	8.7	1,558	22.0	9,443	9.6
Q2	9,656	10.6	1,330	18.8	10,987	11.2
Q3	31,305	34.4	2,950	41.7	34,255	35.0

Q4	19,321	21.3	877	12.4	20,198	20.6
Q5 (Most advantaged)	22,742	25.0	364	5.1	23,106	23.6

^a Includes pre-existing and gestational-onset diabetes and hypertension

^b Accessibility/Remoteness Index of Australia (ARIA+) based on mother's statistical local area of residence at the time of birth

^c Socio-Economic Indices for Areas (SEIFA) Index of Relative Socio-economic Advantage and Disadvantage population quintiles based on mother's statistical local area of residence at the time of birth.

Table 2. Risk difference (RD) and relative risk (RR) of developmental vulnerability according to gestational age and Aboriginality

		Total Births	Developmental vulnerability/special needs				
			n (%)	RD (95% CI)	aRD ^a (95% CI)	RR (95% CI)	aRR ^a (95% CI)
Gestational Age							
	Early/moderate preterm (≤33 weeks)	1117	362 (32.4)	12.6 (9.8, 15.3)	7.5 (5.1, 9.9)	1.63 (1.49, 1.77)	1.37 (1.25, 1.49)
	Late preterm (34-36 weeks)	3932	1062 (27.0)	7.2 (5.7, 8.6)	3.9 (2.6, 5.1)	1.36 (1.29, 1.44)	1.19 (1.13, 1.26)
	Early term (37-38 weeks)	20951	4654 (22.2)	2.4 (1.7, 3.0)	1.8 (1.1, 2.4)	1.12 (1.08, 1.15)	1.09 (1.05, 1.12)
	Full term (39-40 weeks)	43199	362 (19.8)	Reference	Reference	Reference	Reference
	Late term (41 weeks)	16636	3326 (20.0)	0.1 (-0.6, 0.8)	0.0 (-0.7, 0.7)	1.01 (0.97, 1.04)	1.00 (0.97, 1.03)
	Postterm (≥42 weeks)	1908	397 (20.8)	1.0 (-1.0, 2.9)	0.5 (-1.3, 2.2)	1.05 (0.95, 1.14)	1.02 (0.93, 1.11)
Aboriginality							
	non-Aboriginal	90910	17866 (19.7)	Reference	Reference	Reference	Reference
	Aboriginal	7079	2543 (35.9)	16.3 (14.7, 17.8)	5.4 (4.3, 6.6)	1.83 (1.74, 1.91)	1.27 (1.21, 1.32)

RD=Risk difference; aRD=Adjusted risk difference; RR=Relative risk; aRR=Adjusted relative risk

^aAdjusted models based on pooled estimates from 10 multiply imputed datasets, controlling for: mother's schooling, parent's occupation, Aboriginal status, child's sex, parity, maternal age, maternal country of birth, maternal partnership status, private insurance status, smoking during pregnancy, antenatal care in the first 20 weeks, maternal comorbidities (pre-existing or gestational-onset diabetes or hypertension), preschool attendance, school enrolment age, ESL status, geographic

Table 3. Absolute risk, risk difference and relative risk estimates from interaction models, comparing risk of developmental vulnerability for Aboriginal and non-Aboriginal children within each gestational age group.

	Risk estimates				Tests of effect modification			
	non-Aboriginal risk (95% CI)	Aboriginal Risk (95% CI)	Risk Difference (95% CI)	Relative Risk (95% CI)	RERI ^a	P	Beta ^b	P
Model 0 ^c								
Early/moderate preterm (≤33 weeks)	30.5 (27.5, 33.4)	45.0 (36.6, 53.4)	14.5 (5.6, 23.4)	1.48 (1.17, 1.78)	0.38	0.551	-0.19	0.308
Late preterm (34-36 weeks)	25.0 (23.4, 26.5)	42.1 (37.8, 46.5)	17.1 (12.6, 21.7)	1.69 (1.48, 1.89)	0.44	0.126	-0.04	0.720
Early term (37-38 weeks)	20.8 (20.0, 21.6)	39.8 (37.1, 42.5)	19.0 (16.3, 21.7)	1.91 (1.77, 2.06)	0.46	0.005	0.11	0.090
Full term (39-40 weeks)	18.8 (18.1, 19.4)	34.3 (32.5, 36.1)	15.6 (13.8, 17.4)	1.83 (1.72, 1.94)	Reference		Reference	
Late term (41 weeks)	19.2 (18.4, 20.0)	32.1 (29.1, 35.2)	12.9 (9.9, 16.0)	1.67 (1.51, 1.84)	-0.24	0.119	-0.13	0.098
Postterm (≥42 weeks)	20.0 (18.0, 22.1)	32.5 (23.9, 41.1)	12.5 (3.8, 21.2)	1.62 (1.17, 2.07)	-0.26	0.527	-0.16	0.426
Model 1 ^d								

	28.4 (25.7,	33.7 (26.3,		1.19 (0.90,				
Early-moderate preterm (≤ 33 weeks)	31.2)	41.2)	5.3 (-2.6, 13.2)	1.47)	-0.02	0.936	-0.16	0.427
	24.5 (23.1,	31.4 (27.6,		1.28 (1.11,				
Late preterm (34-36 weeks)	26.0)	35.1)	6.8 (2.9, 10.8)	1.45)	0.06	0.734	-0.06	0.558
	21.7 (20.9,	30.5 (28.2,		1.41 (1.30,				
Early term (37-38 weeks)	22.4)	32.8)	8.8 (6.5, 11.1)	1.52)	0.19	0.086	0.06	0.323
	19.5 (18.9,	26.6 (25.1,		1.36 (1.28,				
Full term (39-40 weeks)	20.1)	28.1)	7.1 (5.6, 8.5)	1.44)	Reference		Reference	
	19.1 (18.4,	24.9 (22.4,		1.31 (1.17,				
Late term (41 weeks)	19.8)	27.4)	5.8 (3.3, 8.3)	1.44)	-0.10	0.339	-0.06	0.425
	19.6 (17.7,	25.3 (18.0,		1.29 (0.91,				
Postterm (≥ 42 weeks)	21.5)	32.6)	5.8 (-1.7, 13.2)	1.68)	-0.10	0.730	-0.07	0.739
Model 2 ^e								
	27.4 (24.8,	31.1 (24.3,		1.14 (0.87,				
Early-moderate preterm (≤ 33 weeks)	29.9)	38.0)	3.8 (-3.5, 11.1)	1.41)	-0.03	0.939	-0.13	0.529
	23.5 (22.1,	28.4 (24.9,		1.21 (1.04,				
Late preterm (34-36 weeks)	24.9)	31.9)	4.9 (1.1, 8.6)	1.37)	0.03	0.885	-0.05	0.655
	21.4 (20.6,	28.4 (26.3,		1.33 (1.23,				
Early term (37-38 weeks)	22.1)	30.5)	7.1 (5.0, 9.2)	1.43)	0.19	0.068	0.09	0.175
Full term (39-40 weeks)	19.7 (19.2,	24.9 (23.5,	5.2 (3.8, 6.5)	1.26 (1.19,	Reference		Reference	

	20.3)	26.3)		1.33)				
	19.7 (19.0,	23.8 (21.4,		1.21 (1.08,				
Late term (41 weeks)	20.4)	26.3)	4.1 (1.6, 6.6)	1.34)	-0.09	0.425	-0.06	0.448
	20.2 (18.3,	24.3 (17.2,		1.20 (0.84,				
Postterm (≥42 weeks)	22.0)	31.4)	4.1 (-3.1, 11.4)	1.57)	-0.08	0.789	-0.07	0.770

^a Relative Excess Risk of Interaction (test of interaction on the additive scale); ^b Coefficient for the interaction dummy variable (test of interaction on the multiplicative scale) ^c Model 0 is unadjusted

^d Model 1 is adjusted for low maternal education, low parental occupation, and private insurance/patient status; ^e Model 2 is adjusted for all available covariates included in Table 1.

ppe_12426_f1.tif

ppe_12426_f2.tif

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Hanly, M;Falster, K;Chambers, G;Lynch, J;Banks, E;Homaira, N;Brownell, M;Eades, S;Jorm, L

Title:

Gestational Age and Child Development at Age Five in a Population-Based Cohort of Australian Aboriginal and Non-Aboriginal Children

Date:

2018-01-01

Citation:

Hanly, M., Falster, K., Chambers, G., Lynch, J., Banks, E., Homaira, N., Brownell, M., Eades, S. & Jorm, L. (2018). Gestational Age and Child Development at Age Five in a Population-Based Cohort of Australian Aboriginal and Non-Aboriginal Children. *PAEDIATRIC AND PERINATAL EPIDEMIOLOGY*, 32 (1), pp.114-125. <https://doi.org/10.1111/ppe.12426>.

Persistent Link:

<http://hdl.handle.net/11343/293907>