Title: Nurses' perceptions of preventing falls for patients with dementia in the acute hospital setting

Authors: Darshini Ayton¹, Penny O'Brien¹, Jonathan Treml², Sze-Ee Soh¹, Renata Morello¹, Anna Barker¹

1. Department of Epidemiology and Preventive Medicine, Monash University, Melbourne, AUSTRALIA

2. Elderly Care, University Hospitals Birmingham NHS Foundation Trust Queen Elizabeth Hospital Birmingham, Edgbaston, Birmingham, UNITED KINGDOM

Corresponding author

Associate Professor Anna Barker

Department of Epidemiology and Preventive Medicine

Monash University

99 Commercial Road, Melbourne VIC 3004, AUSTRALIA

Telephone: +61 3 9903 0946

Email: anna.barker@monash.edu

Acknowledgements

This study was funded by the National Health & Medical Research Council, Australia (APP1007627). The authors would like to acknowledge the contribution of the 6-PACK investigator team and the nurses who participated. The authors declare there are no conflicts of interest in relation to this research.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/ajag.12474](https://doi.org/10.1111/ajag.12474)

This article is protected by copyright. All rights reserved

MISS PENNY O'BRIEN (Orcid ID : 0000-0002-0334-5288)

Article type : Brief Report

Nurses' perceptions of preventing falls for patients with dementia in the acute hospital setting

Abstract

Objectives: Safe and high-quality care for patients with dementia is a key priority area of the Australian Commission on Safety and Quality in Health Care, however caring for patients with dementia in the acute hospital setting is perceived as challenging. The aim of this analysis was to explore nurses' perspectives regarding falls prevention for patients with dementia in an acute care setting.

Methods: Secondary analysis of focus group data. Focus groups were conducted with nurses (n=96) across six hospitals in New South Wales and Victoria.

Results: Nurses frequently reported issues relating to the physical environment of the acute care setting, competing priorities in a complex care setting and the need for one-on-one supervision for patients with dementia.

Conclusions: Nurses report that one-on-one supervision is required to keep patients safe. Future research examining the acceptability and cost effectiveness of volunteers providing this supervision is warranted in Australian hospitals.

Keywords: accidental falls, dementia, health services, nursing

Short Report (1089 words)

Introduction

People with dementia are high users of acute hospitals with between 20-25% of in-patients aged ≥ 70 and 47% of in-patients aged ≥ 90 having a diagnosis of dementia [1]. This is a relatively conservative assessment due to the known inaccuracy of coding dementia in hospital administrative data [2,3]. For a person with dementia, the impact of hospitalisation can be profound. These individuals may be more likely to experience preventable adverse events such as infections, dehydration, malnutrition and falls [1]. Impaired cognition has been shown to be the most commonly identified risk factor in patients who fall while in hospital [4]. Hence, safe and high-quality care for patients with dementia is a key priority area of the Australian Commission on Safety and Quality in Health Care [5].

The aim of this analysis was to explore nurses' perspectives regarding falls prevention for patients with dementia in an acute care setting.

Methods

This analysis was part of the 6-PACK project – a multi-centre cluster RCT testing the effectiveness of the 6-PACK falls prevention intervention on falls and fall injuries in six acute hospitals in New South Wales and Victoria [3]. In the pre-implementation phase of the project, nursing staff from 16 medical and 8 surgical wards were invited to participate in focus groups to explore current falls prevention practices [6]. Participants were 96 nurses working on participating wards. A member of the research team invited nurses to attend focus groups at handover times. The recruitment process has been described in detail elsewhere [7]. A total of 12 focus groups were conducted [6, 7]. The discussion guide for the focus groups was initially designed to understand hospital practices, policies and the organisational context influencing falls prevention interventions. The results of the original analysis have been published [7]. An emerging theme of the complexity of preventing falls for patients with dementia was identified. This paper reports on the secondary analysis conducted to examine this theme.

The research questions guiding this analysis were:

1. How do nurses describe the problem of falls for patients with dementia?
2. What strategies do nurses propose to prevent falls in patients with dementia?
3. What challenges do nurses describe in implementing strategies to prevent falls in patients with dementia?

Qualitative analysis of focus group data was continuous with deductive coding guided by the research questions being applied by two researchers who independently coded transcripts using Nvivo.

This study was approved by Monash University Human Research Ethics Committee – CF11/0229 – 2011000072 and relevant hospital ethics committees.

Results

Nurses believed that implementing falls prevention interventions for patients with dementia was challenging. Two key barriers were identified: the perception that standard falls prevention interventions were not effective and competing patient priorities in a complex care setting.

Participants in this study recognised that preventing falls amongst patients with dementia was difficult, with some believing that falls were inevitable in this patient group.

Yeah, we've got dementia patients, you can do as much as you can, and [falls are] still just going to happen, I don't think it's going to be able to be stopped. I don't think falls can be prevented ... it's just the way the patients are I think. (Nurse, Hospital 3)

Nurses felt that standard falls prevention strategies, such as patient education about the risk of falls while in hospital, diversion tactics and use of a gait aid, were not effective for patients with dementia.

People with dementia in particular, you can tell them a million, squillion times what to do but because of their physiological...mental condition, they're just not able to comprehend. (Nurse, Hospital 1)

Because they're not cognitively on the ball we can't give them a magazine to read or a newspaper. So entertaining them that way becomes a difficult issue. (Nurse, Hospital 1)

If they've got dementia then they're just as likely to fall over with the gait aid as without. (Nurse, Hospital 3)

The acute setting was recognised by nurses as a challenging and multifaceted work environment, particularly when it came to caring for patients with dementia.

We have aged care, dementia, falls risks, we have medical [issues]... We had a patient with acute chest pain, in that same room I've got a patient with dementia that's a falls risk... So what do you do? You've got to deal with the acute case first.
(Nurse, Hospital 5)

Some nurses felt that needing to address the acute care concerns first led to high risk patients being 'neglected'. Hence, nurses felt one-to-one supervision in the form of specifically trained patient sitters (also termed constant observers or patient watch) was the most effective way to care for patients with dementia.

There's a lack of manpower... Sometimes you need to have one on one [supervision]... If you got three confused high risk falls [patients], you don't know who to attend to. If you went to the other one, then the other one will fall. (Nurse, Hospital 5)

Discussion and Conclusion

Nurses in this study recognised that preventing falls in patients with dementia in the acute hospital setting is challenging. Many falls prevention interventions have been found to be ineffective for people with dementia [8, 9, 10]. Nurses perceived falls in this patient group as inevitable and believed that one-on-one observation (use of patient sitters) was an effective method to keep patients with dementia safe. However, patient sitters are a significant cost to hospitals and may not be feasible within resource and financial constraints.

Hospitals are increasingly adopting innovative ways to mitigate resource restrictions. For example, since 1995, hospitals in the United Kingdom and United States have been engaging volunteers to provide basic care and supervision to patients with dementia [11, 12]. Volunteers receive training and are able to provide emotional support, supervision and practical assistance such as fluid and nutrition assistance and supervised mobilisation to reduce the risk of adverse outcomes for these patients. Research describing volunteer programs for patients with dementia or to prevent delirium in Australian hospitals is emerging [13]. Many Australian hospitals have existing volunteer networks which could be utilised to ensure the safety of patients with dementia in the acute setting. Research is required to examine whether volunteer programs to prevent falls and provide person-centred

care for patients with dementia is effective and acceptable in the acute hospital setting in Australia.

Impact Statement

This analysis highlights that there are constraints in providing patient centred care to people with dementia in the acute hospital setting, especially in the context of falls prevention. Using volunteers to provide patient centred care may be an innovative solution to overcoming resource restrictions in hospitals.

References:

1. Australian Commission on Safety and Quality in Health Care. Evidence for the safety and quality issues associated with the care of patients with cognitive impairment in acute care settings: a rapid review. 2010. Sydney: ACSQHC.
2. Cummings E, Maher R, Showell CM et al. Hospital coding of dementia: is it accurate? *HIM J* 2011;40: 5-11. 22006431
3. Barker AL, Morello RT, Wolfe R, et al. 6-PACK programme to decrease fall injuries in acute hospitals: cluster randomised controlled trial. *BMJ*. 2016 Jan 26;352:h6781.
4. Härlein J, Halfens RJ, Dassen T, Lahmann NA. Falls in older hospital inpatients and the effect of cognitive impairment: a secondary analysis of prevalence studies. *J Clin Nurs*. 2011; Jan 1;20(1-2):175-83.
5. Australian Commission on Safety and Quality in Health Care. Delirium Critical Care Standard. 2016. Sydney: ACSQHC
6. Barker AL, Morello, RT, Ayton DR, Hill KD, Landgren FS, Brand CA. Development of an implementation plan for the 6-PACK falls prevention programme as part of a randomised controlled trial: protocol for a series of preimplementation studies. *Inj Prev*. 2015.
7. Ayton DR, Barker AL, Morello RT, Brand CA, Talevski J, Landgren FS, Melhem MM, Bian E, Brauer SG, Hill KD, Livingston PM. Barriers and enablers to the implementation of the 6-PACK falls prevention program: A pre-implementation study

in hospitals participating in a cluster randomised controlled trial. PLoS ONE. 2017 Feb 16;12(2):e0171932.

8. Shaw FE, Bond J, Richardson DA, Dawson P, Steen IN, McKeith IG, Kenny RA. Multifactorial intervention after a fall in older people with cognitive impairment and dementia presenting to the accident and emergency department: randomised controlled trial. BMJ. 2003 Jan 11;326(7380):73.
9. Haines TP, Hill AM, Hill KD, McPhail S, Oliver D, Brauer S, Hoffmann T, Beer C. Patient education to prevent falls among older hospital inpatients: a randomized controlled trial. Arch Intern Med. 2011 Mar 28;171(6):516-24.
10. Cameron ID, Murray GR, Gillespie LD, Cumming RG, Robertson MC, Hill KD, Kerse N. Interventions for preventing falls in older people in residential care facilities and hospitals. The Cochrane Library. 2005.
11. Inouye SK, Bogardus ST, Baker DI, Leo-Summers, L, Cooney LM. The Hospital Elder Life Program: A Model of Care to Prevent Cognitive and Functional Decline in Older Hospitalized Patients. J. Am Geriatr. Soc 2000; 48(12):1697-1706.
12. Dementia Action Alliance. (2016) http://www.dementiaaction.org.uk/joint_work/dementia_friendly_hospitals (Accessed Aug 2016).
13. Bateman C, Anderson K, Bird M, Hungerford C. Volunteers improving person-centred dementia and delirium care in a rural Australian hospital. Rural and Remote Health. 2016. 16(3667).

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Ayton, D;O'Brien, P;Trembl, J;Soh, S-E;Morello, R;Barker, A

Title:

Nurses' perceptions of preventing falls for patients with dementia in the acute hospital setting

Date:

2017-12-01

Citation:

Ayton, D., O'Brien, P., Trembl, J., Soh, S. -E., Morello, R. & Barker, A. (2017). Nurses' perceptions of preventing falls for patients with dementia in the acute hospital setting. AUSTRALASIAN JOURNAL ON AGEING, 36 (4), pp.E70-E72. <https://doi.org/10.1111/ajag.12474>.

Persistent Link:

<http://hdl.handle.net/11343/293914>