Apolipoprotein E gene polymorphism, trauma burden, and posttraumatic stress symptoms in U.S. military veterans: Results from the National Health and Resilience in Veterans Study

Natalie P. Mota¹, Shizhong Han², Ilan Harpaz-Rotem^{3,4}, Paul Maruff⁵, John H. Krystal^{3,4}, Steven M. Southwick^{3,4}, Joel Gelernter^{3,4}, Robert H. Pietrzak^{3,4}

¹Department of Clinical Health Psychology, Max Rady College of Medicine, University of Manitoba, Manitoba, Canada

²Department of Psychiatry, University of Iowa Carver College of Medicine, Iowa City, IA, USA

³Clinical Neurosciences Division, U.S. Department of Veterans Affairs National Center for PTSD, VA Connecticut Healthcare System, West Haven, CT, USA

⁴Department of Psychiatry, Yale University School of Medicine, New Haven, CT, USA

⁵Florey Institute of Neuroscience and Mental Health, University of Melbourne and Cogstate, Ltd., Melbourne, Australia

This work took place at the Clinical Neurosciences Division, U.S. Department of Veterans Affairs National Center for PTSD, VA Connecticut Healthcare System, West Haven, CT, USA.

Key words:

apolipoprotein, posttraumatic stress disorder, trauma, social support, veterans

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/da.22698](#).

Correspondence: Natalie Mota, Department of Clinical Health Psychology, University of Manitoba, PZ-350, 771 Bannatyne Ave., Winnipeg, MB R3E 0Y1, Canada. Email: Natalie.Mota@umanitoba.ca

ABSTRACT

Background: Previous research examining the association between apolipoprotein E (*APOE*) gene polymorphism and risk for posttraumatic stress disorder (PTSD) has been inconsistent due to the use of small and select samples. This study examined the relation between *APOE* genotype and PTSD symptoms in two nationally representative samples of U.S. military veterans. The potential effect of cumulative trauma burden and social support in moderating this association was also evaluated.

Methods: The main sample consisted of 1,386 trauma-exposed European American (EA) veterans (mean age: 62--63 years) who participated in the National Health and Resilience in Veterans Study (NHRVS) in 2011. The independent replication sample consisted of 509 trauma-exposed EA veterans from the 2013 NHRVS.

Results: *APOE* ϵ 4 allele carriers reported significantly greater severity of PTSD symptoms than noncarriers in the main, but not the replication, sample. In both samples, the interaction of *APOE* ϵ 4 carrier status and cumulative trauma burden was associated with greater severity of PTSD symptoms (F range = 2.53--8.09, all P 's < .01), particularly re-experiencing/intrusion symptoms (F range = 3.59--4.24, P 's < .001). Greater social support was associated with lower severity of PTSD symptoms among *APOE* ϵ 4 allele carriers with greater cumulative trauma burden (β range -.27 to -.60, P 's < .05).

Conclusion: U.S. military veterans who are *APOE* ϵ 4 allele carriers and exposed to a high number of traumas may be at increased risk for developing PTSD symptoms than ϵ 4

noncarriers. Greater social support may moderate this association, thereby highlighting the potential importance of social support promoting interventions in mitigating the effect of $\epsilon 4 \times$ cumulative trauma burden on PTSD risk.

1 INTRODUCTION

Posttraumatic stress disorder (PTSD) affects 6.4--7.8% of trauma-exposed adults (Kessler, 2000; Pietrzak, Goldstein, Southwick, & Grant, 2011; Wisco et al., 2014). Myriad environmental and psychosocial risk and protective factors for PTSD have been identified in trauma-exposed populations, including military veterans (Andersen, Karstoft, Bertelsen, & Madsen, 2014; Wisco et al., 2014; Xue et al., 2015). There is also increasing interest in identifying genetic risk factors for PTSD and how these markers interact with environmental factors to predict PTSD risk (e.g., Grabe et al., 2009; Kilpatrick et al., 2007). Examining the interactive effects of candidate genes and environmental factors on mental disorders, as opposed to investigating genetic or environmental influences independently, can substantially further our understanding of the etiology of these conditions (Dick et al., 2015). To date, candidate gene studies have identified possible risk alleles mapped to genes implicated in PTSD (Almli, Fani, Smith, & Ressler, 2014), and several genome-wide significant associations between genes such as *AC068718* (rs10170218), *RORA* (rs8042149), *ANKRD55* (rs159572), and *ZNF626* (rs11085374) and PTSD (Guffanti et al., 2013; Logue et al., 2013; Stein et al., 2016).

c a growing body of research has evaluated whether polymorphisms in the apolipoprotein E (*APOE*) gene may contribute to PTSD risk (e.g., Kimbrel et al., 2015; Lyons et al., 2013), most studies have been small and focused on select samples of trauma survivors (e.g., Vietnam or Iraq/Afghanistan-era veterans). *APOE* is a protein-coding gene with three

functionally different alleles ($\epsilon 2$, $\epsilon 3$, $\epsilon 4$), with $\epsilon 3$ being the most frequently occurring in the general population. The *APOE* gene encodes the APOE protein, which plays a role in neuronal repair through its involvement in cholesterol metabolism, and transportation of cholesterol and other lipids to neurons. The *APOE* $\epsilon 4$ allele, however, has been linked to hippocampal atrophy and memory impairment (Pievani et al., 2011; Small, Rosnick, Fratiglioni, & Bäckman, 2004), reduced synaptic plasticity (Chen, Durakoglugil, Xian, & Herz, 2010), and a more robust systemic and central nervous system inflammatory response (Lynch et al., 2003). It has also been associated with greater likelihood of developing several neurological and neuropsychiatric disorders (e.g., Skoog et al., 2015), especially Alzheimer's disease (Bekris, Yu, Bird, & Tsuang, 2010; Kim, Basak, & Holtzman, 2009).

Studies examining the relation between *APOE* gene and PTSD have been mixed. In a sample of 172 Vietnam veterans, Lyons et al. (2013) found that the presence of an *APOE* $\epsilon 4$ allele was associated with a greater number of PTSD symptoms, but not with PTSD diagnosis. They also observed that $\epsilon 4$ carriers with high combat exposure had greatest severity and risk for PTSD. More recently, a study of 859 non-Hispanic black veterans found that each additional *APOE* $\epsilon 4$ allele was associated with 60% greater likelihood of having a PTSD diagnosis and greater severity of PTSD symptoms, and that this finding was only observed among veterans with high levels of combat exposure; these effects were not observed in non-Hispanic white veterans (Kimbrel et al., 2015). *APOE* genotype was also unrelated to severity of postdeployment PTSD symptoms in 230 U.S. army soldiers (Dretsch et al., 2016). Other studies have found the *APOE* $\epsilon 2$ allele, but not $\epsilon 4$, to be associated with memory impairment and re-experiencing symptoms of PTSD (Freeman, Roca, Guggenheim, Kimbrell, & Griffin, 2005), PTSD diagnosis (Kim et al., 2013), and PTSD symptom severity

(Johnson et al., 2015), which is the opposite of the expectation based on the literature related to Alzheimer's disease.

Gene $\times$ environment ($G \times E$) effects of *APOE* $\epsilon 4$ and environmental exposures are also poorly understood. Aside from combat exposure severity (e.g., Kimbrel et al., 2015; Lyons et al., 2013), it remains unknown whether $\epsilon 4$ carrier status may interact with cumulative trauma burden to predict PTSD risk, as has been observed for other genetic polymorphisms (e.g., *SLC6A4*: Grabe et al., 2009). Further, no known study has evaluated potentially modifiable moderators of the *APOE* $\epsilon 4 \times$ trauma burden interaction, such as social support, a robust protective factor for PTSD (Ozer, Best, Lipsey, & Weiss, 2003). A small number of studies have found low social support to interact with high trauma exposure and genetic risk factors to increase the likelihood of PTSD (Kilpatrick et al., 2007; Lian et al., 2014), suggesting that high levels of social support may buffer against genetic risk among trauma-affected individuals.

To address these gaps, we analyzed data from two nationally representative samples of trauma-exposed European American (EA) U.S. military veterans to evaluate two aims: (1) examine the relation between *APOE* $\epsilon 4$ carrier status, alone and interactively with cumulative trauma burden, and overall severity of PTSD symptoms and symptom clusters; and (2) determine whether levels of perceived social support may moderate any effects of *APOE* $\epsilon 4$ carrier status and the interaction of $\epsilon 4 \times$ trauma burden in predicting severity of PTSD symptoms.

2 MATERIALS AND METHODS

2.1 Participants

Participants were recruited from a research panel of over 50,000 U.S. households developed and maintained by GfK Knowledge Networks, Inc. (Menlo Park, CA), and representing approximately 98% of U.S. households. Participants in the research panel who answered affirmatively to the question, “Have you ever served on active duty in the U.S. Armed Forces, Military Reserves, or National Guard?” were eligible to participate in the National Health and Resilience in Veterans Study (NHRVS), a nationally representative study of U.S. veterans. The main sample consisted of 1,386 trauma-exposed EA U.S. military veterans from the NHRVS, conducted in 2011. Participants in the replication sample were an independent sample of 509 trauma-exposed EA U.S. military veterans from a second baseline cohort survey of the NHRVS conducted in 2013. Poststratification weights were applied based on the demographic distribution of veterans (age, sex, education, race/ethnicity, metropolitan area, and Census region) in the GfK Knowledge Networks survey panel and calibrated against U.S. Census data. The NHRVS was approved by the Veterans Affairs (VA) Connecticut Healthcare System and the VA Office of Research & Development.

2.2 Assessments

2.2.1 APOE genotyping

Participants provided saliva for DNA extraction. Saliva was collected using Oragene DNA (OG-250) kits. DNA was extracted using prepIT-L2P reagent (DNA Genotek, Ontario, Canada) according to manufacturer's directions. Samples were genotyped with the PsychChip GWAS array. Genotypes were called using GenomeStudio software V2011.1 and genotyping module V1.8.4 (Illumina, San Diego, CA). Ninety samples with missing genotyping rate $\geq 5\%$ were excluded from analysis. The following criteria were used for including SNPs: minor allele frequency (MAF) $> .01\%$, missing genotyping rate per SNP $< 5\%$, and Hardy-Weinberg equilibrium (HWE) P -value $> 10^{-5}$. This resulted in 423,415 autosomal SNPs and 2,737 samples. Duplicates were detected by estimating the genome-wide identity-by-descent (IBD) sharing for all pairwise samples in PLINK (Purcell et al., 2007) using 93,814 independent SNPs with MAF $> .01$. Nine duplicate pairs and 12 additional pairs with a high level of IBD sharing ($> .1$) were detected. We randomly removed one subject of the duplicate or related pairs, retaining 2,718 independent samples (2,270 EAs). We computed principal components (PC) for the GWAS data using EIGENSOFT (Price et al., 2006) based on a common set SNPs (64,219) with Hapmap3, which were in low linkage disequilibrium (LD) with one another and have a MAF $> .01$. We detected and removed 95 outlier EA samples from the PC analysis, defined as samples whose ancestry was at least three standard deviations from the mean on one of the two largest PCs. We then imputed 1,000 genomes variants into the EA samples following the best practice guidelines of IMPUTE2 (Howie, Donnelly, & Marchini, 2009). Prephasing was first performed with SHAPEIT (Delaneau, Marchini, & Zagury, 2012) to infer haplotypes for the EA samples based on 295,837 autosomal SNPs with MAF $> .01$. Imputation was then carried out on prephased haplotypes using IMPUTE2 against reference data from the 1000 Genomes Phase III integrated variant set. After postimputation QC (SNP missing rate $< .05$, MAF $> .005$, imputation quality score (info) $> .5$, and HWE $> 10^{-6}$), 10,377,932 SNPs remained. We extracted two *APOE* SNPs—rs429358 and rs7412—from the imputed data of EA samples

for a candidate gene analysis of *APOE* ϵ 4 carrier status. These two SNPs define ApoE status and were selected in accordance with previous literature (e.g., Kim et al., 2013; Kimbrel et al., 2015). One SNP (rs7412) was directly genotyped by the PsychChip array and the other (rs429358) was imputed with high-quality score (info = .99). To determine *APOE* ϵ 4 carrier status, we made hard genotype calls of the imputed SNP by applying a posterior genotype probability threshold of .9. There was no evidence of deviation from Hardy--Weinberg expectations for the *APOE* genotype in either the main ($P = .14$) or replication ($P = .74$) sample. A dichotomous variable of 0 versus 1 or 2 *APOE* ϵ 4 alleles was created.

2.2.2 Sociodemographic and military characteristics

Age, sex, household income, education, employment status, marital status, combat veteran status, and number of years of military service were assessed.

2.2.3 Cumulative trauma burden

The *Trauma History Screen* (THS) assessed lifetime exposure to 13 potentially traumatic events (Carlson et al., 2011), including child and adult physical and sexual assault, natural disaster, and unexpected loss of a loved one. Events were summed to yield a measure of cumulative trauma burden.

2.2.4 PTSD symptoms

The PTSD Checklist (PCL) was used to assess lifetime and past-month PTSD symptoms based on respondents' worst reported traumatic event on the THS. The DSM-IV version

(PCL-Specific Stressor [PCL-S]) was used in the main sample ($\alpha = .94$) and the DSM-5 version (PCL-5) was administered in the replication sample ($\alpha = .95$). Veterans were classified as having probable PTSD if their PCL score was ≥ 50 on the DSM-IV version (Weathers, Litz, Herman, Huska, & Keane, 1993) and ≥ 38 on the DSM-5 version (Hoge, Riviere, Wilk, Herrell, & Weathers, 2014). Comparability of the symptom clusters on the two PCL versions was achieved by using the four-factor DSM-IV model of re-experiencing, avoidance, emotional numbing, and hyperarousal symptoms (King, Leskin, King, & Weathers, 1998), and the DSM-5 model of intrusions, avoidance, negative cognitions and mood, and alterations in arousal and reactivity. Responses on items comprising each symptom cluster were summed to yield severity measures.

2.2.5 Social support

Social support was assessed using a 5-item version of the Medical Outcomes Study Social Support Scale (Amstadter et al., 2010); items assessed emotional and instrumental support ($\alpha = .90$ in main sample; $\alpha = .87$ in replication sample). Veterans were asked how often each kind of support was available when needed, and items included, “Someone to get together with for relaxation” and “Someone to love and make you feel wanted.”

2.2.6 Other psychiatric disorders

The Mini International Neuropsychiatric Interview adapted for self-report was used to assess lifetime DSM-IV diagnoses of major depressive, alcohol, and drug use disorders (Lecrubier et al., 1997).

2.3 Data analysis

Descriptive statistics were used to summarize sociodemographic, trauma, and clinical variables of trauma-exposed veterans. A series of univariate analyses of covariance (ANCOVAs) were conducted to evaluate the relation between *APOE* $\epsilon 4$ allele carrier status, trauma burden, and their interaction, and severity of lifetime and past-month PTSD symptoms in the main and replication samples. Covariates included age, sex, top 10 PCs from population stratification analysis, combat exposure (i.e., combat veteran vs. noncombat veteran), and nature of “worst” traumatic event (i.e., assaultive vs. nonassaultive). To evaluate the role of social support in moderating the interaction between *APOE* $\epsilon 4$ allele carrier status $\times$ lifetime trauma burden, we incorporated an $\epsilon 4 \times$ trauma burden $\times$ social support interaction term into the ANCOVAs. To evaluate the relationship between *APOE* $\epsilon 4$ allele carrier status, cumulative trauma burden, and their interaction, and lifetime PTSD symptom clusters, we conducted a parallel series of univariate ANCOVAs with scores on each of the PTSD symptom clusters entered as dependent variables in separate analyses and other PTSD symptom clusters entered as additional covariates. A statistical significance threshold of $\alpha = .01$ was employed in analyses to reduce the likelihood of Type I error. These post-hoc analyses were limited to lifetime PTSD symptom clusters due to the low prevalence and variance of past-month PTSD symptom clusters. Reported raw frequencies are unweighted; means, percentages, and inferential statistics are poststratification weighted to reflect the general population of U.S. veterans. Analyses were conducted using SPSS version 22.

3 RESULTS

Table 1 shows sociodemographic, military, and clinical characteristics of trauma-exposed veterans in the main and replication samples. On average, participants in both samples were 62--63 years of age, predominantly male, some college or higher educated, married/cohabitating, retired, had household incomes <\$60,000, were noncombat veterans, and spent an average of 7 years in the military. On average, veterans in the samples reported experiencing 3.7--3.8 traumatic life events, with 7.1--10.0% screening positive for lifetime PTSD and 3.5--3.8% for past-month PTSD.

Table 2 shows results of analyses evaluating the relation between *APOE* ϵ 4 carrier status, trauma burden, and their interaction, and PTSD symptoms.

In the main sample, *APOE* ϵ 4 carrier status, number of traumas, and their interaction were associated with greater severity of lifetime and past-month PTSD symptoms. Figure 1 illustrates the interaction of *APOE* ϵ 4 and trauma burden on severity of lifetime PTSD symptoms in the main sample. Among ϵ 4 carriers, those with a greater number of traumas had greater PTSD symptoms. Incorporation of an *APOE* ϵ 4 $\times$ lifetime traumas $\times$ social support interaction term into this ANCOVA revealed a significant association with lifetime (three-way interaction, $F = 3.01$, $P < .001$) and past-month (three-way interaction, $F = 4.35$, $P < .001$) PTSD symptoms. Adjusted post-hoc analyses among ϵ 4 carriers with greater cumulative trauma burden revealed that greater levels of social support were associated with significantly reduced likelihood of lifetime ($\beta = -.48$, $t = 6.21$, $P \leq .001$) and past-month ($\beta = -.60$, $t = 7.89$, $P < .001$) PTSD symptoms. Among ϵ 4 carriers with greater cumulative trauma burden (i.e., greater than the median number of three events, $n = 602$), those with

higher levels of perceived social support were less likely to screen positive for current PTSD (2.4% in highest tertile vs. 4.8% in middle tertile vs. 20.5% in lowest tertile; see Fig. 2).

In the replication sample, the interaction of *APOE* $\epsilon 4$ carrier status $\times$ cumulative trauma burden was significantly associated with lifetime and past-month PTSD symptoms; *APOE* $\epsilon 4$ carrier status was unrelated to these outcomes. Incorporation of an *APOE* $\epsilon 4 \times$ lifetime traumas $\times$ social support interaction term revealed a significant association with lifetime ($F = 3.86, P < .001$) and past-month ($F = 4.55, P < .001$) PTSD symptoms. Adjusted post-hoc multivariable linear regression analyses in $\epsilon 4$ carriers with greater cumulative trauma burden revealed that greater levels of social support were associated with significantly reduced likelihood of lifetime ($\beta = -.27, t = 2.32, P \leq .024$) and past-month ($\beta = -.37, t = 3.07, P = .003$) PTSD symptoms. Among $\epsilon 4$ allele carriers with greater cumulative trauma burden, those with higher levels of social support were less likely to screen positive for current PTSD (0% in highest tertile vs. 8.3% in middle tertile vs. 11.8% in lowest tertile).

Adjustment for MDD and SUDs did not substantively change the results. In the main sample, the main and interactive effects of $\epsilon 4$ allele and $\epsilon 4$ allele $\times$ cumulative trauma burden remained significant for both lifetime PTSD symptoms: $F = 20.77, P < .001$ and $F = 36.13, P < .001$, respectively; and past-month PTSD symptoms: $F = 22.76, P < .001$; $F = 30.10, P < .001$, respectively. In the replication sample, the interaction of $\epsilon 4$ allele $\times$ trauma burden also remained significant for past-month ($F = 2.78, P = .003$) but not for lifetime ($F = 1.30, P = .23$) PTSD symptoms.

Table 3 shows results of analyses examining the relationship between *APOE* ϵ 4 carrier status, cumulative trauma burden, and the interaction of these variables, and lifetime PTSD symptom clusters. In the main sample, the interaction of *APOE* ϵ 4 carrier status $\times$ lifetime traumas was significant for re-experiencing and avoidance symptoms. In the replication sample, ϵ 4 carrier status was associated with greater severity of intrusions; and the interaction of *APOE* ϵ 4 $\times$ cumulative trauma burden was significant for intrusion symptoms and alterations in arousal and reactivity.

4 DISCUSSION

This study examined the association between *APOE* genotype, trauma exposure, and PTSD symptoms in two contemporary, nationally representative cohorts of EA military veterans. *APOE* ϵ 4 allele carriers reported greater lifetime and past-month PTSD symptoms than noncarriers in the main sample, however, this association was not significant in our replication sample. This finding aligns with a recent meta-analysis supporting an association between ϵ 4 carrier status and higher risk for combat-related PTSD (Roby, 2017). It is possible that the lack of a main effect in the replication sample was due to small sample size, and differences in sociodemographic factors and environmental exposures across samples may also partly account for the inconsistent findings. It is also possible that the discrepancy in findings between the main and replication samples may be attributable in part to the different PCL versions (DSM-IV vs. DSM-5) used; however, after recomputing a PCL-5 summary score by removing the new DSM-5 PTSD symptoms from the PCL-5 such that it more closely aligned with the PCL-S (DSM-IV), the main effect of ϵ 4 allele carrier status remained nonsignificant for both lifetime ($F = .44$, $P = .51$) and past-month ($F = .01$, $P = .92$)

symptoms. Further evaluation of a possible association between *APOE* ϵ 4 status and PTSD symptoms is needed in different trauma-exposed samples.

In both samples, *APOE* ϵ 4 carriers with greater trauma burden reported greater lifetime and past-month PTSD symptoms. This finding is consistent with previous work linking ϵ 4 allele carrier status to PTSD among veterans with high combat exposure (Lyons et al., 2013; Kimbrel et al., 2015), and extends it to demonstrate that cumulative trauma burden may also moderate the effect of ϵ 4 allele carriage on PTSD symptoms. The *APOE* ϵ 4 allele may hinder the neuronal repair and recovery that is necessary after injury related to extensive stress exposure (Lyons et al., 2013; Kimbrel et al., 2015). Trauma exposure across the lifespan can deleteriously affect brain structure and function, including reductions in hippocampal volume (Bremner, 2006; Woon, Sood, & Hedges, 2010) and dysregulation of the hypothalamic--pituitary--adrenal (HPA) axis (Bremner, 2006; Carpenter et al., 2007; Elzinga et al., 2008), which can increase risk for developing PTSD symptoms (Gilbertson et al., 2002; Pitman et al., 2012). *APOE* ϵ 4 allele carriers are similarly at increased risk of several deficits relevant to PTSD, including hippocampal volume loss and amygdala atrophy (den Heijer et al., 2002; Goni et al., 2012) and greater cortisol levels (Gill-Bea et al., 2010), possibly further exacerbating PTSD vulnerability. Given that individuals with PTSD are at higher risk of dementia (Yaffe et al., 2010), it is also possible that the ϵ 4 allele contributes to a shared mechanistic pathway for the development of both conditions, or that the development of dementia symptoms plays a role in the ϵ 4-PTSD association. For example, a recent study found that Alzheimer's model mice displayed more exaggerated and frequent responses to a PTSD-like induction than control mice (Justice et al., 2015). Additional research utilizing prospective designs is needed to elucidate possible mechanisms linking ϵ 4, PTSD, and cognitive decline and dementia in trauma-affected individuals.

Analyses of PTSD symptom clusters revealed that the *APOE* $\epsilon 4$ $\times$ trauma burden association was linked to greater re-experiencing and avoidance symptoms in the main sample, and greater intrusion symptoms and alterations in arousal and reactivity in the replication sample. In light of prior studies showing higher levels of cerebrospinal fluid cortisol among *APOE* $\epsilon 4$ carriers (Gil-Bea et al., 2010; Peskind, Wilkinson, Petrie, Schellenberg, & Raskind, 2001), it is possible that greater HPA axis reactivity among $\epsilon 4$ carriers may exacerbate hyperarousal symptoms among highly trauma-exposed individuals, which may in turn trigger re-experiencing/intrusion symptoms. Greater hippocampal atrophy and volume loss in $\epsilon 4$ carriers may also contribute to deficits in memory for fear extinction (den Heijer et al., 2002; Fanselow, 2000; Pievani et al., 2011), which can maintain re-experiencing/intrusion symptoms. Villasana, Weber, Akinyeke, and Raber (2016) found that mice carrying the $\epsilon 4$ allele had higher levels of heme oxygenase-1, a marker of stress, in the hippocampus, and also exhibited greater anxiety-related behaviors and conditioned fear responding. These findings suggest a possible mechanism linking $\epsilon 4$ carriage and intrusive symptoms among highly trauma-exposed veterans. Clearly, more research is needed to elucidate mechanisms underlying this association.

Of note, *APOE* $\epsilon 2$ carriage has been found to be associated with memory impairment, more re-experiencing symptoms, and greater overall PTSD symptoms in small samples of humans (Freeman et al., 2005; Johnson et al., 2015), while stress-exposed mice carrying the $\epsilon 2$ allele have been found to have impairments in fear memory extinction (Johnson et al., 2015; Olsen, Agam, Davis, & Raber, 2012). However, a meta-analysis did not find support for an effect of the $\epsilon 2$ variant on risk for combat-related PTSD (Roby, 2017). Further research in

larger samples is needed to help resolve these inconsistent findings regarding *APOE* genotype and PTSD (Johnson et al., 2015; Rogers & Weeber, 2008).

Higher levels of social support were found to be associated with lower PTSD symptoms among *APOE* $\epsilon 4$ allele carriers with greater cumulative trauma burden. This finding adds to a small literature demonstrating an effect of social support in moderating the association between other genetic polymorphisms (Kilpatrick et al., 2007; Lian et al., 2014) and PTSD, depressive symptoms, and suicidal ideation (Chen et al., 2011; Kilpatrick et al., 2007; Kim et al., 2014). Social support has been linked with lower cortisol levels (McQuaid et al., 2016; Rosal, King, Ma, & Reed, 2004), improved cardiovascular and immune functioning (Uchino, 2006), and less-threatening appraisal of stressful events (Sippel, Pietrzak, Charney, Mayes, & Southwick, 2015), suggesting a potential role in buffering the impact of trauma-related stress responses. It is also possible, however, that veterans with more PTSD symptoms engage in behaviors that lead to less social support (e.g., avoidance, social isolation) or may inaccurately perceive their social support (e.g., Platt, Lowe, Galea, Norris, & Koenen, 2016). Nonetheless, assuming that social support may help buffer against the development of PTSD symptoms, facilitating the enhancement of social support networks (e.g., one-to-one mentorship programs, peer support groups, social/relationship skills interventions, Vet-to-Vet programs) among highly trauma-exposed veterans at elevated genetic risk for PTSD may be an important aspect of prevention initiatives (e.g., Hogan, Linden, & Najarian, 2002; Pietrzak et al., 2009; Resnick & Rosenheck, 2008; Williams, Bambara, & Turner, 2012).

The current findings should be considered in light of several limitations. First, PTSD symptoms were assessed via self-report, and it is unclear whether results would be

comparable if PTSD symptoms were assessed using a structured clinical interview such as the Clinician Administered PTSD Scale (CAPS). However, statistically significant and large magnitude associations have been observed between total scores on the PCL and on the CAPS (Macdonald, Greene, Torres, Frueh, & Morland, 2013; Monson et al., 2008). Second, the main sample was larger than the replication sample and different versions of the PCL (i.e., DSM-IV vs. DSM-5) were used in these samples; thus, it is unclear whether differences in patterns of associations between the samples may be related to reduced statistical power or versions of the PCL. Third, the sample was comprised predominantly of male EA veterans. Further research is needed to examine the role of *APOE* ϵ 4 carrier status on risk for PTSD among more diverse samples of veterans and other trauma-affected populations. Fourth, given the strong association between *APOE* ϵ 4 carrier status and poorer cognitive functioning and dementia (e.g., Engelborghs et al., 2003; Wisdom et al., 2011), it remains to be determined whether cognitive impairment might influence the association between ϵ 4 and PTSD, as formal neuropsychological testing was not conducted in this cohort of veterans. Fifth, there have been a number of limitations raised with respect to $G \times E$ research with candidate genes, including the lack of replication of some findings, publication bias, and the frequent use of underpowered samples (e.g., Dick et al., 2015). However, the current study incorporated a number of recommendations made for improving the rigor of $G \times E$ studies, including having an adequately powered main sample, an independent replication sample, and a priori selection of candidate genes (i.e., *APOE*) and environmental factors (i.e., trauma exposure) based on theoretical rationale and existing literature (Dick et al., 2015). A recent review also recommended additional research examining the effect of positive exposures (i.e., social support) on psychopathology (Leighton, Botto, Silva, Jimenez, & Luyten, 2017).

5 CONCLUSION

Results of the present study suggest that *APOE* ϵ 4 carrier status interacts with cumulative trauma burden to predict severity of PTSD symptoms, particularly re-experiencing/intrusive symptoms, in U.S. military veterans. Further, greater social support is associated with lower severity of PTSD symptoms in ϵ 4 allele carriers with greater cumulative trauma burden. Further research is needed to examine the complex relationship and mechanistic pathways between *APOE* genotype, PTSD, and cognitive decline and dementia (Lee et al., 2008; Meziab et al., 2014; Peavy et al., 2007; Yaffe et al., 2010); evaluate how *APOE* and other gene polymorphisms are linked to transdiagnostic aspects of trauma-related psychopathology (e.g., Mota et al., 2015); and examine the efficacy of interventions to enhance social support in mitigating PTSD symptoms among at-risk trauma survivors.

FUNDING AND DISCLOSURE

The U.S. Department of VA supported this work through its support of the VA National Center for PTSD and the Consortium to Alleviate PTSD. Dr. Krystal received additional support from a National Institute on Alcohol Abuse and Alcoholism grant (2P50AA012870-14) and National Center for Advancing Translational Science grants (UL1RR024139, principal investigator: R. Sherwin; UH2TR000960, principal investigator: Dr. Krystal). No author has any relevant conflict of interest to disclose.

REFERENCES

- Almli, L. M., Fani, N., Smith, A. K., & Ressler, K. J. (2014). Genetic approaches to understanding post-traumatic stress disorder. *International Journal of Neuropsychopharmacology*, 17(2), 355--370.
- Amstadter, A. B., Begle, A. M., Cisler, J. M., Hernandez, M. A., Muzzy, W., & Acierno, R. (2010). Prevalence and correlates of poor self-rated health in the United States: The national elder mistreatment study. *American Journal of Geriatric Psychiatry*, 18(7), 615--623.
- Andersen, S. B., Karstoft, K. I., Bertelsen, M., & Madsen, T. (2014). Latent trajectories of trauma symptoms and resilience: The 3-year longitudinal prospective USPER study of Danish veterans deployed in Afghanistan. *Journal of Clinical Psychiatry*, 75(9), 1001--1008.
- Bekris, L. M., Yu, C. E., Bird, T. D., & Tsuang, D. W. (2010). Genetics of Alzheimer disease. *Journal of Geriatric Psychiatry and Neurology*, 23(4), 213--227.
- Bremner, J. D. (2006). Traumatic stress: Effects on the brain. *Dialogues in Clinical Neuroscience*, 8(4), 445.
- Carlson, E. B., Smith, S. R., Palmieri, P. A. et al. (2011). Development and validation of a brief self-report measure of trauma exposure: The Trauma History Screen. *Psychological Assessment*, 23(2), 463--477.

Carpenter, L. L., Carvalho, J. P., Tyrka, A. R., Wier, L. M., Mello, A. F., Mello, M. F., et al. (2007). Decreased adrenocorticotrophic hormone and cortisol responses to stress in healthy adults reporting significant childhood maltreatment. *Biological Psychiatry*, 62(10), 1080--1087.

Chen, F. S., Kumsta, R., von Dawans, B., Monakhov, M., Ebstein, R. P., & Heinrichs, M. (2011). Common oxytocin receptor gene (OXTR) polymorphism and social support interact to reduce stress in humans. *Proceedings of the National Academy of Sciences of the United States of America*, 108(50), 19937--19942.

Chen, Y., Durakoglugil, M. S., Xian, X., & Herz, J. (2010). ApoE4 reduces glutamate receptor function and synaptic plasticity by selectively impairing ApoE receptor recycling. *Proceedings of the National Academy of Sciences of the United States of America*, 107(26), 12011--12016.

Delaneau, O., Marchini, J., & Zagury, J. F. (2012). A linear complexity phasing method for thousands of genomes. *Nature Methods*, 9(2), 179--181.

den Heijer, T., Oudkerk, M., Launer, L. J., Van Duijn, C. M., Hofman, A., & Breteler, M. M. B. (2002). Hippocampal, amygdalar, and global brain atrophy in different apolipoprotein E genotypes. *Neurology*, 59(5), 746--748.

Dick, D. M., Agrawal, A., Keller, M. C., Adkins, A., Aliev, F., Monroe, S., et al. (2015). Candidate gene-environment interaction research: Reflections and recommendations. *Perspectives on Psychological Science*, 10(1), 37--59.

Dretsch, M. N., Williams, K., Emmerich, T., Crynen, G., Ait-Ghezala, G., Chaytow, H., et al. (2016). Brain-derived neurotrophic factor polymorphisms, traumatic stress, mild traumatic brain injury, and combat exposure contribute to postdeployment traumatic stress. *Brain and Behavior*, 6(1).

Elzinga, B.M., Roelofs, K., Tollenaar, M. S., Bakvis, P., van Pelt, J., & Spinhoven, P. (2008). Diminished cortisol responses to psychosocial stress associated with lifetime adverse events: A study among healthy young subjects. *Psychoneuroendocrinology*, 33(2), 227--237.

Engelborghs, S., Dermaut, B., Goeman, J., Saerens, J., Marien, P., Pickut, B. A., et al. (2003). Prospective Belgian study of neurodegenerative and vascular dementia: APOE genotype effects. *Journal of Neurology, Neurosurgery & Psychiatry*, 74, 1148--1151.

Fanselow, M. S. (2000). Contextual fear, gestalt memories, and the hippocampus. *Behavioural Brain Research*, 110, 73--81.

Freeman, T., Roca, V., Guggenheim, F., Kimbrell, T., & Griffin, W. S. T. (2005). Neuropsychiatric associations of apolipoprotein E alleles in subjects with combat-related

posttraumatic stress disorder. *Journal of Neuropsychiatry and Clinical Neurosciences*, 17(4), 541--543.

Gil-Bea, F. J., Aisa, B., Solomon, A., Solas, M., del Carmen Mugueta, M., Winblad, B., et al. (2010). HPA axis dysregulation associated to apolipoprotein E4 genotype in Alzheimer's disease. *Journal of Alzheimer's Disease*, 22(3), 829--838.

Gilbertson, M. W., Shenton, M. E., Ciszewski, A., Kasai, K., Lasko, N. B., Orr, S. P., et al. (2002). Smaller hippocampal volume predicts pathologic vulnerability to psychological trauma. *Nature Neuroscience*, 5(11), 1242--1247.

Goni, J., Cervantes, S., Arrondo, G., Lamet, I., Pastor, P., & Pastor, M. A. (2013). Selective brain gray matter atrophy associated with APOE ϵ 4 and MAPT H1 in subjects with mild cognitive impairment. *Journal of Alzheimer's Disease*, 33(4), 1009--1019.

Grabe, H. J., Spitzer, C., Schwahn, C., Marcinek, A., Frahnw, A., & Barnow, S. (2009). Serotonin transporter gene (SLC6A4) promoter polymorphisms and the susceptibility to posttraumatic stress disorder in the general population. *American Journal of Psychiatry*, 166(8), 926--933.

Guffanti, G., Galea, S., Yan, L., Roberts, A. L., Solovieff, N., Aiello, A. E., et al. (2013). Genome-wide association study implicates a novel RNA gene, the lincRNA AC068718. 1, as a risk factor for post-traumatic stress disorder in women. *Psychoneuroendocrinology*, 38(12), 3029--3038.

Hogan, B. E., Linden, W., & Najarian, B. (2002). Social support interventions: Do they work? *Clinical Psychology Review, 22*(3), 381--440.

Hoge, C. W., Riviere, L. A., Wilk, J. E., Herrell, R. K., & Weathers, F. W. (2014). The prevalence of post-traumatic stress disorder (PTSD) in US combat soldiers: A head-to-head comparison of DSM-5 versus DSM-IV-TR symptom criteria with the PTSD checklist. *Lancet Psychiatry, 1*(4), 269--77.

Howie, B. N., Donnelly, P., & Marchini, J. (2009). A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS Genetics, 5*(6), e1000529.

Johnson, L. A., Zuloaga, D. G., Bidiman, E., Marzulla, T., Weber, S., Wahbeh, H., et al. (2015). ApoE2 exaggerates PTSD-related behavioral, cognitive, and neuroendocrine alterations. *Neuropsychopharmacology, 40*(10), 2443--2453.

Kessler, R. C. (2000). Posttraumatic stress disorder: The burden to the individual and to society. *Journal of Clinical Psychiatry, 61*, 4--14.

Kilpatrick, D. G., Koenen, K. C., Ruggiero, K. J., Acierno, R., Galea, S., Resnick, H. S., et al. (2007). The serotonin transporter genotype and social support and moderation of

posttraumatic stress disorder and depression in hurricane-exposed adults. *American Journal of Psychiatry*, 164(11), 1693--1699.

Kim, J., Basak, J. M., & Holtzman, D. M. (2009). The role of apolipoprotein E in Alzheimer's disease. *Neuron*, 63(3), 287--303.

Kim, J. M., Stewart, R., Kim, S. W., Kang, H. J., Kim, S. Y., Lee, J. Y., et al. (2014). Interactions between a serotonin transporter gene, life events and social support on suicidal ideation in Korean elders. *Journal of Affective Disorders*, 160, 14--20.

Kim, T. Y., Chung, H. G., Shin, H. S., Kim, S. J., Choi, J. H., Chung, M. Y., et al. (2013). Apolipoprotein E gene polymorphism, alcohol use, and their interactions in combat-related posttraumatic stress disorder. *Depression & Anxiety*, 30(12), 1194--1201.

Kimbrel, N. A., Hauser, M. A., Garrett, M., Ashley-Koch, A., Liu, Y., Dennis, M. F., et al. (2015). Effect of the APOE ϵ 4 allele and combat exposure on PTSD among Iraq/Afghanistan-era veterans. *Depression & Anxiety*, 32(5), 307--315.

King, D. W., Leskin, G. A., King, L. A., & Weathers, F. W. (1998). Confirmatory factor analysis of the clinician-administered PTSD Scale: Evidence for the dimensionality of posttraumatic stress disorder. *Psychological Assessment*, 10(2), 90--96.

Lecrubier, Y., Sheehan, D. V., Weiller, E., Amorim, P., Bonora, I., Sheehan, K. H., et al. (1997). The Mini International Neuropsychiatric Interview (MINI). A short diagnostic

structured interview: Reliability and validity according to the CIDI. *European Psychiatry*, 12(5), 224--231.

Lee, B. K., Glass, T. A., Wand, G. S., McAtee, M. J., Bandeen-Roche, K., Bolla, K. I., & Schwartz, B. S. (2008). Apolipoprotein e genotype, cortisol, and cognitive function in community-dwelling older adults. *American Journal of Psychiatry*, 165(11), 1456--1464.

Leighton, C., Botto, A., Silva, J. R., Jimenez, J. P., & Luyten, P. (2017). Vulnerability or sensitivity to the environment? Methodological issues, trends, and recommendations in gene-environment interactions research in human behavior. *Frontiers in Psychiatry*, 8(106).

Lian, Y., Xiao, J., Wang, Q., Ning, L., Guan, S., Ge, H., et al. (2014). The relationship between glucocorticoid receptor polymorphisms, stressful life events, social support, and post-traumatic stress disorder. *BMC Psychiatry*, 14(1), 232.

Logue, M. W., Baldwin, C., Guffanti, G., Melista, E., Wolf, E. J., Reardon, A. F., et al. (2013). A genome-wide association study of post-traumatic stress disorder identifies the retinoid-related orphan receptor alpha (RORA) gene as a significant risk locus. *Molecular Psychiatry*, 18(8), 937--942.

Lynch, J. R., Tang, W., Wang, H., Vitek, M. P., Bennett, E. R., Sullivan, P. M., et al. (2003). APOE genotype and an ApoE-mimetic peptide modify the systemic and central nervous system inflammatory response. *Journal of Biological Chemistry*, 278(49), 48529--48533.

Lyons, M. J., Genderson, M., Grant, M. D., Logue, M., Zink, T., McKenzie, R., et al. (2013). Gene-environment interaction of ApoE genotype and combat exposure on PTSD. *American Journal of Medical Genetics*, 162(7), 762--769.

Macdonald, A., Greene, C. J., Torres, J. G., Frueh, B. C., & Morland, L. A. (2013). Concordance between clinician-assessed and self-reported symptoms of posttraumatic stress disorder across three ethnoracial groups. *Psychological Trauma: Theory, Research, Practice, and Policy*, 5(3), 201--208.

McQuaid, R. J., McInnis, O. A., Paric, A., Al-Yawer, F., Matheson, K., & Anisman, H. (2016). Relations between plasma oxytocin and cortisol: The stress buffering role of social support. *Neurobiology of Stress*, 3, 52--60.

Meziab, O., Kirby, K. A., Williams, B., Yaffe, K., Byers, A. L., & Barnes, D. E. (2014). Prisoner of war status, posttraumatic stress disorder, and dementia in older veterans. *Alzheimer's & Dementia*, 10(3), S236--S241.

- Monson, C. M., Gradus, J. L., Young-Xu, Y., Schnurr, P. P., Price, J. L., & Schumm, J. A. (2008). Change in posttraumatic stress disorder symptoms: Do clinicians and patients agree? *Psychological Assessment, 20*(2), 131--138.
- Mota, N., Sumner, J. A., Lowe, S. R., Neumeister, A., Uddin, M., Aiello, A. E., et al. (2015). The rs1049353 polymorphism in the CNR1 gene interacts with childhood abuse to predict posttraumatic threat symptoms. *Journal of Clinical Psychiatry, 76*(12), e1622--e1623.
- Olsen, R. H., Agam, M., Davis, M. J., & Raber, J. (2012). ApoE isoform-dependent deficits in extinction of contextual fear conditioning. *Genes, Brain and Behavior, 11*(7), 806--812.
- Ozer, E. J., Best, S. R., Lipsey, T. L., & Weiss, D. S. (2003). Predictors of posttraumatic stress disorder and symptoms in adults: A meta-analysis. *Psychological Bulletin, 129*(1), 52--73.
- Peavy, G. M., Lange, K. L., Salmon, D. P., Patterson, T. L., Goldman, S., Gamst, A. C., et al. (2007). The effects of prolonged stress and APOE genotype on memory and cortisol in older adults. *Biological Psychiatry, 62*(5), 472--478.
- Peskind, E. R., Wilkinson, C. W., Petrie, E. C., Schellenberg, G. D., & Raskind, M. A. (2001). Increased CSF cortisol in AD is a function of APOE genotype. *Neurology, 56*(8), 1094--1098.

Pietrzak, R. H., Goldstein, R. B., Southwick, S. M., & Grant, B. F. (2011). Medical comorbidity of full and partial posttraumatic stress disorder in United States adults: Results from wave 2 of the National Epidemiologic Survey on Alcohol and Related Conditions. *Psychosomatic Medicine*, 73(8), 697--707.

Pietrzak, R. H., Johnson, D. C., Goldstein, M. B., Malley, J. C., & Southwick, S. M. (2009). Psychological resilience and postdeployment social support protect against traumatic stress and depressive symptoms in soldiers returning from Operations Enduring Freedom and Iraqi Freedom. *Depression & Anxiety*, 26(8), 745--751.

Pievani, M., Galluzzi, S., Thompson, P. M., Rasser, P. E., Bonetti, M., & Frisoni, G. B. (2011). APOE4 is associated with greater atrophy of the hippocampal formation in Alzheimer's disease. *Neuroimage*, 55(3), 909--919.

Pitman, R. K., Rasmusson, A. M., Koenen, K. C., Shin, L. M., Orr, S. P., Gilbertson, M. W., et al. (2012). Biological studies of post-traumatic stress disorder. *Nature Reviews Neuroscience*, 13(11), 769--787.

Platt, J. M., Lowe, S. R., Galea, S., Norris, F. H., & Koenen, K. C. (2016). A longitudinal study of the bidirectional relationship between social support and posttraumatic stress following a natural disaster. *Journal of Traumatic Stress*, 29(3), 205--213.

Price, A. L., Patterson, N. J., Plenge, R. M., et al. (2006). Principal components analysis corrects for stratification in genome-wide association studies. *Nature Genetics*, 38(8), 904--909.

Purcell, S., Neale, B., Todd-Brown, K., et al. (2007). PLINK: A tool set for whole-genome association and population-based linkage analyses. *American Journal of Human Genetics*, 81(3), 559--575.

Resnick, S. G., & Rosenheck, R. A. (2008). Integrating peer-provided services: A quasi-experimental study of recovery orientation, confidence, and empowerment. *Psychiatric Services*, 59(11), 1307--1314.

Roby, Y. (2017). Apolipoprotein E variants and genetic susceptibility to combat-related post-traumatic stress disorder: A meta-analysis. *Psychiatric Genetics*. Epub ahead of print. doi: 10.1097/YPG.0000000000000174

Rogers, J. T., & Weeber, E. J. (2008). Reelin and apoE actions on signal transduction, synaptic function and memory formation. *Neuron Glia Biology*, 4(03), 259--270.

Rosal, M. C., King, J., Ma, Y., & Reed, G. W. (2004). Stress, social support, and cortisol: Inverse associations? *Behavioral Medicine* 30(1), 11--22.

Sippel, L. M., Pietrzak, R. H., Charney, D. S., Mayes, L. C., & Southwick, S. M. (2015). How does social support enhance resilience in the trauma-exposed individual? *Ecology & Society*, 20(4), 136--145.

Skoog, I., Waern, M., Duberstein, P., Blennow, K., Zetterberg, H., Börjesson-Hanson, A., et al. (2015). A 9-year prospective population-based study on the association between the APOE* E4 allele and late-life depression in Sweden. *Biological Psychiatry*, 78(10), 730--736.

Small, B. J., Rosnick, C. B., Fratiglioni, L., & Bäckman, L. (2004). Apolipoprotein E and cognitive performance: A meta-analysis. *Psychology & Aging*, 19(4), 592--600.

Stein, M. B., Chen, C. Y., Ursano, R. J., Cai, T., Gelernter, J., Heeringa, S. G., et al. (2016). Genome-wide association studies of posttraumatic stress disorder in 2 cohorts of US Army soldiers. *JAMA Psychiatry*, 73(7), 695--704.

Sureshkumar, R., Bharath, S., Jain, S., Prakash, O., Purushottam, M., Thennarasu, K., et al. (2012). ApoE4 and late onset depression in Indian population. *Journal of Affective Disorders*, 136(3), 244--248.

Uchino, B. N. (2006). Social support and health: A review of physiological processes potentially underlying links to disease outcomes. *Journal of Behavioral Medicine*, 29(4), 377-387.

Villasana, L. E., Weber, S., Akinyeke, T., & Raber, J. (2016). Genotype differences in anxiety and fear learning and memory of WT and ApoE4 mice associated with enhanced generation of hippocampal reactive oxygen species. *Journal of Neurochemistry*, 138(6), 896--908.

Weathers, F. W., Litz, B. T., Herman, D. S., Huska, J. A., & Keane, T. M. (1993). The PTSD Checklist (PCL): Reliability, validity, and diagnostic utility. Annual convention of the international society for traumatic stress studies, San Antonio, TX (Vol. 462).

Williams, R. M., Bambara, J., & Turner, A. P. (2012). A scoping study of one-to-one peer mentorship interventions and recommendations for application with veterans with postdeployment syndrome. *Journal of Head Trauma Rehabilitation*, 27(4), 261--273.

Wisco, B. E., Marx, B. P., Wolf, E. J., Miller, M. W., Southwick, S. M., Pietrzak, & R. H. (2014). Posttraumatic stress disorder in the US veteran population: Results from the National Health and Resilience in Veterans Study. *Journal of Clinical Psychiatry*, 75(12), 1338--1346.

Woon, F. L., Sood, S., & Hedges, D. W. (2010). Hippocampal volume deficits associated with exposure to psychological trauma and posttraumatic stress disorder in adults: A meta-analysis. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, 34(7), 1181--1188.

Xue, C., Ge, Y., Tang, B., Liu, Y., Kang, P., Wang, M., & Zhang, L. (2015). A meta-analysis of risk factors for combat-related PTSD among military personnel and veterans. *PLoS One*, 10(3), e0120270.

Yaffe, K., Vittinghoff, E., Lindquist, K., Barnes, D., Covinsky, K. E., Neylan, T., et al. (2010). Posttraumatic stress disorder and risk of dementia among US veterans. *Archives of General Psychiatry*, 67(6), 608--613.

Table 1. Sociodemographic, Military, and Trauma and Clinical Characteristics of the Main and Replication Samples of Trauma-Exposed U.S. European-Ancestry Military Veterans

	Main Sample (<i>n</i> = 1,386)	Replication Sample (<i>n</i> = 509)
	Weighted Mean (<i>SD</i>) or <i>n</i> (Weighted %)	Weighted Mean (<i>SD</i>) or <i>n</i> (Weighted %)
Sociodemographic characteristics		
Age	62.6 (14.3)	62.4 (15.6)
Male sex	1,260 (92.8%)	457 (90.7%)
Some college or higher education	1,183 (66.0%)	428 (65.4%)
Married/living with partner	1,084 (74.9%)	379 (72.3%)
Currently employed	536 (36.1%)	152 (30.3%)
Household income ≥\$60,000/year	728 (42.4%)	255 (44.0%)
Military characteristics		
Combat veteran	479 (32.7%)	213 (42.0%)
Number of years in military	6.8 (7.5)	7.0 (7.2)

Trauma and clinical characteristics

Number of lifetime traumatic events	3.8 (2.5)	3.7 (2.5)
Index traumatic event		
Sudden death of close family member or friend	451 (34.4%)	151 (32.7%)
Life-threatening illness or injury	226 (16.6%)	84 (14.2%)
Military-related trauma	113 (7.9%)	42 (9.5%)
Child physical or sexual abuse	59 (3.2%)	23 (5.4%)
Other traumatic event	537 (37.9%)	209 (38.2%)
Lifetime PCL score ^a	28.4 (11.4)	15.2 (15.2)
Positive screen for lifetime PTSD	95 (7.1%)	44 (10.0%)
Past-month PCL score ^a	24.2 (10.1)	9.4 (12.1)
Positive screen for past-month PTSD	41 (3.8%)	15 (3.5%)
Lifetime major depressive disorder	244 (18.2%)	60 (25.6%)
Lifetime alcohol use disorder	659 (48.6%)	212 (41.7%)
Lifetime drug use disorder	200 (14.6%)	63 (12.3%)
APOE ε4 allele carrier	334 (24.1%)	133 (25.6%)

0 alleles	1,052 (75.9%)	376 (74.4%)
1 allele	318 (22.9%)	124 (23.7%)
2 alleles	16 (1.2%)	9 (1.9%)

Note. PCL, PTSD Checklist; PTSD, posttraumatic stress disorder.

^aThe DSM-IV version of the PCL was used in the main sample (score range = 17--85) and the DSM-5 version of the PCL was used in the replication sample (score range = 0--80).

Table 2. Results of Analyses Evaluating Relation Between *APOE* ϵ 4 Allele Carrier Status, Cumulative Trauma Burden, and Lifetime and Past-Month PTSD Symptoms

Main Sample ($n = 1,386$)				
	Lifetime		Past-Month	
	PTSD Symptoms		PTSD Symptoms	
	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>
<i>APOE</i> ϵ 4 carrier	16.11	<.001	6.75	.010
Cumulative trauma burden	28.31	<.001	22.48	<.001
<i>APOE</i> ϵ 4 $\times$ cumulative trauma burden	6.08	<.001	8.09	<.001
Replication Sample ($n = 509$)				
	Lifetime		Past-Month	
	PTSD Symptoms		PTSD Symptoms	
	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>
<i>APOE</i> ϵ 4 carrier	.41	.52	.04	.84
Cumulative trauma burden	11.97	<.001	8.66	<.001

<i>APOE</i> ε4 × cumulative trauma burden	2.53	.006	3.06	.001
---	------	------	------	------

Note. PTSD, posttraumatic stress disorder; *APOE* ε4, apolipoprotein epsilon 4.

Analyses are adjusted for age, sex, ancestral proportion scores, combat veteran status, and type of index trauma (assaultive vs. nonassaultive).

Author Manuscript

Table 3. Results of Analyses Evaluating Relation Between *APOE* ϵ 4 Allele Carrier Status, Cumulative Trauma Burden, and Lifetime PTSD Symptom Clusters

Main Sample ($n = 1,386$)								
	Re-experiencing		Avoidance		Emotional Numbing		Hyperarousal	
	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>
<i>APOE</i> ϵ 4 carrier	4.12	.043	5.98	.015	2.32	.13	4.11	.043
Cumulative trauma burden	9.08	<.001	3.30	<.001	4.05	<.001	3.09	<.001
<i>APOE</i> ϵ 4 $\times$ cumulative trauma burden	4.24	<.001	3.01	.001	1.63	.085	2.14	.016
Replication Sample ($n = 509$)								
	Intrusions		Avoidance		Negative Cognitions and Mood		Alterations in Arousal and Reactivity	
	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>
<i>APOE</i> ϵ 4 carrier	14.99	<.001	.22	.64	5.29	.022	1.96	.16
Cumulative trauma burden	3.89	<.001	.74	.71	1.12	.34	2.96	.001

<i>APOE</i> ε4 × cumulative trauma burden	3.59	<.001	.86	.57	1.47	.15	5.30	<.001
--	------	-------	-----	-----	------	-----	------	-------

Note. PTSD, posttraumatic stress disorder; *APOE* ε4, apolipoprotein epsilon 4.

Analyses are adjusted for age, sex, ancestral proportion scores, combat veteran status, nature of index trauma

(assaultive vs. nonassaultive), and other PTSD symptom clusters.

The DSM-IV version of the PCL was used in the main sample and the DSM-5 version of the PCL was used in the replication sample.

Figure 1. Interaction of *APOE* $\epsilon 4$ allele carrier status and cumulative trauma burden in predicting lifetime severity of PTSD symptoms in the main sample ($n = 1,386$).

Note. PTSD, posttraumatic stress disorder; *APOE* $\epsilon 4$, apolipoprotein epsilon 4.

PTSD Checklist score range = 17--85. Lines represent fitted regression lines adjusted for age, sex, ancestral proportion scores, combat veteran status, and nature of index trauma (assaultive vs. nonassaultive). Slopes and 95% confidence intervals for $\epsilon 4$ allele noncarriers = 1.93 (1.72--2.15) and 3.31 (2.96--3.66) for $\epsilon 4$ allele carriers.

Figure 2. Prevalence of current PTSD in highly trauma-exposed veterans by $\epsilon 4$ carrier status and level of perceived social support in the main sample ($n = 602$).

Note. Current PTSD was identified as a score ≥ 50 on the DSM-IV version of the PTSD Checklist. Low, moderate, and high levels of perceived social support reflect tertiles of total scores on the 5-item version of the Medical Outcomes Study Social Support Scale.

*Statistically significantly greater prevalence than $\epsilon 4$ allele carriers with moderate and high levels of perceived social support, $\chi^2(2) = 9.83$, $P = .007$.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Mota, NP;Han, S;Harpaz-Rotem, I;Maruff, P;Krystal, JH;Southwick, SM;Gelernter, J;Pietrzak, RH

Title:

Apolipoprotein E gene polymorphism, trauma burden, and posttraumatic stress symptoms in US military veterans: Results from the National Health and Resilience in Veterans Study

Date:

2018-02-01

Citation:

Mota, N. P., Han, S., Harpaz-Rotem, I., Maruff, P., Krystal, J. H., Southwick, S. M., Gelernter, J. & Pietrzak, R. H. (2018). Apolipoprotein E gene polymorphism, trauma burden, and posttraumatic stress symptoms in US military veterans: Results from the National Health and Resilience in Veterans Study. *DEPRESSION AND ANXIETY*, 35 (2), pp.168-177. <https://doi.org/10.1002/da.22698>.

Persistent Link:

<http://hdl.handle.net/11343/293917>