MISS HANNA SARAH ROSINGER (Orcid ID : 0000-0001-8453-0532)

DR. ARMANDO GERALDES (Orcid ID : 0000-0001-7239-4529)

Article type : Original Article

The tip of the iceberg: genome wide marker analysis reveals hidden hybridization during invasion

Running title: Hybridization of co-invaders

Hanna S. Rosinger¹, Armando M. Geraldes^{2,4}, Kristin A. Nurkowski^{1,4}, Paul Battlay¹, Roger D. Cousens³, Loren H. Rieseberg⁴, Kathryn A. Hodgins¹

¹School of Biological Sciences, Monash University, Melbourne, VIC, Australia

²Department of Zoology, University of British Columbia, Vancouver, BC, Canada,

³School of BioSciences, University of Melbourne, Melbourne, VIC, Australia

⁴Department of Botany and Biodiversity Research Centre, University of British Columbia, Vancouver, BC, Canada

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/MEC.15768](https://doi.org/10.1111/MEC.15768)

This article is protected by copyright. All rights reserved

19 * **Correspondence:**

20 Hanna S. Rosinger

21 Hanna.Rosinger@monash.edu

22 **Abstract**

23 Biological invasions are accelerating, and invasive species can have large economic impacts as well
24 as severe consequences for biodiversity. During invasions, species can interact, potentially resulting
25 in hybridization. Here, we examined two *Cakile* species, *C. edentula* and *C. maritima*
26 (Brassicaceae), that co-occur and may hybridize during range expansion in separate regions of the
27 globe. *Cakile edentula* invaded each location first, while *C. maritima* established later, apparently
28 replacing the former. We assessed the evidence for hybridization in western North America and
29 Australia, where both species have been introduced, and identified source populations with 4561
30 SNPs using Genotype-by-Sequencing. Our results indicate that *C. edentula* in Australia originated
31 from one region of eastern North America while in western North America it is likely from multiple
32 sources. *Cakile maritima* in Australia is derived from at least two different parts of Europe while the
33 introduction in western North America is from one. Although morphological evidence of
34 hybridization is generally limited to mixed species populations in Australia and virtually absent
35 elsewhere, our genetic analysis revealed relatively high levels of hybridization in Australia (58%
36 hybrids using Admixture) and supported the presence of hybrids in western North America (16%
37 hybrids using Admixture) and New Zealand. Hybrids might be commonly overlooked in invaders, as
38 identification based solely on morphological traits may represent only the tip of the iceberg. Our
39 study reveals a repeated pattern of invasion, hybridization and apparent replacement of one species
40 by another, which offers an opportunity to investigate the role of hybridization and introgression
41 during invasion.

42 **Keywords:** invasion, hybridization, *Cakile edentula*, *Cakile maritima*, Genotype-by-Sequencing
43 (GBS), range expansion

44 **1 Introduction**

45 Biogeographic barriers on a global, regional and local scale are often overcome by human activities,
46 leading to biological invasions (Sax & Gaines, 2003; Simberloff, 2013; Vilatersana, Sanz, Galian, &
47 Castells, 2016). Biological invasions can have a large economic impact (Hoffmann & Broadhurst,
48 2016; Pimentel, Zuniga, & Morrison, 2005), as well as severe negative consequences for biodiversity

49 and ecosystems (Sakai et al., 2001). Most long-distance introductions of invasive species in historic
50 times are directly (e.g. ornamentals) or indirectly the result of anthropogenic activities (e.g. via
51 ballast on ships) (Baker, 1974; Ruiz et al., 2000; Sakai et al., 2001). Invasions can also lead to novel
52 interactions between species that previously had not co-occurred and, where there are no strong
53 reproductive barriers, this may lead to instances of hybridization (Abbott, 1992; Ellstrand &
54 Schierenbeck, 2000; Vallejo- Marín & Hiscock, 2016).

55 Rather than hybridization just being an incidental event, it could actually facilitate the success of
56 invasive plant species, as invasive hybrid lineages can have increased fecundity and size (Hovick &
57 Whitney, 2014). Various hypotheses have been proposed by which hybridization facilitates rapid
58 range expansion (Bock et al., 2015; Ellstrand & Schierenbeck, 2000), including evolutionary novelty,
59 increased genetic variation, heterosis, dumping genetic load (i.e. genetic rescue) (Ellstrand &
60 Schierenbeck, 2000) and demographic rescue (Mesgaran et al., 2016). But convincing empirical data
61 are limited. Hybridization is certainly not the sole evolutionary pathway to invasiveness, but can
62 catalyze its evolution (Ellstrand & Schierenbeck, 2000). Not all of the potential consequences of
63 hybridization are beneficial, however, and there can be significant costs associated with the
64 phenomenon, such as outbreeding depression (Baack, Melo, Rieseberg, & Ortiz-Barrientos, 2015)
65 and genetic swamping (Todesco et al., 2016). Our capacity to assess the role of hybridization during
66 any particular invasion is hampered by the fact that it can be difficult to identify, especially when
67 repeated backcrossing with one parental species has occurred rendering morphological identification
68 difficult (Ward, Gaskin, & Wilson, 2008). However, genome-wide molecular markers can provide
69 estimates of the extent of past hybridization and introgression across the genome (Payseur &
70 Rieseberg, 2016).

71 On the beaches of Australia, the North Island of New Zealand and western North America a repeated
72 pattern of invasion by two species of sea-rocket with contrasting mating systems (Barbour &
73 Rodman, 1970; Cousens, Ades, Mesgaran, & Ohadi, 2013; Cousens & Cousens, 2011; Rodman,
74 1974, 1986) offers a rare opportunity to investigate the role of hybridization during invasion in
75 distinct, geographically isolated regions. *Cakile edentula* (American sea-rocket), native to eastern
76 North America, invaded each location first, while *Cakile maritima* (European sea-rocket)
77 (Brassicaceae), native to Europe and northern Africa, arrived later. The invasion and replacement
78 history in western North America and Australia are reviewed elsewhere (Barbour & Rodman, 1970;
79 Cousens et al., 2013; Rodman, 1986), but we briefly outline it below.

In Australia, *C. edentula* was first recorded in Victoria in 1863 and subsequently spread along the coastline of Australia (Rodman, 1986). In 1897, *C. maritima* was recorded for the first time in Western Australia, and a second introduction into South Australia (1918: see Cousens et al., 2013; Ohadi et al., 2016) spread from there to the east (Heyligers, 1984; Rodman, 1986). In contrast to *C. edentula*, *C. maritima* seems still to be actively spreading in Australia and appears to have replaced *C. edentula* throughout much of its initial introduced range (Cousens et al., 2013; Rodman, 1986). In western North America, a similar pattern of replacement occurred. *Cakile edentula* was found near San Francisco around 1880 (Barbour & Rodman, 1970), while *C. maritima* reached western North America by 1935 where it was found sympatric with *C. edentula* near San Francisco. The most recent published field study showed that *C. maritima* had replaced *C. edentula* throughout most of coastal California but not Oregon or Washington (Boyd & Barbour, 1993). In each case, there has been complete replacement of *C. edentula* by *C. maritima* over wide geographic areas (Barbour & Rodman, 1970; Cousens et al., 2013; Rodman, 1986), which was originally assumed to involve either direct or indirect competition (Rodman, 1986), although several additional mechanisms have been proposed such as disease (Bock, 2008; Cousens et al., 2013; Thrall, Young, & Burdon, 2000), coincidence (Cousens et al., 2013; Rodman, 1986), or greater lifetime fecundity of *C. maritima* (Boyd & Barbour, 1993). However, the mechanism of the replacement remains unclear.

Cakile edentula and *C. maritima* are closely related and cross-compatible (Li, Cousens, & Mesgaran, 2019; Mesgaran et al., 2016; Rodman, 1974). Both species are found in coastal strandline habitat, providing opportunities for hybridization in regions where they co-occur, but the species exhibit contrasting mating systems (Rodman, 1974). *Cakile edentula* (self-compatible) benefits from high levels of reproductive assurance as it is able to set seeds autonomously at high rates (Li, Mesgaran, Ades, & Cousens, 2020); one of Baker's (1965) ideal weed traits. In contrast, the establishment of *C. maritima* (self-incompatible) may be initially hindered (during both initial establishment as well as subsequent range expansion) by a lack of compatible mates limiting sexual reproduction and resulting in strong Allee effects. The apparent presence of hybrids, based on an intermediate leaf and fruit shape of both parental species, in some sites in Australia led Mesgaran et al., (2016) to develop a model for the interacting species, with the novel outcome that transient hybridization could overcome Allee effects in *C. maritima*. As a consequence, we hypothesized that past hybridization with *C. edentula* could be a common feature of *C. maritima*'s establishment and range expansion in western North America, Australia and New Zealand.

We used genome-wide markers derived from Genotype-by-Sequencing (GBS) to examine the invasion history of these two species in Australia and western North America and quantify the extent and distribution of hybridization. There have been several previous studies examining the population genetic structure of *C. edentula* and *C. maritima* in their native ranges in Europe (Clausing, Vickers, Kadereit, 2000; Kadereit, Arafah, Somogyi, & Westberg, 2005; Westberg, 2005), Africa (Gandour, Hessini, & Abdelly, 2008), eastern and western North America (Gormally, Hamrick, & Donovan, 2011) as well as in the introduced range of Australia (Ohadi et al., 2016). However, no study of the invasion history on two continents has been attempted nor has the extent of hybridization across multiple introductions been quantified. Specifically, we aimed to (1) identify probable source regions (from Europe and eastern North America); (2) determine whether both recent and advanced generation hybrids occur in the introduced ranges and the extent of their geographic distribution; and (3) determine if the change in levels of species ancestry post-invasion reflects a chronosequence along the direction of invasion of *C. maritima*. We predicted that early generation hybrids should be present at the leading edge of *C. maritima*'s invasion into *C. edentula*-occupied areas, but later generation backcrosses with *C. maritima* should be more common in areas closer to where *C. maritima* first established. This should contribute to a gradient in species ancestry whereby *C. maritima* ancestry will be dominant in hybrids near the invasion source, while *C. edentula* ancestry will be more prevalent in hybrids identified in areas recently invaded by *C. maritima*. We predicted high levels of *C. maritima* ancestry in hybrids near the invasion source because *C. maritima* phenotypes are now exclusively present in the regions surrounding the invasion source, and studies of pollinators suggest preferential visitation of both hybrids and *C. maritima* over *C. edentula* which should facilitate backcrossing to *C. maritima* (Mesgaran et al., 2016).

2 Methods

2.1 Study species

Cakile maritima's native range extends over a wide climatic range from northern Norway to northern Africa. Current taxonomy recognizes subsp. *maritima* (Mediterranean), subsp. *baltica* (Baltic), subsp. *integrifolia* (Atlantic coast), subsp. *islandica* (Northern Europe and Northwestern Russia) and subsp. *euxina* (Black Sea) (Marhold, 2011). This is paralleled in the western Atlantic by *C. edentula*, for which two subspecies are recognized in its native range (Rodman, 1974) subsp. *edentula* (Labrador to North Carolina) and subsp. *harperi* (North Carolina to Florida). Although *C. maritima* has a sporophytic self-incompatibility system, the level of self-incompatibility varies among plants (Thrall

et al., 2000). *Cakile edentula* is self-compatible and can set seed autonomously at a high rate (Barbour, 1970; Rodman, 1974), although field estimates are suggestive of intermediate levels of autonomous selfing (Li et al., 2020). Both species are diploid ($2n = 18$) (Rodman, 1974). Hybrids are readily produced through artificial pollination (Rodman, 1974) with either parent as the pollen donor when emasculated (Li et al., 2019; Mesgaran et al., 2016), although crosses are more successful when *C. edentula* acts as the pollen recipient, consistent with the SI x SC rule (Harrison & Darby, 1955).

2.2 Samples

Samples of *Cakile spp.* were obtained from the native ranges (Europe and northern Africa, eastern North America) and the two introduced ranges (Australasia, western North America). We collected four of the five subspecies (subsp. *baltica*, subsp. *maritima*, subsp. *integrifolia* and subsp. *islandica*) of *C. maritima*. In the native range of *C. edentula* we sampled only *C. edentula* subsp. *edentula* as this subspecies is most likely the source of invasions in Australia and western North America (Cousens et al., 2013; Rodman, 1974). We obtained 214 samples of *C. maritima*, 137 samples of *C. edentula*, 17 putative hybrids (identified by morphology in the field) and two *C. lanceolata* samples from 92 locations in total (Figure S1; Table 1 & S1). Most samples were our own field collections of silica dried leaf tissue although a few samples were purified DNA from colleagues. We collected our samples along a transect through a population, ensuring that individuals were at least 2 m apart to avoid sampling close relatives or the same individual and collected individuals randomly with respect to their putative species based on morphology.

2.3 DNA extraction and Genotype-by-sequencing

We performed DNA extractions from dried leaf material using a modified CCDB DNA Extraction Protocol following Whitlock, Hipperson, Mannarelli, and Burke (2008). DNA quantity was assessed using a QuBit broad-sensitivity DNA quantification system (Invitrogen, Carlsbad, CA, USA) and a double-digest GBS library preparation was carried out (using PstI-HF (NEB) and MspI (NEB) enzymes, see Supplementary Information for details). Sequencing (125bp PE) was conducted on an Illumina HiSeq2500 (McGill University and Genome Quebec Innovation Centre) on two lanes.

2.4 SNP calling

Quality statistics of raw reads were assessed through FastQC (http://hannonlab.cshl.edu/fastx_toolkit) and the reads were demultiplexed using STACKS process_radtags (Catchen, Amores, Hohenlohe, Cresco, & Postlethwait, 2011). We removed adapter sequences and trimmed the reads using Sickle (Joshi & Fass, 2011) with a Q-score of ≥ 20 and read length of ≥ 20 base pair. FASTQ quality filter (http://hannonlab.cshl.edu/fastx_toolkit) was then used to filter for reads with a Q-score of 20 or greater for $\geq 90\%$ of the read length. The filtered reads were aligned using the Burrows-Wheeler Aligner (BWA) (Li & Durbin, 2009) to a *C. maritima* draft genome. Early access to the draft genome was provided by S.I. Wright, University of Toronto (<https://genome.jgi.doe.gov/portal/CakmarStandDraft/CakmarStandDraft.info.html>, GenBank: MK637688.1). The reference genome is found in 26,153 scaffolds with a scaffold N50 of 85,425. We assessed if there was a bias when mapping the reads of *C. edentula* to the reference genome of *C. maritima* but found limited evidence for such a bias (see Supplementary Information, Figure S2 & S3).

We called variants with GATK HaplotypeCaller (Poplin et al., 2017). We refer to this as the *unfiltered dataset* (Rosinger et al., 2020). Using VCFtools (Danecek et al., 2011) we removed individuals with fewer than 25000 reads, removed indels and restricted individual genotypes to have a depth between 5- 100,000. Furthermore, we filtered for a minimum quality score of 20, a genotype quality of 20, and a minor allele frequency of 0.05. Subsequently, we kept only bi-allelic variants that were successfully genotyped in more than 50% of individuals and removed individuals that had more than 50% missing data. The above filtering steps resulted in 18,573 SNPs from 258 individuals. Additionally, we removed 121 SNPs which showed $> 80\%$ observed heterozygosity, because such high observed heterozygosity could be caused by paralogues. We refer to this as the *filtered dataset* (Rosinger et al., 2020), which had a mean coverage of 39.21 (minimum coverage 9.18, maximum coverage 504.73).

2.5 Genetic clustering

Population genetic structure was inferred using Admixture (Alexander, Novembre, & Lange, 2009). For Admixture and most of our analysis we thinned our *filtered dataset* for linkage using a single SNP per 1kb window, resulting in a reduction to 4561 SNPs from 257 individuals (excluding the outgroup *C. lanceolata*). We will refer to this as the *global thinned dataset*. We ran Admixture using the *global thinned dataset* with a major termination criterion of 1×10^{-9} , 1000 bootstraps and ten-fold cross-validation for $K=1-10$, where K equals the number of genetic groups. The K that produced the lowest cross-validation error was selected as the best K value. We refer to this as the *unsupervised run*. All following analyses were conducted in R v.3.5.2 (R Core team, 2018) except where otherwise stated. The output of Admixture visualized with pophelper v.2.3.0 (Francis, 2017) and pie charts.

To complement the population clustering analysis and to provide further insight in the population differentiation, we conducted a principal component analysis (PCA) and an unrooted phylogenetic network analysis. Genetic differentiation between native and introduced populations was summarized in a PCA with an 95% confidence ellipse using the R package SNPRelate (Zheng et al., 2012), tidyverse (Wickham, Francois, Henry, & Müller, 2019) and car (Fox & Weisberg, 2019) on the *global thinned dataset*. We used SPLITSTREE5 (Huson & Bryant, 2006) to visualize the overall sample relatedness with an unrooted phylogenetic network. To do this, we created two datasets from our unfiltered dataset (see details in Supplementary Information); (1) a global dataset containing all samples (*global Splitstree dataset*) and (2) a native range dataset containing samples from Europe and eastern North America (*native range Splitstree dataset*).

2.6 Hybrid identification

We used three different approaches to identify hybrids using genetic data:

(1) A *supervised run* of Admixture for $K=2$ using the *global thinned dataset*, by setting the samples from the two native ranges as reference individuals. Providing known ancestries allows the program to set some rows in the matrix Q to known constants and provides a more accurate estimation of the ancestries of the remaining individuals, and of the ancestral allele frequencies (Alexander et al., 2009). The other settings were retained from the *unsupervised run*. We refer to this as the *supervised run* and used this run to classify individuals by their Q -scores as hybrid, or pure species. We used the highest standard error from the Q scores, resulting in individuals classified as hybrids if $0.025 < Q < 0.975$ of their genome was assigned to the *C. edentula* cluster.

(2) We used the program NewHybrids (Anderson & Thompson, 2002) to identify early generation hybrids. It classifies their generation using a Bayesian model-based clustering framework to compute, by Markov chain Monte Carlo, the posterior probability that each individual belongs to each of the distinct first two generation hybrid classes (parental species, F1, F2, BC to species 1, BC to species 2). As the program is unable to deal with a large dataset, we restricted our data to 63 SNPs that showed fixed differences between the two species obtained from individuals classified as parental species using the *supervised run* of Admixture. Details of the settings used are provided in the Supplementary Information.

(3) We used the R package Hlest (Fitzpatrick, 2012), which uses maximum likelihood to estimate ancestry and heterozygosity. For this package, we used the 471 loci that showed fixed differences between the individuals of the native ranges. Because it is possible that there is a low level of segregating variation within each species for these loci due to sampling error, particularly for *C. maritima* where the sample size is lower, we set the allele frequencies as 0.99 for *C. edentula* and 0.03 for *C. maritima*. We also tested other SNP sets and allele frequencies. The details of the settings used and the hybrid assignments along with the results are provided in the Supplementary Information.

We tested for a chronosequence by assessing if there was a correlation between the distance of each population from the first entry point of *C. maritima* (Adelaide in Australia, San Francisco in western North America) and the level of *C. maritima* and *C. edentula* ancestry using a Spearman's rank correlation test in R using the ggpubr package (Kassambara, 2020). We used the ranked order of populations from this origin point along the coastline for each range. In Australia, we only used the south-east mainland individuals (see Supplementary Information for details). We tested the correlation between the Q value of the *C. edentula* cluster of the *supervised run* for each population and the rank order of the sampling locations along the coastline to the first entry point of *C. maritima*. We used individuals that were classified as hybrids by Admixture or all samples (including the parental species). We repeated this analysis using the S value from Hlest and the hybrid classifications of this program.

Additionally, we used the program TreeMix (Pickrell & Pritchard, 2012) to identify evidence for hybridization in the introduced ranges using the *global thinned dataset* for which we constructed maximum likelihood trees and calculated the f_3 statistic (for details see Supplementary Information).

260 2.7 Genetic diversity and differentiation

261 Genetic diversity and differentiation within the two native ranges and two introduced ranges were
 262 assessed for the 256 individuals (the New Zealand and *C. lanceolata* samples were excluded) using
 263 the *global thinned dataset*. We calculated observed heterozygosity (H_O) and allelic richness (A_R)
 264 with the *diveR*sity package (Keenan, McGinnity, Cross, Crozier, & Prodöhl, 2013). The 95%
 265 confidence intervals of A_R were calculated with 1000 bootstraps. We estimated differences in genetic
 266 diversity between the species and ranges because we expected self-fertilization in *C. edentula* and
 267 bottlenecks potentially experienced during introduction would reduce diversity. Because sampling at
 268 individual locations was limited in the native ranges, we grouped individuals based on their range,
 269 and their hybrid ancestry (pure parental or hybrid) using the *supervised run* Q-value assignments of
 270 the *global thinned dataset* into eight groups. We used the Q value assignment of the *C. edentula*
 271 cluster and the highest standard error (0.024) of the *supervised run* to classify individuals. To
 272 determine regional differentiation we calculated Weir and Cockerham's (1984) pairwise F_{ST} between
 273 the above eight groups using the *global thinned dataset* with VCFtools (Danecek et al., 2011).
 274 Additionally, we calculated the F_{ST} for pure parental individuals, grouping individuals according to
 275 their Admixture cluster from the *unsupervised run* and range (see Supplementary Information).

276

277 3 Results

278 3.1 Genetic structuring and differentiation

279 The Admixture analysis of the *unsupervised run* showed genetic structuring of *C. maritima*, *C.*
 280 *edentula* and hybrids with an optimal K value of 8 (Figure 1 A, B & S4). Genetic structure was
 281 present in the native range of *C. edentula*, where single samples from Lake Michigan and Rhode
 282 Island constituted one group, samples from New Brunswick within the Gulf of St. Lawrence a second
 283 group, samples from Newfoundland and Quebec (along the St Lawrence River) a third group and
 284 samples from Nova Scotia a final group. As expected, for *C. maritima*, there were two main groups:
 285 one group was largely from the Baltic and Atlantic coasts, which we term the "Atlantic" group
 286 (comprising mainly the dark blue cluster, Figure 1 A, B) and a second admixed group was associated
 287 with the Mediterranean, that we term the "Mediterranean" group (comprising mainly the light and

medium blue clusters, Figure 1 A, B). In Australia, several genetic clusters were identified. First, in Queensland, New South Wales and Tasmania we identified pure *C. edentula* individuals. Second, for populations along the west coast of Australia, we identified a *C. maritima* cluster associated with the Atlantic coast in the native range. Third, in South Australia, genetic clusters associated with the Mediterranean were found. In the south-east of Australia there was evidence of hybrids between *C. maritima* and *C. edentula* (see below). In the introduced range of western North America, we identified pure *C. edentula* along with pure *C. maritima* (Figure 1 A, B). A small number of samples from Washington, Oregon and California showed evidence of hybridization (see below).

The PCA and SPLITSTREE5 analyses confirmed the findings of Admixture. There was clear differentiation of *C. maritima* and *C. edentula* in the *global thinned dataset*. The first eigenvector (EV) (Figure 2 A & S5 A) explained 33.17% of the variation and clearly delineated the species. The *C. edentula* group showed less variation than the *C. maritima* group along the first two EVs. Two *C. maritima* groupings were also evident with one representing *C. maritima* from Europe and Australia (EV1<0, EV2<0) and the other representing exclusively *C. maritima* from western North America (EV1<0, EV2>0). In the SPLITSTREE5 network, using the *global Splitstree dataset*, *C. edentula* (as identified by the *supervised run*) formed a monophyletic group without admixture. *C. maritima* samples were split into three groups (Figure 2 B, C): *C. maritima* (Mediterranean group), *C. maritima* (Atlantic group) and *C. maritima* in western North America. Hybrids of the two species were scattered in between the *C. maritima* groups or between the two-parental species along the network. The additional native range SPLITSTREE5 analysis (Figure S6) mirrored this pattern but provides clearer *C. edentula* grouping in the native range.

Pairwise F_{ST} (Table S2) using the *global thinned dataset* revealed clear genetic differentiation between the two-parental species originating from the native range ($F_{ST} > 0.527$). Within the introduced ranges the pairwise F_{ST} between the two species was similar to the comparison of the native ranges. Hybrids identified using Admixture in the introduced ranges showed higher genetic differentiation from *C. edentula* than from *C. maritima* (Table S2).

3.2 Genetic diversity

Population statistics revealed that in their native ranges, *C. edentula*, the self-compatible species, has considerably less H_O than *C. maritima* and the hybrids of the two species (Table S3). A_R was

significantly reduced in *C. edentula* in comparison to *C. maritima*, the largely self-incompatible species. In the introduced ranges, no clear reduction of H_O or A_R was observed in either of the species. Hybrids of the two-species had higher H_O and A_R compared to both parental species.

3.3 Hybrid classification

The three approaches classified different proportions of individuals as hybrids, as expected due to their ability to detect recent hybrids (NewHybrid, Hiest), versus hybrid ancestry (Admixture, Hiest). All hybrids identified by NewHybrids were also identified as hybrids with Hiest and Admixture (Table S4 & S5). The fourteen putative hybrids included in the samples as a result of morphological identification were assigned by all analyses as hybrids, providing evidence of the accuracy of the assignments. Furthermore, the NewHybrid analysis confirmed that these hybrids were likely the product of the first two generations of interbreeding. NewHybrids analysis revealed 19 hybrids (Figure 3; Table S4) with 17 hybrids in Australia (13.49%), one in western North America (1.47%) and one in New Zealand. In Australia, F1 and F2 hybrids were detected in the current sympatric zones where individuals with both species' phenotypic traits were clearly identifiable in the populations. Hybrids (Figure S5 B) grouped in the PCA according to their generation, with F1 and F2 hybrids grouped between the parental species, and backcrosses grouped closer to species they backcrossed to. In this same PCA the advanced generation hybrids identified with the *supervised run* of Admixture as well as Hiest frequently grouped with *C. maritima*, suggestive of further backcrossing to that species.

Classification of hybrids using the *supervised run* of Admixture revealed 73 hybrids in Australia (57.94%) from 15 locations, 11 hybrids in western North America (16.18%) from five locations and one hybrid from New Zealand (Figure 1; Table S4). In western North America hybrids were found in each of two locations in California and Oregon and in one location in Washington.

All Admixture hybrids were also identified as hybrids in Hiest and the ancestry assignments were highly correlated between the programs (Figure S7). When the 471 loci that are fixed between native range samples were used, and we allowed for a low level of polymorphism within each species (0.99 *C. edentula*, 0.03 *C. maritima*), a larger number of hybrids were identified using Hiest than Admixture (138 versus 85, Table S6). Changing the allele frequencies and SNP set impacted the number of hybrids identified (see Supplementary Information), but this only influenced the

classification of individuals with an apparent low level of ancestry from the alternate species. In all the runs, advanced generation hybrids were identified in this analysis with many in regions where *C. maritima* has not been recorded for many decades, but also in the current sympatric zone (New South Wales, Queensland and Tasmania).

We then examined if patterns of ancestry in Australia and western North America reflected the likely invasion route of *C. maritima*. Specifically, we tested if low levels of *C. edentula* ancestry were found in areas where *C. maritima* first arrived, and if high levels of *C. edentula* ancestry were found in regions *C. maritima* has more recently invaded and where *C. edentula* is still present. Using the supervised Admixture analysis, the mean *C. edentula* ancestry of hybrids at each location was correlated with the ranked distance from where *C. maritima* first arrived in south-eastern mainland Australia ($\rho = 0.82$, $p < 0.01$) (Table 2). This pattern was also significant when testing across all samples, including individuals identified as parental species ($\rho = 0.89$, $p < 0.05$). However, in western North America, although the direction of the correlation was as predicted, a geographic pattern in ancestry was only significant when using locations north of San Francisco as well as parental and hybrid individuals ($\rho = 0.72$, $p < 0.05$). The same pattern of significance was found when using the results of the Hltest (Figure 4 & 5 & S8; Table 2).

We used TreeMix to assess geneflow between *C. edentula* and *C. maritima* within each introduced range. The maximum likelihood tree in both invasive ranges showed bidirectional gene flow (Figure 6). In Australia geneflow occurred from the *C. edentula* branch into Australian *C. maritima* (Mediterranean); a migration event also occurred from this group into the *C. edentula* branch (Figure 6 B). In western North America the same pattern occurs. There is evidence of a migration event from the *C. edentula* branch into western North American *C. maritima* as well as a migration event from the western North American *C. maritima* branch into the western North American *C. edentula* (Figure 6 A). The f_3 statistic of TreeMix (Table 3) confirmed that the hybrids (identified by the supervised Admixture run) in the introduced range are admixed from the *C. edentula* and *C. maritima* parental individuals within both introduced ranges (Australia $f_3 = -0.006$, $Z = -31.97$; western North America $f_3 = -0.005$, $Z = -23.22$).

4 Discussion

Our analysis sheds light on the origin and extent of hybridization of two introduced species in two separate invasions, which experienced a parallel pattern of invasion and apparent replacement of one species by another. Except at places where the two species are currently sympatric and new hybrids are still being formed, it would be difficult to determine morphologically that hybridization has ever taken place, since backcrossing soon hides its phenotypic evidence. *Cakile maritima* is highly variable within and between populations in its native range and hybrids in the introduced range could easily be overlooked (e.g. Cousens et al., 2013) without the use of molecular methods. However, our analysis identified extensive hybrid ancestry in the introductions, particularly in Australia. It is therefore an intriguing possibility that hybridization may be commonly overlooked in a much wider range of invasive taxa, especially where morphological trait indicators of hybridization are more cryptic. Alien floras commonly include many congeneric species whose capacity for interbreeding is yet to be established. While previous authors (Ellstrand & Schierenbeck, 2000) have raised our attention to obvious hybrid species and allopolyploids, perhaps the impacts of hybridization are often more insidious. It is thus important – though not an easy task – to determine in future the extent to which such non-apparent introgression has been beneficial during invasion.

4.1 Native range patterns

One of our primary goals was to identify the source regions for the invasions for each species and our analysis provided evidence of geographic structuring in the *C. edentula* native range, at a much finer grain than currently recognized taxonomically (Figure 1). Samples from Quebec, Newfoundland, Nova Scotia and New Brunswick contain separate Admixture clusters, likely within *C. edentula* subsp. *edentula* var. *edentula* as this subspecies is the only one described in this region of the North American Atlantic coast (Rodman, 1974). Two single samples from Lake Michigan and Rhode Island grouped together in one cluster of the Admixture analysis; those samples might belong to the Atlantic coast variety of *C. edentula* subsp. *edentula* var. *edentula* as it is known to have invaded Lake Michigan in historical times (Huebner, 2009; Rodman, 1974), where it now coexists with the Great Lakes endemic var. *lacustris*. A second possibility, suggested by Gormally et al., (2011), but without morphological evidence, is that var. *lacustris* has dispersed to the Atlantic. Genetically distinct regional variation is not surprising, as the directions of currents and the influences of geological features on seed dispersal can be highly predictable (Lapointe, 2000). Similar conclusions have been reached in the Mediterranean by Westberg (2005) and Gandour et al. (2008). *Cakile edentula* subsp. *harperi* occurs in areas south of the populations sampled in our study (Rodman, 1974), but comprehensive studies of herbarium samples by Rodman (1974) and Cousens et al.,

(2013) have found no morphological evidence that subsp. *harperi* has been introduced anywhere outside its native range.

Our analyses revealed clustering of *C. maritima* in its native Europe largely consistent with the accepted taxonomic distributions (Ball, 1964; Marhold, 2011; Rodman, 1974) as well as one previous population genetic analysis (Clausing et al., 2000). Other genetic studies with greater sampling intensity, however, showed more differentiation on a local level (Kadereit et al., 2005; Westberg, 2005). The absence of fine-grain local differentiation in our study might be driven by the limited number of native range samples for this species and restricted sampling of the Baltic area.

Cakile edentula showed lower genetic diversity than *C. maritima* in their native ranges as measured by A_R and H_O (Table S3) and showed less variation along the EVs and in the SPLITSTREE network analysis (Figure 2). Higher selfing rates in *C. edentula* would be expected to reduce the effective population size compared to the largely self-incompatible *C. maritima* (Pollak, 1987).

4.2 Introduced range patterns

4.2.1 Australia and New Zealand

Although *C. edentula* has now disappeared from much of its original introduced range in Australia, some pure *C. edentula* populations still remain. Our analyses show that they likely originate from populations located in Nova Scotia as they contained an Admixture cluster found exclusively in this region of the native range and showed the lowest genetic differentiation from this region (Figure 1; Table S7). *Cakile edentula*'s A_R and H_O did not change considerably in Australia compared to the native range (Table S3), which is inconsistent with a strong invasion bottleneck. The genetic structure of the Australian *C. maritima* samples is consistent with a history of multiple introductions. This is in accordance with previous morphological and genetic studies of invasion history in Australia (Cousens et al., 2013; Ohadi et al., 2016; Rodman, 1976, 1986). In particular, the cluster associated with the Atlantic European group is found in western Australia, while a Mediterranean cluster predominates in southern and eastern Australia (Figure 1; Table S8). Similarly, analysis of microsatellite markers indicated that that western and south-eastern populations of *C. maritima* in Australia were genetically distinct and most likely resulted from independent introductions with severely limited gene flow from west to east (Ohadi et al., 2016). Finally, Australian *C. maritima* showed higher A_R and H_O values than its native range, consistent with admixture of multiple source populations and/or hybridization with *C. edentula*. Many successful invasions are sourced from

multiple introductions (e.g., Vallejo- Marín et al., 2020; van Boheemen et al. 2017) and both hybridization and multiple introductions and admixture may spur successful invasions (Ellstrand & Schierenbeck, 2006; Dlugosch & Parker, 2008; Hodgins, Bock, Rieseberg, 2018).

Our data provides substantial evidence for extensive hybridization in Australia between the two species. TreeMix supported bidirectional gene flow between the parental species (identified morphologically) (Figure 6). This was confirmed by the Admixture global analysis (Figure 1), the PCA and SplitsTree analysis, as many Australian samples fell in-between the native range samples of both species (Figure 2), and the f_3 test (Table 3). Further support is provided by three separate analyses which specifically detect hybrid individuals (Figure 1 & 3 & 4 & 5 & S8; Table S4). As expected, Australian hybrids (*supervised Admixture run*) had higher genetic diversity than both parental species (Table S3). Furthermore, the pattern of hybrid ancestry was geographically structured and reflected the historical invasion route of *C. maritima* in south-eastern Australia. This pattern was consistent across two separate approaches (*supervised Admixture run*, H1est) to identify hybrid ancestry (Figure 1 & 4; Table 2). NewHybrids confirmed the presence of a small number of early generation hybrids (within two generations) where both species still co-occur and some mixed populations show pure genotypes of both parental species and early generation hybrids, demonstrating on-going hybridization of the two taxa (Figure 3). In areas where *C. edentula* still persists, backcrossing to *C. edentula* has also occurred, but is rare, and recent backcrosses to *C. maritima* appear to be more common. In those parts of Australia where *C. maritima* has already appeared to have replaced *C. edentula* (i.e., where no *C. edentula* phenotypes remain; Cousens et al., 2013; Rodman, 1986), evidence is consistent with past hybridization between the species and repeated backcrossing to *C. maritima* (Figure 1 & 4 & 6). In areas of Western Australia, where *C. edentula* has never been identified, evidence of hybridization with *C. edentula* was also found, confirming a previous observation by Ohadi et al., (2016). The sample from New Zealand was identified as a hybrid where the same replacement of *C. edentula* by *C. maritima* has also taken place (Cousens & Cousens, 2011).

4.2.2 Western North America

Our results revealed that *C. edentula* in western North America most likely originated from two sources in eastern North America. We also found that western North American *C. maritima* potentially originated from the Mediterranean region, as *C. maritima* in western North America contained the same Admixture clusters as the Mediterranean and showed the lowest differentiation

from this region (Figure 1; Table S7 & S8). However, these populations were genetically distinct (Figure 2 & S5) suggesting the possibility of an unknown source for this invasion, or the impact of an invasion bottleneck. *Cakile edentula* and *C. maritima* in western North America showed, as in Australia, no reduction of H_O and A_R , which may reflect the impacts of undetected hybridization, large founding populations, or multiple introductions.

Like Australia, hybridization was identified between the two species in western North America, although the proportion of hybrids was less (e.g., 58% versus 16% using the *supervised Admixture* run). TreeMix identified bidirectional gene flow between the species in western North America (Figure 6; Table 3), and evidence consistent with hybridization was apparent in the global Admixture analysis (Figure 1), the PCA and Splitstree analysis (Figure 2). Furthermore, we employed three independent methods to specifically identify hybrid individuals and their likely generation. From this we identified 11 hybrid samples (all 11 were identified by both H1est and Admixture and one as an F2 by NewHybrids) from five locations in western North America. Specimens of hybrids based on morphological identification are largely unknown for this region, either in herbaria or in the field (Rodman, 1974). But more recently, Cody and Cody (2004) reported a small percentage of hybrids in a population from British Columbia. Although the fitness and demographic consequences of hybridization during introduction require further investigation, the lower incidence of hybrids in western North America compared to Australia suggests that hybridization could have facilitated the establishment and rapid spread of *C. maritima* to a greater degree in Australia. In support of this hypothesis, the complete replacement of *C. edentula* by *C. maritima* phenotypes has not progressed as far north in western North America compared to Australia, where few northern populations of *C. edentula* remain. Indeed, although the introduction of *C. maritima* in western North America is more recent than Australia, migration rates for this species based on herbarium records are much lower in western North America (Barbour & Rodman, 1970; Rodman, 1986). However, the mechanism driving differences in hybridization rates in western North America compared to Australia is unclear and requires further investigation.

4.3 Hybrid identification and significance

The pattern of invasion first by *C. edentula*, then by *C. maritima*, has been repeated in three regions. Prior to this study, hybrids were known only from Australia. However, we also identified clear evidence of hybridization in western North America and in New Zealand. Hybrids between the two species can be produced readily by handcrossing (e.g. Li et al., 2019; Mesgaran et al., 2016; Rodman,

1974) and our data demonstrate that recent and advanced generation hybrids are at least partially fertile in natural populations. Our results show backcrossing to both parental species, although backcrossing to *C. maritima* was much more frequent. This pattern of biased backcrossing towards *C. maritima* was predicted based on field observations of pollinator visitations (Mesgaran et al., 2016), the morphological replacement of *C. edentula* by *C. maritima*, and previous genetic studies (Mesgaran et al., 2016; Ohadi et al., 2016). It is also consistent with expected mating asymmetries between these species and their hybrids caused by the inheritance of the self-incompatibility system and traits associated with pollinator attraction in hybrids (Li et al., 2019). In artificial crosses, early generation hybrids inherited mostly (but not exclusively) self-incompatibility, as well as larger floral displays, similar to *C. maritima* (Li et al., 2019). This suggests that F1 hybrids will often need to rely on outcrossing, and that larger floral displays should facilitate this. Consequently, these traits in the hybrids should further contribute to backcrossing to the self-incompatible parent (*C. maritima*). A similar asymmetric pattern of species ancestry has been identified in hybrids of other species with such differences in mating system (Brandvain, Kenney, Flagel, Coop, & Sweigart, 2014; Pickup et al., 2019; Ruhsam, Hollingsworth, & Ennos, 2011).

Our identification of advanced generation backcrosses to *C. maritima* means that portions of the *C. edentula* genome have been retained in a largely *C. maritima* background (i.e. introgression), long after morphological evidence of hybridization has gone from a population. The role of selection and neutral evolutionary processes in governing patterns of introgression across the genome, however, remains to be investigated in this system. Theory suggests that regions of the genome that are not introgressed will harbour incompatibilities or a high number of additive deleterious alleles in the introgressing species (Harris & Nielsen, 2016; Juric, Aeschbacher, & Coop 2016). A greater fixation rate of weakly deleterious alleles is predicted in the *C. edentula* due to its higher level of inbreeding, and indeed, the low levels of genetic variability in this species relative to *C. maritima* support a lower effective population size in this species. Selection against a higher genetic load originating from *C. edentula* in hybrids should more rapidly lead to the reconstitution of a *C. maritima* genome following transient hybridization during range expansion. In line with the expectation of selection against selfing ancestry in outcrossers, in *Mimulus guttatus* (outcrossing) genomic regions with high recombination rates have reduced levels of ancestry from the selfing species *Mimulus nasutus* (Brandvain et al., 2014). However, several remarkable examples in plants have demonstrated the infusion of favorable alleles via hybridization (adaptive introgression), including the transfer of herbivore resistance in *Helianthus* (Whitney, Randell, & Rieseberg, 2006). Indeed, Cody and Cody

(2004) proposed the intriguing possibility of adaptive introgression in the *Cakile* system but this remains to be investigated. Our identification of replicated patterns of hybridization, replacement and invasion in *Cakile* provide an exciting opportunity for further investigation of the beneficial and detrimental consequences of hybridization during range expansion.

5 Conclusion

Here we confirm that, particularly in Australia, the apparent replacement of *C. edentula* by *C. maritima* is not complete and remnants of the *C. edentula* genome are evident in contemporary *C. maritima* populations. Furthermore, it appears that both early and later generation hybrids are at least partially fertile in natural populations and that there is a higher frequency of backcrossing to *C. maritima*. The patterns of hybridization we identified is consistent with the hypothesis that mating among these cross-compatible invaders has facilitated the establishment of the self-incompatible *C. maritima* whose range expansion may otherwise be limited due to Allee effects, as has been observed in other potential self-incompatible invaders (Uesugi, Baker, de Silva, Nurkowski, & Hodgins, 2020). The evolutionary consequence of hybridization for both species remains unclear, as is its role, if any, in the rapid expansion of one invader at the expense of another.

6 Author Contributions

KH, RC and LR conceived of and designed the study. KH, KN and RC carried out sampling. KN conducted the molecular laboratory work. HR carried out the bioinformatics analyses with significant input from AG, PB and KH. AG, KH, LR, PB, RC and HR contributed to the writing and approved the final manuscript.

7 Funding

KH, RC and LR received funding from the Australian Research Council (Grant ID: DP180102531). HR was supported by Monash Graduate Scholarship (MGS).

8 Acknowledgments

We would like to thank all collectors of this study, Sarah Bou-assi for help with molecular laboratory work and Lotte van Boheemen for help with the initial bioinformatics analysis.

9 Data Availability Statement

Sequence data are available at the National Center for Biotechnology Information Sequence Read Archive under Bioproject PRJNA637114. Unfiltered dataset available on https://bridges.monash.edu/articles/dataset/GBSCAK_vcf_gz/12526220/1; filtered dataset on https://bridges.monash.edu/articles/dataset/filtered_dataset_GBS_Cakile/12996854. Scripts are available on <https://github.com/HannaRos/Cakile-GBS-scripts>.

10 Data reference

[dataset]. Rosinger, H.S., Geraldles, A. M., Nurkowski, K. A., Battlay, P., Cousens, R. D., Rieseberg, L. H., Hodgins, K. A.; 2020; GBSCAK.vcf.gz; Monash University Bridges; <https://doi.org/10.26180/5ef01e7a6359b>

[dataset]. Rosinger, H.S., Geraldles, A. M., Nurkowski, K. A., Battlay, P., Cousens, R. D., Rieseberg, L. H., Hodgins, K. A.; 2020; filtered dataset GBS Cakile; Monash University Bridges; <https://doi.org/10.26180/5f6be2a84dc4d>

11 References

Abbott, R. J. (1992). Plant invasions, interspecific hybridization and the evolution of new plant taxa. *Trends in Ecology & Evolution*, 7(12), 401-405.

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, 19(9), 1655-1664. <https://doi.org/10.1101/gr.094052.109>

Anderson, E. C., & Thompson, E. A. (2002). A model-based method for identifying species hybrids using multilocus genetic data. *Genetics*, 160(3), 1217-1229.

Baack, E., Melo, M. C., Rieseberg, L. H., & Ortiz-Barrientos, D. (2015). The origins of reproductive isolation in plants. *New Phytologist*, 207(4), 968-984. <https://doi.org/10.1111/nph.13424>

Baker, H. G. (1965) Characteristics and modes of origin of weeds. In: The Genetics of Colonizing Species (eds Baker H, Stebbins G), pp. 147–168. Academic Press, New York.

Baker, H. G. (1974). The evolution of weeds. *Annual Review of Ecology and Systematics*, 5(1), 1-24. <https://doi.org/10.1146/annurev.es.05.110174.000245>

Ball, P. W. (1964). A revision of *Cakile* in Europe. *Feddes Repertorium*, 69, 35-40.

Barbour, M. G. (1970). Seedling ecology of *Cakile maritima* along the California coast. *Bulletin of the Torrey Botanical Club*, 280-289. <https://doi.org/10.2307/2483647>

Barbour, M. G., & Rodman, J. E. (1970). Saga of the West Coast sea-rockets: *Cakile edentula* ssp. *californica* and *C. maritima*. *Rhodora*, 72(791), 370-386.

590 Bock, C.H. (2008) The effect of *Alternaria brassicicola* infection on the reproductive fitness of the
 591 naturally occurring littoral ruderals *Cakile maritima* and *C. edentula*. *Australian Plant Pathology*
 592 37(6), 569-580. <https://doi.org/10.1071/AP08057>
 593 Bock, D. G., Caseys, C., Cousens, R. D., Hahn, M. A., Heredia, S. M., Hübner, S., ... Rieseberg, L.
 594 H. (2015). What we still don't know about invasion genetics. *Molecular Ecology*, 24(9), 2277-2297.
 595 <https://doi.org/10.1111/mec.13032>
 596 Brandvain, Y., Kenney, A. M., Flagel, L., Coop, G., & Sweigart, A. L. (2014). Speciation and
 597 Introgression between *Mimulus nasutus* and *Mimulus guttatus*. *PLoS Genet* 10(6) : e1004410.
 598 <https://doi.org/10.1371/journal.pgen.1004410>
 599 Catchen, J. M., Amores, A., Hohenlohe, P., Cresko, W., & Postlethwait, J. H. (2011). Stacks:
 600 building and genotyping loci de novo from short-read sequences. *G3: Genes, Genomes, Genetics*,
 601 1(3), 171-182. <https://doi.org/10.1534/g3.111.000240>
 602 Boyd, R. S., & Barbour, M. G. (1993). Replacement of *Cakile edentula* by *C. maritima* in the strand
 603 habitat of California. *American Midland Naturalist*, 209-228. <https://doi.org/10.2307/2426122>
 604 Clausen, G., Vickers, K., & Kadereit, J. W. (2000). Historical biogeography in a linear system:
 605 genetic variation of Sea Rocket (*Cakile maritima*) and Sea Holly (*Eryngium maritimum*) along
 606 European coasts. *Molecular Ecology*, 9(11), 1823-1833. <https://doi.org/10.1046/j.1365-294x.2000.01083.x>
 607
 608 Cody, T. W., & Cody, M. L. (2004). Morphology and spatial distribution of alien sea-rockets (*Cakile*
 609 *spp.*) on South Australian and Western Canadian beaches. *Australian Journal of Botany*, 52(2), 175-
 610 183.
 611 Cousens, R. D., Ades, P. K., Mesgaran, M. B., & Ohadi, S. (2013). Reassessment of the invasion
 612 history of two species of *Cakile* (Brassicaceae) in Australia. *Cunninghamia*, 13, 275-290.
 613 Cousens, R. D., & Cousens, J. M. (2011). Invasion of the New Zealand coastline by European sea-
 614 rocket (*Cakile maritima*) and American sea-rocket (*Cakile edentula*). *Invasive Plant Science and*
 615 *Management*, 4(2), 260-263.
 616 Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., ... McVean, G.
 617 (2011). The variant call format and VCFtools. *Bioinformatics*, 27(15), 2156-2158.
 618 <https://doi.org/10.1093/bioinformatics/btr330>
 619 Dlugosch, K. M., & Parker, I. M. (2008). Founding events in species invasions: genetic variation,
 620 adaptive evolution, and the role of multiple introductions. *Molecular Ecology*, 17(1), 431-
 621 449. doi:10.1111/j.1365-294X.2007.03538.x

622 Ellstrand, N. C., & Schierenbeck, K. A. (2000). Hybridization as a stimulus for the evolution of
623 invasiveness in plants?. *Proceedings of the National Academy of Sciences*, 97(13), 7043-7050.
624 <https://doi.org/10.1073/pnas.97.13.7043>

625 Fitzpatrick, B. M. (2012). Estimating ancestry and heterozygosity of hybrids using molecular
626 markers. *BMC Evolutionary Biology*, 12(1), 131. <https://doi.org/10.1186/1471-2148-12-131>

627 Fox, J. & Weisberg S. (2019). An R companion to applied regression, third Edition. Thousand Oaks
628 CA: Sage. URL: <https://socialsciences.mcmaster.ca/jfox/Books/Companion/>

629 Francis, R. M. (2017). pophelper: an R package and web app to analyse and visualize population
630 structure. *Molecular Ecology Resources*, 17(1), 27-32. <https://doi.org/10.1111/1755-0998.12509>

631 Gandour, M., Hessini, K., & Abdelly, C. (2008). Understanding the population genetic structure of
632 coastal species (*Cakile maritima*): seed dispersal and the role of sea currents in determining
633 population structure. *Genetics Research*, 90(2), 167-178.
634 <https://doi.org/10.1017/S0016672308009269>

635 Gormally, C. L., Hamrick, J.L., & Donovan, J. H. A. (2011). Genetic structure of a widely dispersed
636 beach annual, *Cakile edentula* (Brassicaceae). *American Journal of Botany*, 98(10), 1657-1662.
637 <https://doi.org/10.3732/ajb.1000499>

638 Harris, K., & Nielsen, R. (2016). The genetic cost of Neanderthal introgression. *Genetics*, 203(2),
639 881-891. <https://doi.org/10.1534/genetics.116.186890>

640 Harrison, B. J., & Darby, L. A. (1955). Unilateral hybridization. *Nature*, 176(4490), 982-982.
641 <https://doi.org/10.1038/176982a0>

642 Heyligers, P. C. (1984). Beach invaders: sea rockets and beach daisies thrive. *Australian Natural*
643 *History*, 21(5), 212-214

644 Hodgins, K. A., Bock, D. G., & Rieseberg, L. H. (2018). Trait evolution in invasive species. *Annual*
645 *Plant Reviews Online*, 459-496. <https://doi.org/10.1002/9781119312994.apr0643>

646 Hoffmann, B. D., & Broadhurst, L. M. (2016). The economic cost of managing invasive species in
647 Australia. *NeoBiota*, 31, 1. <https://doi.org/10.3897/neobiota.31.6960>

648 Hovick, S. M., & Whitney, K. D. (2014). Hybridisation is associated with increased fecundity and
649 size in invasive taxa: meta-analytic support for the hybridisation-invasion hypothesis. *Ecology*
650 *Letters*, 17(11), 1464-1477. <https://doi.org/10.1111/ele.12355>

651 Huebner, D. C. (2009). A morphological, molecular and experimental assessment of the conservation
652 status of great lakes sea-rocket (*Cakile edentula* var. *lacustris*, Brassicaceae) (Unpublished doctoral
653 dissertation). Northwestern University.

654 Huson, D. H., & Bryant, D. (2006). Application of phylogenetic networks in evolutionary studies.
 655 *Molecular Biology and Evolution*, 23(2), 254-267. <https://doi.org/10.1093/molbev/msj030>
 656 Joshi, N. A. and Fass, J. N. (2011). Sickel: a Sliding-Window, Adaptive, Quality-Based Trimming
 657 Tool for FastQ Files (Version 1.33) [Software]. Retrieved from <https://github.com/najoshi/sickle>
 658 Juric, I., Aeschbacher, S., & Coop, G. (2016). The strength of selection against Neanderthal
 659 introgression. *PLoS Genetics*, 12(11), e1006340. <https://doi.org/10.1371/journal.pgen.1006340>
 660 Kadereit, J. W., Arafah, R., Somogyi, G., & Westberg, E. (2005). Terrestrial growth and marine
 661 dispersal? Comparative phylogeography of five coastal plant species at a European scale. *Taxon*,
 662 54(4), 861-876. <https://doi.org/10.2307/25065567>
 663 Kassambara, A. (2020). Ggpubr: ggplot2 Based Publication Ready Plots. R package version 0.2.5 [R
 664 package]. A Retrieved from <https://cran.r-project.org/package=ggpubr>
 665 Keenan, K., McGinnity, P., Cross, T. F., Crozier, W. W., & Prodöhl, P. A. (2013). DiveRsity: An R
 666 package for the estimation of population genetics parameters and their associated errors. *Methods*
 667 *Ecol Evolution*, 4(8), 782-788. doi:10.1111/2041-210X.12067
 668 Lapointe, M. (2000). Modern diatom assemblages in surface sediments from the Maritime Estuary
 669 and the Gulf of St. Lawrence, Québec (Canada). *Marine Micropaleontology*, 40(1-2), 43-65.
 670 [https://doi.org/10.1016/S0377-8398\(00\)00031-1](https://doi.org/10.1016/S0377-8398(00)00031-1)
 671 Li, C., Cousens, R. D., & Mesgaran, M. B. (2019). How can natural hybridisation between
 672 self-compatible and self-incompatible species be bidirectional?. *Weed Research*, 59(5), 339-348.
 673 <https://doi.org/10.1111/wre.12372>
 674 Li, H. & Durbin, R. (2009). Fast and accurate short read alignment with Burrows- Wheeler
 675 transform. *Bioinformatics* 25, 1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>
 676 Li, C., Mesgaran, M. B., Ades, P. K., & Cousens, R. D. (2020). Inheritance of breeding system in
 677 *Cakile* (Brassicaceae) following hybridization: implications for plant invasions. *Annals of Botany*,
 678 125(4), 639-650. <https://doi.org/10.1093/aob/mcz198>
 679 Maddison, W. P. and Maddison, D.R. (2019). Mesquite: A Modular System for Evolutionary
 680 Analysis [Software]. Retrieved from <http://www.mesquiteproject.org>
 681 Marhold, K. (2011). Brassicaceae. In: Euro+Med Plantbase- the Information Resource for Euro-
 682 Mediterranean Plant Diversity. Retrieved from
 683 <http://ww2.bgbm.org/EuroPlusMed/PTaxonDetail.asp?NameId=14225&PTRefFk=7200000>
 684 Mesgaran, M. B., Lewis, M. A., Ades, P. K., Donohue, K., Ohadi, S., Li, C., & Cousens, R. D.
 685 (2016). Hybridization can facilitate species invasions, even without enhancing local adaptation.

686 *Proceedings of the National Academy of Sciences*, 113(36), 10210-10214.
687 <https://doi.org/10.1073/pnas.1605626113>
688 Ohadi, S., Ades, P. K., Ford, R., Strand, A. E., Tibbits, J., Mesgaran, M. B., & Cousens, R. D.
689 (2016). Genetic structure along the strandline: Unravelling invasion history in a one-dimensional
690 system. *Journal of Biogeography*, 43(3), 451-460. <https://doi.org/10.1111/jbi.12640>
691 Payseur, B. A., & Rieseberg, L. H. (2016). A genomic perspective on hybridization and speciation.
692 *Molecular Ecology*, 25(11), 2337-2360. <https://doi.org/10.1111/mec.13557>
693 Pickrell, J.K., & Pritchard, J.K. (2012). Inference of population splits and mixtures from genome-
694 wide allele frequency data. *PLoS Genetics*, In Press.
695 Pickup, M., Brandvain, Y., Fraïsse, C., Yakimowski, S., Barton, N. H., Dixit, T., ... Field, D. L.
696 (2019). Mating system variation in hybrid zones: facilitation, barriers and asymmetries to gene flow.
697 *New Phytologist*, 224(3), 1035-1047. <https://doi.org/10.1111/nph.16180>
698 Pimentel, D., Zuniga, R., & Morrison, D. (2005). Update on the environmental and economic costs
699 associated with alien-invasive species in the United States. *Ecological Economics*, 52(3), 273-288.
700 <https://doi.org/10.1016/j.ecolecon.2004.10.002>
701 Pollak, E. (1987). On the theory of partially inbreeding finite populations. I. Partial selfing. *Genetics*,
702 117(2), 353-360.
703 Poplin, R., Ruano-Rubio, V., DePristo, M. A., Fennell, T. J., Carneiro, M. O., Van der Auwera, G.
704 A., ... Shakir, K. (2017). Scaling accurate genetic variant discovery to tens of thousands of samples.
705 *BioRxiv*, 201178. <https://doi.org/10.1101/201178>
706 Reich, D., Thangaraj, K., Patterson, N., Price, A. L., & Singh, L. (2009). Reconstructing Indian
707 population history. *Nature*, 461(7263), 489-494. <https://doi.org/10.1038/nature08365>
708 Rodman, J. E. (1974). Systematics and evolution of the genus *Cakile* (Cruciferae). *Contributions*
709 *from the Gray Herbarium of Harvard University*, (205), 3-146.
710 Rodman, J. E. (1976). Differentiation and migration of *Cakile* (Cruciferae): seed glucosinolate
711 evidence. *Systematic Botany*, 137-148.
712 Rodman, J. E. (1986). Introduction, establishment and replacement of sea-rockets (*Cakile*,
713 Cruciferae) in Australia. *Journal of Biogeography*, 159-171. <https://doi.org/10.2307/2844990>
714 R Core Team (2018). R: A language and environment for statistical computing. R Foundation for
715 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
716 Ruiz, G. M., Rawlings, T. K., Dobbs, F. C., Drake, L. A., Mullady, T., Huq, A., & Colwell, R. R.
717 (2000). Global spread of microorganisms by ships. *Nature*, 408(6808), 49-50.
718 <https://doi.org/10.1038/35040695>

719 Ruhsam, M., Hollingsworth, P. M., & Ennos, R. A. (2011). Early evolution in a hybrid swarm
 720 between outcrossing and selfing lineages in *Geum*. *Heredity*, 107(3), 246-255.
 721 <https://doi.org/10.1038/hdy.2011.9>
 722 Sakai, A. K., Allendorf, F. W., Holt, J. S., Lodge, D. M., Molofsky, J., With, K. A., ... McCauley, D.
 723 E. (2001). The population biology of invasive species. *Annual review of Ecology and Systematics*,
 724 32(1), 305-332. <https://doi.org/10.1146/annurev.ecolsys.32.081501.114037>
 725 Sax, D. F., & Gaines, S. D. (2003). Species diversity: from global decreases to local increases.
 726 *Trends in Ecology & Evolution*, 18(11), 561-566.
 727 Simberloff, D. (2013). Biological invasions: Prospects for slowing a major global change. *Elementa*
 728 *Science of the Anthropocene*, 1, p.000008. doi: <http://doi.org/10.12952/journal.elementa.000008>
 729 Sedlazeck, F. J., Rescheneder, P., & von Haeseler, A. (2013). NextGenMap: fast and accurate read
 730 mapping in highly polymorphic genomes. *Bioinformatics*, 29(21), 2790-2791.
 731 <https://doi.org/10.1093/bioinformatics/btt468>
 732 Thrall, P. H., Young, A. G., & Burdon, J. J. (2000). An analysis of mating structure in populations of
 733 the annual sea rocket, *Cakile maritima* (Brassicaceae). *Australian Journal of Botany*, 48(6), 731-738.
 734 <https://doi.org/10.1071/BT99060>
 735 Todesco, M., Pascual, M. A., Owens, G. L., Ostevik, K. L., Moyers, B. T., Hübner, S., ... Rieseberg,
 736 L. H. (2016). Hybridization and extinction. *Evolutionary Applications*, 9(7), 892-908.
 737 <https://doi.org/10.1111/eva.12367>
 738 Uesugi, A., Baker, D. J., de Silva, N., Nurkowski, K., & Hodgins, K. A. (2020). A lack of genetically
 739 compatible mates constrains the spread of an invasive weed. *New Phytologist*, 226(6), 1864-1872.
 740 <https://doi.org/10.1111/nph.16496>
 741 Vallejo-Marín, M., Friedman, J., Twyford, A. D., Lepais, O., Ickert-Bond, S. M., Streisfeld, M. A., ...
 742 & Puzey, J. R. (2020). Population genomic and historical analysis reveals a global invasion by
 743 bridgehead processes in *Mimulus guttatus*. *bioRxiv*. <https://doi.org/10.1101/2020.06.26.173286>
 744 Vallejo-Marín, M., & Hiscock, S. J. (2016). Hybridization and hybrid speciation under global
 745 change. *New Phytologist*, 211(4), 1170-1187. <https://doi.org/10.1111/nph.14004>
 746 van Boheemen, L. A., Lombaert, E., Nurkowski, K. A., Gauffre, B., Rieseberg, L. H., & Hodgins, K.
 747 A. (2017). Multiple introductions, admixture and bridgehead invasion characterize the introduction
 748 history of *Ambrosia artemisiifolia* in Europe and Australia. *Molecular Ecology*, 26(20), 5421-5434.
 749 <https://doi.org/10.1111/mec.14293>

- Vilatersana, R., Sanz, M., Galian, A., & Castells, E. (2016). The invasion of *Senecio pterophorus* across continents: multiple, independent introductions, admixture and hybridization. *Biological invasions*, 18(7), 2045-2065. <https://doi.org/10.1007/s10530-016-1150-1>
- Ward, S. M., Gaskin, J. F., & Wilson, L. M. (2008). Ecological genetics of plant invasion: what do we know?. *Invasive Plant Science and Management*, 1(1), 98-109. <https://doi.org/10.1614/IPSM-07-022.1>
- Weir, B. S., & Cockerham, C. C. (1984). Estimating F-statistics for the analysis of population structure. *Evolution*, 38, 1358-1370. <https://doi.org/10.2307/2408641>
- Westberg, E. D. (2005). European phylogeography of the coastal plants *Cakile maritima* Scop. (Brassicaceae) and *Eryngium maritimum* L. (Apiaceae) (Unpublished doctoral dissertation). Johannes Gutenberg-Universität.
- Whitlock, R., Hipperson, H., Mannarelli, M., & Burke, T. (2008). A high-throughput protocol for extracting high-purity genomic DNA from plants and animals. *Molecular Ecology Resources*, 8(4), 736-741. <https://doi.org/10.1111/j.1755-0998.2007.02074.x>
- Whitney, K. D., Randell, R. A., & Rieseberg, L. H. (2006). Adaptive introgression of herbivore resistance traits in the weedy sunflower *Helianthus annuus*. *The American Naturalist*, 167(6), 794-807. <https://doi.org/10.1086/504606>
- Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., ... Kuhn, M. (2019). Welcome to the Tidyverse. *Journal of Open Source Software*, 4(43), 1686. <https://doi.org/10.21105/joss.01686>
- Zheng, X., Levine, D., Shen, J., Gogarten, S. M., Laurie, C., & Weir, B. S. (2012). A high-performance computing toolset for relatedness and principal component analysis of SNP data. *Bioinformatics*, 28(24), 3326-3328. <https://doi.org/10.1093/bioinformatics/bts606>

Figures and figure captions

Figure 1 Admixture results of the *unsupervised run* of the *global thinned dataset*. (A) A distruct plot for K=8. Individuals are ordered according to their cluster association of the *supervised run*. AUS=Australia, eNA= eastern North America, EU= Europe and northern Africa, NZ= New Zealand, wNA=western North America. E= *C. edentula*, M= *C. maritima*, H= Hybrids. (B) Population pie charts for K=8, Admixture proportions for each population are displayed. A global map is displayed as well as close ups of western North America, Europe, the Australian mainland and Tasmania. Colours correspond to the clusters in the distruct plot. Arrows indicates direction of invasion and direction of Spearman test.

792

793 Figure 3 Geographic distribution of the hybrid assignment test by NewHybrid. Individuals are
 794 coloured according to their NewHybrid classification. A global map and close-ups of western North
 795 America, the Australian mainland and Tasmania are presented. BC-E= backcross to *C. edentula*, BC-
 796 M= backcross to *C. maritima*.

797

(A)

(B)

Figure 4 Results of a hybridization assignment test implemented by Hlest using 471 SNPs (0.99, 0.03). (A) Association of ancestry index (S) and interclass heterozygosity (H) are given for western North America (left) and Australia (right). Individuals are coloured according to their Hlest classification. For hybrids the continuous model was a better fit than the hybrid classes. (B) The geographic distribution of individuals and their S index; yellow= *C. edentula* proportion, blue= *C. maritima* proportion. A global map and close-ups of western North America, the Australian mainland and Tasmania are presented. Arrows indicates direction of invasion and direction of Spearman's test.

Figure 5 Results of the Spearman correlation test displayed (Table 2). The associations between population mean Q values of hybrids identified using the *supervised* Admixture *run* and the ranked order of populations from the first entry point of *C. maritima* (A) in eastern Australia and (B) western North America.

Figure 6 Maximum likelihood trees with two migration events generated by TreeMix. Native ranges and (A) western North America, (B) Australia. Individuals are grouped by species (identified morphologically), likely subspecies and geographic origin.

819 Table 1 Number of individuals and sampling locations as well as mean number of individuals sampled per sampling location in each range is
820 presented.

821

Range	Phenotype	Number of individuals	Number of sampling locations	Mean number of individuals sampled per sampling location
Eastern North America	<i>C. edentula</i>	55	26	2.03
	<i>C. lanceolata</i>	2	2	1
Europe and northern Africa	<i>C. maritima</i> subsp. <i>integrifolia</i> and <i>baltica</i>	12	12	1
	<i>C. maritima</i> subsp. <i>maritima</i>	12	12	1
	<i>C. maritima</i> subsp. <i>islandica</i>	1	1	1
Western North America	<i>C. edentula</i>	39	4	4
	<i>C. maritima</i>	79	10	5.9
	Hybrids	2	1 (in mixed)	/

	Unknown	1	0 (in <i>C. edentula</i>)	/
	Mixed populations		3	15.6
	Total	120	17	7.05
New Zealand	Unknown	1	1	1
Australia	<i>C. edentula</i>	43	3	7.33
	<i>C. maritima</i>	110	11	8
	Hybrids	14	5 (in mixed)	/
	Mixed population		7	8.4
	Total	167	21	7.95

822

823

824 Table 2 Results of the Spearman's rank correlation test in the introduced ranges examining the association between species ancestry for *C.*
825 *edentula*, *C. maritima* and hybrids or hybrids and the rank order of sampling locations based on the distance along the coastline from the
826 first recorded case of *C. maritima* in western North America (San Francisco) or south-east mainland of Australia (Adelaide). Spearman's
827 Rank Correlation Coefficient ρ and p values are presented for correlation between Q-value of the *supervised run* of the *C. edentula* cluster

828 for each population in western North America and Australia and correlation between ancestry index (S) (Figure 4) and rank order of
829 sampling locations.

830

			Q		S		
Range	Species	# populations (# individuals)	ρ	p	# populations (# individuals)	ρ	p
south-east Australia	<i>C. edentula</i> , <i>C. maritima</i> , hybrids	10 (65)	0.815	0.004	10 (65)	0.815	0.004
	Hybrids	7 (30)	0.893	0.012	8 (38)	0.905	0.005
western North America	<i>C. edentula</i> , <i>C. maritima</i> , hybrids	10 (68)	0.511	0.132	10 (68)	0.576	0.088
all sampling locations	Hybrids	5 (11)	0.300	0.683	10 (50)	0.467	0.213
western North America	<i>C. edentula</i> , <i>C. maritima</i> , hybrids	8 (47)	0.719	0.045	8 (47)	0.810	0.022
north of San Francisco	Hybrids	5 (11)	0.300	0.683	7 (30)	0.679	0.110

831

832 Table 3 Results of the f_3 statistic using TreeMix. Tests of admixture in the invasive range of Australia and western North America were done
833 separately and both were based on three groups (hybrids, *C. edentula*, *C. maritima*). Hybrid classification was done according to the
834 supervised run of Admixture. The f_3 statistic, the standard error of f_3 and the Z-score are reported.

835

Range	Target	Source 1	Source 2	f_3	Standard error	Z-score
-------	--------	----------	----------	-------	----------------	---------

					of f_3	
Australia	Australian hybrids	Australian <i>C. edentula</i>	Australian <i>C. maritima</i>	-0.0058	0.0002	-31.9723
w. North America	w. North American hybrids	w. North America <i>C. edentula</i>	w. North American <i>C. maritima</i>	-0.0049	0.0002	-23.2228

(A)

(B)

(C)

- invasive wNA *C. maritima*
- ◆ native atl. *C. maritima*
- ◆ native *C. maritima*
- invasive med. *C. maritima*
- hybrid
- ◆ native *C. edentula* New Brunswick
- ◆ native *C. edentula* Quebec & Newfoundland & Nova Scotia
- invasive *C. edentula* Nova Scotia
- ◆ native *C. edentula* Lake Michigan & Rhode Island
- invasive *C. edentula* Lake Michigan & Rhode Island

This article is protected by copyright. All rights reserved

Author Manuscript

- *C. maritima*
- *C. edentula*
- BC-M
- BC-E
- F1
- F2

(A) $\rho = 0.89, p = 0.012$ mec_15768.f5.pdf

(B) $\rho = 0.3, p = 0.68$

(A)

mec_15768_f6.pdf

(B)

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Rosinger, HS;Geraldes, AM;Nurkowski, KA;Battlay, P;Cousens, RD;Rieseberg, LH;Hodgins, KA

Title:

The tip of the iceberg: Genome wide marker analysis reveals hidden hybridization during invasion

Date:

2021-01-23

Citation:

Rosinger, H. S., Geraldes, A. M., Nurkowski, K. A., Battlay, P., Cousens, R. D., Rieseberg, L. H. & Hodgins, K. A. (2021). The tip of the iceberg: Genome wide marker analysis reveals hidden hybridization during invasion. MOLECULAR ECOLOGY, 30 (3), pp.810-825. <https://doi.org/10.1111/mec.15768>.

Persistent Link:

<http://hdl.handle.net/11343/298149>

License:

[CC BY-NC-ND](#)