Hematopoietic stem cells, progenitor cells and leukemic stem cells in adult myeloproliferative neoplasms

Ashley P. Ng^{1,2,3}

¹The Walter and Eliza Hall Institute of Medical Research, Parkville, Victoria 3052, Australia

²The Department of Medical Biology, University of Melbourne, Parkville, Victoria, 3010, Australia

³The Royal Melbourne Hospital, Parkville, Victoria, 3052, Australia

Correspondence:

Ashley P. Ng

Walter and Eliza Hall Institute of Medical Research

1G Royal Parade, Parkville, Victoria, 3052, Australia

Email : ang@wehi.edu.au

Running Title : HSCs, Progenitors and LSCs in MPNs

Keywords : Myeloproliferative Neoplasms, Stem and Primitive Progenitor Cells, Signal transduction

Abstract

The understanding of myeloproliferative neoplasms has changed dramatically since Dameshek proposed his classification over fifty years ago. Our knowledge of the types of cells which comprise the hematopoietic system and of how they are regulated has also appreciated significantly over this time. This review will relate what is currently known about the acquired genetic mutations associated with adult myeloproliferative neoplasms to how they lead to the hematopoietic perturbations of myeloproliferative disease. There will be a particular focus on how stem and progenitor cell compartments are affected by *BCR-ABL1* and *JAK2V617F* mutations, and the particular issue of resistance of leukemic stem cells to conventional and targeted therapies.

Introduction

Myeloproliferative neoplasms are a class of hematopoietic neoplasms characterised by an overproduction of cells of mature myeloid lineages which may lead to complications of rheostasis, thrombosis and hemostasis, splenomegaly, bone marrow failure and leukemic transformation in affected patients (DAMESHEK, 1951).

The Philadelphia chromosome with the *BCR-ABL1* fusion gene arising from the 9;22 chromosomal translocation in chronic myeloid leukaemia (CML) was the first example of an acquired causative mutation that could be associated with a myeloproliferative disorder (Chandra et al., 2011). Several gene mutations were subsequently associated with Philadelphia/*BCR-ABL1* negative (Ph-) myeloproliferative neoplasms (reviewed in (Tefferi, 2010) (Vainchenker et al., 2011)). Most notable of those is the valine to phenylalanine substitution in the *JAK2* gene (*JAK2V617F*) that occurs in ~95% of polycythemia vera (PV), ~ 60% of essential thrombocythemia (ET) and ~50% of idiopathic myelofibrosis (IMF) (James et al., 2005) (Levine et al., 2005) (Delhommeau et al., 2006) (Barosi et al., 2012). These acquired genetic mutations have been incorporated into the diagnostic criteria of myeloproliferative neoplasms as proposed by the World Health Organisation (Swerdlow, 2008), integrating molecular and cytogenetic findings with traditional morphologic pathology in recognition of the importance of specific mutations which lead to constitutively activated kinases that drive disease phenotypes in most myeloproliferative neoplasms (Delhommeau et al., 2006) (Gupta et al., 2002) (Vardiman and Hyjek, 2011) (Anastasi, 2011).

The realisation that *BCR-ABL1* and *JAK2V617F* mutations lead to specific tyrosine kinase activation resulted in the development of targeted therapies directed at kinase inhibition

(O'Brien et al., 2003) (Cools et al., 2003) (Tefferi, 2012). Kinases, such as ABL1 or JAK2, are the targets of choice for inhibition by small molecules, and were therefore attractive targets for pharmaceutical intervention with kinase inhibitors (Arora and Scholar, 2005). In the case of CML, this has resulted in the *BCR-ABL1* kinase inhibitor, imatinib, becoming the standard of therapy for treatment of CML especially for those patients diagnosed in chronic phase (Hochhaus et al., 2009). This can be attributed to the reduction in the incidence of CML blast crisis, improvement in overall survival and the tolerability of imatinib in the vast majority of patients with CML (Hehlmann, 2012). The superiority of imatinib (O'Brien et al., 2003), and more recently, second generation tyrosine kinase inhibitors (TKIs) (reviewed (Cortes and Kantarjian, 2012)) over previous therapy for CML is evidenced and related to their dramatic ability to reduce the molecularly detectable *BCR-ABL1* positive disease (Cross et al., 2012). JAK2 kinase inhibitors have also entered into clinical practice for patients with IMF. In recent seminal phase III trials, JAK1/2 inhibition was very successful at decreasing splenomegaly and disease related constitutional symptoms with significant improvement in quality of life scores (Verstovsek et al., 2012) (Harrison et al., 2012), but did not show survival benefits compared to best available therapy (Harrison et al., 2012). Inhibition of JAK2 as a therapeutic target is also unlikely have the same magnitude of benefit in reducing disease burden as has been seen for BCR-ABL1 inhibition in CML, with dose limiting toxicity of endogenous JAK2 inhibition apparent in hemopoietic cells as well as off-target effects being noted (Tefferi, 2012).

The success of BCR-ABL1 kinase inhibitors in a significant majority of patients with CML raised the initial hope that these TKIs may provide potentially definitive curative therapy with complete eradication of *BCR-ABL1* disease. Despite these advances it has been shown

that TKIs may not completely eradicate the disease clone in CML, even in patients who obtain an apparently complete molecular response (Branford et al., 2007) (Ross et al., 2010). This observation can be explained in significant part by the unique physiological characteristics of hematopoietic stem cells harboring the *BCR-ABL1* mutation. Similarly, current approaches to treatment of Ph- myeloproliferative neoplasms, although successful at preventing and managing several complications of PV, ET and IMF, do not for the most part eradicate the underlying disease clone. Allogeneic transplantation is current the only strategy whereby this may occur for CML and Ph- myeloproliferative neoplasms. Allogeneic transplantation however, carries significant cost in terms of non-relapse and relapse related morbidity and mortality for both CML (Goldman et al., 2010) and the select group of patients with Ph- myeloproliferative neoplasms who have transformed to acute leukaemia or have IMF and who are suitable for the procedure and have appropriately matched donors (Barosi et al., 2012) (Gupta et al., 2012).

This review will focus on hematopoietic stem cell and progenitor cell biology and relate what is currently known about acquired genetic mutations associated with adult myeloproliferative neoplasms to how they perturb hematopoiesis to generate myeloproliferative phenotypes. There will be a particular focus on how stem and progenitor cell compartments are affected by *BCR-ABL1* and *JAK2V617F* mutations and the issue of resistance of leukemic stem cells to conventional and targeted therapies.

Hematopoietic Stem Cell Biology

Mammalian hematopoiesis has been well characterised. Multiple specialised cellular compartments and their important molecular regulators have been identified. Hematopoietic stem cells (HSCs) are rare and specialised residents of the bone marrow and underpin the development and maintenance of hematopoiesis throughout the lifetime of an organism. Located in highly specialised and relatively hypoxic bone marrow micro-environments (Parmar et al., 2007) (reviewed in (Adams and Scadden, 2006)) adjacent to N-cadherin⁺ osteoblasts (Zhang et al., 2003) in endochondral ossified endosteal bone niches (Chan et al., 2009), HSCs are maintained in a relatively quiescent state where they retain the characteristics of *pluripotency*, the "potential" of a single cell to generate all the cell lineages of the hematopoietic system, and *self-renewal*, the ability to divide "assymmetrically" to sustain a pool of daughter stem cells which possess the pluripotency and self-renewal abilities of the parent HSC (Tajbakhsh et al., 2009) (Ting et al., 2012) (Figure 1).

HSC quiescence and self-renewal are maintained by several molecular regulators acting via specific cell signaling pathways (reviewed in (Blank et al., 2008)). Thrombopoietin/Mpl receptor (Qian et al., 2007) (Yoshihara et al., 2007), Wnt/ β -catenin (Fleming et al., 2008), Notch1/Jagged1 (Stier et al., 2002) (Calvi et al., 2003), osteopontin (Nilsson et al., 2005), TGF- β /Smad4 (Karlsson et al., 2007), Tie2/Angiopoietin1 (Arai et al., 2004) and stem cell factor/cKit signaling (Kent et al., 2008), have all been demonstrated to be important in regulating HSC quiescence/self-renewal.

Importantly, quiescent HSCs have been shown to be resistant to conventional chemotherapeutic agents (Hodgson and Bradley, 1979) and γ radiation (Ploemacher et al., 1992). The resistance of quiescent HSCs to mechanisms which can trigger cell death by apoptosis is an im-

portant defining characteristic of this population and is mediated by specific molecular regulators, such as the prosurvival Bcl-2 family protein, Mcl-1 (Opferman et al., 2005).

Notably, HSCs may be stimulated into cell cycle by other regulators. Inflammatory mediators such as interferon alpha (Essers et al., 2009) and interferon gamma (Baldrige et al., 2010) may induce "quiescent" HSCs into cell cycle, whilst Prostaglandin E2 generated by the cyclooxygenase (COX) enzyme in HSCs, may modulate HSC self-renewal acting via EP2 and EP4 receptors (North et al., 2007) (Frisch et al., 2009) (Durand and Zon, 2010).

Hematopoietic progenitor cells are the progeny of HSCs. Progenitors comprise a significant proportion of adult bone marrow, coordinate the production of precise numbers of mature blood cells of diverse functional lineages, and expand in times of hematopoietic stress to attempt to meet physiologic requirements. The steady state production of mature functional cell lineages and the hematopoietic stress response is mediated in large part by hematopoietic cytokines, such as erythropoietin (EPO) for erythroid (Miyake et al., 1977) (Gasson et al., 1985), granulocyte colony stimulating factor (G-CSF) for granulocyte (Nicola et al., 1983) (Metcalf and Nicola, 1983), and thrombopoietin (TPO) for the megakaryocyte lineage and platelet production (Kaushansky et al., 1994) (Lok et al., 1994) (Ng et al., 2012a).

Hematopoietic cytokines, cell signaling molecules and mutations of epigenetic regulators associated with myeloproliferative neoplasms

The discovery of hematopoietic cytokines, their cognate receptors and the characterisation of the molecules which comprise the intracellular signaling pathways has led directly to significant therapeutic advances in supportive hematological care and peripheral stem cell mobilisation (reviewed in (Metcalf, 2010) (Robb, 2007)).

The use of genetically manipulated murine models has allowed the precise characterisation of the roles of specific signaling molecules in physiological hematopoiesis. For instance, during EPO and EPO receptor signaling (reviewed in (Constantinescu et al., 1999), Figure 2), JAK2, a member of the Janus (JAK) family of non-receptor protein tyrosine kinases (Wilks et al., 1991), was shown to be an essential down-stream mediator that was required for definitive fetal erythropoiesis (Parganas et al., 1998) (Neubauer et al., 1998). STAT5, which is recruited to the EPO receptor and is phosphorylated by JAK2 to induce its activation, nuclear translocation and transcription (Constantinescu et al., 1999), was shown to be important during fetal erythropoiesis through analysis of mice lacking both STAT5a and STAT5b redundant isoforms (Teglund et al., 1998). The role of STAT5 in fetal erythropoiesis was attributed in significant part, to its transcription of the pro-survival BCL-X_L protein and inhibition of apoptosis of fetal erythrocytes (Socolovsky et al., 1999).

Murine models have also permitted dissection of intracellular signaling molecules required for manifestations of disease phenotypes associated with *BCR-ABL1* and *JAK2V617* mutations (Figure 3). Both *BCR-ABL1* and *JAK2V617F* appear dependent on STAT5 signaling for features of myeloproliferation in murine models (Funakoshi-Tago et al., 2010) (Yan et al., 2012) (Walz et al., 2012). Endogenous JAK2 may serve as an important mediator of the disease phenotype associated with the *FIP1L1-PDGFRα* fusion identified in a proportion of patients with chronic eosinophilic leukaemia (Li et al., 2012a). Of particular note, although *BCR-ABL1* appears dependent on STAT5 for the development of chronic myeloid leukaemia-like myeloproliferative disease, endogenous JAK2 may be required for initiation of transformation to B- acute lymphoblastic leukaemia in these models (Hantschel et al., 2012). This last finding highlights that not only the acquired genetic lesion, but specific interactions with

endogenous signaling molecules and the cell types in which this occurs may all play an important role in the specific manifestations of myeloproliferative neoplasms.

Similarly, other mutations identified in a minority of cases of Ph- myeloproliferative neoplasms, such as gain-of-function mutations in *JAK2* exon 12 (Scott et al., 2007) and *MPL* (Pardanani et al., 2006), and loss-of-function mutations for negative regulators such as *CBL* (Aranaz et al., 2012) and *LNK* (Pardanani et al., 2010), may all directly perturb cell signaling pathways.

A subset of patients with Ph- myeloproliferative neoplasms and interestingly CML patients with advanced phases of disease (Makishima et al., 2011), have had mutations identified in epigenetic regulators such as *ASXL1*, *DNMT3A*, *IDH1*, *IDH2*, *TET2*, *SUZ12* (Martinez-Aviles et al., 2012) and *EZH2* (Guglielmelli et al., 2011), of which several have been associated with significant differences in disease outcome for Ph- myeloproliferative neoplasms. Recently, recurrent mutations in the serine/arginine-rich splicing factor 2 (*SRSF2*), relatively uncommon in patients with chronic myeloproliferative neoplasms (Yoshida et al., 2011), were identified in ~18.9% of patients with myeloproliferative neoplasms which had undergone leukemic transformation and conferred a worse overall survival in this group of transformed myeloid leukemia (tAML) by univariate analysis (Zhang et al., 2012). *DNMT3A* mutations have also been found to occur more frequently in IMF and transformed acute myeloid leukemia associated with the *JAK2V617F* mutation when compared to ET and PV cohorts (Stegelmann et al., 2011).

Although regulators such as the polycomb group proteins *SUZ12* and *EZH2* have been demonstrated to have direct roles in the regulation of HSCs and thrombopoiesis through their

involvement in the Polycomb Repressor Complex 2 (PRC2) (Majewski et al., 2008) (Majewski et al., 2010), and the *TET2* DNA methylation enzyme has been shown to modulate HSC self-renewal (Moran-Crusio et al., 2011) (Ko et al., 2011) (Shide et al., 2012), it remains unclear as to whether this occurs via regulation of intracellular cell signaling molecule expression or by other mechanisms. Taken together however, these findings suggest that mutations in epigenetic regulators may modulate the emergence of specific phenotypes associated with Ph- myeloproliferative neoplasms, including those associated with the JAK2V617F mutation (reviewed (Vainchenker et al., 2011)). Of note, ectopic expression of the histone-methyltransferase EZH2 in hematopoietic stem cells was recently shown to induce a myeloproliferative disease (Herrera-Merchan et al., 2012). Interestingly, mutant JAK2V617F has also been shown to lead to aberrant phosphorylation and reduction of the methyl transferase activity of protein arginine methyltransferase 5 (PRMT5) and subsequent H2A and H4 histone arginine methylation. PRMT5 activity was demonstrated to have direct effects on colony formation and erythroid differentiation *in vivo* (Liu et al., 2011), suggesting perturbation of this epigenetic regulator may also be a mechanism by which the JAK2V617F mutation can lead to myeloproliferation. These findings also suggest that for a subset of patients, the efficacy of therapy targeted to specific signaling molecules, such as with JAK2 inhibitors for IMF or BCR-ABL1 TKIs for patients with advanced phases of CML, may be limited, especially if other mutations are present that may drive the disease in HSCs and progenitor cells.

Expression of *JAK2V617F* in HSCs and Progenitors

Knowledge of the biology of HSC and progenitor cells would suggest that maintenance of a myeloproliferative phenotype requires acquired mutations to be harboured in cell populations with the ability to self-renew in order to allow ongoing generation of progeny containing the

disease-associated mutations. Importantly, expression of the *JAK2V617F* mutation been demonstrated in bone marrow cell populations expressing surface markers associated with HSCs in patients with PV (Jamieson et al., 2006) (Butcher et al., 2007). Interestingly however, X-chromosome inactivation analysis in females affected by PV (Adamson et al., 1976) (Briere and el-Kassar, 1998) (Mitterbauer et al., 1999) (Zamora et al., 2005) and molecular profiling of *JAK2V617F* mutations in peripheral blood leukocyte populations, noted this mutation was detected in lymphoid cells only in a minority of patients (Ishii et al., 2006) (Butcher et al., 2007). One interpretation of these data is that the *JAK2V617F* mutation does not lead to a proliferative advantage to lymphocytes. Indeed, global gene expression profiling of HSC and progenitor cells of patients carrying *JAK2V617F* demonstrated a relative reduction in genes associated with B- and T- cells (Berkofsky-Fessler et al., 2010). This suggests that rather than conferring an abnormal self-renewal ability on early myeloid progenitor populations, the *JAK2V617F* mutation may lead to a passive selectivity against lymphoid lineage formation from HSCs resulting from no lymphoid acquired growth advantage, with the circulating lymphocytes in the majority of patients with PV representing pre-existing lymphoid cells prior to acquisition of the mutation.

Supporting this contention are murine models of PV in which HSC and progenitor populations expressing *JAK2V617F* were transplanted into lethally irradiated recipients. These experiments demonstrated that the HSC population, but not the progenitor cell population, was capable of inducing and maintaining a PV-like disease in transplanted recipients (Mullally et al., 2010) (Mullally et al., 2012). These findings argue against *JAK2V617F* having a fully transformative effect on progenitor cells unlike some other leukaemia oncogenes (Zuber et al., 2009). Rather, it appears progenitor cell populations with cytokine signaling hypersensi-

tivity still require derivation from HSCs with the *JAK2V617F* mutation in order for the myeloproliferative phenotype to develop (Dupont et al., 2007).

The physiologic effect of the *JAK2V617F* mutation on HSC function remains unresolved. Interpretation of studies of xenotransplantation of human stem and progenitor cells harbouring the *JAK2V617F* mutation into immunodeficient mice has been limited by the poor rates of cross-species engraftment (Ishii et al., 2007) (James et al., 2008). Analysis of one murine model in which the human *JAK2V617F* mutation is expressed at endogenous levels demonstrated that this mutation did not confer functional advantages and even impaired HSCs containing the mutation when compared to HSCs expressing the wild-type JAK2 (Li et al., 2010). This finding, however, is in contrast with other murine models in which the stem and early progenitor compartment is relatively expanded in mice carrying a *JAK2V617F* mutation expressed from the endogenous JAK2 promoter (Akada et al., 2010) (Yan et al., 2012), with HSCs with the *JAK2V617F* mutation in one model demonstrating a weak competitive advantage relative to wild-type HSCs (Mullally et al., 2012).

Both human and murine *JAK2V617F* expression can lead to erythroid expansion in several mouse models (reviewed in (Li et al., 2011)), with an increase of erythroid progenitors (Mullally et al., 2010) (Akada et al., 2010) (Akada et al., 2012) and cytokine-independent erythroid progenitor colony growth (Akada et al., 2010) (Yan et al., 2012) (Akada et al., 2012). These findings phenocopy erythroid progenitors in human disease which demonstrated enhanced sensitivity to cytokine signaling especially to EPO (Dai et al., 1991) (Dai et al., 1992) (Dupont et al., 2007), as well as EPO independent colony growth (Prchal and Axelrad, 1974). These data demonstrate a causal link between the *JAK2V617F* mutation and the PV phenotype *in vivo*. Not all murine models expressing human *JAK2V617F* however lead to

erythroid expansion. A human *JAK2V617F* transgenic model demonstrated a dose-dependent JAK2V617F to murine wild-type JAK2 ratio may underlie expression of a PV or ET phenotype, with the ET phenotype associated with a lower JAK2V617F to wild-type JAK2 ratio of < 0.4 (Tiedt et al., 2008). Interestingly, this finding was not observed in a model using murine JAK2V617F which also had lower expression of JAK2V617F when compared to wild-type JAK2 (Mullally et al., 2010). These findings suggest that factors such as the JAK2V617F to wild-type JAK2 ratio, JAK2V617F species specific differences and mouse genetic background may all play a role in the manifestation of disease phenotype in these models.

Importantly, selective expression of the *JAK2V617F* in progenitor cells expressing the EPO receptor, thus restricting *JAK2V617F* expression to predominantly erythroid progenitor cells, was shown to be sufficient for producing a PV-like phenotype which in addition to the pathological features of PV, was able to recapitulate erythroid progenitor expansion and EPO independent progenitor colony growth as had been seen in murine models in which *JAK2V617F* was expressed in HSCs (Akada et al., 2012) (Mullally et al., 2012). These findings highlight both the importance of EPO signaling as well as EPO receptor-expressing progenitor cells for the manifestation of the PV phenotype associated with the *JAK2V617F* mutation.

Expression of BCR-ABL1 in HSCs and Progenitors

Characterisation of patients with CML has demonstrated *BCR-ABL1* expression in bone marrow cell populations enriched for HSCs and progenitor cells (Jamieson et al., 2004) (Hope et al., 2003) (Warner et al., 2004). In CML patients, it was noted there was an expansion of the progenitor pool particularly in patients with accelerated phase or blast crisis, with notable

amplification of *BCR-ABL1* transcripts in myeloid progenitors in the latter phase. The granulocyte-macrophage progenitors from CML patients also appeared to demonstrate an *in vitro* self-renewal capacity not seen in controls, which was related to increased expression of beta-catenin in CML progenitor cells relative to control progenitor cells (Jamieson et al., 2004). This latter finding was not replicated in committed murine hematopoietic progenitors in a murine model of *BCR-ABL1*-transduced CML (Huntly et al., 2004). Notably however, progenitors from CML patients and murine models in which *BCR-ABL1* expression had been transduced into hematopoietic stem and progenitor cells, demonstrated growth factor-independent colony formation which was not seen in normal controls (Graham et al., 2002) (Yan et al., 2012).

Global gene expression profiling of CML CD34⁺ bone marrow cells enriched for HSC and progenitor cells has demonstrated an up-regulation of cell cycle-associated genes when compared to control populations (Graham et al., 2007) (Bruns et al., 2009) (Benito et al., 2012). These findings suggest that *BCR-ABL1* induces proliferation within the HSC and progenitor cell compartment. These changes were shown to resolve with imatinib therapy (Benito et al., 2012). This last finding may be interpreted as either imatinib inhibition of *BCR-ABL1*-induced cellular proliferation, or a restoration of the composition of the HSC/progenitor cell compartment with *BCR-ABL1* tyrosine kinase inhibition (Jamieson et al., 2004) resulting in imatinib-induced apoptosis of proliferating *BCR-ABL1*-expressing progenitor cells (Kuroda et al., 2006).

The leukemic stem cell and CML

A population of highly quiescent HSCs expressing *BCR-ABL1* can be isolated from untreated CML patients (Holyoake et al., 1999) and from CML patients treated with imatinib (Hamilton et al., 2012). These quiescent, non-proliferating CD34⁺ CML cells have been shown to be resistant to a range of pro-apoptotic stimuli including chemotherapy and tyrosine kinase inhibition with imatinib (Graham et al., 2002) (Holtz et al., 2005) (Figure 4). The quiescent *BCR-ABL1*-expressing HSCs can be regarded as leukemic stem cells (LSCs). By way of comparison, proliferating CD34⁺ progenitors in CML are sensitive to imatinib-induced apoptosis that is mediated in significant part by the BCL-2 family pro-apoptotic proteins Bim and Bad (Kuroda et al., 2006). Interestingly, a BIM deletion polymorphism was recently shown to confer intrinsic imatinib resistance and inferior responses in patients with CML treated with imatinib (Ng et al., 2012b). It has also been suggested that BCL-2 family pro-survival proteins may play a role in resistance and survival of CML LSCs, although these data remains unpublished (Crews and Jamieson, 2012).

Several signaling pathways may allow survival of the CML LSCs in spite of BCR-ABL1 tyrosine kinase inhibition, such as Hedgehog pathway signaling (Dierks et al., 2008), and Wnt/beta-catenin signaling, the latter which may in part be regulated via Prostaglandin E2 (Heidel et al., 2012). Recently, induced pluripotent stem cells derived from human CML patients by transfection with factors Oct3/4, Sox2, KLF4 and c-myc (CML-iPSCs) were shown to express BCR-ABL1 and demonstrated a decrease in STAT5 and CRKL phosphorylation upon treatment with imatinib. Notably however, phosphorylation of ERK1/2, AKT and JNK were unchanged after imatinib treatment suggesting CML-iPSCs were capable of utilizing alternative signaling pathways other than STAT5 for survival. This was in contrast to imatinib-sensitive hematopoietic cells derived from CML-iPSCs which were shown to be BCR-ABL1 and

STAT5 dependent (Kumano et al., 2012). Similar conclusions have been reached using primary CML CD34+ bone marrow cells and transgenic murine models of CML LSCs (Hamilton et al., 2012).

In summary, HSCs and progenitor cells are distinct components of the hematopoietic system, each with unique biological properties. Acquired genetic mutations associated with myeloproliferative neoplasms may affect HSCs and progenitors differently and may perturb different cellular processes. Examination of *BCR-ABL1* and *JAK2V617F* gene mutations in human and mouse studies highlight the importance of deregulation of cytokine and intracellular cell signaling pathways in specific progenitor cell populations in order to produce the myeloproliferative phenotypes associated with these mutations, and for which STAT5 has emerged as a critical player in both *BCR-ABL1*- and *JAK2V617F*- associated myeloid disease. Analysis of human samples and murine models have demonstrated that although *BCR-ABL1* and *JAK2V617F* mutations significantly perturb the hematopoietic progenitor composition, as single mutations, they do not appear to fully transform progenitor cells into LSCs that can initiate and sustain disease *in vivo*. In addition, several other mutations in signaling molecules and epigenetic regulators have been characterized in myeloproliferative neoplasms which may play a role in the emergence of specific disease phenotypes.

Persistence of the leukemic stem cell in patients with CML treated with TKIs

Studies of human CML have demonstrated the presence of chemotherapy- radiotherapy- and targeted TKI therapy-resistant quiescent HSCs which harbour the causative mutation and which can maintain the *de novo* disease. These HSCs are the LSCs in CML, the unique prop-

erties of which reflect properties of HSCs in general, namely cell cycle quiescence and resistance to a variety of apoptotic stimuli.

Collectively, experimental data suggests CML LSCs are detectable by highly sensitive PCR in patients who have apparently achieved a stable complete molecular response (CMR) (Ross et al., 2011b) to the TKIs imatinib (Ross et al., 2010) and dasatinib (Ross et al., 2011a). Two prospective clinical trials, the French intergroup STIM study (Mahon et al., 2010) and Australasian Leukemia and Lymphoma Group CML8 study (Ross et al., 2010) have enrolled patients who had achieved stable CMR with undetectable peripheral blood *BCR-ABL1* transcripts for at least 2 years while receiving imatinib, to assess the incidence and types of relapse upon cessation of the TKI. Eight of 18 patients on the CML8 trial maintained undetectable *BCR-ABL1* after a median follow up of 2 years (Ross et al., 2010), while interim 12 month analysis for the STIM study demonstrated 41% molecular relapse free survival (Mahon et al., 2010). The latest update of the STIM study showed the probability of maintaining of CMR at 24 and 36 months in the STIM study remained ~ 39% (Mahon et al., 2011). Interestingly, 5/39 patients below a defined threshold for confirmed molecular relapse demonstrated fluctuating *BCR-ABL1* transcript levels over a median follow up of 22 months, suggesting molecular detection of transcript by highly sensitive PCR may not predict imminent cytogenetic or hematologic relapse when remaining at less than 1 log rise in *BCR-ABL1* transcript level. This observation may have basis in HSC biology. Observation of hematopoietic reconstitution in murine models (Capel et al., 1990) (Drize et al., 2001) (Cao et al., 2004) and non-human primates (Laukkanen et al., 2005) have demonstrated complex recruitment and extinction of individual HSC clones when reconstitution was observed closely over time. These observations suggest the fluctuating low level *BCR-ABL1* transcript that are detected may re-

flect this phenomenon of clonal HSC recruitment and extinction. These findings also recall the concept a "functional cure" proposed for patients with CML who were assessed more than 18 months after allogeneic bone marrow transplant in whom low levels of transcript were detected, but remained in cytogenetic and hematologic remission despite not having definitive eradication of the leukemic clone (Radich et al., 2001).

Given the data related to the allogeneic transplant experience in CML and the quiescent nature of imatinib-resistant CML LSCs, close monitoring and longer term follow up will be required for patients enrolled in the CML8 and STIM trials to further elucidate the patterns of relapse over the coming years. Evidence from long term follow up of allogeneic transplantation for CML clearly demonstrates the occurrence of late CML relapse beyond five years (Radich et al., 2001) (Goldman et al., 2010). *BCR-ABL1* is also believed to be the driving mutation for CML blast transformation, such that patients may still be at risk of this event in the absence of definitive eradication of CML clone. Although the incidence of blast crisis has been markedly reduced in the imatinib era (Hehlmann, 2012), this has been in the context where therapy has been intended to be uninterrupted. Patients with molecular relapse enrolled in CML8 study have to date remained sensitive to reintroduction of imatinib (Ross et al., 2011b), although in the latest update of the STIM study, not all patients were able to return to an imatinib induced CMR. This included one patient who experienced a loss of complete cytogenetic response (Mahon et al., 2011).

Interferon in CML and Ph- myeloproliferative neoplasms

Ultimately, despite the success of BCR-ABL1 TKI treatment for the majority of patients with CML, elimination of LSCs in CML will be needed to be able to establish a definitive cure of

the disease which would permit the cessation of therapy. This is likely to require targeting signaling pathways and molecular regulators other than the BCR-ABL1 tyrosine kinase, either through inhibiting non BCR-ABL1 molecular regulators implicated in CML LSC survival, or through the use of therapeutic agents which may sensitise or synergise with imatinib to induce apoptosis in the LSC. Synergistic action of combination therapy had been suggested in a retrospective analysis of patients undergoing induction treatment with interferon alpha and imatinib (Palandri et al., 2010). In prospective studies, the use of interferon in combination with imatinib was suggested to improve the rate of molecular response when compared to imatinib alone (Simonsson et al., 2011) (Preudhomme et al., 2010). Improvements in molecular response rates however were not observed in the German CML IV study (Hehlmann et al., 2011). Importantly, these prospective studies did not show benefit in event free survival or overall survival with combination therapy over imatinib alone. Putative mechanisms by which interferon may synergise with imatinib are via interferon alpha-induced Fas-mediated apoptosis (Selleri et al., 1997), interferon-induced immune modulation (Burchert et al., 2010), or induction of LSC clones into cell cycle whereupon they may become sensitive to imatinib (Essers et al., 2009).

For Ph- myeloproliferative neoplasms, Pegylated interferon alfa-2a (PEG-IFN- α -2a) in particular, has been shown to have significant activity in two phase II trials in patients with PV, or PV and ET, when used at relatively low and well tolerated doses. Significant hematological responses and progressive molecular responses for JAK2V617 assessable disease were noted in both studies. Interestingly, a small cohort of patients in both studies were shown to remain in sustained complete molecular response with undetectable JAK2V617F even after cessation of interferon therapy (Kiladjian et al., 2008) (Quintas-Cardama et al., 2009). Activi-

ty of low dose recombinant interferon alpha has additionally been shown for IMF, with clear bone marrow responses demonstrated for patients with early primary IMF (Silver et al., 2011). Taken together, these studies suggest Ph- myeloproliferative neoplasms demonstrate particular sensitivity to IFN alpha signaling. IFN alpha was shown to have anti-proliferative effects especially on human cell lines and patient derived CD34+ progenitors harboring the JAK2V617F mutation (Pasquier et al., 2009). PEG-IFN- α -2a has also been shown to induce apoptosis of hematopoietic progenitor cells derived from normal bone marrow and JAK2V617F positive CD34+ cells from patients with PV *in vitro* (Lu et al., 2010b). In this latter study, this effect was partially inhibited by a p38 MAP kinase inhibitor suggesting interferon signaling may in part act via this signaling pathway to increase p53 transcription. Consequently, combination treatment with low dose interferon alpha with the Bcl-xL inhibitor ABT-737 (Lu et al., 2010a) or the MDM2 antagonist Nutlin-3 which promotes apoptosis by retarding p53 degradation (Lu et al., 2012), was demonstrated to have synergistic and selective activity against JAK2V617F positive progenitors compared to controls *in vitro*. Relative selectivity of interferon alpha *in vivo* against amplifying hematopoietic stem and progenitor cells with the JAK2V617F mutation has been reported in a chimeric mouse model, including activity against disease-initiating LSC (Marty et al., 2011). To build on these findings, a phase III trial of PEG-IFN- α -2a in Ph- myeloproliferative neoplasms is required.

Other potential drug targets against the leukemic stem cell in CML and JAK2V617F driven myeloproliferative neoplasms

Pre-clinical data continue to suggest that modulation of CML LSCs may be feasible with other agents, such as indomethacin which in combination with imatinib, may aid in targeting CML LSCs (Heidel et al., 2012). Similarly, pan histone deacetylase inhibitors (Zhang et al.,

2010) and more specific inhibitors of NAD-dependent histone deacetylase SIRT1 (Li et al., 2012b) in combination with imatinib, have shown promising activity in the ability to potentially target the CML LSC. Whether these approaches will provide additional benefit to patients with CML in chronic phase or the more advanced accelerated phase or blast crisis disease in which mutations in genes involved in cell signaling pathways other than *BCR-ABL1* or epigenetic regulators can be found, remain to be seen. Importantly, these additional non *BCR-ABL1* mutations may account for the relative resistance to TKIs in these advanced stages of CML (Makishima et al., 2011).

Similarly, conventional chemotherapeutic approaches to Ph- myeloproliferative neoplasms are not curative and targeted therapy with JAK2 inhibitors are unlikely to drive a paradigm shift in the treatment of these neoplasms in the way that BCR-ABL1 TKIs have done for CML. Dose-limiting toxicity of JAK2 inhibitor therapy can be directly related to inhibition of wild-type JAK2. The maximal tolerated dose of the JAK1/2 inhibitor Ruxolitinib (Verstovsek et al., 2010) and selective JAK2 inhibitor TG101348 (Pardanani et al., 2011) was defined by hematologic toxicity and could be predicted given the importance of JAK2 signaling in mediating cytokine response in normal hematopoiesis (Khwaja, 2006). At selected therapeutic doses, thrombocytopenia and anemia are frequent side effects of the JAK1/2 inhibitor Ruxolitinib (INCB018424) (Harrison et al., 2012) and the selective JAK2 inhibitor TG101348 (Pardanani et al., 2011) with no clinically meaningful reduction in allele burden for JAK2V617F positive disease (Verstovsek et al., 2010) (Pardanani et al., 2011), induction of disease remission, or survival benefits over best supportive therapies observed (Harrison et al., 2012) (Tefferi, 2012). In contrast, ABL1 kinase does not appear to be required for normal myelopoiesis (Tybulewicz et al., 1991) (Schwartzberg et al., 1991) with the majority of patients with CML

in chronic phase treated with *BCR-ABL1* inhibitors not experiencing dose-limiting hematologic toxicity at therapeutic doses capable of significant activity against *BCR-ABL1* leukemic cells (Kantarjian et al., 2002) (Jabbour et al., 2011) (Kantarjian et al., 2012). It is therefore likely conventional therapies will remain as the mainstay of treatment for the majority of patients with Ph- myeloproliferative neoplasms, with allogeneic transplantation for select and suitable patients currently the only therapy that has the potential to eliminate the LSC clone, although with significant potential cost in terms of treatment-related, graft versus host disease-related, and disease-related morbidity and mortality.

Other pharmacological agents continue to be evaluated to address this clinical need for Ph- myeloproliferative neoplasms with the JAK2V617F mutation. Inhibitors of the HSP90 chaperone which associates and stabilizes JAK2 (Marubayashi et al., 2010), have been shown to promote degradation of wild-type and mutant JAK2V617F (Bareng et al., 2007) (Wang et al., 2009) (Proia et al., 2011) and inhibit JAK/STAT signaling *in vitro* and *in vivo*. Interestingly, the HSP90 inhibitor PU-H71 may have potentially greater selectivity for JAK2V617F expressing cells (Marubayashi et al., 2010). This selectivity also been suggested for parbinostat, a histone deacetylase/HSP90 inhibitor when used in combination with a JAK2 inhibitor (Wang et al., 2009). Taken together, these preliminary findings suggest there may be strategies in the future with which to be able to selectively target JAKV617F disease, with an improved therapeutic index over current JAK2 inhibitor therapy.

Conclusion

As further insights are gained into the biological consequences of mutations associated with myeloproliferation and therapies are developed which aim to target critical molecules such as

JAK2V617F in Ph- myeloproliferative neoplasms, BCR-ABL1 in CML, and other strategies against Ph- myeloproliferative neoplasms and the LSCs in CML emerge, we can continue to address the challenges that remain in the treatment of patients with myeloproliferative neoplasms. This is especially important for those patients in whom conventional and currently available targeted therapies are inadequate.

Acknowledgements

The author's studies are supported by Australian National Health and Medical Research Council Program Grant 461219; Independent Research Institutes Infrastructure Support Scheme; Cure Cancer Australia/Leukaemia Foundation Australia Post Doctoral Fellowship; Lion's Fellowship, Cancer Council of Victoria; the Australian Cancer Research Fund; and a grant from the Victorian State Government Operational Infrastructure Support.

Conflicts of Interest

There are no conflicts of interest.

References

- Adams, G. B., and Scadden, D. T. (2006). The hematopoietic stem cell in its place. *Nat Immunol* 7, 333-337.
- Adamson, J. W., Fialkow, P. J., Murphy, S., Prchal, J. F., and Steinmann, L. (1976). Polycythemia vera: stem-cell and probable clonal origin of the disease. *New England Journal of Medicine* 295, 913-916.
- Akada, H., Akada, S., Hutchison, R. E., and Mohi, G. (2012). Erythroid lineage-restricted expression of Jak2V617F is sufficient to induce a myeloproliferative disease in mice. *Haematologica*
- Akada, H., Yan, D., Zou, H., Fiering, S., Hutchison, R. E., and Mohi, M. G. (2010). Conditional expression of heterozygous or homozygous Jak2V617F from its endogenous promoter induces a polycythemia vera-like disease. *Blood* 115, 3589-3597.
- Anastasi, J. (2011). The myeloproliferative neoplasms including the eosinophilia-related myeloproliferations associated with tyrosine kinase mutations: changes and issues in classification and diagnosis criteria. *Semin Diagn Pathol* 28, 304-313.
- Arai, F., Hirao, A., Ohmura, M., Sato, H., Matsuoka, S., Takubo, K., Ito, K., Koh, G. Y., and Suda, T. (2004). Tie2/angiopoietin-1 signaling regulates hematopoietic stem cell quiescence in the bone marrow niche. *Cell* 118, 149-161.
- Aranaz, P., Hurtado, C., Erquiaga, I., Migueliz, I., Ormazabal, C., Cristobal, I., Garcia-Delgado, M., Novo, F. J., and Vizmanos, J. L. (2012). CBL mutations in myeloproliferative neoplasms are also found in its proline-rich domain and in patients with the V617FJAK2. *Haematologica* 97, 1234-1241.
- Arora, A., and Scholar, E. M. (2005). Role of tyrosine kinase inhibitors in cancer therapy. *J Pharmacol Exp Ther* 315, 971-979.
- Baldrige, M. T., King, K. Y., Boles, N. C., Weksberg, D. C., and Goodell, M. A. (2010). Quiescent haematopoietic stem cells are activated by IFN-gamma in response to chronic infection. *Nature* 465, 793-797.
- Bareng, J., Jilani, I., Gorre, M., Kantarjian, H., Giles, F., Hannah, A., and Albitar, M. (2007). A potential role for HSP90 inhibitors in the treatment of JAK2 mutant-positive diseases as demonstrated using quantitative flow cytometry. *Leuk Lymphoma* 48, 2189-2195.
- Barosi, G., Lupo, L., and Rosti, V. (2012). Management of myeloproliferative neoplasms: from academic guidelines to clinical practice. *Curr Hematol Malig Rep* 7, 50-56.
- Benito, R., Lumberras, E., Abaigar, M., Gutierrez, N. C., Delgado, M., Robledo, C., Garcia, J. L., Rodriguez-Vicente, A. E., Canizo, M. C., and Rivas, J. M. (2012). Imatinib therapy of chronic myeloid leukemia restores the expression levels of key genes for DNA damage and cell-cycle progression. *Pharmacogenet Genomics* 22, 381-388.
- Berkofsky-Fessler, W., Buzzai, M., Kim, M. K., Fruchtman, S., Najfeld, V., Min, D. J., Costa, F. F., Bischof, J. M., Soares, M. B., McConnell, M. J., Zhang, W., Levine, R., Gilliland, D. G., Calogero, R., and Licht, J. D. (2010). Transcriptional profiling of polycythemia vera identifies gene expression patterns both dependent and independent from the action of JAK2V617F. *Clin Cancer Res* 16, 4339-4352.
- Blank, U., Karlsson, G., and Karlsson, S. (2008). Signaling pathways governing stem-cell fate. *Blood* 111, 492-503.
- Branford, S., Seymour, J. F., Grigg, A., Arthur, C., Rudzki, Z., Lynch, K., and Hughes, T. (2007). BCR-ABL messenger RNA levels continue to decline in patients with chronic phase chronic myeloid leukemia treated with imatinib for more than 5 years and approximately half of all first-line treated patients have stable undetectable BCR-ABL using strict sensitivity criteria. *Clin Cancer Res* 13, 7080-7085.
- Briere, J., and el-Kassar, N. (1998). Clonality markers in polycythaemia and primary thrombocythaemia. *Baillieres Clin Haematol* 11, 787-801.
- Bruns, I., Czibere, A., Fischer, J. C., Roels, F., Cadeddu, R. P., Buest, S., Bruennert, D., Huenerlituerkoglu, A. N., Stoecklein, N. H., Singh, R., Zerbini, L. F., Jager, M., Kobbe, G.,

Gattermann, N., Kronenwett, R., Brors, B., and Haas, R. (2009). The hematopoietic stem cell in chronic phase CML is characterized by a transcriptional profile resembling normal myeloid progenitor cells and reflecting loss of quiescence. *Leukemia* 23, 892-899.

Burchert, A., Muller, M. C., Kostrewa, P., Erben, P., Bostel, T., Liebler, S., Hehlmann, R., Neubauer, A., and Hochhaus, A. (2010). Sustained molecular response with interferon alfa maintenance after induction therapy with imatinib plus interferon alfa in patients with chronic myeloid leukemia. *J Clin Oncol* 28, 1429-1435.

Butcher, C. M., Hutton, J. F., Hahn, U., To, L. B., Bardy, P., Lewis, I., and D'Andrea, R. J. (2007). Cellular origin and lineage specificity of the JAK2(V617F) allele in polycythemia vera. *Blood* 109, 386-387.

Calvi, L. M., Adams, G. B., Weibrecht, K. W., Weber, J. M., Olson, D. P., Knight, M. C., Martin, R. P., Schipani, E., Divieti, P., Bringham, F. R., Milner, L. A., Kronenberg, H. M., and Scadden, D. T. (2003). Osteoblastic cells regulate the haematopoietic stem cell niche. *Nature* 425, 841-846.

Cao, Y. A., Wagers, A. J., Beilhack, A., Dusich, J., Bachmann, M. H., Negrin, R. S., Weissman, I. L., and Contag, C. H. (2004). Shifting foci of hematopoiesis during reconstitution from single stem cells. *Proc Natl Acad Sci U S A* 101, 221-226.

Capel, B., Hawley, R. G., and Mintz, B. (1990). Long- and short-lived murine hematopoietic stem cell clones individually identified with retroviral integration markers. *Blood* 75, 2267-2270.

Chan, C. K., Chen, C. C., Luppen, C. A., Kim, J. B., DeBoer, A. T., Wei, K., Helms, J. A., Kuo, C. J., Kraft, D. L., and Weissman, I. L. (2009). Endochondral ossification is required for haematopoietic stem-cell niche formation. *Nature* 457, 490-494.

Chandra, H. S., Heisterkamp, N. C., Hungerford, A., Morrisette, J. J. D., Nowell, P. C., Rowley, J. D., and Testa, J. R. (2011). Philadelphia Chromosome Symposium: commemoration of the 50th anniversary of the discovery of the Ph chromosome. *Cancer Genetics* 204, 171-179.

Constantinescu, S. N., Ghaffari, S., and Lodish, H. F. (1999). The Erythropoietin Receptor: Structure, Activation and Intracellular Signal Transduction. *Trends Endocrinol Metab* 10, 18-23.

Cools, J., DeAngelo, D. J., Gotlib, J., Stover, E. H., Legare, R. D., Cortes, J., Kutok, J., Clark, J., Galinsky, I., and Griffin, J. D. (2003). A tyrosine kinase created by fusion of the PDGFRA and FIP1L1 genes as a therapeutic target of imatinib in idiopathic hypereosinophilic syndrome. *New England Journal of Medicine* 348, 1201-1214.

Cortes, J., and Kantarjian, H. (2012). How I treat newly diagnosed chronic phase CML. *Blood* 120, 1390-1397.

Crews, L. A., and Jamieson, C. H. (2012). Chronic myeloid leukemia stem cell biology. *Curr Hematol Malig Rep* 7, 125-132.

Cross, N. C., White, H. E., Muller, M. C., Saglio, G., and Hochhaus, A. (2012). Standardized definitions of molecular response in chronic myeloid leukemia. *Leukemia*

Dai, C. H., Krantz, S. B., Dessypris, E. N., Means, R. T. J., Horn, S. T., and Gilbert, H. S. (1992). Polycythemia vera. II. Hypersensitivity of bone marrow erythroid, granulocyte-macrophage, and megakaryocyte progenitor cells to interleukin-3 and granulocyte-macrophage colony-stimulating factor. *Blood* 80, 891-899.

Dai, C. H., Krantz, S. B., Means, R. T. J., Horn, S. T., and Gilbert, H. S. (1991). Polycythemia vera blood burst-forming units-erythroid are hypersensitive to interleukin-3. *J Clin Invest* 87, 391-396.

DAMESHEK, W. (1951). Some speculations on the myeloproliferative syndromes. *Blood* 6, 372-375.

Delhommeau, F., Pisani, D. F., James, C., Casadevall, N., Constantinescu, S., and Vainchenker, W. (2006). Oncogenic mechanisms in myeloproliferative disorders. *Cell Mol Life Sci* 63, 2939-2953.

Dierks, C., Beigi, R., Guo, G. R., Zirlik, K., Stegert, M. R., Manley, P., Trussell, C., Schmitt-Graeff, A., Landwerlin, K., Veelken, H., and Warmuth, M. (2008). Expansion of Bcr-Abl-positive leukemic stem cells is dependent on Hedgehog pathway activation. *Cancer Cell* 14, 238-249.

Drize, N. J., Olshanskaya, Y. V., Gerasimova, L. P., Manakova, T. E., Samoylina, N. L., Todria, T. V., and Chertkov, J. L. (2001). Lifelong hematopoiesis in both reconstituted and sublethally irradiated mice is provided by multiple sequentially recruited stem cells. *Exp Hematol* 29, 786-794.

Dupont, S., Massé, A., James, C., Teyssandier, I., Lécluse, Y., Larbret, F., Ugo, V., Saulnier, P., Koscielny, S., and Le Couédic, J. P. (2007). The JAK2 617V>F mutation triggers erythropoietin hypersensitivity and terminal erythroid amplification in primary cells from patients with polycythemia vera. *Blood* 110, 1013-1021.

Durand, E. M., and Zon, L. I. (2010). Newly emerging roles for prostaglandin E2 regulation of hematopoiesis and hematopoietic stem cell engraftment. *Curr Opin Hematol* 17, 308-312.

Essers, M. A., Offner, S., Blanco-Bose, W. E., Waibler, Z., Kalinke, U., Duchosal, M. A., and Trumpp, A. (2009). IFN α activates dormant haematopoietic stem cells in vivo. *Nature* 458, 904-908.

Fleming, H. E., Janzen, V., Lo Celso, C., Guo, J., Leahy, K. M., Kronenberg, H. M., and Scadden, D. T. (2008). Wnt signaling in the niche enforces hematopoietic stem cell quiescence and is necessary to preserve self-renewal in vivo. *Cell Stem Cell* 2, 274-283.

Frisch, B. J., Porter, R. L., Gigliotti, B. J., Olm-Shipman, A. J., Weber, J. M., O'Keefe, R. J., Jordan, C. T., and Calvi, L. M. (2009). In vivo prostaglandin E2 treatment alters the bone marrow microenvironment and preferentially expands short-term hematopoietic stem cells. *Blood* 114, 4054-4063.

Funakoshi-Tago, M., Tago, K., Abe, M., Sonoda, Y., and Kasahara, T. (2010). STAT5 activation is critical for the transformation mediated by myeloproliferative disorder-associated JAK2 V617F mutant. *Journal of Biological Chemistry* 285, 5296.

Gasson, J. C., Golde, D. W., Kaufman, S. E., Westbrook, C. A., Hewick, R. M., Kaufman, R. J., Wong, G. G., Temple, P. A., Leary, A. C., Brown, E. L., and et, a. (1985). Molecular characterization and expression of the gene encoding human erythroid-potentiating activity. *Nature* 315, 768-771.

Goldman, J. M., Majhail, N. S., Klein, J. P., Wang, Z., Sobocinski, K. A., Arora, M., Horowitz, M. M., and Rizzo, J. D. (2010). Relapse and late mortality in 5-year survivors of myeloablative allogeneic hematopoietic cell transplantation for chronic myeloid leukemia in first chronic phase. *J Clin Oncol* 28, 1888-1895.

Graham, S. M., Jorgensen, H. G., Allan, E., Pearson, C., Alcorn, M. J., Richmond, L., and Holyoake, T. L. (2002). Primitive, quiescent, Philadelphia-positive stem cells from patients with chronic myeloid leukemia are insensitive to STI571 in vitro. *Blood* 99, 319-325.

Graham, S. M., Vass, J. K., Holyoake, T. L., and Graham, G. J. (2007). Transcriptional analysis of quiescent and proliferating CD34+ human hemopoietic cells from normal and chronic myeloid leukemia sources. *Stem Cells* 25, 3111-3120.

Guglielmelli, P., Biamonte, F., Score, J., Hidalgo-Curtis, C., Cervantes, F., Maffioli, M., Fanelli, T., Ernst, T., Winkelman, N., Jones, A. V., Zoi, K., Reiter, A., Duncombe, A., Villani, L., Bosi, A., Barosi, G., Cross, N. C., and Vannucchi, A. M. (2011). EZH2 mutational status predicts poor survival in myelofibrosis. *Blood* 118, 5227-5234.

Gupta, R., Knight, C. L., and Bain, B. J. (2002). Receptor tyrosine kinase mutations in myeloid neoplasms. *Br J Haematol* 117, 489-508.

Gupta, V., Hari, P., and Hoffman, R. (2012). Allogeneic hematopoietic cell transplantation for myelofibrosis in the era of JAK inhibitors. *Blood* 120, 1367-1379.

Hamilton, A., Helgason, G. V., Schemionek, M., Zhang, B., Myssina, S., Allan, E. K., Nicolini, F. E., Muller-Tidow, C., Bhatia, R., Brunton, V. G., Koschmieder, S., and Holyoake, T. L. (2012). Chronic myeloid leukemia stem cells are not dependent on Bcr-Abl

kinase activity for their survival. *Blood* 119, 1501-1510.

Hantschel, O., Warsch, W., Eckelhart, E., Kaupe, I., Grebien, F., Wagner, K. U., Superti-Furga, G., and Sexl, V. (2012). BCR-ABL uncouples canonical JAK2-STAT5 signaling in chronic myeloid leukemia. *Nat Chem Biol* 8, 285-293.

Harrison, C., Kiladjian, J. J., Al-Ali, H. K., Gisslinger, H., Waltzman, R., Stalbovskaya, V., McQuitty, M., Hunter, D. S., Levy, R., Knoops, L., Cervantes, F., Vannucchi, A. M., Barbui, T., and Barosi, G. (2012). JAK inhibition with ruxolitinib versus best available therapy for myelofibrosis. *N Engl J Med* 366, 787-798.

Hehlmann, R. (2012). How I treat CML blast crisis. *Blood* 120, 737-747.

Hehlmann, R., Lauseker, M., Jung-Munkwitz, S., Leitner, A., Muller, M. C., Pletsch, N., Proetel, U., Haferlach, C., Schlegelberger, B., Balleisen, L., Hanel, M., Pffirmann, M., Krause, S. W., Nerl, C., Pralle, H., Gratwohl, A., Hossfeld, D. K., Hasford, J., Hochhaus, A., and Saussele, S. (2011). Tolerability-adapted imatinib 800 mg/d versus 400 mg/d versus 400 mg/d plus interferon-alpha in newly diagnosed chronic myeloid leukemia. *J Clin Oncol* 29, 1634-1642.

Heidel, F. H., Bullinger, L., Feng, Z., Wang, Z., Neff, T. A., Stein, L., Kalaitzidis, D., Lane, S. W., and Armstrong, S. A. (2012). Genetic and Pharmacologic Inhibition of beta-Catenin Targets Imatinib-Resistant Leukemia Stem Cells in CML. *Cell Stem Cell* 10, 412-424.

Herrera-Merchan, A., Arranz, L., Ligos, J. M., de Molina, A., Dominguez, O., and Gonzalez, S. (2012). Ectopic expression of the histone methyltransferase Ezh2 in haematopoietic stem cells causes myeloproliferative disease. *Nat Commun* 3, 623.

Hochhaus, A., O'Brien, S. G., Guilhot, F., Druker, B. J., Branford, S., Foroni, L., Goldman, J. M., Muller, M. C., Radich, J. P., Rudoltz, M., Mone, M., Gathmann, I., Hughes, T. P., and Larson, R. A. (2009). Six-year follow-up of patients receiving imatinib for the first-line treatment of chronic myeloid leukemia. *Leukemia* 23, 1054-1061.

Hodgson, G. S., and Bradley, T. R. (1979). Properties of haematopoietic stem cells surviving 5-fluorouracil treatment: evidence for a pre-CFU-S cell? *Nature* 281, 381-382.

Holtz, M. S., Forman, S. J., and Bhatia, R. (2005). Nonproliferating CML CD34+ progenitors are resistant to apoptosis induced by a wide range of proapoptotic stimuli. *Leukemia* 19, 1034-1041.

Holyoake, T., Jiang, X., Eaves, C., and Eaves, A. (1999). Isolation of a highly quiescent subpopulation of primitive leukemic cells in chronic myeloid leukemia. *Blood* 94, 2056-2064.

Hope, K. J., Jin, L., and Dick, J. E. (2003). Human acute myeloid leukemia stem cells. *Arch Med Res* 34, 507-514.

Huntly, B. J., Shigematsu, H., Deguchi, K., Lee, B. H., Mizuno, S., Duclos, N., Rowan, R., Amaral, S., Curley, D., Williams, I. R., Akashi, K., and Gilliland, D. G. (2004). MOZ-TIF2, but not BCR-ABL, confers properties of leukemic stem cells to committed murine hematopoietic progenitors. *Cancer Cell* 6, 587-596.

Ishii, T., Bruno, E., Hoffman, R., and Xu, M. (2006). Involvement of various hematopoietic-cell lineages by the JAK2V617F mutation in polycythemia vera. *Blood* 108, 3128-3134.

Ishii, T., Zhao, Y., Sozer, S., Shi, J., Zhang, W., Hoffman, R., and Xu, M. (2007). Behavior of CD34+ cells isolated from patients with polycythemia vera in NOD/SCID mice. *Exp Hematol* 35, 1633-1640.

Jabbour, E., Kantarjian, H. M., O'Brien, S., Shan, J., Quintas-Cardama, A., Garcia-Manero, G., Rios, M. B., and Cortes, J. E. (2011). Front-line therapy with second-generation tyrosine kinase inhibitors in patients with early chronic phase chronic myeloid leukemia: what is the optimal response? *J Clin Oncol* 29, 4260-4265.

James, C., Mazurier, F., Dupont, S., Chaligne, R., Lamrissi-Garcia, I., Tulliez, M., Lippert, E., Mahon, F. X., Pasquet, J. M., and Etienne, G. (2008). The hematopoietic stem cell compartment of JAK2V617F-positive myeloproliferative disorders is a reflection of disease heterogeneity. *Blood* 112, 2429-2438.

James, C., Ugo, V., Le Couédic, J. P., Staerk, J., Delhommeau, F., Lacout, C., Garçon, L., Raslova, H., Berger, R., and Bennaceur-Griscelli, A. (2005). A unique clonal JAK2 mutation leading to constitutive signalling causes polycythaemia vera. *Nature* *434*, 1144-1148.

Jamieson, C. H., Ailles, L. E., Dylla, S. J., Muijtjens, M., Jones, C., Zehnder, J. L., Gotlib, J., Li, K., Manz, M. G., Keating, A., Sawyers, C. L., and Weissman, I. L. (2004). Granulocyte-macrophage progenitors as candidate leukemic stem cells in blast-crisis CML. *N Engl J Med* *351*, 657-667.

Jamieson, C. H., Gotlib, J., Durocher, J. A., Chao, M. P., Mariappan, M. R., Lay, M., Jones, C., Zehnder, J. L., Lilleberg, S. L., and Weissman, I. L. (2006). The JAK2 V617F mutation occurs in hematopoietic stem cells in polycythemia vera and predisposes toward erythroid differentiation. *Proc Natl Acad Sci U S A* *103*, 6224-6229.

Kantarjian, H., Sawyers, C., Hochhaus, A., Guilhot, F., Schiffer, C., Gambacorti-Passerini, C., Niederwieser, D., Resta, D., Capdeville, R., Zoellner, U., Talpaz, M., Druker, B., Goldman, J., O'Brien, S. G., Russell, N., Fischer, T., Ottmann, O., Cony-Makhoul, P., Facon, T., Stone, R., Miller, C., Tallman, M., Brown, R., Schuster, M., Loughran, T., Gratwohl, A., Mandelli, F., Saglio, G., Lazzarino, M., Russo, D., Baccarani, M., and Morra, E. (2002). Hematologic and cytogenetic responses to imatinib mesylate in chronic myelogenous leukemia. *N Engl J Med* *346*, 645-652.

Kantarjian, H. M., Shah, N. P., Cortes, J. E., Baccarani, M., Agarwal, M. B., Undurraga, M. S., Wang, J., Ipina, J. J., Kim, D. W., Ogura, M., Pavlovsky, C., Junghanss, C., Milone, J. H., Nicolini, F. E., Robak, T., Van Droogenbroeck, J., Vellenga, E., Bradley-Garelik, M. B., Zhu, C., and Hochhaus, A. (2012). Dasatinib or imatinib in newly diagnosed chronic-phase chronic myeloid leukemia: 2-year follow-up from a randomized phase 3 trial (DASISION). *Blood* *119*, 1123-1129.

Karlsson, G., Blank, U., Moody, J. L., Ehinger, M., Singbrant, S., Deng, C. X., and Karlsson, S. (2007). Smad4 is critical for self-renewal of hematopoietic stem cells. *J Exp Med* *204*, 467-474.

Kaushansky, K., Lok, S., Holly, R. D., Broudy, V. C., Lin, N., Bailey, M. C., Forstrom, J. W., Buddle, M. M., Oort, P. J., Hagen, F. S., and et, a. (1994). Promotion of megakaryocyte progenitor expansion and differentiation by the c-Mpl ligand thrombopoietin. *Nature* *369*, 568-571.

Kent, D., Copley, M., Benz, C., Dykstra, B., Bowie, M., and Eaves, C. (2008). Regulation of hematopoietic stem cells by the steel factor/KIT signaling pathway. *Clin Cancer Res* *14*, 1926-1930.

Khwaja, A. (2006). The role of Janus kinases in haemopoiesis and haematological malignancy. *Br J Haematol* *134*, 366-384.

Kiladjian, J. J., Cassinat, B., Chevret, S., Turlure, P., Cambier, N., Roussel, M., Bellucci, S., Grandchamp, B., Chomienne, C., and Fenaux, P. (2008). Pegylated interferon-alfa-2a induces complete hematologic and molecular responses with low toxicity in polycythemia vera. *Blood* *112*, 3065-3072.

Ko, M., Bandukwala, H. S., An, J., Lamperti, E. D., Thompson, E. C., Hastie, R., Tsangaratou, A., Rajewsky, K., Koralov, S. B., and Rao, A. (2011). Ten-Eleven-Translocation 2 (TET2) negatively regulates homeostasis and differentiation of hematopoietic stem cells in mice. *Proc Natl Acad Sci U S A* *108*, 14566-14571.

Kumano, K., Arai, S., Hosoi, M., Taoka, K., Takayama, N., Otsu, M., Nagae, G., Ueda, K., Nakazaki, K., Kamikubo, Y., Eto, K., Aburatani, H., Nakauchi, H., and Kurokawa, M. (2012). Generation of induced pluripotent stem cells from primary chronic myelogenous leukemia patient samples. *Blood*

Kuroda, J., Puthalakath, H., Cragg, M. S., Kelly, P. N., Bouillet, P., Huang, D. C., Kimura, S., Ottmann, O. G., Druker, B. J., Villunger, A., Roberts, A. W., and Strasser, A. (2006). Bim and Bad mediate imatinib-induced killing of Bcr/Abl+ leukemic cells, and resistance due to their loss is overcome by a BH3 mimetic. *Proc Natl Acad Sci U S A* *103*, 14907-14912.

Laukkanen, M. O., Kuramoto, K., Calmels, B., Takatoku, M., von Kalle, C., Donahue, R. E., and Dunbar, C. E. (2005). Low-dose total body irradiation causes clonal fluctuation of primate hematopoietic stem and progenitor cells. *Blood* 105, 1010-1015.

Levine, R. L., Wadleigh, M., Cools, J., Ebert, B. L., Wernig, G., Huntly, B. J. P., Boggon, T. J., Wlodarska, I., Clark, J. J., and Moore, S. (2005). Activating mutation in the tyrosine kinase JAK2 in polycythemia vera, essential thrombocythemia, and myeloid metaplasia with myelofibrosis. *Cancer cell* 7, 387-397.

Li, B., Zhang, G., Li, C., He, D., Li, X., Zhang, C., Tang, F., Deng, X., Lu, J., Tang, Y., Li, R., Chen, Z., and Duan, C. (2012a). Identification of JAK2 as a Mediator of FIP1L1-PDGFR α -Induced Eosinophil Growth and Function in CEL. *PLoS One* 7, e34912.

Li, J., Kent, D. G., Chen, E., and Green, A. R. (2011). Mouse models of myeloproliferative neoplasms: JAK of all grades. *Dis Model Mech* 4, 311-317.

Li, J., Spensberger, D., Ahn, J. S., Anand, S., Beer, P. A., Ghevaert, C., Chen, E., Forrai, A., Scott, L. M., Ferreira, R., Campbell, P. J., Watson, S. P., Liu, P., Erber, W. N., Huntly, B. J., Ottersbach, K., and Green, A. R. (2010). JAK2 V617F impairs hematopoietic stem cell function in a conditional knock-in mouse model of JAK2 V617F-positive essential thrombocythemia. *Blood* 116, 1528-1538.

Li, L., Wang, L., Li, L., Wang, Z., Ho, Y., McDonald, T., Holyoake, T. L., Chen, W., and Bhatia, R. (2012b). Activation of p53 by SIRT1 inhibition enhances elimination of CML leukemia stem cells in combination with imatinib. *Cancer Cell* 21, 266-281.

Liu, F., Zhao, X., Perna, F., Wang, L., Koppikar, P., Abdel-Wahab, O., Harr, M. W., Levine, R. L., Xu, H., and Tefferi, A. (2011). JAK2V617F-mediated phosphorylation of PRMT5 downregulates its methyltransferase activity and promotes myeloproliferation. *Cancer cell* 19, 283-294.

Lok, S., Kaushansky, K., Holly, R. D., Kuijper, J. L., Lofton-Day, C. E., Oort, P. J., Grant, F. J., Heipel, M. D., Burkhead, S. K., Kramer, J. M., and et, a. (1994). Cloning and expression of murine thrombopoietin cDNA and stimulation of platelet production in vivo. *Nature* 369, 565-568.

Lu, M., Wang, J., Li, Y., Berenzon, D., Wang, X., Mascarenhas, J., Xu, M., and Hoffman, R. (2010a). Treatment with the Bcl-xL inhibitor ABT-737 in combination with interferon alpha specifically targets JAK2V617F-positive polycythemia vera hematopoietic progenitor cells. *Blood* 116, 4284-4287.

Lu, M., Wang, X., Li, Y., Tripodi, J., Mosoyan, G., Mascarenhas, J., Kremyanskaya, M., Najfeld, V., and Hoffman, R. (2012). Combination treatment in vitro with Nutlin, a small-molecule antagonist of MDM2, and pegylated interferon alpha 2a specifically targets JAK2V617F positive polycythemia vera cells. *Blood*

Lu, M., Zhang, W., Li, Y., Berenzon, D., Wang, X., Wang, J., Mascarenhas, J., Xu, M., and Hoffman, R. (2010b). Interferon-alpha targets JAK2V617F-positive hematopoietic progenitor cells and acts through the p38 MAPK pathway. *Exp Hematol* 38, 472-480.

Mahon, F. X., Rea, D., Guilhot, J., Guilhot, F., Huguet, F., Nicolini, F., Legros, L., Charbonnier, A., Guerci, A., Varet, B., Etienne, G., Reiffers, J., and Rousselot, P. (2010). Discontinuation of imatinib in patients with chronic myeloid leukaemia who have maintained complete molecular remission for at least 2 years: the prospective, multicentre Stop Imatinib (STIM) trial. *Lancet Oncol* 11, 1029-1035.

Mahon, F. X., Rea, D., Guilhot, J., Guilhot, F., Huguet, F., Nicolini, F., Legros, L., Charbonnier, A., Guerci, A., Varet, B., Etienne, G., Reiffers, J., and Rousselot, P. (2011). Discontinuation of imatinib in patients with chronic myeloid leukemia who have maintained complete molecular response: update results of the STIM study [abstract]. *Blood (ASH Annual Meeting Abstracts)* 118 (21), Abstract 603.

Majewski, I. J., Blewitt, M. E., de Graaf, C. A., McManus, E. J., Bahlo, M., Hilton, A. A., Hyland, C. D., Smyth, G. K., Corbin, J. E., Metcalf, D., Alexander, W. S., and Hilton, D. J. (2008). Polycomb repressive complex 2 (PRC2) restricts hematopoietic stem cell activity.

PLoS Biol 6, e93.

- Majewski, I. J., Ritchie, M. E., Phipson, B., Corbin, J., Pakusch, M., Ebert, A., Busslinger, M., Koseki, H., Hu, Y., Smyth, G. K., Alexander, W. S., Hilton, D. J., and Blewitt, M. E. (2010). Opposing roles of polycomb repressive complexes in hematopoietic stem and progenitor cells. *Blood* 116, 731-739.
- Makishima, H., Jankowska, A. M., McDevitt, M. A., O'Keefe, C., Dujardin, S., Cazzolli, H., Przychodzen, B., Prince, C., Nicoll, J., Siddaiah, H., Shaik, M., Szpurka, H., Hsi, E., Advani, A., Paquette, R., and Maciejewski, J. P. (2011). CBL, CBLB, TET2, ASXL1, and IDH1/2 mutations and additional chromosomal aberrations constitute molecular events in chronic myelogenous leukemia. *Blood* 117, e198-206.
- Martinez-Aviles, L., Besses, C., Alvarez-Larran, A., Torres, E., Serrano, S., and Bellosillo, B. (2012). TET2, ASXL1, IDH1, IDH2, and c-CBL genes in JAK2- and MPL-negative myeloproliferative neoplasms. *Ann Hematol* 91, 533-541.
- Marty, C., Lacout, C., Cuingnet, M., Hasan, S., Solary, E., Vainchenker, W., and Villeval, J. L. (2011). JAK2V617F Promotes Stem Cell Amplification Driving MPN Clonal Dominance in Mice and Treatment by IFN α Prevents This Effect. *Blood* (ASH Annual Meeting Abstracts) 118(21), Abstract 616.
- Marubayashi, S., Koppikar, P., Taldone, T., Abdel-Wahab, O., West, N., Bhagwat, N., Caldas-Lopes, E., Ross, K. N., Gonen, M., Gozman, A., Ahn, J. H., Rodina, A., Ouerfelli, O., Yang, G., Hedvat, C., Bradner, J. E., Chiosis, G., and Levine, R. L. (2010). HSP90 is a therapeutic target in JAK2-dependent myeloproliferative neoplasms in mice and humans. *J Clin Invest* 120, 3578-3593.
- Metcalf, D., and Nicola, N. A. (1983). Proliferative effects of purified granulocyte colony-stimulating factor (G-CSF) on normal mouse hemopoietic cells. *J Cell Physiol* 116, 198-206.
- Metcalf, D. (2010). The colony-stimulating factors and cancer. *Nature Reviews Cancer* 10, 425-434.
- Mitterbauer, G., Winkler, K., Gisslinger, H., Geissler, K., Lechner, K., and Mannhalter, C. (1999). Clonality analysis using X-chromosome inactivation at the human androgen receptor gene (Humara). Evaluation of large cohorts of patients with chronic myeloproliferative diseases, secondary neutrophilia, and reactive thrombocytosis. *Am J Clin Pathol* 112, 93-100.
- Miyake, T., Kung, C. K., and Goldwasser, E. (1977). Purification of human erythropoietin. *J Biol Chem* 252, 5558-5564.
- Moran-Crusio, K., Reavie, L., Shih, A., Abdel-Wahab, O., Ndiaye-Lobry, D., Lobry, C., Figueroa, M. E., Vasanthakumar, A., Patel, J., Zhao, X., Perna, F., Pandey, S., Madzo, J., Song, C., Dai, Q., He, C., Ibrahim, S., Beran, M., Zavadil, J., Nimer, S. D., Melnick, A., Godley, L. A., Aifantis, I., and Levine, R. L. (2011). Tet2 loss leads to increased hematopoietic stem cell self-renewal and myeloid transformation. *Cancer Cell* 20, 11-24.
- Mullally, A., Lane, S. W., Ball, B., Megerdichian, C., Okabe, R., Al-Shahrour, F., Paktinat, M., Haydu, J. E., Housman, E., Lord, A. M., Wernig, G., Kharas, M. G., Mercher, T., Kutok, J. L., Gilliland, D. G., and Ebert, B. L. (2010). Physiological Jak2V617F expression causes a lethal myeloproliferative neoplasm with differential effects on hematopoietic stem and progenitor cells. *Cancer Cell* 17, 584-596.
- Mullally, A., Poveromo, L., Schneider, R. K., Al-Shahrour, F., Lane, S. W., and Ebert, B. L. (2012). Distinct roles for long-term hematopoietic stem cells and erythroid precursor cells in a murine model of Jak2V617F polycythemia vera. *Blood* 120, 166-172.
- Neubauer, H., Cumano, A., Muller, M., Wu, H., Huffstadt, U., and Pfeffer, K. (1998). Jak2 deficiency defines an essential developmental checkpoint in definitive hematopoiesis. *Cell* 93, 397-409.
- Ng, A. P., Kauppi, M., Metcalf, D., Di Rago, L., Hyland, C. D., and Alexander, W. S. (2012a). Characterization of thrombopoietin (TPO)-responsive progenitor cells in adult mouse bone marrow with in vivo megakaryocyte and erythroid potential. *Proc Natl Acad Sci U S A* 109, 2364-2369.

Ng, K. P., Hillmer, A. M., Chuah, C. T., Juan, W. C., Ko, T. K., Teo, A. S., Ariyaratne, P. N., Takahashi, N., Sawada, K., Fei, Y., Soh, S., Lee, W. H., Huang, J. W., Allen, J. C. J., Woo, X. Y., Nagarajan, N., Kumar, V., Thalamuthu, A., Poh, W. T., Ang, A. L., Mya, H. T., How, G. F., Yang, L. Y., Koh, L. P., Chowbay, B., Chang, C. T., Nadarajan, V. S., Chng, W. J., Than, H., Lim, L. C., Goh, Y. T., Zhang, S., Poh, D., Tan, P., Seet, J. E., Ang, M. K., Chau, N. M., Ng, Q. S., Tan, D. S., Soda, M., Isobe, K., Nothen, M. M., Wong, T. Y., Shahab, A., Ruan, X., Cacheux-Rataboul, V., Sung, W. K., Tan, E. H., Yatabe, Y., Mano, H., Soo, R. A., Chin, T. M., Lim, W. T., Ruan, Y., and Ong, S. T. (2012b). A common BIM deletion polymorphism mediates intrinsic resistance and inferior responses to tyrosine kinase inhibitors in cancer. *Nat Med* 18, 521-528.

Nicola, N. A., Metcalf, D., Matsumoto, M., and Johnson, G. R. (1983). Purification of a factor inducing differentiation in murine myelomonocytic leukemia cells. Identification as granulocyte colony-stimulating factor. *J Biol Chem* 258, 9017-9023.

Nilsson, S. K., Johnston, H. M., Whitty, G. A., Williams, B., Webb, R. J., Denhardt, D. T., Bertoncello, I., Bendall, L. J., Simmons, P. J., and Haylock, D. N. (2005). Osteopontin, a key component of the hematopoietic stem cell niche and regulator of primitive hematopoietic progenitor cells. *Blood* 106, 1232-1239.

North, T. E., Goessling, W., Walkley, C. R., Lengerke, C., Kopani, K. R., Lord, A. M., Weber, G. J., Bowman, T. V., Jang, I. H., Grosser, T., Fitzgerald, G. A., Daley, G. Q., Orkin, S. H., and Zon, L. I. (2007). Prostaglandin E2 regulates vertebrate haematopoietic stem cell homeostasis. *Nature* 447, 1007-1011.

O'Brien, S. G., Guilhot, F., Larson, R. A., Gathmann, I., Baccarani, M., Cervantes, F., Cornelissen, J. J., Fischer, T., Hochhaus, A., Hughes, T., Lechner, K., Nielsen, J. L., Rousselot, P., Reiffers, J., Saglio, G., Shepherd, J., Simonsson, B., Gratwohl, A., Goldman, J. M., Kantarjian, H., Taylor, K., Verhoef, G., Bolton, A. E., Capdeville, R., and Druker, B. J. (2003). Imatinib compared with interferon and low-dose cytarabine for newly diagnosed chronic-phase chronic myeloid leukemia. *N Engl J Med* 348, 994-1004.

Opferman, J. T., Iwasaki, H., Ong, C. C., Suh, H., Mizuno, S., Akashi, K., and Korsmeyer, S. J. (2005). Obligate role of anti-apoptotic MCL-1 in the survival of hematopoietic stem cells. *Science* 307, 1101-1104.

Palandri, F., Castagnetti, F., Iacobucci, I., Martinelli, G., Amabile, M., Gugliotta, G., Poerio, A., Testoni, N., Breccia, M., Bocchia, M., Crugnola, M., Rege-Cambrin, G., Martino, B., Pierri, I., Radaelli, F., Specchia, G., Pane, F., Saglio, G., Rosti, G., and Baccarani, M. (2010). The response to imatinib and interferon-alpha is more rapid than the response to imatinib alone: a retrospective analysis of 495 Philadelphia-positive chronic myeloid leukemia patients in early chronic phase. *Haematologica* 95, 1415-1419.

Pardanani, A., Gotlib, J. R., Jamieson, C., Cortes, J. E., Talpaz, M., Stone, R. M., Silverman, M. H., Gilliland, D. G., Shorr, J., and Tefferi, A. (2011). Safety and efficacy of TG101348, a selective JAK2 inhibitor, in myelofibrosis. *J Clin Oncol* 29, 789-796.

Pardanani, A., Lasho, T., Finke, C., Oh, S. T., Gotlib, J., and Tefferi, A. (2010). LNK mutation studies in blast-phase myeloproliferative neoplasms, and in chronic-phase disease with TET2, IDH, JAK2 or MPL mutations. *Leukemia* 24, 1713-1718.

Pardanani, A. D., Levine, R. L., Lasho, T., Pikman, Y., Mesa, R. A., Wadleigh, M., Steensma, D. P., Elliott, M. A., Wolanskyj, A. P., and Hogan, W. J. (2006). MPL515 mutations in myeloproliferative and other myeloid disorders: a study of 1182 patients. *Blood* 108, 3472-3476.

Parganas, E., Wang, D., Stravopodis, D., Topham, D. J., Marine, J. C., Teglund, S., Vanin, E. F., Bodner, S., Colamonici, O. R., and van Deursen, J. M. (1998). Jak2 is essential for signaling through a variety of cytokine receptors. *Cell* 93, 385-395.

Parmar, K., Mauch, P., Vergilio, J. A., Sackstein, R., and Down, J. D. (2007). Distribution of hematopoietic stem cells in the bone marrow according to regional hypoxia. *Proc Natl Acad Sci U S A* 104, 5431-5436.

Pasquier, F., Tonetti, C., Besancenot, R., Cassinat, B., Kiladjian, J. J., Herault, O., Remeau, P., Lacout, C., Villeval, J. L., Vainchenker, W., Chaligne, R., and Giraudier, S. (2009). Interferon-Alpha Preferentially Targets JAK2V617F-Positive Rather Than Wild-Type Early Progenitor Cells in Myeloproliferative Disorders. *Blood (ASH Annual Meeting Abstracts)* 114(22), Abstract 436.

Ploemacher, R. E., van Os, R., van Beurden, C. A., and Down, J. D. (1992). Murine haemopoietic stem cells with long-term engraftment and marrow repopulating ability are more resistant to gamma-radiation than are spleen colony forming cells. *Int J Radiat Biol* 61, 489-499.

Prchal, J. F., and Axelrad, A. A. (1974). Letter: Bone-marrow responses in polycythemia vera. *N Engl J Med* 290, 1382.

Preudhomme, C., Guilhot, J., Nicolini, F. E., Guerci-Bresler, A., Rigal-Huguet, F., Maloisel, F., Coiteux, V., Gardembas, M., Berthou, C., Vekhoff, A., Rea, D., Jourdan, E., Allard, C., Delmer, A., Rousselot, P., Legros, L., Berger, M., Corm, S., Etienne, G., Roche-Lestienne, C., Eclache, V., Mahon, F. X., and Guilhot, F. (2010). Imatinib plus peginterferon alfa-2a in chronic myeloid leukemia. *N Engl J Med* 363, 2511-2521.

Proia, D. A., Foley, K. P., Korb, T., Sang, J., Smith, D., Bates, R. C., Liu, Y., Rosenberg, A. F., Zhou, D., Koya, K., Barsom, J., and Blackman, R. K. (2011). Multifaceted intervention by the Hsp90 inhibitor ganetespib (STA-9090) in cancer cells with activated JAK/STAT signaling. *PLoS One* 6, e18552.

Qian, H., Buza-Vidas, N., Hyland, C. D., Jensen, C. T., Antonchuk, J., Mansson, R., Thoren, L. A., Eklom, M., Alexander, W. S., and Jacobsen, S. E. (2007). Critical role of thrombopoietin in maintaining adult quiescent hematopoietic stem cells. *Cell Stem Cell* 1, 671-684.

Quintas-Cardama, A., Kantarjian, H., Manshouri, T., Luthra, R., Estrov, Z., Pierce, S., Richie, M. A., Borthakur, G., Konopleva, M., Cortes, J., and Verstovsek, S. (2009). Pegylated interferon alfa-2a yields high rates of hematologic and molecular response in patients with advanced essential thrombocythemia and polycythemia vera. *J Clin Oncol* 27, 5418-5424.

Radich, J. P., Gooley, T., Bryant, E., Chauncey, T., Clift, R., Beppu, L., Edmands, S., Flowers, M. E., Kerkof, K., Nelson, R., and Appelbaum, F. R. (2001). The significance of bcr-abl molecular detection in chronic myeloid leukemia patients "late," 18 months or more after transplantation. *Blood* 98, 1701-1707.

Robb, L. (2007). Cytokine receptors and hematopoietic differentiation. *Oncogene* 26, 6715-6723.

Ross, D. M., Bartley, P. A., Goyne, J., Morley, A. A., Seymour, J. F., and Grigg, A. P. (2011a). Durable complete molecular remission of chronic myeloid leukemia following dasatinib cessation, despite adverse disease features. *Haematologica* 96, 1720-1722.

Ross, D. M., Branford, S., Seymour, J. F., Schwarzer, A. P., Arthur, C., Bartley, P. A., Slader, C., Field, C., Dang, P., and Filshie, R. J. (2010). Patients with chronic myeloid leukemia who maintain a complete molecular response after stopping imatinib treatment have evidence of persistent leukemia by DNA PCR. *Leukemia* 24, 1719-1724.

Ross, D. M., Hughes, T. P., and Melo, J. V. (2011b). Do we have to kill the last CML cell? *Leukemia* 25, 193-200.

Schwartzberg, P. L., Stall, A. M., Hardin, J. D., Bowdish, K. S., Humaran, T., Boast, S., Harbison, M. L., Robertson, E. J., and Goff, S. P. (1991). Mice homozygous for the abm1 mutation show poor viability and depletion of selected B and T cell populations. *Cell* 65, 1165-1175.

Scott, L. M., Tong, W., Levine, R. L., Scott, M. A., Beer, P. A., Stratton, M. R., Futreal, P. A., Erber, W. N., McMullin, M. F., and Harrison, C. N. (2007). JAK2 exon 12 mutations in polycythemia vera and idiopathic erythrocytosis. *New England Journal of Medicine* 356, 459-468.

Selleri, C., Sato, T., Del Vecchio, L., Luciano, L., Barrett, A. J., Rotoli, B., Young, N. S., and Maciejewski, J. P. (1997). Involvement of Fas-mediated apoptosis in the inhibitory effects of interferon-alpha in chronic myelogenous leukemia. *Blood* 89, 957-964.

Shide, K., Kameda, T., Shimoda, H., Yamaji, T., Abe, H., Kamiunten, A., Sekine, M., Hidaka, T., Katayose, K., Kubuki, Y., Yamamoto, S., Miike, T., Iwakiri, H., Hasuike, S., Nagata, K., Marutsuka, K., Iwama, A., Matsuda, T., Kitanaka, A., and Shimoda, K. (2012). TET2 is essential for survival and hematopoietic stem cell homeostasis. *Leukemia*

Silver, R. T., Vandris, K., and Goldman, J. J. (2011). Recombinant interferon-alpha may retard progression of early primary myelofibrosis: a preliminary report. *Blood* 117, 6669-6672.

Simonsson, B., Gedde-Dahl, T., Markevarn, B., Remes, K., Stentoft, J., Almqvist, A., Bjoreman, M., Flogegard, M., Koskenvesa, P., Lindblom, A., Malm, C., Mustjoki, S., Myhr-Eriksson, K., Ohm, L., Rasanen, A., Sinisalo, M., Sjalander, A., Stromberg, U., Bjerrum, O. W., Ehrencrona, H., Gruber, F., Kairisto, V., Olsson, K., Sandin, F., Nagler, A., Nielsen, J. L., Hjorth-Hansen, H., and Porkka, K. (2011). Combination of pegylated IFN-alpha2b with imatinib increases molecular response rates in patients with low- or intermediate-risk chronic myeloid leukemia. *Blood* 118, 3228-3235.

Socolovsky, M., Fallon, A. E., Wang, S., Brugnara, C., and Lodish, H. F. (1999). Fetal anemia and apoptosis of red cell progenitors in Stat5a-/-5b-/- mice: a direct role for Stat5 in Bcl-X(L) induction. *Cell* 98, 181-191.

Stegelmann, F., Bullinger, L., Schlenk, R. F., Paschka, P., Griesshammer, M., Blersch, C., Kuhn, S., Schauer, S., Dohner, H., and Dohner, K. (2011). DNMT3A mutations in myeloproliferative neoplasms. *Leukemia* 25, 1217-1219.

Stier, S., Cheng, T., Dombkowski, D., Carlesso, N., and Scadden, D. T. (2002). Notch1 activation increases hematopoietic stem cell self-renewal in vivo and favors lymphoid over myeloid lineage outcome. *Blood* 99, 2369-2378.

Swerdlow, S. H. (2008). WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues (Lyon, France: International Agency for Research on Cancer).

Tajbakhsh, S., Rocheteau, P., and Le Roux, I. (2009). Asymmetric cell divisions and asymmetric cell fates. *Annu Rev Cell Dev Biol* 25, 671-699.

Tefferi, A. (2010). Novel mutations and their functional and clinical relevance in myeloproliferative neoplasms: JAK2, MPL, TET2, ASXL1, CBL, IDH and IKZF1. *Leukemia* 24, 1128-1138.

Tefferi, A. (2012). JAK inhibitors for myeloproliferative neoplasms: clarifying facts from myths. *Blood* 119, 2721-2730.

Teglund, S., McKay, C., Schuetz, E., van Deursen, J. M., Stravopodis, D., Wang, D., Brown, M., Bodner, S., Grosveld, G., and Ihle, J. N. (1998). Stat5a and Stat5b proteins have essential and nonessential, or redundant, roles in cytokine responses. *Cell* 93, 841-850.

Tiedt, R., Hao-Shen, H., Sobas, M. A., Looser, R., Dirnhofer, S., Schwaller, J., and Skoda, R. C. (2008). Ratio of mutant JAK2-V617F to wild-type Jak2 determines the MPD phenotypes in transgenic mice. *Blood* 111, 3931-3940.

Ting, S. B., Deneault, E., Hope, K., Cellot, S., Chagraoui, J., Mayotte, N., Dorn, J. F., Laverdure, J.-P., Harvey, M., Hawkins, E. D., Russell, S. M., Maddox, P. S., Iscove, N. N., and Sauvageau, G. (2012). Asymmetric segregation and self-renewal of hematopoietic stem and progenitor cells with endocytic Ap2a2. *Blood* 119, 2510-2522.

Tybulewicz, V. L., Crawford, C. E., Jackson, P. K., Bronson, R. T., and Mulligan, R. C. (1991). Neonatal lethality and lymphopenia in mice with a homozygous disruption of the c-abl proto-oncogene. *Cell* 65, 1153-1163.

Vainchenker, W., Delhommeau, F., Constantinescu, S. N., and Bernard, O. A. (2011). New mutations and pathogenesis of myeloproliferative neoplasms. *Blood* 118, 1723-1735.

Vardiman, J., and Hyjek, E. (2011). World health organization classification, evaluation, and genetics of the myeloproliferative neoplasm variants. *Hematology Am Soc Hematol Educ*

Program 2011, 250-256.

- Verstovsek, S., Kantarjian, H., Mesa, R. A., Pardanani, A. D., Cortes-Franco, J., Thomas, D. A., Estrov, Z., Fridman, J. S., Bradley, E. C., Erickson-Viitanen, S., Vaddi, K., Levy, R., and Tefferi, A. (2010). Safety and efficacy of INCB018424, a JAK1 and JAK2 inhibitor, in myelofibrosis. *N Engl J Med* 363, 1117-1127.
- Verstovsek, S., Mesa, R. A., Gotlib, J., Levy, R. S., Gupta, V., DiPersio, J. F., Catalano, J. V., Deininger, M., Miller, C., and Silver, R. T. (2012). A double-blind, placebo-controlled trial of ruxolitinib for myelofibrosis. *New England Journal of Medicine* 366, 799-807.
- Walz, C., Ahmed, W., Lazarides, K., Betancur, M., Patel, N., Hennighausen, L., Zaleskas, V. M., and Van Etten, R. A. (2012). Essential role for Stat5a/b in myeloproliferative neoplasms induced by BCR-ABL1 and JAK2V617F in mice. *Blood* 119, 3550-3560.
- Wang, Y., Fiskus, W., Chong, D. G., Buckley, K. M., Natarajan, K., Rao, R., Joshi, A., Balusu, R., Koul, S., Chen, J., Savoie, A., Ustun, C., Jillella, A. P., Atadja, P., Levine, R. L., and Bhalla, K. N. (2009). Cotreatment with panobinostat and JAK2 inhibitor TG101209 attenuates JAK2V617F levels and signaling and exerts synergistic cytotoxic effects against human myeloproliferative neoplastic cells. *Blood* 114, 5024-5033.
- Warner, J. K., Wang, J. C., Hope, K. J., Jin, L., and Dick, J. E. (2004). Concepts of human leukemic development. *Oncogene* 23, 7164-7177.
- Wilks, A. F., Harpur, A. G., Kurban, R. R., Ralph, S. J., Zurcher, G., and Ziemiecki, A. (1991). Two novel protein-tyrosine kinases, each with a second phosphotransferase-related catalytic domain, define a new class of protein kinase. *Mol Cell Biol* 11, 2057-2065.
- Yan, D., Hutchison, R. E., and Mohi, G. (2012). Critical requirement for Stat5 in a mouse model of polycythemia vera. *Blood* 119, 3539-3549.
- Yoshida, K., Sanada, M., Shiraishi, Y., Nowak, D., Nagata, Y., Yamamoto, R., Sato, Y., Sato-Otsubo, A., Kon, A., Nagasaki, M., Chalkidis, G., Suzuki, Y., Shiosaka, M., Kawahata, R., Yamaguchi, T., Otsu, M., Obara, N., Sakata-Yanagimoto, M., Ishiyama, K., Mori, H., Nolte, F., Hofmann, W. K., Miyawaki, S., Sugano, S., Haferlach, C., Koeffler, H. P., Shih, L. Y., Haferlach, T., Chiba, S., Nakauchi, H., Miyano, S., and Ogawa, S. (2011). Frequent pathway mutations of splicing machinery in myelodysplasia. *Nature* 478, 64-69.
- Yoshihara, H., Arai, F., Hosokawa, K., Hagiwara, T., Takubo, K., Nakamura, Y., Gomei, Y., Iwasaki, H., Matsuoka, S., Miyamoto, K., Miyazaki, H., Takahashi, T., and Suda, T. (2007). Thrombopoietin/MPL signaling regulates hematopoietic stem cell quiescence and interaction with the osteoblastic niche. *Cell Stem Cell* 1, 685-697.
- Zamora, L., Espinet, B., Florensa, L., Besses, C., Bellosillo, B., and Sole, F. (2005). Clonality analysis by HUMARA assay in Spanish females with essential thrombocythemia and polycythemia vera. *Haematologica* 90, 259-261.
- Zhang, B., Strauss, A. C., Chu, S., Li, M., Ho, Y., Shiang, K. D., Snyder, D. S., Huettner, C. S., Shultz, L., Holyoake, T., and Bhatia, R. (2010). Effective targeting of quiescent chronic myelogenous leukemia stem cells by histone deacetylase inhibitors in combination with imatinib mesylate. *Cancer Cell* 17, 427-442.
- Zhang, J., Niu, C., Ye, L., Huang, H., He, X., Tong, W. G., Ross, J., Haug, J., Johnson, T., Feng, J. Q., Harris, S., Wiedemann, L. M., Mishina, Y., and Li, L. (2003). Identification of the haematopoietic stem cell niche and control of the niche size. *Nature* 425, 836-841.
- Zhang, S. J., Rampal, R., Manshouri, T., Patel, J., Mensah, N., Kayserian, A., Hricik, T., Heguy, A., Hedvat, C., Gonen, M., Kantarjian, H., Levine, R. L., Abdel-Wahab, O., and Verstovsek, S. (2012). Genetic analysis of patients with leukemic transformation of myeloproliferative neoplasms shows recurrent SRSF2 mutations that are associated with adverse outcome. *Blood* 119, 4480-4485.
- Zuber, J., Radtke, I., Pardee, T. S., Zhao, Z., Rappaport, A. R., Luo, W., McCurrach, M. E., Yang, M. M., Dolan, M. E., Kogan, S. C., Downing, J. R., and Lowe, S. W. (2009). Mouse models of human AML accurately predict chemotherapy response. *Genes Dev* 23, 877-889.

Figure Legends

Figure 1. Schematic diagram of hematopoietic stem cell regulation by signaling pathways. Several ligands (blue box) and their cognate receptors (white box) implicated in the regulation of hematopoietic stem cell (HSC) quiescence and self-renewal within the HSC osteoblastic niche are shown. Interferon alpha and interferon gamma have been demonstrated to induce quiescent HSC into cell cycle.

Figure 2. Schematic representation of cytokine signal transduction. Ligand binding by erythropoietin (EPO) to its cognate EPO receptor results in receptor dimerisation, cross phosphorylation and activation of Jak2 molecules associated with the cytoplasmic region of the EPO receptor. Phosphorylation of Stat5 molecules by activated Jak2 kinases, leads to nuclear translocation of Stat5 molecules and binding to gene promoter regions with initiation of gene transcription. RNA Pol; RNA polymerase.

Figure 3. Schematic representation of intracellular signaling molecules important in the development of myeloproliferative disorder phenotypes. Stat5 appears to be required for erythroid and granulocytic myeloproliferation associated with *JAK2V617F* and *BCR-ABL1* mutations, respectively. Endogenous Jak2 appears to be important for initiation of B lineage acute lymphoblastic leukaemia (B-ALL) associated with *BCR-ABL1* mutations.

Figure 4. Schematic diagram of cell populations in chronic myeloid leukaemia (CML), and their sensitivity to the tyrosine kinase inhibitor, imatinib. Dotted arrows indicate strategies proposed to target quiescent leukaemic stem cells (LSC), such as inducing quiescent LSC into cell cycle or targeting non-BCR-ABL1 signaling pathways in the LSC.

Osteoblast HSC niche

Figure 2

Figure 3

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Ng, Ashley P.

Title:

Hematopoietic stem cells, progenitor cells and leukemic stem cells in adult myeloproliferative neoplasms

Date:

2012

Citation:

Ng, A. P. (2012). Hematopoietic stem cells, progenitor cells and leukemic stem cells in adult myeloproliferative neoplasms. Leukemia and Lymphoma. doi: 10.3109/10428194.2012.734615

Publication Status:

In Press

Persistent Link:

<http://hdl.handle.net/11343/32701>