**The Joint Distribution of the Surplus
Prior to Ruin and the Deficit at Ruin
in Some Sparre Andersen Models**

by

David C M Dickson
The University of Melbourne

and

Steve Drekić
University of Waterloo

RESEARCH PAPER NUMBER 108

AUGUST 2003

Centre for Actuarial Studies
Department of Economics
The University of Melbourne
Victoria 3010
Australia

The joint distribution of the surplus prior to ruin and the deficit at ruin in some Sparre Andersen models

David C M Dickson and Steve Drekic

Abstract

In the Sparre Andersen risk model, we derive a general expression for $h(u, x, y)$, the joint density function of the surplus prior to ruin and the deficit at ruin when the initial surplus is u . This density function is expressed in terms of the corresponding density function when the initial surplus is 0. We apply a known result for $h(0, x, y)$ in the situation when claim inter-arrival times follow a generalised Erlang distribution to derive expressions for $h(u, x, y)$ when individual claims have a phase-type(m) distribution, $m \in \mathbb{Z}^+$. We also consider the case when claim inter-arrival times follow a phase-type(2) distribution and derive an expression for $h(0, x, y)$. Finally, we illustrate our results with examples.

1 Introduction

In this paper, we consider the Sparre Andersen risk model. Thus we assume that the time until the first claim and the times between claims form a sequence of independent and identically distributed (i.i.d.) continuous random variables, $\{T_i\}_{i=1}^\infty$, with common density function k . Let $\{X_i\}_{i=1}^\infty$ be a sequence of i.i.d. random variables, where X_i denotes the amount of the i th claim. Let $P(x) = 1 - \bar{P}(x) = \Pr(X_1 \leq x)$, $x \geq 0$, denote the distribution function of X_1 . We assume that $P(0) = 0$ and that X_1 is a continuous random variable with density function p . We further assume that the sequences $\{T_i\}_{i=1}^\infty$ and $\{X_i\}_{i=1}^\infty$ are independent. Let $\kappa = E(X_1)$ and $\mu = E(T_1)$.

Let c denote the insurer's premium income per unit time, which we assume to be received continuously. We define the probability of ultimate ruin for

this risk process by

$$\psi(u) = \Pr \left(u + \sum_{i=1}^n (cT_i - X_i) < 0 \text{ for some } n, n = 1, 2, 3, \dots \right)$$

and define $\delta(u) = 1 - \psi(u)$. It is well known (see, for example, Rolski et al (1999, Section 6.5)) that $\delta(u) = \Pr(L \leq u)$ where the random variable L denotes the maximum of the aggregate loss process. We assume that $c\mu - \kappa > 0$, so that $\psi(u) < 1$. The implied loading factor is $\theta = c\mu/\kappa - 1$.

Let $\{U(t)\}_{t \geq 0}$ denote the surplus process starting from $U(0) = u \geq 0$, and define the time of ruin, T , as

$$T = \begin{cases} \inf\{t: U(t) < 0\} \\ \infty \text{ if } U(t) \geq 0 \text{ for all } t > 0 \end{cases}$$

so that $\psi(u) = \Pr(T < \infty)$. Let T^- denote the time immediately prior to ruin.

We now define three defective distribution functions. First of all, let

$$\begin{aligned} H(u, x, y) &= \int_0^x \int_0^y h(u, r, s) ds dr \\ &= \Pr(U(T^-) \leq x, |U(T)| \leq y, T < \infty \mid U(0) = u) \end{aligned}$$

denote the joint distribution function of the surplus prior to ruin and the deficit at ruin, where h denotes the joint density function. Then we can define the marginal distribution functions

$$F(u, x) = \lim_{y \rightarrow \infty} H(u, x, y) \quad \text{and} \quad G(u, y) = \lim_{x \rightarrow \infty} H(u, x, y)$$

so that F is the distribution function of the surplus prior to ruin, and G is the distribution function of the deficit at ruin. Let their respective defective density functions be denoted by

$$f(u, x) = \frac{d}{dx} F(u, x) \quad \text{and} \quad g(u, y) = \frac{d}{dy} G(u, y).$$

It is convenient to introduce the proper density function of the deficit at ruin when $u = 0$, which we denote γ , and define by $\gamma(y) = g(0, y)/\psi(0)$. Finally, define the function ϕ by

$$\phi(u) = E[w(U(T^-), |U(T)|) I(T < \infty) \mid U(0) = u] \quad (1.1)$$

where w is an arbitrary bounded function and $I(A)$ denotes the indicator of an event A .

The remainder of the paper is as follows. In Section 2, we provide a general solution for $H(u, x, y)$, and consequently its joint density function, $h(u, x, y)$. In Section 3, we consider the case in which the density function k has a generalised Erlang distribution (as considered by Gerber and Shiu (2003a, 2003b)) in combination with claim amount density function p being phase-type(m), $m \in \mathbb{Z}^+$. In Section 4, we consider the case when the density function k is phase-type(2), or, equivalently, Coxian(2). As the phase-type(2) class is not a special case of the generalised Erlang family of distributions, the approach we take in Section 4 is different to that of Section 3. In fact, our approach leads to a much simpler method of finding $h(u, x, y)$ than that of Cheng and Tang (2003) who consider the situation when k is an Erlang(2) density function, a density function which belongs to both the phase-type(2) and generalised Erlang classes. In Section 5, we give some examples.

2 A formula for $H(u, x, y)$

To derive an expression for $H(u, x, y)$, we apply arguments given in Section 6.5 of Rolski et al (1999). We consider first the case when $u \geq x$. In order for the surplus immediately prior to ruin to be less than or equal to x , the surplus cannot fall below 0 on the first occasion that it drops below its initial level u . Hence it follows that

$$H(u, x, y) = \int_0^u g(0, z)H(u - z, x, y)dz = \psi(0) \int_0^u \gamma(z)H(u - z, x, y)dz.$$

Taking partial derivatives with respect to x and y yields

$$h(u, x, y) = \psi(0) \int_0^u \gamma(z)h(u - z, x, y)dz.$$

In the case when $0 \leq u < x$, it is possible for ruin to occur at the time the surplus first falls below its initial level u , and for the surplus prior to ruin to be less than or equal to x , and for the deficit at ruin to be less than or equal to y . The probability of this event is

$$J(u, x, y) = \int_0^{x-u} \int_u^{u+y} h(0, r, s)dsdr \quad (2.1)$$

as the event is equivalent to ruin occurring from initial surplus 0 with a surplus immediately prior to ruin less than or equal to $x - u$ and a deficit at ruin between u and $u + y$. Hence, for $0 \leq u < x$, we have

$$H(u, x, y) = \psi(0) \int_0^u \gamma(z)H(u - z, x, y)dz + J(u, x, y)$$

and

$$h(u, x, y) = \psi(0) \int_0^u \gamma(z) h(u - z, x, y) dz + h(0, x - u, u + y).$$

Therefore, for $u \geq 0$, we have

$$H(u, x, y) = \psi(0) \int_0^u \gamma(z) H(u - z, x, y) dz + Z(u, x, y)$$

where $Z(u, x, y) = I(u < x) J(u, x, y)$, and

$$h(u, x, y) = \psi(0) \int_0^u \gamma(z) h(u - z, x, y) dz + I(u < x) h(0, x - u, u + y).$$

Let

$$H^*(s, x, y) = \int_0^\infty e^{-su} H(u, x, y) du$$

be the Laplace transform of H , with similar notation for Laplace transforms of other functions. Then by basic properties of Laplace transforms,

$$H^*(s, x, y) = \frac{Z^*(s, x, y)}{1 - \psi(0)\gamma^*(s)}.$$

From Rolski et al (1999, Section 6.5), L has a compound geometric distribution, and so its Laplace-Stieltjes transform is given by

$$E(e^{-sL}) = \int_0^\infty e^{-su} d\delta(u) = \frac{\delta(0)}{1 - \psi(0)\gamma^*(s)}$$

which implies that

$$H^*(s, x, y) = \frac{E(e^{-sL})}{\delta(0)} Z^*(s, x, y).$$

Since the product of two transforms is the transform of a convolution, it immediately follows that

$$\begin{aligned} H(u, x, y) &= \frac{1}{\delta(0)} \int_0^u Z(u - z, x, y) d\delta(z) \\ &= \frac{1}{\delta(0)} \int_0^u I(u - z < x) J(u - z, x, y) d\delta(z). \end{aligned} \quad (2.2)$$

Hence the above equation provides a means of finding $H(u, x, y)$ provided that we know both δ and J . In a similar fashion, we also have the corresponding analogue for $h(u, x, y)$, namely

$$h(u, x, y) = \frac{1}{\delta(0)} \int_0^u I(u - z < x) h(0, x - u + z, u - z + y) d\delta(z). \quad (2.3)$$

We remark that equation (6.4) of Dickson and Hipp (2000) for $G(u, y)$ can be obtained from equation (2.2) as

$$\lim_{x \rightarrow \infty} J(u, x, y) = G(0, u + y) - G(0, u).$$

See also Willmot and Lin (1998, p.92). Moreover, Dufresne and Gerber (1988) show that in the classical Poisson risk model

$$h(0, x, y) = f(0, x) \frac{p(x + y)}{\bar{P}(x)}. \quad (2.4)$$

As the arguments underlying this formula equally apply in the Sparre Andersen model, finding J reduces to finding the density function of the surplus prior to ruin when the initial surplus is 0.

We end this section by remarking that in the classical model $f(0, x) = (\lambda/c)\bar{P}(x)$, so that $h(0, x, y) = (\lambda/c)p(x + y)$ (see, for example, Dufresne and Gerber (1988)). Hence, for $0 \leq u < x$, equation (2.3) yields

$$h(u, x, y) = \frac{1}{\delta(0)} \int_0^u \frac{\lambda}{c} p(x + y) d\delta(z) = \frac{\lambda}{c} p(x + y) \frac{1 - \psi(u)}{1 - \psi(0)}, \quad (2.5)$$

and for $u > x$ it yields

$$h(u, x, y) = \frac{1}{\delta(0)} \int_{u-x}^u \frac{\lambda}{c} p(x + y) d\delta(z) = \frac{\lambda}{c} p(x + y) \frac{\psi(u - x) - \psi(u)}{1 - \psi(0)}, \quad (2.6)$$

results first given by Dickson (1992).

3 Generalised Erlang inter-claim times

In this section we consider the situation where k has a generalised Erlang density function. In particular, we assume that k is the density function of $T_1 = \sum_{j=1}^n W_j$ where $\{W_j\}_{j=1}^n$ are n independent, exponentially distributed random variables with $E(W_j) = 1/\lambda_j$. This generalised Erlang distribution belongs to the phase-type(n) family of distributions and as such, k can be written as

$$k(t) = \underline{\beta} \exp\{tS\} \underline{S}^o, \quad t > 0 \quad (3.1)$$

where $\underline{\beta} = (1, 0, 0, \dots, 0, 0)$ is a $1 \times n$ row vector,

$$S = \begin{bmatrix} -\lambda_1 & \lambda_1 & 0 & \dots & 0 & 0 \\ 0 & -\lambda_2 & \lambda_2 & \dots & 0 & 0 \\ 0 & 0 & -\lambda_3 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & -\lambda_{n-1} & \lambda_{n-1} \\ 0 & 0 & 0 & \dots & 0 & -\lambda_n \end{bmatrix}$$

is an $n \times n$ transition rate matrix, and $\underline{S}^o = (0, 0, 0, \dots, 0, \lambda_n)'$ is an $n \times 1$ column vector. Also, the matrix exponential in equation (3.1) is defined by

$$\exp\{tS\} = \sum_{j=0}^{\infty} \frac{t^j}{j!} S^j.$$

For a detailed treatment of phase-type distributions, we refer the reader to Neuts (1981), Latouche and Ramaswami (1999), Rolski et al (1999), and Asmussen (2000).

For this choice of k , Gerber and Shiu (2003b) show that

$$h(0, x, y) = \mathcal{C} p(x + y) \sum_{j=1}^n a_j e^{-\rho_j x} \quad (3.2)$$

where $\mathcal{C} = \prod_{r=1}^n (\lambda_r / c)$, $a_j = \prod_{r=1, r \neq j}^n (\rho_r - \rho_j)^{-1}$, and $\{\rho_r\}_{r=1}^n$ are the n roots in the right half of the complex plane (assumed to be distinct) satisfying the following equation for ξ :

$$\prod_{j=1}^n \left(1 - \frac{c}{\lambda_j} \xi\right) - p^*(\xi) = 0. \quad (3.3)$$

We remark that equation (3.3) may be alternatively expressed as $k^*(-c\xi)p^*(\xi) = 1$. Furthermore, it is clear from equation (3.3) that $\xi = 0$ is always one of these n roots. Substituting equation (3.2) into equation (2.3), we obtain

$$h(u, x, y) = \frac{\mathcal{C}}{\delta(0)} p(x + y) \sum_{j=1}^n a_j e^{-\rho_j(x-u)} \int_0^u I(u - z < x) e^{-\rho_j z} d\delta(z). \quad (3.4)$$

Let us now assume that claim amounts have a general phase-type(m) distribution, $m \in \mathbb{Z}^+$, with density function $p(x) = \underline{\alpha} \exp\{xA\} \underline{A}^o$ and corresponding distribution function $P(x) = 1 - \underline{\alpha} \exp\{xA\} \underline{e}_m$ where $\underline{e}_m$ denotes

an $m \times 1$ column vector of ones and $\underline{A}^o = -A\underline{e}_m$. From Theorem 4.4 in Asmussen (2000), the distribution function of the maximal aggregate loss L in the general Sparre Andersen model with phase-type(m) distributed claim amounts is given by

$$\delta(u) = 1 - \underline{\eta} \exp\{u(A + \underline{A}^o \underline{\eta})\} \underline{e}_m, \quad u \geq 0 \quad (3.5)$$

where the defective row vector $\underline{\eta}$ is the unique solution satisfying the relation

$$\underline{\eta} = (1/c) \underline{\alpha} \int_0^\infty \exp\{t(A + \underline{A}^o \underline{\eta})\} k(t/c) dt. \quad (3.6)$$

However, since k has the phase-type form given by equation (3.1), an explicit solution for $\underline{\eta}$ may actually be obtained. Omitting the tedious mathematical details, it can be verified that

$$\underline{\eta} = \underline{C} \underline{\alpha} \sum_{j=1}^n a_j (\rho_j I_m - A)^{-1} \quad (3.7)$$

satisfies equation (3.6) where I_m denotes the $m \times m$ identity matrix. Furthermore, it follows from equation (3.5) and the fact that $\underline{A}^o = -A\underline{e}_m$ that the density function of L is given by $\delta'(u) = \delta(0) \underline{\eta} \exp\{u(A + \underline{A}^o \underline{\eta})\} \underline{A}^o$, $u > 0$, with $\delta(0) = 1 - \underline{\eta} \underline{e}_m$.

Consider first the case when $0 \leq u < x$. In this case, $I(u - z < x) = 1$ for all $z \in [0, u]$, and so equation (3.4) becomes

$$\begin{aligned} h(u, x, y) &= \frac{c}{\delta(0)} p(x + y) \sum_{j=1}^n a_j e^{-\rho_j(x-u)} \left[\delta(0) + \int_0^u e^{-\rho_j z} \delta'(z) dz \right] \\ &= \underline{C} \underline{\alpha} \exp\{(x + y)A\} \underline{A}^o \sum_{j=1}^n a_j e^{-\rho_j(x-u)} \left(1 + \int_0^u e^{-\rho_j z} \underline{\eta} \exp\{zD\} \underline{A}^o dz \right) \end{aligned} \quad (3.8)$$

where we define $D = A + \underline{A}^o \underline{\eta}$ for the sake of notational convenience. However, note that

$$\begin{aligned} \int_0^u e^{-\rho_j z} \underline{\eta} \exp\{zD\} \underline{A}^o dz &= \underline{\eta} \left(\int_0^u \exp\{zD\} e^{-\rho_j z} dz \right) \underline{A}^o \\ &= \underline{\eta} \left(\int_0^u \exp\{z(D - \rho_j I_m)\} dz \right) \underline{A}^o \\ &= \underline{\eta} B_j^{-1} (\exp\{u B_j\} - I_m) \underline{A}^o \end{aligned} \quad (3.9)$$

where we define $B_j = D - \rho_j I_m$. Substituting equation (3.9) into equation (3.8) immediately yields for $0 \leq u < x$

$$h(u, x, y) = C \underline{\alpha} \exp\{(x+y)A\} \underline{A}^o \sum_{j=1}^n a_j e^{-\rho_j(x-u)} [1 + \underline{\eta} B_j^{-1} (\exp\{u B_j\} - I_m) \underline{A}^o]. \quad (3.10)$$

On the other hand, if $u > x$, then $I(u - z < x) = 1$ for all $z \in (u - x, u]$, and so equation (3.4) becomes

$$h(u, x, y) = C \underline{\alpha} \exp\{(x+y)A\} \underline{A}^o \sum_{j=1}^n a_j e^{\rho_j(u-x)} \int_{u-x}^u e^{-\rho_j z} \underline{\eta} \exp\{z D\} \underline{A}^o dz. \quad (3.11)$$

In a similar fashion which led to equation (3.9), we have

$$\int_{u-x}^u e^{-\rho_j z} \underline{\eta} \exp\{z D\} \underline{A}^o dz = \underline{\eta} B_j^{-1} [\exp\{u B_j\} - \exp\{(u-x) B_j\}] \underline{A}^o.$$

Therefore, for $u > x$, equation (3.11) becomes after some straightforward manipulation

$$h(u, x, y) = C \underline{\alpha} \exp\{(x+y)A\} \underline{A}^o \sum_{j=1}^n a_j \underline{\eta} B_j^{-1} [e^{-\rho_j x} \exp\{u D\} - \exp\{(u-x) D\}] \underline{A}^o. \quad (3.12)$$

If we now integrate equations (3.10) and (3.12) from $y = 0$ to $y = \infty$, we immediately obtain the following marginal density function of the surplus prior to ruin. For $0 \leq u < x$,

$$f(u, x) = C \underline{\alpha} \exp\{xA\} \underline{e}_m \sum_{j=1}^n a_j e^{-\rho_j(x-u)} [1 + \underline{\eta} B_j^{-1} (\exp\{u B_j\} - I_m) \underline{A}^o], \quad (3.13)$$

and for $u > x$

$$f(u, x) = C \underline{\alpha} \exp\{xA\} \underline{e}_m \sum_{j=1}^n a_j \underline{\eta} B_j^{-1} [e^{-\rho_j x} \exp\{u D\} - \exp\{(u-x) D\}] \underline{A}^o. \quad (3.14)$$

The marginal density function of the deficit at ruin can also be obtained by integrating equation (3.12) from $x = 0$ to $x = u$ followed by integrating equation (3.10) from $x = u$ to $x = \infty$. This is equivalent to integrating equation (3.4) from $x = 0$ to $x = \infty$. After a fair bit of matrix algebra and simplification, it can be shown that

$$g(u, y) = \underline{\eta} \exp\{u D\} \exp\{y A\} \underline{A}^o, \quad y > 0, \quad (3.15)$$

a result which agrees with that of Drekić et al (2001, p.12).

We conclude this section with a few remarks. First of all, by comparing equations (3.13) and (3.14) as $x \rightarrow u^-$ and $x \rightarrow u^+$, respectively, the density function $f(u, x)$ has a jump discontinuity at $x = u$ unless $\sum_{j=1}^n a_j = 0$, in which case $f(u, x)$ is strictly continuous. Furthermore, in the classical model (i.e. when $k(t) = \lambda e^{-\lambda t}$), we have $n = 1$, $C = \lambda/c$, $\rho_1 = 0$, $a_1 = 1$, $B_1 = D$, and $\eta = -(\lambda/c)\alpha A^{-1}$. With these values, it is a straightforward exercise to show that (i) equations (3.13) and (3.14) reduce to give equations (3.1) and (3.2), respectively, in Dickson (1992), and (ii) equation (3.15) agrees with the result given in Drekić et al (2001, p.13).

Finally we remark that the most obvious, and we believe useful, application of the generalised Erlang distribution is the Erlang distribution itself. Whilst there are many distributions in the class, the class is restrictive in the sense that the generalised Erlang density is a combination of densities and the parameters $\{\lambda_j\}_{j=1}^n$ determine the coefficients of the densities in the combination.

4 Phase-type(2) inter-claim times

In this section, we consider the case when k is a phase-type(2) density function. In what follows, we adopt the notation and approach of Dickson and Hipp (2000) who observed that when k is a phase-type(2) density function, it satisfies the differential equation

$$k(t) + A_1 k'(t) + A_2 k''(t) = 0 \quad (4.1)$$

for $t > 0$, where A_1 and A_2 are constants with $A_2 > 0$. Cheng and Tang (2003) considered the Erlang(2, σ) case with $k(t) = \sigma^2 t e^{-\sigma t}$, so that $A_1 = 2/\sigma$ and $A_2 = 1/\sigma^2$.

We begin by deriving an expression for $h(0, x, y)$ which is required to use equation (2.3). In order to do so, we use the function ϕ defined in equation (1.1) with $w(x_1, x_2) = I(x_1 \leq x)I(x_2 \leq y)$. However, we will initially work with a general function w . We start from equation (2.3) of Cheng and Tang (2003) which applies to any Sparre Andersen model:

$$\begin{aligned} c\phi(u) &= \int_u^\infty k\left(\frac{s-u}{c}\right) \int_0^s \phi(s-x)p(x)dxds \\ &\quad + \int_u^\infty k\left(\frac{s-u}{c}\right) \int_s^\infty w(s, x-s)p(x)dxds \\ &= \int_u^\infty k\left(\frac{s-u}{c}\right) [\tau(s) + \omega(s)] ds \end{aligned} \quad (4.2)$$

where $\tau(u) = \int_0^u \phi(u-x)p(x)dx$ and $\omega(u) = \int_u^\infty w(u, x-u)p(x)dx$. Differentiation of equation (4.2) then yields

$$c\phi'(u) = -k(0)[\tau(u) + \omega(u)] - \frac{1}{c} \int_u^\infty k' \left(\frac{s-u}{c} \right) [\tau(s) + \omega(s)] ds$$

and hence

$$\begin{aligned} c\phi''(u) &= -k(0)[\tau'(u) + \omega'(u)] + \frac{1}{c}k'(0)[\tau(u) + \omega(u)] \\ &\quad + \frac{1}{c^2} \int_u^\infty k'' \left(\frac{s-u}{c} \right) [\tau(s) + \omega(s)] ds. \end{aligned}$$

Following Dickson and Hipp (2000, p.152), we use equation (4.1) to obtain

$$\begin{aligned} &A_2c^2\phi''(u) - A_1c\phi'(u) + \phi(u) \\ &= A_2k'(0)[\tau(u) + \omega(u)] - A_2ck(0)[\tau'(u) + \omega'(u)] + k(0)A_1[\tau(u) + \omega(u)] \\ &= \tau(u) + \omega(u) - A_2ck(0)[\tau'(u) + \omega'(u)] \end{aligned}$$

since $A_1k(0) + A_2k'(0) = 1$ (see Dickson and Hipp (2000, equation (2.1))). Taking the Laplace transform of this equation, and noting that τ is a convolution, we obtain

$$\begin{aligned} &A_2c^2[s^2\phi^*(s) - s\phi(0) - \phi'(0)] - A_1c[s\phi^*(s) - \phi(0)] + \phi^*(s) \\ &= \phi^*(s)p^*(s) + \omega^*(s) - A_2ck(0)[s\phi^*(s)p^*(s) + s\omega^*(s) - \omega(0)] \end{aligned}$$

so that

$$\phi^*(s) = \frac{A_2c^2[s\phi(0) + \phi'(0)] - A_1c\phi(0) + \omega^*(s) - A_2ck(0)[s\omega^*(s) - \omega(0)]}{A_2c^2s^2 - A_1cs + 1 - p^*(s) + A_2ck(0)sp^*(s)},$$

or, equivalently

$$s\phi^*(s) = \frac{A_2c^2[s\phi(0) + \phi'(0)] - A_1c\phi(0) + \omega^*(s) - A_2ck(0)[s\omega^*(s) - \omega(0)]}{A_2c^2s - A_1c + \kappa q^*(s) + A_2ck(0)p^*(s)} \quad (4.3)$$

where we define $q^*(s) = [1 - p^*(s)]/(\kappa s)$. Dickson and Hipp (2000, Section 3) show that there is a unique positive number s_0 such that

$$A_2c^2s_0 - A_1c + \kappa q^*(s_0) + A_2ck(0)p^*(s_0) = 0.$$

As s_0 is a zero of the denominator in equation (4.3), it must also be a zero of the numerator. Making use of this observation, we write

$$s\phi^*(s) = \frac{A_2c^2(s - s_0)\phi(0) + \omega^*(s) - \omega^*(s_0) - A_2ck(0)[s\omega^*(s) - s_0\omega^*(s_0)]}{A_2c^2s - A_1c + \kappa q^*(s) + A_2ck(0)p^*(s)}.$$

If we now set $s = 0$ in the denominator, we get $-A_1c + \kappa + A_2ck(0) = \kappa - c\mu \neq 0$ (see Dickson and Hipp (2000, equation (2.3))). Hence the numerator must be zero with $s = 0$ as $\lim_{s \rightarrow 0^+} s\phi^*(s) = 0$. Thus

$$-A_2c^2s_0\phi(0) + \omega^*(0) - \omega^*(s_0) + A_2ck(0)s_0\omega^*(s_0) = 0$$

giving

$$\phi(0) = \frac{1}{A_2c^2s_0}[\omega^*(0) - \omega^*(s_0)] + \frac{k(0)}{c}\omega^*(s_0). \quad (4.4)$$

We remark that equation (4.4) generalises Corollary 2.3 of Cheng and Tang (2003) as their r_2 is our s_0 .

Now let $w(x_1, x_2) = I(x_1 \leq x)I(x_2 \leq y)$, so that $\phi(0) = H(0, x, y)$. It then follows that

$$\begin{aligned} \omega(u) &= \int_u^\infty I(u \leq x)I(r - u \leq y)p(r)dr \\ &= I(u \leq x) \int_u^\infty I(r \leq u + y)p(r)dr \\ &= I(u \leq x) [\bar{P}(u) - \bar{P}(u + y)]. \end{aligned}$$

This gives

$$\omega^*(0) = \int_0^\infty I(u \leq x) [\bar{P}(u) - \bar{P}(u + y)] du = \int_0^x [\bar{P}(u) - \bar{P}(u + y)] du$$

and

$$\omega^*(s_0) = \int_0^x e^{-s_0u} [\bar{P}(u) - \bar{P}(u + y)] du,$$

thereby yielding

$$\begin{aligned} H(0, x, y) &= \frac{1}{A_2c^2s_0} \int_0^x [\bar{P}(u) - \bar{P}(u + y)] du \\ &\quad - \frac{1 - A_2ck(0)s_0}{A_2c^2s_0} \int_0^x e^{-s_0u} [\bar{P}(u) - \bar{P}(u + y)] du. \end{aligned} \quad (4.5)$$

Differentiating equation (4.5) with respect to x and then y gives

$$\begin{aligned} h(0, x, y) &= \frac{1}{A_2c^2s_0} \{p(x + y) - [1 - A_2ck(0)s_0]e^{-s_0x}p(x + y)\} \\ &= \frac{1}{A_2c^2s_0} (1 - e^{-s_0x})p(x + y) + \frac{k(0)}{c}e^{-s_0x}p(x + y). \end{aligned} \quad (4.6)$$

We can easily verify that

$$h(0, x, y) = f(0, x) \frac{p(x + y)}{\bar{P}(x)}$$

by first letting $y \rightarrow \infty$ in equation (4.5) and then differentiating the result with respect to x .

We remark that equation (4.6) generalises equation (3.9) of Cheng and Tang (2003) with $u = 0$, as their $k(0)$ is equal to 0. Also, note that equation (4.6) may be expressed in the form of equation (3.2) by setting $n = 2$, $\mathcal{C} = 1$, $\rho_1 = 0$, $\rho_2 = s_0$, $a_1 = (A_2 c^2 s_0)^{-1}$, and $a_2 = k(0)/c - (A_2 c^2 s_0)^{-1}$. Moreover, it is readily verified that $\xi = 0$ and $\xi = s_0$ are the two distinct roots in the right half of the complex plane satisfying $k^*(-c\xi)p^*(\xi) = 1$. Therefore, if claim amounts have a phase-type(m) distribution, then equations (3.4) through to (3.15) continue to hold (with constants assigned as above) in the case when inter-claim times are phase-type(2) distributed.

5 Examples

In this section, we present three examples to illustrate the results of the previous sections. The results in Examples 5.2 and 5.3 were obtained with the aid of the symbolic computational package *Mathematica*.

Example 5.1 *We start with a straightforward example for which the phase-type machinery is not required. Let $p(x) = \nu e^{-\nu x}$, $x > 0$, and let $k(t) = 4te^{-2t}$, $t > 0$ (i.e. exponential claim amounts and Erlang(2,2) inter-claim times). In this case, $k(0) = 0$ and $A_2 = 1/4$, so that equation (4.6) becomes*

$$h(0, x, y) = \frac{4\nu}{c^2 s_0} (1 - e^{-s_0 x}) e^{-\nu(x+y)}.$$

It is well known (see, for example, Rolski et al (1999, Section 6.5.1)) that

$$\delta(u) = 1 - (1 - R/\nu)e^{-Ru}$$

where R is the adjustment coefficient. Hence, for $0 \leq u < x$,

$$\begin{aligned} h(u, x, y) &= \frac{1}{\delta(0)} \int_0^u h(0, x - u + z, u - z + y) d\delta(z) \\ &= \frac{4\nu e^{-\nu(x+y)}}{c^2 s_0} \left[(1 - e^{-s_0(x-u)}) + \frac{\nu - R}{R} (1 - e^{-Ru}) \right. \\ &\quad \left. - \frac{\nu - R}{s_0 + R} e^{-s_0(x-u)} (1 - e^{-(s_0+R)u}) \right] \end{aligned}$$

and integrating out y we obtain

$$\begin{aligned} f(u, x) &= \frac{4e^{-\nu x}}{c^2 s_0} \left[(1 - e^{-s_0(x-u)}) + \frac{\nu - R}{R} (1 - e^{-Ru}) \right. \\ &\quad \left. - \frac{\nu - R}{s_0 + R} e^{-s_0(x-u)} (1 - e^{-(s_0+R)u}) \right]. \end{aligned}$$

Similarly, for $u > x$,

$$\begin{aligned} h(u, x, y) &= \frac{1}{\delta(0)} \int_{u-x}^u h(0, x-u+z, u-z+y) d\delta(z) \\ &= (\nu - R) \frac{4\nu e^{-\nu(x+y)}}{c^2 s_0} e^{-Ru} \left[\frac{1}{R} (e^{Rx} - 1) - \frac{e^{Rx} - e^{-s_0 x}}{s_0 + R} \right] \end{aligned}$$

and

$$f(u, x) = (\nu - R) \frac{4e^{-\nu x}}{c^2 s_0} e^{-Ru} \left[\frac{1}{R} (e^{Rx} - 1) - \frac{e^{Rx} - e^{-s_0 x}}{s_0 + R} \right].$$

We remark that these results can also be obtained from Example 1 of Li and Garrido (2003) by setting $n = 2$ there.

Example 5.2 Consider an inter-claim time distribution that is an equal mixture of two exponentials at rates $1/3$ and 1 respectively, so that

$$k(t) = \frac{1}{2}e^{-t} + \frac{1}{6}e^{-t/3}, \quad t > 0.$$

We remark that this distribution belongs to the phase-type(2) class but is not a member of the generalised Erlang family. Moreover, we have $\mu = 2$, $k(0) = 2/3$, $A_1 = 4$, and $A_2 = 3$. Suppose that the individual claim amount distribution is Erlang(3, 1.5), so that

$$p(x) = \frac{27}{16}x^2 e^{-1.5x}, \quad x > 0, \quad \text{and} \quad \bar{P}(x) = e^{-1.5x} \left(1 + \frac{3}{2}x + \frac{9}{8}x^2 \right), \quad x \geq 0.$$

We note that $\kappa = 2$. Written in phase-type form, we have $\underline{\alpha} = (1, 0, 0)$, $\underline{A}^o = (0, 0, 1.5)'$, and

$$A = \begin{bmatrix} -1.5 & 1.5 & 0 \\ 0 & -1.5 & 1.5 \\ 0 & 0 & -1.5 \end{bmatrix}.$$

Let $\theta = 0.1$ so that $c = 1.1$. Then we obtain (to 5 decimal places of accuracy) $s_0 = 0.79184$, $\underline{\eta} = (0.34458, 0.30566, 0.28019)$, and

$$D = \begin{bmatrix} -1.50000 & 1.50000 & 0 \\ 0 & -1.50000 & 1.50000 \\ 0.51687 & 0.45849 & -1.07972 \end{bmatrix}.$$

As D has one real eigenvalue (-0.05110) and two complex ones $(-2.0143 \pm 0.73357i)$, expressions for $h(u, x, y)$ will involve trigonometric functions. For

$0 \leq u < x$, we obtain

$$\begin{aligned} h(u, x, y) = & p(x+y) \{ 5 - (4.7147 + 0.21209e^{-0.79184x}) e^{-0.0511u} \\ & + 0.43824e^{-0.79184(x-u)} \\ & + (0.06263 + 0.032e^{-0.79184x}) e^{-2.0143u} \text{Cos}[0.73357u] \\ & + (0.04288 + 0.02663e^{-0.79184x}) e^{-2.0143u} \text{Sin}[0.73357u] \} \end{aligned}$$

and for $u > x$

$$\begin{aligned} h(u, x, y) = & p(x+y) \{ - (4.7147 + 0.21209e^{-0.79184x}) e^{-0.0511u} \\ & + 4.92682e^{-0.0511(u-x)} \\ & + (0.06263 + 0.032e^{-0.79184x}) e^{-2.0143u} \text{Cos}[0.73357u] \\ & + (0.04288 + 0.02663e^{-0.79184x}) e^{-2.0143u} \text{Sin}[0.73357u] \\ & - 0.09464e^{-2.0143(u-x)} \text{Cos}[0.73357(u-x)] \\ & - 0.06951e^{-2.0143(u-x)} \text{Sin}[0.73357(u-x)] \}. \end{aligned}$$

The density function $f(u, x)$ has the same form as the two equations above, except that $p(x+y)$ is replaced by $\bar{P}(x)$. Figures 1 and 2 display the shape of $f(u, x)$ for $u = 1$ and $u = 3$, respectively. The density functions are very similar to those illustrated by Dufresne and Gerber (1988) and Dickson (1992). Since $k(0) = 2/3 \neq 0$, we have $a_1 + a_2 \neq 0$. Based on our earlier remark at the end of Section 3, this implies that $f(u, x)$ possesses a jump discontinuity at $x = u$. This is indeed the case as evidenced by Figures 1 and 2, although it becomes more difficult to tell for larger values of u . Finally, the density function $g(u, y)$ is

$$\begin{aligned} g(u, y) = & (0.6925 + 0.71232y + 0.27792y^2) e^{-0.0511u-1.5y} \\ & - (0.27222 + 0.02459y - 0.30356y^2) e^{-2.0143u-1.5y} \text{Cos}[0.73357u] \\ & - (0.38035 - 0.55516y + 0.00296y^2) e^{-2.0143u-1.5y} \text{Sin}[0.73357u]. \end{aligned}$$

Example 5.3 Suppose inter-claim times have a generalised Erlang distribution with $n = 3$, $\lambda_1 = \lambda_2 = 0.5$, and $\lambda_3 = 2$. With these values, equation (3.1) simplifies to give

$$k(t) = \frac{2}{9}e^{-2t} - \frac{2}{9}e^{-0.5t} + \frac{1}{3}te^{-0.5t}, \quad t > 0$$

with $\mu = 4.5$. Let the individual claim amounts be distributed with phase-type representation $\underline{\alpha} = (0.1, 0.1, 0.3, 0.5)$, $\underline{A}^o = (0, 0, 0, 0.5)'$, and

$$A = \begin{bmatrix} -1/10 & 1/10 & 0 & 0 \\ 0 & -1/6 & 1/6 & 0 \\ 0 & 0 & -1/3 & 1/3 \\ 0 & 0 & 0 & -1/2 \end{bmatrix}.$$

Then the mean claim amount is $\kappa = 5.7$. Moreover, we have

$$p(x) = \frac{5}{112}e^{-x/10} - \frac{1}{40}e^{-x/6} + \frac{17}{70}e^{-x/3} - \frac{1}{80}e^{-x/2}, \quad x > 0,$$

and

$$\bar{P}(x) = \frac{25}{56}e^{-x/10} - \frac{3}{20}e^{-x/6} + \frac{51}{70}e^{-x/3} - \frac{1}{40}e^{-x/2}, \quad x \geq 0.$$

Let $\theta = 0.2$ so that $c = 1.52$. Then we obtain (to 5 decimal places of accuracy) $C = 0.14238$, $\rho_1 = 0$, $\rho_2 = 0.56407$, $\rho_3 = 1.29160$,

$$\underline{\eta} = (0.15407, 0.19102, 0.19612, 0.23601),$$

and

$$D = \begin{bmatrix} -0.10000 & 0.10000 & 0 & 0 \\ 0 & -0.16667 & 0.16667 & 0 \\ 0 & 0 & -0.33333 & 0.33333 \\ 0.07703 & 0.09551 & 0.09806 & -0.38199 \end{bmatrix}.$$

We note that D has real eigenvalues, namely -0.50122 , -0.27485 , -0.18112 , and -0.02480 . Therefore, for $0 \leq u < x$, we obtain

$$\begin{aligned} h(u, x, y) = & 0.14238p(x+y)[6.1611 + 0.00217e^{-0.50122u} - 0.23444e^{-0.27485u} \\ & + 0.09197e^{-0.18112u} - 4.6482e^{-0.0248u} + 1.1547e^{-1.2916(x-u)} - 2.8792e^{-0.56407(x-u)} \\ & + (0.00047e^{-0.50122u} - 0.03189e^{-0.27485u} + 0.00877e^{-0.18112u} \\ & - 0.0679e^{-0.0248u})e^{-1.2916x} - (0.00182e^{-0.50122u} - 0.13636e^{-0.27485u} \\ & + 0.03969e^{-0.18112u} - 0.34755e^{-0.0248u})e^{-0.56407x}] \end{aligned}$$

and for $u > x$

$$\begin{aligned} h(u, x, y) = & 0.14238p(x+y)[0.00217e^{-0.50122u} - 0.23444e^{-0.27485u} + 0.09197e^{-0.18112u} \\ & - 4.6482e^{-0.0248u} + 4.3685e^{-0.0248(u-x)} - 0.06106e^{-0.18112(u-x)} \\ & + 0.12998e^{-0.27485(u-x)} - 0.00083e^{-0.50122(u-x)} + (0.00047e^{-0.50122u} \\ & - 0.03189e^{-0.27485u} + 0.00877e^{-0.18112u} - 0.0679e^{-0.0248u})e^{-1.2916x} \\ & - (0.00182e^{-0.50122u} - 0.13636e^{-0.27485u} + 0.03969e^{-0.18112u} \\ & - 0.34755e^{-0.0248u})e^{-0.56407x}]. \end{aligned}$$

As we commented in Example 5.2, the density function $f(u, x)$ has the same form as the two equations above, except that $p(x+y)$ is replaced by $\bar{P}(x)$. Figures 3 and 4 display the shape of $f(u, x)$ for $u = 20$ and $u = 40$, respectively. Since $a_1 + a_2 + a_3 = 1.37257 - 2.43676 + 1.06419 = 0$, $f(u, x)$ is

strictly continuous for all $x > 0$, as evidenced by Figures 3 and 4. For the sake of completeness, the density function $g(u, y)$ is given by

$$\begin{aligned}
g(u, y) = & (0.00014e^{-0.50122u} - 0.01847e^{-0.27485u} + 0.01029e^{-0.18112u} + 0.07682e^{-0.0248u})e^{-y/10} \\
& - (0.00005e^{-0.50122u} - 0.0087e^{-0.27485u} + 0.01683e^{-0.18112u} + 0.01187e^{-0.0248u})e^{-y/6} \\
& + (0.00034e^{-0.50122u} + 0.05711e^{-0.27485u} - 0.00567e^{-0.18112u} + 0.01936e^{-0.0248u})e^{-y/3} \\
& - (0.00122e^{-0.50122u} + 0.00039e^{-0.27485u} - 0.00007e^{-0.18112u} + 0.00033e^{-0.0248u})e^{-y/2}.
\end{aligned}$$

References

- [1] Asmussen, S. (2000) *Ruin Probabilities*. World Scientific Publishing, Singapore.
- [2] Cheng, Y. and Tang, Q. (2003) Moments of the surplus before ruin and the deficit at ruin in the Erlang(2) risk process. *North American Actuarial Journal* 7, 1-12.
- [3] Dickson, D. C. M. (1992) On the distribution of the surplus prior to ruin. *Insurance: Mathematics & Economics* 11, 191-207.
- [4] Dickson, D. C. M. and Hipp, C. (2000) Ruin problems for phase-type(2) risk processes. *Scandinavian Actuarial Journal*, 147-167.
- [5] Drekić, S., Dickson, D. C. M., Stanford, D. A., and Willmot, G. E. (2001) On the distribution of the deficit at ruin when claims are phase-type. IIPR Research Report 01-12, Department of Statistics and Actuarial Science, University of Waterloo. To appear in the *Scandinavian Actuarial Journal*.
- [6] Dufresne, F. and Gerber, H. U. (1988) The surpluses immediately before and at ruin, and the amount of the claim causing ruin. *Insurance: Mathematics & Economics* 7, 193-199.
- [7] Gerber, H. U. and Shiu, E. S. W. (1998) On the time value of ruin. *North American Actuarial Journal* 2, 48-78.
- [8] Gerber, H. U. and Shiu, E. S. W. (2003a) First discussion of Y. Cheng and Q. Tang's "Moments of the surplus before ruin and the deficit at ruin in the Erlang(2) risk process". To appear in the *North American Actuarial Journal*.

- [9] Gerber, H. U. and Shiu, E. S. W. (2003b) Second discussion of Y. Cheng and Q. Tang's "Moments of the surplus before ruin and the deficit at ruin in the Erlang(2) risk process". To appear in the *North American Actuarial Journal*.
- [10] Latouche, G. and Ramaswami, V. (1999) *Introduction to Matrix Analytic Methods in Stochastic Modeling*. ASA SIAM, Philadelphia.
- [11] Li, S. and Garrido, J. (2003) On ruin for the Erlang(n) risk process. Technical Report No. 1/03, Department of Mathematics and Statistics, Concordia University.
- [12] Neuts, M. F. (1981) *Matrix-geometric Solutions in Stochastic Models: An Algorithmic Approach*. John Hopkins University Press, Baltimore.
- [13] Rolski, T., Schmidli, H., Schmidt, V., and Teugels, J. (1999) *Stochastic Processes for Insurance and Finance*. John Wiley, Chichester.
- [14] Willmot, G. E. and Lin, X. (1998) Exact and approximate properties of the distribution of the surplus before ruin. *Insurance: Mathematics & Economics* 23, 91-110.

David C M Dickson
 Centre for Actuarial Studies
 Department of Economics
 University of Melbourne
 Victoria 3010
 Australia
 email: dcmd@unimelb.edu.au

Steve Drekić
 Department of Statistics and
 Actuarial Science
 University of Waterloo
 Waterloo, Ontario
 Canada N2L 3G1
 email: sdrekić@math.uwaterloo.ca

Figure 1

Figure 2

Figure 3

Figure 4

RECENT ADDITIONS TO THE RESEARCH PAPER SERIES

No.	Date	Subject	Author
84	FEBRUARY 2001	DISCRETE TIME RISK MODELS UNDER STOCHASTIC FORCES OF INTEREST	Jun Cai
85	FEBRUARY 2001	MODERN LANDMARKS IN ACTUARIAL SCIENCE Inaugural Professorial Address	David C M Dickson
86	JUNE 2001	LUNDBERG INEQUALITIES FOR RENEWAL EQUATIONS	Gordon E Willmot Jun Cai X Sheldon Lin
87	SEPTEMBER 2001	VOLATILITY, BETA AND RETURN WAS THERE EVER A MEANINGFUL RELATIONSHIP?	Richard Fitzherbert
88	NOVEMBER 2001	EXPLICIT, FINITE TIME RUIN PROBABILITIES FOR DISCRETE, DEPENDENT CLAIMS	Zvetan G Ignatov Vladimir K Kaishev Rossen S Krachunov
89	NOVEMBER 2001	ON THE DISTRIBUTION OF THE DEFICIT AT RUIN WHEN CLAIMS ARE PHASE-TYPE	Steve Drekcic David C M Dickson David A Stanford Gordon E Willmot
90	NOVEMBER 2001	THE INTEGRATED SQUARE-ROOT PROCESS	Daniel Dufresne
91	NOVEMBER 2001	ON THE EXPECTED DISCOUNTED PENALTY FUNCTION AT RUIN OF A SURPLUS PROCESS WITH INTEREST	Jun Cai David C M Dickson
92	JANUARY 2002	CHAIN LADDER BIAS	Greg Taylor
93	JANUARY 2002	FURTHER OBSERVATIONS ON CHAIN LADDER BIAS	Greg Taylor
94	JANUARY 2002	A GENERAL CLASS OF RISK MODELS	Daniel Dufresne
95	JANUARY 2002	THE DISTRIBUTION OF THE TIME TO RUIN IN THE CLASSICAL RISK MODEL	David C M Dickson Howard R Waters
96	MAY 2002	A NOTE ON THE MAXIMUM SEVERITY OF RUIN AND RELATED PROBLEMS	David C M Dickson
97	JUNE 2002	UPPER BOUNDS FOR ULTIMATE RUIN PROBABILITIES IN THE SPARRE ANDERSEN MODEL WITH INTEREST	Jun Cai David C M Dickson
98	JUNE 2002	CONTINUOUS COMPOUNDING, VOLATILITY AND THE EQUITY PREMIUM	Richard Fitzherbert

No.	Date	Subject	Author
99	JUNE 2002	THE DEFICIT AT RUIN IN THE STATIONARY RENEWAL RISK MODEL	Gordon E Willmot David C M Dickson Steve Drekić David A Stanford
100	AUGUST 2002	ASIAN AND BASKET ASYMPTOTICS	Daniel Dufresne
101	AUGUST 2002	RUIN PROBABILITIES WITH A MARKOV CHAIN INTEREST MODEL	Jun Cai David C M Dickson
102	AUGUST 2002	THE GERBER-SHIU DISCOUNTED PENALTY FUNCTION IN THE STATIONARY RENEWAL RISK MODEL	Gordon E Willmot David C M Dickson
103	NOVEMBER 2002	INITIAL CAPITAL AND MARGINS REQUIRED TO SECURE A JAPANESE LIFE INSURANCE POLICY PORTFOLIO UNDER VARIABLE INTEREST RATES	Manabu Sato David C M Dickson Richard M Fitzherbert
104	NOVEMBER 2002	STATISTICAL CASE ESTIMATION	Greg Taylor Mireille Campbell
105	MARCH 2003	ACTUARIAL PRACTICE AND CONTROL: OBJECTIVES AND CAPABILITIES	Julian D Gribble
106	MARCH 2003	THE IDENTIFICATION AND MEASUREMENT OF SPECULATIVE RISK	Richard Fitzherbert
107	AUGUST 2003	SOME OPTIMAL DIVIDENDS PROBLEMS	David C M Dickson Howard R Waters
108	AUGUST 2003	THE JOINT DISTRIBUTION OF THE SURPLUS PRIOR TO RUIN AND THE DEFICIT AT RUIN IN SOME SPARRE ANDERSEN MODELS	David C M Dickson Steve Drekić

A complete list of papers can be found at <http://www.economics.unimelb.edu.au/actwww/papers.html>

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

DICKSON, DCM;DREKIC, S

Title:

The joint distribution of the surplus prior to ruin and the deficit at ruin in some Sparre Andersen models

Date:

2004

Citation:

DICKSON, D. C. M. & DREKIC, S. (2004). The joint distribution of the surplus prior to ruin and the deficit at ruin in some Sparre Andersen models. Insurance: Mathematics & Economics, 34 (1), pp.97-107. <https://doi.org/10.1016/j.insmatheco.2003.11.003>.

Persistent Link:

<http://hdl.handle.net/11343/33767>